

Total Synthesis of Schilancitrilactones B and C**

Liang Wang, Hengtao Wang, Yihang Li, and Pingping Tang*

Abstract: The first total syntheses of schilancitrilactones B and C have been accomplished in 17 steps (longest linear sequence) from commercially available materials. Key steps include an intramolecular radical cyclization to provide the seven-membered ring, late-stage iodination, and an intermolecular radical addition reaction to complete the total synthesis.

Schilancitrilactones B and C (**1** and **2**; Figure 1)^[1] were isolated in 2012 by Sun and co-workers from the stems of *Schisandra Lancifolia*, which have been used in traditional Chinese medicine for the treatment of neurasthenia and related diseases.^[2] Preliminary biological assays indicated that





Figure 1. Schilancitrilactones B and C.

schilancitrilactone C showed biological activities for inhibiting HIV-1 while schilancitrilactone B was not bioactive. The structures of these compounds were striking in that they contain a 5/7/5/5/5-fused pentacyclic ring system bearing nine stereogenic centers. In addition, the three *cis*-fused five-membered rings (rings C–E), all with the envelope conformations, and seven contiguous chiral centers (including two quaternary centers) form a structurally rigid tricyclic ring system. Construction of these highly oxygenated unique motifs remains challenging. Herein, we present the first total synthesis of schilancitrilactones B and C. The key steps include the successful implementation of an intramolecular radical cyclization to prepare a seven-membered ring, late-stage iodination, and an intermolecular radical C–C bond formation.

Recently, the total synthesis of *Schisandraceae* triterpenoids has been of great interest to synthetic organic chemists because of the intriguing structures and diverse biological activities.^[3] In 2011, Yang and co-workers reported the first total synthesis of schindilactone A.^[4] Recently, the group of Li disclosed the first asymmetric total synthesis of rubrifloridilactone A.^[5] Herein we report our efforts on developing a new strategy to solve the chemical synthesis of **1** and **2**, and a pathway for the synthesis of their analogues and derivatives for medicinal studies. Our retrosynthetic analysis is shown in Scheme 1. It was hypothesized that **1** and **2** might be



Scheme 1. Retrosynthetic analysis of schilancitrilactones B and C.


accessible by an intermolecular radical addition reaction between the alkyl iodide **3** and vinyl stannane **4**. The alkyl iodide **3** was expected to arise by late-stage iodination at C20 from the compound **5**, which in turn could be prepared from the compound **6** by a series of steps including an intramolecular radical cyclization at the C7–C8 bond to prepare the seven-membered ring.^[6] The compound **6** was further deconstructed at the C10–C19 bond into the two simple building blocks **7** and **8**, which could be put together by an aldol reaction. The building blocks **4**, **7**, and **8** could be prepared from the commercially available compounds citraconic anhydride (**16**), L-carvone (**9**), and 1,3-cyclohexadiene (**19**), respectively.

Our work began with the synthesis of the alkyl iodide **7** (Scheme 2). Following the procedure by Fukuyama and co-workers,^[7] L-carvone (**9**) was converted into the correspond-

[*] L. Wang, H. Wang, Y. Li, Prof. Dr. P. Tang
State Key Laboratory and Institute of Elemento-Organic Chemistry
Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Nankai University
Tianjin 300071 (China)
E-mail: ptang@nankai.edu.cn

[**] We gratefully acknowledge the State Key Laboratory of Elemento-Organic Chemistry for generous start-up financial support. This work was supported by the “1000 Youth Talents Plan”, NSFC (21402098, 21421062), and the Natural Science Foundation of Tianjin (13JCYBJC36500). Prof. Biao Yu and Prof. Guangxin Liang are greatly acknowledged for helpful comments.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201501169>.



Scheme 2. Reagents and conditions: a) 30% H_2O_2 , NaOH (aq), MeOH , 0°C ; b) H_2SO_4 , THF/ H_2O (5:1), reflux; c) NaIO_4 , $i\text{PrOH}/\text{H}_2\text{O}$ (1:1), 0°C to RT; d) I_2 , KI , NaHCO_3 , $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ (1:3), 0°C , 51% for 4 steps; e) NaBH_4 , MeOH , 0°C , 85%; f) AIBN , Bu_3SnH , toluene, 100°C , 90%; g) PPh_3 , I_2 , imidazole, 0°C to RT, THF, 84%. AIBN = 2,2'-azobis(2-methylpropionitrile), THF = tetrahydrofuran.

ing aldehyde **13** in a four-step sequence involving epoxidation, epoxide hydrolysis, oxidative cleavage of diols, and iodolactonization in 51% overall yield (4 steps). The aldehyde **13** was selectively reduced with NaBH_4 to provide the alcohol **14** in 85% yield. The deiodination of **14** with AIBN and Bu_3SnH afforded the compound **15**, which was converted into the corresponding **7** with I_2 in the presence of Ph_3P and imidazole in 84% yield.^[8]


Depicted in Scheme 3 is the construction of the vinyl stannane compound **4**. The vinyl bromide **18** was prepared



Scheme 3. Reagents and conditions: a) $\text{PPh}_3\text{CHCO}_2t\text{Bu}$, toluene, RT, 54%; b) TFA , CH_2Cl_2 , 0°C ; c) Br_2 , TFA , $\text{CDCl}_3/\text{CCl}_4$ (1:1), RT; d) Et_3N , DMF , 0°C to RT, 76% for 3 steps; e) $[\text{Pd}(\text{allyl})\text{Cl}]_2$ (5 mol %), $(\text{Bu}_3\text{Sn})_2$, LiCl , 1,4-dioxane, RT, 49%. DMF = N,N -dimethylformamide, TFA = trifluoromethanesulfonfyl.


from the commercially available compound citraconic anhydride (**16**) in a reported four-step process in a 41% overall yield.^[9] Stannylation of **18** was achieved and afforded **4** with $[\text{Pd}(\text{allyl})\text{Cl}]_2$ and $(\text{Bu}_3\text{Sn})_2$ in 49% yield.^[10] It is noteworthy that **4** is not stable during purification, thus resulting in a low yield.

We then moved on to construct the aldehyde compound **8** (Scheme 4). By using the reaction conditions developed by



Scheme 4. Reagents and conditions: a) TPP , O_2 , $h\nu$, CCl_4 , -10°C then thiourea, MeOH ; b) BzCl , Et_3N , DMAP , CH_2Cl_2 , RT, 73% for 2 steps; c) $[\text{Pd}(\text{allyl})\text{Cl}]_2$ (3 mol %), ligand **A**, **22**, K_2CO_3 , MeOH , THF, 0°C ; then DIPEA , 55°C ; 70%; d) NaH , CH_3I , DMF , 0°C , 90%, d.r. (at C13) = 4:1; e) NaBr , DMF , 180°C , 88%, d.r. (at C13) = 1:1; f) LDA , $\text{BrCH}_2\text{CO}_2t\text{Bu}$, THF, -78°C ; g) TFA , CH_2Cl_2 , 0°C to RT, 93% for 2 steps; h) EtMgBr , THF/ Et_2O (1:1), -78°C to RT, 80%; i) O_3 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1:1), -78°C , then Me_2S , -78°C to RT; j) $(\text{Bn}_2\text{NH}_2)(\text{OCOCF}_3)$, toluene, 63°C , 80% for 2 steps. Bz = benzoyl, DIPEA = diisopropylethylamine, DMAP = 4-(N,N -dimethylamino)pyridine, LDA = lithium-diisopropyl amide, TPP = 5,10,15,20-tetraphenyl-21*H*,23*H*-porphine. Thermal ellipsoids are shown at 50% probability.^[11]

Trost and co-workers,^[11] the lactone **23** was obtained in a reported three-step process from the commercially available 1,3-cyclohexadiene (**19**). The steps included asymmetric palladium-catalyzed allylic alkylation. Methylation of **23** with NaH and CH_3I provided the compound **24** in 90% yield with 4:1 diastereoselectivity at C13, and was then subjected to decarboxylation mediated by NaBr to produce a 1:1 mixture of the lactone **25** in 88% yield. Alkylation of **25** with *tert*-butyl bromoacetate gave the single diastereomer **26**. Deprotection of **26** was achieved using trifluoroacetic acid and gave the acid **27** in 93% yield (two steps). Addition of ethyl magnesium bromide followed by acidic workup gave rise to the tricycle **28**, having an ethyl group installed stereoselectively onto the tricyclic framework.^[12] The absolute configuration of **28** was determined by X-ray crystallographic analysis. The cyclohexene ring in **28** was oxidatively cleaved by ozonolysis and the resulting dialdehyde **29** was directly subjected to intramolecular aldol condensation, thus yielding the ring-closed unsaturated aldehyde **8** (80% yield for two steps).^[13]



Scheme 5. Reagents and conditions: a) LDA, THF, -78°C , then **8**, 86%, d.r. (at C19) = 17:1; b) CuCl_2 , EDC, toluene, 80°C , 83%; c) CuI , Zn , $\text{Pyr}/\text{H}_2\text{O}$ (1:4), ultrasound, RT, 55% for **31**, 4% for **31'**; d) *m*CPBA, NaHCO_3 , CH_2Cl_2 , -15°C , 51%; e) NaOMe , MeOH , RT; then $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, NaBH_4 , MeOH/THF (1:5), -15°C , 73%; f) ICl , THF, RT, 63%, d.r. (at C20) = 1.5:1; (g) **4**, AIBN, Bu_3SnH , toluene, 4 Å M.S., 100°C , 9% for **1**, 36% for **2**. EDC = 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide *m*CPBA = *m*-chloroperbenzoic acid. Thermal ellipsoids are shown at 50% probability.^[21]

With the key intermediates **4**, **7**, and **8** in hand, we finished the total synthesis of schilancitrilactones B and C as shown in Scheme 5. The iodo compound **7** was converted into the lithium enolate with LDA at -78°C and then reacted with **8** to give the aldol adduct **30** in 86% yield (d.r. = 17:1 at C19). Dehydration of **30** with 2 equivalents of EDC and a catalytic amount of CuCl_2 provided a 2:1 mixture of the inseparable diene lactone **6** in 83% yield.^[14] The structure of the *E* isomer was confirmed by X-ray crystallographic analysis. Then we investigated the intramolecular radical cyclization to form the seven-membered ring. Initially, the conventional radical conditions (AIBN, Bu_3SnH) led to rapid decomposition of **6** and trace amounts of cyclization product was observed. Photoredox catalysis^[15] was also evaluated and no desired product was found. By using the method (CuI , Zn under ultrasound) for conjugate additions in aqueous media discovered by Luche et al.,^[16] we were pleased to observe the cyclization product **31** in 55% yield, together with the isomer **31'** in 4% yield. The structure of **31'** was confirmed by X-ray crystallographic analysis. We reasoned that the conformation of **6** was suited for cyclization to give the seven-membered ring over the five-membered ring.^[17] Epoxidation of **31** with *m*CPBA provided the epoxide **32** in 51% yield, and underwent ring opening with $\text{NaOMe}/\text{NiCl}_2 \cdot 6\text{H}_2\text{O}/\text{NaBH}_4$ to give the alcohol **5** in 73%.^[18] During this transformation, the epoxide **32** was converted into the intermediate **33** with NaOMe and then further reduced to give the desired product **5** with $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ and NaBH_4 . Finally, we investigated the late-stage iodination and intermolecular radical addition reaction. It was found that treatment of **5** with ICl delivered

the iodo compound **3** as a mixture of diastereomers (d.r. = 1.5:1 at C20) in 63% yield,^[18] and when **3** was heated with the vinyl stannane **4**, AIBN, and Bu_3SnH provided the schilancitrilactones B (**1**, 9%) and C (**2**, 36%) in 45% total yield. Around 25% yield of other isomers were observed based on the ^1H NMR analysis of the crude reaction mixture.^[20] The characterization data obtained for synthetic **1** and **2** were in accord with the reported data for the natural products.

In summary, the first total synthesis of schilancitrilactones B and C has been accomplished by employing an intramolecular radical cyclization, late-stage iodination, and intermolecular radical addition as key steps in the 17 step synthesis (longest linear sequence) from commercially available materials. This strategy opens a pathway for the syntheses of other compounds related to schilancitrilactones B and C, as well as their derivatives and analogues.

Keywords: cyclizations · natural products · radical chemistry · total synthesis · terpenoids

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 5732–5735
Angew. Chem. **2015**, *127*, 5824–5827

- [1] X. Luo, Y.-M. Shi, R.-H. Luo, S.-H. Luo, X.-N. Li, R.-R. Wang, S.-H. Li, Y.-T. Zheng, X. Du, W.-L. Xiao, J.-X. Pu, H.-D. Sun, *Org. Lett.* **2012**, *14*, 1286.
- [2] a) W.-L. Xiao, R.-T. Li, S.-X. Huang, J.-X. Pu, H.-D. Sun, *Nat. Prod. Rep.* **2008**, *25*, 871; b) Y.-M. Shi, W.-L. Xiao, J.-X. Pu, H.-D. Sun, *Nat. Prod. Rep.* **2015**, *32*, 367.
- [3] Selected examples of synthetic efforts on Schisandraceae triterpenoids: a) Y. Tang, L. Deng, Y. Zhang, G. Dong, J.

- Chen, Z. Yang, *Org. Lett.* **2005**, *7*, 593; b) Y. Tang, Y. Zhang, M. Dai, T. Luo, L. Deng, J. Chen, Z. Yang, *Org. Lett.* **2005**, *7*, 885; c) Y. Zhang, Y. Tang, T. Luo, J. Shen, J. Chen, Z. Yang, *Org. Lett.* **2006**, *8*, 107; d) D. Fischer, E. A. Theodorakis, *Eur. J. Org. Chem.* **2007**, 4193; e) Y. Zhang, W. Ren, Y. Lan, Q. Xiao, K. Wang, J. Xu, J. Chen, Z. Yang, *Org. Lett.* **2008**, *10*, 665; f) Q. Wang, C. Chen, *Org. Lett.* **2008**, *10*, 1223; g) L. A. Paquette, K. W. Lai, *Org. Lett.* **2008**, *10*, 2111; h) K. W. Lai, L. A. Paquette, *Org. Lett.* **2008**, *10*, 2115; i) L. A. Paquette, K. W. Lai, *Org. Lett.* **2008**, *10*, 3781; j) M. F. Hossain, K. Matcha, S. Ghosh, *Tetrahedron Lett.* **2011**, *52*, 6473; k) A. Bartoli, G. Chouraqi, J.-L. Parrain, *Org. Lett.* **2012**, *14*, 122; l) V. A. Ignatenko, Y. Han, G. P. Tochtrop, *J. Org. Chem.* **2013**, *78*, 12229; m) B. Gockel, S. S. Goh, E. J. Puttock, H. Baars, G. Chaubet, E. A. Anderson, *Org. Lett.* **2014**, *16*, 4480.
- [4] a) Q. Xiao, W.-W. Ren, Z.-X. Chen, T.-W. Sun, Y. Li, Q.-D. Ye, J.-X. Gong, F.-K. Meng, L. You, Y.-F. Liu, M.-Z. Zhao, L.-M. Xu, Z.-H. Shan, Y. Shi, Y.-F. Tang, J.-H. Chen, Z. Yang, *Angew. Chem. Int. Ed.* **2011**, *50*, 7373; *Angew. Chem.* **2011**, *123*, 7511; b) T.-W. Sun, W.-W. Ren, Q. Xiao, Y.-F. Tang, Y.-D. Zhang, Y. Li, F.-K. Meng, Y.-F. Liu, M.-Z. Zhao, L.-M. Xu, J.-H. Chen, Z. Yang, *Chem. Asian J.* **2012**, *7*, 2321; c) Y. Li, Z.-X. Chen, Q. Xiao, Q.-D. Ye, T.-W. Sun, F.-K. Meng, W.-W. Ren, L. You, L.-M. Xu, Y.-F. Wang, J.-H. Chen, Z. Yang, *Chem. Asian J.* **2012**, *7*, 2334; d) W.-W. Ren, Z.-X. Chen, D. Xiao, Y. Li, T.-W. Sun, Z.-Y. Zhang, Q.-D. Ye, F.-K. Meng, L. You, M.-Z. Zhao, L.-M. Xu, Y.-F. Tang, J.-H. Chen, Z. Yang, *Chem. Asian J.* **2012**, *7*, 2341.
- [5] J. Li, P. Yang, M. Yao, J. Deng, A. Li, *J. Am. Chem. Soc.* **2014**, *136*, 16477.
- [6] Selected reviews for radical cyclization: a) J. M. Cuerva, J. Justicia, J. L. Oller-López, J. E. Oltra, *Top. Curr. Chem.* **2006**, *264*, 63; b) J. Justicia, L. Cienfuegos, A. G. Campaña, D. Miguel, V. Jakoby, A. Gansäuer, J. M. Cuerva, *Chem. Soc. Rev.* **2011**, *40*, 3525; c) T. Taniguchi, H. Ishibashi, *Heterocycles* **2013**, *87*, 527. For selected example, see: J. Justicia, J. L. Oller-López, A. G. Campaña, J. E. Oltra, J. M. Cuerva, E. Buñuel, D. J. Cárdenas, *J. Am. Chem. Soc.* **2005**, *127*, 14911.
- [7] a) S. Takita, S. Yokoshima, T. Fukuyama, *Org. Lett.* **2011**, *13*, 2068; b) S. Takita, S. Yokoshima, T. Fukuyama, *Synthesis* **2011**, *23*, 3848.
- [8] E. J. Corey, S. G. Pyne, W.-G. Su, *Tetrahedron Lett.* **1983**, *24*, 4881.
- [9] T. Benneche, Z. Hussain, A. A. Scheie, J. Lönn-Stensrud, *New J. Chem.* **2008**, *32*, 1567.
- [10] W. D. Wulff, G. A. Peterson, W. E. Bauta, K.-S. Chan, K.-L. Faron, S. R. Gilbertson, R. W. Kaesler, D. C. Yang, C. K. Murray, *J. Org. Chem.* **1986**, *51*, 277.
- [11] a) B. M. Trost, S. Tanimori, P. T. Dunn, *J. Am. Chem. Soc.* **1997**, *119*, 2735; b) Y.-H. Wang, L.-L. Zhu, Y.-X. Zhang, Z. Chen, *Chem. Commun.* **2010**, *46*, 577.
- [12] K. Harrar, O. Reiser, *Chem. Commun.* **2012**, *48*, 3457.
- [13] a) E. J. Corey, R. L. Danheiser, S. Chandrasekaran, P. Siret, G. E. Keck, J.-L. Gras, *J. Am. Chem. Soc.* **1978**, *100*, 8031; b) G. Coulthard, W. Erb, V. K. Aggarwal, *Nature* **2012**, *489*, 278.
- [14] a) H. Sai, H. Ohmizu, *Tetrahedron Lett.* **1999**, *40*, 5019; b) H. Sai, T. Ogiku, H. Ohmizu, *Tetrahedron* **2007**, *63*, 10345.
- [15] For selected reviews, see: a) K. Zeitler, *Angew. Chem. Int. Ed.* **2009**, *48*, 9785; *Angew. Chem.* **2009**, *121*, 9969; b) T. P. Yoon, M. A. Ischay, J. Du, *Nat. Chem.* **2010**, *2*, 527; c) J. M. R. Narayanam, C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, *40*, 102; d) J. Xuan, W. Xiao, *Angew. Chem. Int. Ed.* **2012**, *51*, 6828; *Angew. Chem.* **2012**, *124*, 6934; e) C. K. Prier, D. A. Rankic, D. W. C. MacMillan, *Chem. Rev.* **2013**, *113*, 5322.
- [16] J. L. Luche, C. Allavena, *Tetrahedron Lett.* **1988**, *29*, 5369.
- [17] a) E. Lee, C. H. Yoon, T. H. Lee, *J. Am. Chem. Soc.* **1992**, *114*, 10981; b) E. Lee, C. H. Yoon, *Tetrahedron Lett.* **1996**, *37*, 5929; c) C. S. Swindell, M. C. Chander, J. M. Heerding, P. G. Klimko, L. T. Rahman, J. V. Raman, H. Venkataraman, *J. Org. Chem.* **1996**, *61*, 1101.
- [18] T. Satoh, K. Nanba, S. Suzuki, *Chem. Pharm. Bull.* **1971**, *19*, 817.
- [19] a) G. Giordano, L. Coppi, A. Restelli, *J. Org. Chem.* **1990**, *55*, 5400; b) P. S. Baran, A. L. Zografos, D. P. O'Malley, *J. Am. Chem. Soc.* **2004**, *126*, 3726.
- [20] J. E. Baldwin, D. R. Kelly, *J. Chem. Soc. Chem. Commun.* **1985**, 682.
- [21] CCDC 1034710 (**28**), 1046754 (**6**), and 1034713 (**31'**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Received: February 6, 2015
Published online: March 13, 2015