

A Highly Efficient Gold-Catalyzed Photoredox α -C(sp³)-H Alkynylation of Tertiary Aliphatic Amines with Sunlight


Jin Xie, Shuai Shi, Tuo Zhang, Nina Mehrkens, Matthias Rudolph, and A. Stephen K. Hashmi*

Abstract: A new α -C(sp³)-H alkynylation of unactivated tertiary aliphatic amines with 1-iodoalkynes as radical alkynylating reagents in the presence of [Au₂(μ -dppm)₂]²⁺ in sunlight provides propargylic amines. Based on mechanistic studies, a C-C coupling of an α -aminoalkyl radical and an alkynyl radical is proposed for the C(sp³)-C(sp) bond formation. The mild, convenient, efficient, and highly selective C(sp³)-H alkynylation reaction shows excellent regioselectivity and good functional-group compatibility. A scale-up to gram quantities is possible with sunlight used as a clean and sustainable energy source.


C-Alkynylation reactions are important for organic synthesis, as alkyne moieties are not only versatile building blocks, but also important structural motives in organic materials and biologically active molecules.^[1] Sonogashira coupling is a very powerful method for the formation of C(sp²)-C(sp) bonds,^[2] but the chemospecific construction of C(sp³)-C(sp) bonds still remains a challenge, which was addressed during the past decade by the development of nucleophilic^[3] and electrophilic^[4] C(sp³)-C(sp) bond formation, but much less by radical alkynylation of C(sp³) centers. The pioneering contributions in this field used alkylmercury halides,^[5] alkyl halides,^[6] and even C-H bonds^[7] as C(sp³) radical precursors and combined them with alkynyl sulfones and radical initiators under UV conditions. In 2006, a radical deboronative alkynylation of *B*-alkylcatecholboranes with alkynyl sulfones using di-*tert*-butylhyponitrite as a radical initiator was reported.^[8] Recently, the silver-catalyzed oxidative decarboxylative alkynylation of aliphatic carboxylic acids^[9] and visible-light-induced deboronative alkynylation of alkyl trifluoroborates were achieved by using ethynylbenziodoxolones (EBX) as a radical alkynylating reagent.^[10] The broad substrate scope is a further advantage of the radical C(sp³)-C(sp) coupling methods—but most of these methods rely on the use of hazardous radical initiators or external oxidants (organostannanes, iodine(III) reagents, AIBN, K₂S₂O₈ etc.) and require prefunctionalization of the sub-

strates or special photoreactors. In a continuation of our work on gold-catalyzed C-H activation,^[12,13] we now present the first example of a gold-catalyzed redox-neutral radical α -C(sp³)-H alkynylation of unactivated tertiary aliphatic amines that is based on photoredox catalysis with sunlight^[11] and thus obviates prefunctionalization of the substrates (Scheme 1). The products are formed by the coupling of an α -aminoalkyl radical^[14] and an alkynyl radical^[15] instead of the previously reported radical addition/elimination pathways.

Previous work:



This work (radical α -C-H alkynylation of tertiary amines):



Scheme 1. Different strategies for radical C(sp³)-C(sp) coupling reactions.

Organic halides and tertiary amines were usually used in photoredox reactions as the oxidative and reductive quencher, respectively, undergoing reductive dehalogenation reactions.^[16] As the radical C(sp³)-H alkynylation of tertiary amines was unprecedented,^[17] we studied the feasibility of radical C(sp³)-C(sp) coupling reactions of α -aminoalkyl radicals and alkynyl radicals generated from tertiary amines and electrophilic alkynes under photoredox conditions. The bench-stable and easily available [Au₂(μ -dppm)₂]²⁺ (dppm = bis(diphenylphosphanyl)methane) is a promising photosensitizer,^[18] which was recently applied by Barriault and co-workers in elegant photoredox reactions with unactivated halogen-substituted starting materials.^[16c] As a consequence, we started to investigate the reaction of *N,N*-diisopropylmethylamine (**2a**) with various electrophilic alkynes **3a-c** in sunlight and using [Au₂(μ -dppm)₂]Cl₂ **1a** as a photocatalyst (Table 1, entries 1–3). Trace amounts of the desired redox-neutral coupling product **4aa** were obtained with **3a** or **3b** as the alkynylating reagents (entries 1 and 2). A 53 % yield of **4aa** (the by-products **5a** and diyne **6a** were observed by GC-MS analysis) was obtained with **3c**^[19] (entry 3). The use of K₂HPO₄ as an additional base delivered slightly higher yields (51 % versus 40 % without K₂HPO₄; entry 4). Of the gold

[*] Dr. J. Xie, Dr. S. Shi, M. Sc. T. Zhang, Dr. M. Rudolph, Prof. Dr. A. S. K. Hashmi
Organisch-Chemisches Institut, Universität Heidelberg
Im Neuenheimer Feld 270, 69120 Heidelberg (Germany)
E-mail: hashmi@hashmi.de
Homepage: <http://www.hashmi.de>


M. Sc. N. Mehrkens
Anorganisch-Chemisches Institut, Universität Heidelberg
Im Neuenheimer Feld 270, 69120 Heidelberg (Germany)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201412399>.

Table 1: Optimization of the reaction conditions.^[a]

Entry	Photocatalyst (mol %)	X	Base (2.0 equiv)	Equiv 2a	Time [h]	Yield of 4aa [%] ^[b]
1	1a (3)	BX	K ₂ HPO ₄	3	6	trace
2	1a (3)	Br	K ₂ HPO ₄	3	6	trace
3	1a (3)	I	K ₂ HPO ₄	3	6	53
4	1a (3)	I	—	3	6	40
5	1b (3)	I	K ₂ HPO ₄	3	6	62
6	1c (3)	I	K ₂ HPO ₄	3	6	60
7	1b (3)	I	—	5	1.5	81
8	1b (1)	I	—	5	1.5	81
9	[Ru(bpy) ₃]Cl ₂ (1)	I	—	5	6	62
10	[Ir(ppy) ₃] (1)	I	—	5	1.5	76
11 ^[c]	1b (1)	I	—	5	1.5	trace
12 ^[d]	1b (1)	I	—	5	2	78
13 ^[e]	1b (1)	I	—	5	1.5	trace
14 ^[f]	1b (1)	I	—	5	1.5	trace
15	—	I	—	5	1.5	28
16 ^[g]	—	H	—	5	6	0

[a] Reaction conditions: **3a–d** (0.1 mmol), **2a** (3–5 equiv), MeCN (0.25 mL), sunlight. [b] Yield of isolated product. [c] UV light ($\lambda = 254$ nm). [d] UVA light ($\lambda = 315$ –400 nm). [e] 35 W fluorescent bulb. [f] In the dark. [g] 10 mol % **1b** and 3.0 equiv TBHP at room temperature. bpy = 2,2'-bipyridine, ppy = 2-phenylpyridine, TBHP = *tert*-butyl hydroperoxide.



photocatalysts screened, **1b** showed the highest catalytic activity (entries 3, 5, and 6). By adding 2.0 equiv *N,N*-diisopropylmethylamine, K₂HPO₄ could not only be substituted as an external base, but the yield of the reaction improved to 81 % of **4aa** in 1.5 h (entry 7). Even the loading of photocatalyst **1b** could be reduced to 1 mol % (entry 8). MeCN was the best solvent.^[20] The use of [Ru(bpy)₃]Cl₂ or *fac*-[Ir(ppy)₃] instead of the gold complex **1b** resulted in a lower yield (entries 9 and 10). Besides sunlight, UVA light ($\lambda = 315$ –400 nm) could also efficiently promote the reaction (entry 12). However, irradiation of the reaction mixture with either UV light ($\lambda = 254$ nm) or a fluorescent bulb delivered only traces of the desired product (entries 11 and 13). The control experiments demonstrated that both light and photocatalyst were necessary for the formation of **4aa** (entries 14 and 15). **3d** and the tertiary amine **2a** did not provide **4aa** with the very efficient I₂-TBHP oxidative catalytic system (entry 16).^[21]

A variety of substituted 1-iodoalkynes were examined under the optimized reaction conditions (Table 2). Both donor and acceptor substituents in the *o*-, *m*-, or *p*-position

Table 2: Reaction scope with regard to the 1-iodoalkynes.^[a]


G	Yield [%]	Time [h]
	4aa , 81%	(1.5 h)
	4ab , 78%	(5 h)
	4ac , 75%	(6 h)
	4ad , 76%	(6 h)
	4ae , 88%	(2 h)
	4af , 80%	(4 h)
	4ag , 91%	(3 h)
	4ah , 83%	(6 h)
	4ai , 88%	(5 h)
	4aj , 80%	(5 h)
	4ak , 82%	(3 h)
	4al , 78%	(4 h)
	4am , 71%	(6 h)
	4an , 75%	(3 h)
	4ao , 81%	(4 h)
	4ap , 51%	(25 min)
	4aq , 18%	(6 h)
	4ar , 66%	(3 h)

[a] Reaction conditions: [Au₂(μ-dppm)₂](OTf)₂ (1 mol %), 1-iodoalkyne (0.2 mmol), **2a** (5.0 equiv), MeCN (0.5 mL), room temperature, sunlight.


of the phenylacetylene derivatives gave the corresponding coupling products **4aa–am** selectively in 71–91 % yield. The heteroaromatic **4an** and **4ao** were isolated in yields of 75 % and 81 %. 3-Iodopropiolate furnished **4ap** in 51 % yield. When an aliphatic group was present, **4aq** was obtained in only 18 % yield; hex-1-yne was the main product—which suggests that a hydrogen abstraction seems to be favored over a radical–radical cross-coupling for alkylalkynyl radicals. 1-(Iodoethynyl)cyclohex-1-ene furnished the desired **4ar** in 66 % yield.

While *N,N*-dialkylanilines^[3d] and *N*-aryltetrahydroisoquinolines^[3p] are more reactive, mild alkynylations of α -C(sp³)-H bonds of tertiary aliphatic amines are relatively rare.^[22] Our new procedure gave moderate to good yields of **4aa–4oa** (Table 3). In the absence of **1b**, only a small amount of **4ba** is obtained, thus emphasizing the necessity of a photocatalyst. For the unsymmetrical tertiary amines bearing various functional groups (-OH, ether, acetal, -CHO, *N*-Boc, etc.) from Table 3, exclusively **4aa**, **4ba** and **4ea–4ga**, or mainly **4ha–4oa** were obtained, which indicated a reaction at the methyl group adjacent to the nitrogen atom instead of the methylene or methine group. The redox-neutral radical alkynylation procedure was applied in the selective C(sp³)-H alkynylation of the antidepressant drug citalopram (**4oa**). *N*-Arylamines failed to undergo the radical alkynylation reaction.^[20] A gram-scale conversion gave a 75 % yield of **4aa** (Scheme 2).

Table 3: Reaction scope with regard to the tertiary aliphatic amines.^[a]




[a] Reaction conditions: $[\text{Au}_2(\mu\text{-dppm})_2](\text{OTf})_2$ (1 mol %), **3c** (0.2 mmol), tertiary amines (5.0 equiv), MeCN (0.5 mL), room temperature, sunlight. The regioisomeric ratio (r.r.) was determined by ^1H NMR. Yields of the isolated product. [b] 1.2 equiv 4-dimethylaminopyridine (DMAP) was added. [c] Without photocatalyst **1b**. [d] UVA light was used instead of sunlight.



Scheme 2. Gram-scale experiment.






The mechanistic experiments are shown in Scheme 3. The addition of the radical inhibitors TEMPO or hydroquinone or the electron-transfer scavenger 1,4-dinitrobenzene significantly inhibits the transformation. This finding implies a radical pathway that proceeds through a single-electron-transfer (SET) process should be involved. Under the optimized reaction conditions, besides the desired **4aa**, **5a** (13%) and diyne **6a** (trace) were also detected as by-products by GC-MS analysis [Eq. (1)]. Lowering the concentration of the tertiary amine slowed down the α -deprotonation of the amino-centered radical, thus favoring the competing homodimerization and hydrogen abstraction. Indeed, **5a** and **6a** were formed in 23% and 28% yield, respectively, with only two equivalents of amine **2a** [Eq. (2)]. Thus, an excess of

a) Inhibition experiments:





Additive	Yield of 4aa
TEMPO	trace
Hydroquinone	trace
1,4-Dinitrobenzene	trace

b) Evidences for alkynyl radical intermediate:




c) Evidence of α -aminoalkyl radical intermediate:



Scheme 3. Mechanistic experiments.

tertiary amines/base is essential for the reaction, as the generation of the α -aminoalkyl radical is favored under basic conditions.^[14g] With K_2HPO_4 instead of the tertiary amine, **6a** was produced in 11 % yield (substoichiometric with respect to the catalyst), despite a low conversion of **3c** [Eq. (3)]. The conversion of **3c** failed in the absence of a photocatalyst or in the dark [Eq. (4)]. No reaction occurred if phenylacetylene was employed instead of the iodoalkyne [Eq. (5)]. Electron-rich furan was used to trap the alkynyl radical,^[15a,b] and the expected $C(sp^2)-C(sp)$ coupling product was observed (see the Supporting Information for details). These control experiments indicate the generation of an alkynyl radical from 1-iodoalkyne in the presence of photocatalyst **1b** in sunlight. The dimerization of **2a** to **7**^[14c] was observed by GC-MS analysis during optimization of the reaction with aliphatic 1-iodohex-1-yne [Eq. (6)].^[20] The nucleophilicity of an α -aminoalkyl radical^[14a-d] results in the addition of Michael acceptor **8** delivering the expected addition product **9** [Eq. (7)]. These results strongly indicate that an α -aminoalkyl radical is another important intermediate. The model reaction (1) needed continuous irradiation with sunlight; no conversion of (iodoethynyl)benzene **3c** was detected in the dark.^[20]



Scheme 4. Possible reaction mechanism.

A possible mechanism is shown in Scheme 4 based on the visible-light-mediated α -C(sp³)-H arylation of tertiary amines with 1,4-dicyanobenzene^[14g] and on our results. First the excited-state form of dinuclear gold complex **10** is generated by irradiation with sunlight. Owing to its powerful reducing ability [$E^\circ(\text{Au}_2^{3+}/\text{Au}_2^{2+}) = -1.5$ to -1.7 V versus SSCE],^[23] it should be possible to donate an electron to **3c**,^[24] which then fragments, thereby forming the alkynyl radical species **11**.^[15a,b,20] Then, the resulting dimeric gold complex **12** would be reduced through a SET process from tertiary amines **2a** to give amine radical cation **13**, which can afford α -aminoalkyl radical **14** by a deprotonation process. Finally, the radical-radical coupling of the α -aminoalkyl radical **14** and alkynyl radical **11** affords the desired product **4aa**.^[25] Alternatively, the abstraction of a hydrogen atom by the alkynyl radical **11** from the α -position of the amino-centered radical **13** or the homocoupling of two alkynyl radicals **11** can generate by-products **5a** and **6a**. The selectivity for the preferred cross-recombination of two different radicals can easily be explained by the “persistent” radical effect.^[26,27]

In summary, a highly efficient gold-catalyzed radical C(sp³)-H alkynylation of tertiary aliphatic amines using readily available 1-iodoalkynes as a radical alkynyating reagent in the presence of $[\text{Au}_2(\mu\text{-dppm})_2]^{2+}$ under mild reaction conditions was developed. Mechanistic studies indicate a coupling of an α -aminoalkyl radical and an alkynyl radical.

Keywords: C–H activation · gold catalysis · photoredox catalysis · radical C–C coupling · sunlight

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 6046–6050
Angew. Chem. **2015**, *127*, 6144–6148

- [1] F. Diederich, P. J. Stang, R. R. Tykwinski, *Acetylene Chemistry: Chemistry, Biology and Material Science*, Wiley-VCH, Weinheim, **2005**.
- [2] a) K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* **1975**, *16*, 4467–4470; b) R. Chinchilla, C. Nájera, *Chem. Rev.* **2007**, *107*, 874–922; c) R. Chinchilla, C. Nájera, *Chem. Soc. Rev.* **2011**, *40*, 5084–5121.
- [3] a) B. M. Trost, A. H. Weiss, *Adv. Synth. Catal.* **2009**, *351*, 963–983; b) C.-J. Li, *Acc. Chem. Res.* **2010**, *43*, 581–590; selected examples: c) M. Eckhardt, G. C. Fu, *J. Am. Chem. Soc.* **2003**, *125*, 13642–13643; d) Z. Li, C.-J. Li, *J. Am. Chem. Soc.* **2004**, *126*, 11810–11811; e) Y. Zhao, H. Wang, X. Hou, Y. Hu, A. Lei, H. Zhang, L. Zhu, *J. Am. Chem. Soc.* **2006**, *128*, 15048–15049; f) D. K. Rayabarapu, J. A. Tunge, *J. Am. Chem. Soc.* **2005**, *127*,

- 13510–13511; g) X. Xu, X. Li, *Org. Lett.* **2009**, *11*, 1027–1029; h) G. Cahiez, O. Gager, J. Buendia, *Angew. Chem. Int. Ed.* **2010**, *49*, 1278–1281; *Angew. Chem.* **2010**, *122*, 1300–1303; i) I. Jovel, S. Prateetongkum, R. Jackstell, N. Vogl, C. Weckbecker, M. Beller, *Chem. Commun.* **2010**, *46*, 1956–1958; j) M. Chen, X. Zheng, W. Li, J. He, A. Lei, *J. Am. Chem. Soc.* **2010**, *132*, 4101–4103; k) C. Zhang, D. Seidel, *J. Am. Chem. Soc.* **2010**, *132*, 1798–1799; l) O. Vechorkin, A. Godinat, R. Scopelliti, X. Hu, *Angew. Chem. Int. Ed.* **2011**, *50*, 11777–11781; *Angew. Chem.* **2011**, *123*, 11981–11985; m) T. Hatakeyama, Y. Okada, Y. Yoshimoto, M. Nakamura, *Angew. Chem. Int. Ed.* **2011**, *50*, 10973–10976; *Angew. Chem.* **2011**, *123*, 11165–11168; n) F. Ye, X. Ma, Q. Xiao, H. Li, Y. Zhang, J. Wang, *J. Am. Chem. Soc.* **2012**, *134*, 5742–5745; o) T. Sugiishi, H. Nakamura, *J. Am. Chem. Soc.* **2012**, *134*, 2504–2507; p) M. Rueping, R. M. Koenigs, K. Poscharny, D. C. Fabry, D. Leonori, C. Vila, *Chem. Eur. J.* **2012**, *18*, 5170–5174.
- [4] J. P. Brand, J. Waser, *Chem. Soc. Rev.* **2012**, *41*, 4165–4179.
- [5] a) G. A. Russell, P. Ngovivatchai, *Tetrahedron Lett.* **1986**, *27*, 3479–3482; b) G. A. Russell, P. Ngovivatchai, H. I. Tashtoush, A. Pla-Dalmau, R. K. Khanna, *J. Am. Chem. Soc.* **1988**, *110*, 3530–3538.
- [6] J. Xiang, P. L. Fuchs, *Tetrahedron Lett.* **1998**, *39*, 8597–8600.
- [7] J. Gong, P. L. Fuchs, *J. Am. Chem. Soc.* **1996**, *118*, 4486–4487.
- [8] A.-P. Schaffner, V. Darmency, P. Renaud, *Angew. Chem. Int. Ed.* **2006**, *45*, 5847–5849; *Angew. Chem.* **2006**, *118*, 5979–5981.
- [9] X. Liu, Z. Wang, X. Cheng, C. Li, *J. Am. Chem. Soc.* **2012**, *134*, 14330–14333.
- [10] H. Huang, G. Zhang, L. Gong, S. Zhang, Y. Chen, *J. Am. Chem. Soc.* **2014**, *136*, 2280–2283.
- [11] a) T. P. Yoon, M. A. Ischay, J. N. Du, *Nat. Chem.* **2010**, *2*, 527–532; b) J. M. R. Narayanam, C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, *40*, 102–113; c) J. Xuan, W.-J. Xiao, *Angew. Chem. Int. Ed.* **2012**, *51*, 6828–6838; *Angew. Chem.* **2012**, *124*, 6934–6944; d) S. Maity, N. Zheng, *Synlett* **2012**, *23*, 1851–1856; e) L. Shi, W. Xia, *Chem. Soc. Rev.* **2012**, *41*, 7687–7697; f) C. K. K. Priy, D. A. Rankic, D. W. C. MacMillan, *Chem. Rev.* **2013**, *113*, 5322–5363; g) D. Ravelli, M. Fagnoni, A. Albini, *Chem. Soc. Rev.* **2013**, *42*, 97–113; h) Y. Xi, H. Yi, A. Lei, *Org. Biomol. Chem.* **2013**, *11*, 2387–2403; i) D. P. Hari, B. König, *Angew. Chem. Int. Ed.* **2013**, *52*, 4734–4743; *Angew. Chem.* **2013**, *125*, 4832–4842; j) M. Reckenthäler, A. G. Griesbeck, *Adv. Synth. Catal.* **2013**, *355*, 2727–2744; k) J. Xie, H. Jin, P. Xu, C. Zhu, *Tetrahedron Lett.* **2014**, *55*, 36–48; l) D. M. Schultz, T. P. Yoon, *Science* **2014**, *343*, 1239176.
- [12] a) A. S. K. Hashmi, S. Schäfer, M. Wölfl, C. Diezgil, P. Fischer, A. Laguna, M. C. Blanco, M. C. Gimeno, *Angew. Chem. Int. Ed.* **2007**, *46*, 6184–6187; *Angew. Chem.* **2007**, *119*, 6297–6300; b) A. S. K. Hashmi, M. Wietek, I. Braun, M. Rudolph, F. Rominger, *Angew. Chem. Int. Ed.* **2012**, *51*, 10633–10637; *Angew. Chem.* **2012**, *124*, 10785–10789; c) A. S. K. Hashmi, I. Braun, P. Nösel, J. Schädlich, M. Wietek, M. Rudolph, F. Rominger, *Angew. Chem.* **2012**, *124*, 4532–4536; *Angew. Chem. Int. Ed.* **2012**, *51*, 4456–4460; d) M. M. Hansmann, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem.* **2013**, *125*, 2653–2659; *Angew. Chem. Int. Ed.* **2013**, *52*, 2593–2598; e) P. Nösel, L. N. dos Santos Comprido, T. Lauterbach, M. Rudolph, F. Rominger, A. S. K. Hashmi, *J. Am. Chem. Soc.* **2013**, *135*, 15662–15666.
- [13] J. Xie, C. Pan, A. Abdukader, C. Zhu, *Chem. Soc. Rev.* **2014**, *43*, 5245–5256.
- [14] a) Y. Miyake, Y. Ashida, K. Nakajima, Y. Nishibayashi, *Chem. Commun.* **2012**, *48*, 6966–6968; b) Y. Miyake, K. Nakajima, Y. Nishibayashi, *J. Am. Chem. Soc.* **2012**, *134*, 3338–3341; c) P. Kohls, D. Jadhav, G. Pandey, O. Reiser, *Org. Lett.* **2012**, *14*, 672–675; d) S. Zhu, A. Das, L. Bui, H. Zhou, D. P. Curran, M. Rueping, *J. Am. Chem. Soc.* **2013**, *135*, 1823–1829; e) H. Zhou, P. Lu, X. Gu, P. Li, *Org. Lett.* **2013**, *15*, 5646–5649; f) L. R.

- Espelt, E. M. Wiensch, T. P. Yoon, *J. Org. Chem.* **2013**, *78*, 4107; g) A. McNally, C. K. Prier, D. W. C. MacMillan, *Science* **2011**, *334*, 1114–1117; h) Z. Zuo, D. Ahneman, L. Chu, J. Terrett, A. G. Doyle, D. W. C. MacMillan, *Science* **2014**, *345*, 437–440; i) during revision of our paper, Xiao and co-workers reported a radical–radical coupling of an α -aminoalkyl radical and allyl radical, see J. Xuan, T.-T. Zeng, Z.-J. Feng, Q.-H. Deng, J.-R. Chen, L.-Q. Lu, W.-J. Xiao, H. Alper, *Angew. Chem.* **2015**, *127*, 1645–1648; *Angew. Chem. Int. Ed.* **2015**, *54*, 1625–1628; j) S. B. Lang, K. M. O'Nele, J. A. Tunge, *J. Am. Chem. Soc.* **2014**, *136*, 13606–13609.
- [15] For rare examples of alkynyl radicals in organic synthesis, see a) G. Martelli, P. Spagnolo, M. Tiecco, *J. Chem. Soc. D* **1969**, 282–283; b) G. Martelli, P. Spagnolo, M. Tiecco, *J. Chem. Soc. B* **1970**, 1413–1418; c) P. Boutillier, S. Z. Zard, *Chem. Commun.* **2001**, 1304–1305.
- [16] For selective light-mediated reductive dehalogenation reactions, see a) J. D. Nguyen, E. M. D'Amato, J. M. R. Narayanam, C. R. J. Stephenson, *Nat. Chem.* **2012**, *4*, 854–859; b) H. Kim, C. Lee, *Angew. Chem. Int. Ed.* **2012**, *51*, 12303–12306; *Angew. Chem.* **2012**, *124*, 12469–12472; c) G. Revol, T. McCallum, M. Morin, F. Gagosz, L. Barriault, *Angew. Chem. Int. Ed.* **2013**, *52*, 13342–13345; *Angew. Chem.* **2013**, *125*, 13584–13587.
- [17] The oxidative α -C(sp³)-H nucleophilic alkynylation of tertiary amines were developed in the last decade, but sacrificial external oxidants were usually required.
- [18] The dinuclear gold complex absorbs light at about 300 nm to form a high-energy and long-lived photoexcited state with a quantum yield of 0.23, see D. Li, C.-M. Che, H.-L. Kwong, V. W.-W. Yam, *J. Chem. Soc. Dalton Trans.* **1992**, 3325–3329.
- [19] The 1-iodoalkynes can be easily prepared in high yields from various commercially available alkynes, see
- [20] See the Supporting Information.
- [21] a) Z.-J. Cai, S.-Y. Wang, S.-J. Ji, *Org. Lett.* **2013**, *15*, 5226–5229; b) T. Nobuta, N. Tada, A. Fujiya, A. Kariya, T. Miura, A. Itoh, *Org. Lett.* **2013**, *15*, 574–577; c) F. Xiao, H. Chen, H. Xie, S. Chen, L. Yang, G.-J. Deng, *Org. Lett.* **2014**, *16*, 50–53.
- [22] A dehydrogenative nucleophilic alkynylation of branched tertiary aliphatic amines catalyzed by a Shvo complex has been achieved in moderate yields, but the harsh reaction conditions (150°C, 24 h) and poor functional-group compatibility seriously limited the utility of this method, see Ref. [3i].
- [23] C.-M. Che, H.-L. Kwong, K.-C. Poon, V. W.-W. Yam, *J. Chem. Soc. Dalton Trans.* **1990**, 3215–3219.
- [24] The cyclic voltammogram of (iodoethynyl)benzene (**3c**) exhibited an irreversible reduction with $E_{1/2} = -1.29$ V versus Fc in MeCN.^[20]
- [25] The SOMO energy of α -aminoalkyl radical **14** and alkynyl radical **11** were calculated as -7.12 eV and -10.07 eV, respectively.
- [26] Although initially discovered with “persistent” radicals, this kinetic phenomenon also applies to short-lived radicals. It is only necessary that the lifetimes of the two radicals involved show a significant difference (which is the case, Ref. [27]): a) A. Studer, *Chem. Eur. J.* **2001**, *7*, 1159–1164; b) H. Fischer, *Chem. Rev.* **2001**, *101*, 3581–3610.
- [27] The lifetime of an α -aminoalkyl radical (Et₂NCHMe) and phenylethynyl radical are about 700 ns and 2 ns, respectively; see a) J. C. Scaiano, *J. Phys. Chem.* **1981**, *85*, 2851–2855; b) X. Gu, Y. Guo, F. Zhang, A. M. Mebel, R. I. Kaiser, *Chem. Phys. Lett.* **2007**, *436*, 7–14. Thus, α -aminoalkyl radical **14** is more long-lived and the initially preferred dimerization of the alkynyl radical is in full accord with the mechanistic model.

Received: December 26, 2014

Revised: February 24, 2015

Published online: March 30, 2015

