

Catalytic Asymmetric Total Synthesis of (–)-Galanthamine and (–)-Lycoramine**

Lei Li, Qiao Yang, Yuan Wang, and Yanxing Jia*


Dedicated to Professor Jieping Zhu on the occasion of his 50th birthday

Abstract: The catalytic asymmetric total syntheses of (–)-galanthamine (**1**) and (–)-lycoramine (**2**) have been achieved by using a conceptually new strategy featuring two metal-catalyzed reactions as the key steps. A new method for the construction of 3,4-fused benzofurans has been developed through a palladium-catalyzed intramolecular Larock annulation reaction, which was successfully applied to the construction of the ABD tricyclic skeleton of **1** and **2**. To achieve the asymmetric synthesis of **1** and **2**, a $\text{Sc}^{\text{III}}/\text{N},\text{N}'$ -dioxide complex was used to catalyze the enantioselective conjugate addition of 3-alkyl-substituted benzofuranone to methyl vinyl ketone for the construction of a chiral quaternary carbon center.

The 3,4-fused (dihydro)benzofuran skeleton, in which the 3-position of the (dihydro)benzofuran is bridged to the 4-position, represents the key structural motif of a diverse group of natural products which exhibit a wide range of biological activities (Figure 1). These natural products include the medicinally important galanthamine-type *Amaryllidaceae*

alkaloids and the *Opium* alkaloids, which could serve as ideal model compounds for the development of new synthetic methods and strategies.^[1]

As a continuation of our ongoing projects focused on the total synthesis of 3,4-fused indole alkaloids,^[2] we have recently developed a general and convenient strategy for the construction of 3,4-fused indoles by an intramolecular Larock indolization reaction (Scheme 1 a).^[3] Considering the



Scheme 1. Synthesis of 3,4-fused indoles and 3,4-fused benzofurans by an intramolecular Larock annulation.




Figure 1. Representative 3,4-fused (dihydro)benzofuran natural products.

structural similarity of 3,4-fused benzofurans with 3,4-fused indoles, we were curious whether the palladium-catalyzed intramolecular Larock annulation could be applied to the preparation of 3,4-fused benzofurans (Scheme 1 b). If this idea could be realized, then we would have opportunity to explore a new strategy for the synthesis of (–)-galanthamine (**1**) and (–)-lycoramine (**2**).


(–)-Galanthamine (**1**), a representative member of the *Amaryllidaceae* alkaloids, has been clinically used for the treatment of Alzheimer's disease and other memory impairments.^[4] (–)-Lycoramine (**2**) has a similar, albeit less potent activity as an acetylcholinesterase inhibitor and an allosteric potentiating ligand.^[5] Because of the high cost of its isolation and the limited natural sources, a number of total syntheses of **1** have been reported.^[6–16] According to the strategy for the formation of the ring system, these syntheses can be divided into two categories: 1) synthesis proceeding from AC ring → ADC ring → ABCD ring (Scheme 2, path a),^[6–8] and it mainly involves intramolecular phenolic oxidative coupling followed by intramolecular oxa-Michael addition to form the benzofuran B ring, and 2) synthesis proceeding from the AC(AB) ring → ABC ring → ABCD ring (Scheme 2, path b),^[9–16] in which the formation of the bridged seven-membered D ring is placed at the late stage of the synthesis. To our knowledge, no synthesis from the ABD ring → ABCD ring (Scheme 2, path c) has been reported.

[*] L. Li,^[†] Q. Yang,^[†] Dr. Y. Wang, Prof. Y. Jia
State Key Laboratory of Natural and Biomimetic Drugs
School of Pharmaceutical Sciences, Peking University
Xue Yuan Rd. 38, Beijing 100191 (China)
E-mail: yxjia@bjmu.edu.cn

[†] These authors contributed equally to this work.

[**] We are grateful to the National Natural Science Foundation of China (Nos. 21402003, and 21290183) for their financial support.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201411338>.



Scheme 2. Strategies for the synthesis of galanthamine (**1**).

Structurally, galanthamine-type alkaloids contain a chiral all-carbon quaternary center, and the enantioselective construction of this sterically congested quaternary center^[17] is the critical step in the total synthesis of these alkaloids. Although numerous asymmetric total syntheses of these alkaloids have been reported,^[7–16] only the groups of Trost,^[9,15] and Fan,^[13] as well as that of Zhou and Xie,^[14] have recently achieved catalytic asymmetric synthesis of galanthamine (**1**) and related alkaloids.


To explore a new strategy for the effective construction of polycyclic ring systems and a new method for the catalytic asymmetric establishment of the crucial chiral all-carbon quaternary stereocenter, a retrosynthetic analysis of **1** and **2** is outlined in Scheme 3. We envisioned that both **1** and **2** could



Scheme 3. Retrosynthetic analysis for **1** and **2**. Boc = *tert*-butoxycarbonyl, TES = triethylsilyl.

be derived from the same advanced intermediate **3**. The tetracyclic compound **3** could be generated from the tricyclic compound **4** by intramolecular ketone–lactone condensation. In turn, **4** could be accessed from **5** by key catalytic asymmetric Michael addition with methyl vinyl ketone (MVK). The benzofuranone **5** should be readily prepared by oxidation of 3,4-fused benzofuran **6**, which could be derived from **7** by a novel palladium-catalyzed annulation reaction.

We firstly examined the feasibility of our strategy for the synthesis of 3,4-fused benzofuran by an intramolecular Larock annulation reaction (Scheme 4).^[18,19] A variety of reaction conditions were screened and we found that under the reaction conditions of [Pd₂(dba)₃]·CHCl₃ (5 mol %) and



Scheme 4. Realization of the intramolecular Larock annulation reaction. dba = dibenzylideneacetone, Ts = 4-toluenesulfonyl, DMF = *N,N*-dimethylformamide.

P(*t*Bu)₃·HBF₄ (20 mol %) at 100 °C, the desired product **9a** was obtained in 95 % yield (see the Supporting Information).^[20] The substrate scope of this reaction was subsequently examined. The transformation was found to be quite general, and a variety of 3,4-fused benzofurans containing either carbon, oxygen, or nitrogen tethers were obtained in reasonable yields (Table 1).

Table 1: Synthesis of 3,4-fused benzofurans.^[a]

$[Pd_2(dba)_3] \cdot CHCl_3$ (5 mol%)
 $P(tBu)_3 \cdot HBF_4$ (20 mol%)
 K_2CO_3 (5 equiv)
 DMF, 100 °C, 1 h
 $c = 0.01$ M

9a, 95%

9b, 88%

9c, 82%

9d, 95%

9e, 44%

9f, 95%

9g, 95%

9h, 67%

9i, 48%

9j, 41%

9k, 68%

9l, 50%

9m, 88%


9n, 27%

9o: R = Boc, 89%

9p: R = CO₂Me, 92%

[a] Yields are those of the isolated products. TMS = trimethylsilyl.

We then set out to apply this method to the assembly of the ABD ring system of **1** and **2** (Scheme 5). Reductive coupling of the known aldehyde **10**^[9c] and amine **11**,^[3a] and subsequent protection with Boc₂O provided **7** in 81 % yield. Treatment of **7** under our aforementioned reaction conditions



Scheme 5. Construction of ring B/D in the tricyclic synthon. *m*-CPBA = *meta*-chloroperoxybenzoic acid, TBAF = tetra-*n*-butylammonium fluoride.


successfully afforded the 3,4-fused benzofuran **12** in 89% yield. Removal of the TES group in **12** with TBAF (95%), and subsequent oxidation with *m*-CPBA, afforded **5** in 56% yield.

After successful construction of the ABD ring system, we turned our attention to the construction of the chiral all-carbon quaternary stereocenter in **1** and **2**. Although a few catalytic enantioselective conjugate additions of 3-substituted benzofuranones were reported,^[21] to the best of our knowledge, the catalytic enantioselective conjugate addition of 3-alkyl-substituted benzofuranone to MVK has not been reported.

Initially, various amine-thiourea or urea bifunctional organocatalysts were screened for the conjugate addition of **5** to MVK. However, the highest enantioselectivity obtained was 55% *ee* (see the Supporting Information). These results prompted us to investigate alternative catalytic reaction systems. Inspired by the recent work of Feng and co-workers,^[22] we turned our attention to examining an asymmetric Michael addition using the chiral metal/*N,N'*-dioxide complexes. As described in Table 2, we initially investigated the Michael addition, in CH₂Cl₂ at room temperature, catalyzed by Yb(OTf)₃/13 complexes (entries 1–3). But only Yb(OTf)₃/13c gave the desired product **4** in 10% yield with 42% *ee* (entry 3). When the solvent was changed to ethanol, **4** was obtained in higher yield (33%) with similar *ee* value (entry 4). However, when Sc(OTf)₃ was used instead of Yb(OTf)₃ at room temperature, the reaction proceeded smoothly to afford **4** in 85% yield with 93% *ee* (entry 5). Additionally, when the reaction was run at 10 °C, **4** was obtained in 85% yield with 94% *ee* (entry 6). Under the aforementioned reaction conditions, conjugate additions of several 3-alkyl-substituted benzofuranones to MVK were investigated. The results showed that both the yield and the *ee* values were excellent (Table 3).

With **4** in hand, we proceeded with the total synthesis of **1** (Scheme 6). Treatment of **4** with LDA yielded the cyclization product **3** in 95% yield. Subsequent direct reduction of the **3** with Et₃SiH provided the β-alcohol **16** in 73% yield and α-alcohol **17** in 12% yield, with the simultaneous reduction of ketone and removal of the Boc group.^[23] Selective protection of the amine in **16** and **17** with methyl chloroformate and subsequent oxidation with Dess–Martin periodinane pro-


Table 2: Michael addition catalyzed by Lewis acid/*N,N'*-dioxide complexes.^[a]



Entry	Lewis acid	Ligand	Solvent	Yield [%] ^[b]	<i>ee</i> [%] ^[c]
1	Yb(OTf) ₃	13a	CH ₂ Cl ₂	n.r.	–
2	Yb(OTf) ₃	13b	CH ₂ Cl ₂	n.r.	–
3	Yb(OTf) ₃	13c	CH ₂ Cl ₂	10	42
4	Yb(OTf) ₃	13c	EtOH	33	38
5	Sc(OTf) ₃	13c	EtOH	85	93
6 ^[d]	Sc(OTf) ₃	13c	EtOH	85	94

[a] Reaction conditions: **5** (0.1 mmol), MVK (0.15 mmol), solvent (0.3 mL). [b] Yield of isolated product. [c] Determined by HPLC using a chiral stationary phase. [d] The reaction was run at 10 °C. n.r. = no reaction, Tf = trifluoromethanesulfonyl.


Table 3: Substrate scope of the asymmetric Michael reaction.^[a]




[a] Reaction conditions: **14** (0.1 mmol), MVK (0.15 mmol), EtOH (0.3 mL). Yields are those of the isolated products. Enantiomeric excess was determined by HPLC using a chiral stationary phase.

vided the known ketone **18**, the key intermediate in Fan's total synthesis of **1**, in 75% yield over three steps. The physical properties (¹H and ¹³C NMR spectra, MS data, and [α]_D) of **18** are consistent with those described in the literature.^[13] Treatment of **18** with TMSOTf and Et₃N provided the corresponding silyl enol ether, which was oxidized under Saegusa conditions to give the enone **19** in 67% yield. Finally, **19** was readily converted into **1** in a two-step sequence.^[13]

After completion of the synthesis of **1**, we turned our attention to the synthesis of **2** (Scheme 7). Considering the lower yield in the conversion of **3** into **17** (Scheme 6), we tried to develop a more efficient approach. Thus, stereoselective reduction of **3** with L-selectride gave the sole α-alcohol **20** in 91% yield. Reduction of **20** with Et₃SiH provided **17** in 88% yield. Reaction of **17** and formaldehyde under standard reaction conditions [NaBH₄, NaBH₃CN or NaBH(OAc)₃] gave **2** in only 30–40% yield. Methylation of **17** under



Scheme 6. Synthesis of **1**. DMP = Dess–Martin periodinane, LDA = lithium diisopropylamide, THF = tetrahydrofuran.



Scheme 7. Synthesis of **2**.

Eschweiler–Clarke conditions (HCO_2H , HCHO , reflux)^[24] and subsequent treatment of the resulting product with $\text{K}_2\text{CO}_3/\text{MeOH}/\text{H}_2\text{O}$ afforded **2** in 84% yield over two steps.

In summary, an asymmetric total synthesis of both (–)-galanthamine (**1**) and (–)-lycoramine (**2**) have been achieved based on a conceptually new strategy by employing two metal-catalyzed reactions. A new method for the construction of 3,4-fused tricyclic benzofurans, the core structure of a variety of bioactive important natural products, has been developed using a palladium-catalyzed intramolecular Larock annulation reaction. In addition, a $\text{Sc}^{\text{III}}/\text{N},\text{N}'$ -dioxide complex catalyzed the enantioselective conjugate addition reaction of 3-alkyl-substituted benzofuranone to MVK for the construction of a quaternary carbon center was developed for the first time.

Keywords: alkaloids · asymmetric catalysis · natural products · palladium · total synthesis

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 6255–6259
Angew. Chem. **2015**, *127*, 6353–6357

[1] For a recent review, see: a) U. Rinner, T. Hudlicky, *Top. Curr. Chem.* **2012**, *309*, 33; for recent syntheses reported after those

cited in Ref. [1a], see: b) M. Ichiki, H. Tanimoto, S. Miwa, R. Saito, T. Sato, N. Chida, *Chem. Eur. J.* **2013**, *19*, 264; c) J. Li, G.-L. Liu, X.-H. Zhao, J.-Y. Du, H. Qu, W.-D. Chu, M. Ding, C.-Y. Jin, M.-X. Wei, C.-A. Fan, *Chem. Asian J.* **2013**, *8*, 1105; d) V. Varghese, T. Hudlicky, *Angew. Chem. Int. Ed.* **2014**, *53*, 4355; *Angew. Chem.* **2014**, *126*, 4444.

[2] For a review for the synthesis of 3,4-fused indoles, see: a) D. Shan, Y. Jia, *Chin. J. Org. Chem.* **2013**, *33*, 1144; b) Q. Liu, Y.-A. Zhang, P. Xu, Y. Jia, *J. Org. Chem.* **2013**, *78*, 10885; c) Y.-A. Zhang, Q. Liu, C. Wang, Y. Jia, *Org. Lett.* **2013**, *15*, 3662; d) Q. Liu, Q. Li, Y. Ma, Y. Jia, *Org. Lett.* **2013**, *15*, 4528; e) L. Guo, F. Zhang, W. Hu, L. Li, Y. Jia, *Chem. Commun.* **2014**, *50*, 3299, and references therein.

[3] a) D. Shan, Y. Gao, Y. Jia, *Angew. Chem. Int. Ed.* **2013**, *52*, 4902; *Angew. Chem.* **2013**, *125*, 5002; b) Y. Gao, D. Shan, Y. Jia, *Tetrahedron* **2014**, *70*, 5136; c) for a similar work, see: S. P. Breazzano, Y. B. Poudel, D. L. Boger, *J. Am. Chem. Soc.* **2013**, *135*, 1600.

[4] For a review, see: a) J. Marco-Contelles, M. C. Carreiras, C. Rodríguez, M. Villarroya, A. G. García, *Chem. Rev.* **2006**, *106*, 116; for racemic syntheses reported after those cited in Ref. [4a], see: b) X.-D. Hu, Y.-Q. Tu, E. Zhang, S. Gao, S. Wang, A. Wang, C.-A. Fan, M. Wang, *Org. Lett.* **2006**, *8*, 1823; c) T. Ishikawa, K. Kudo, K. Kuroyabu, S. Uchida, T. Kudoh, S. Saito, *J. Org. Chem.* **2008**, *73*, 7498; d) M. G. Banwell, X. Ma, O. P. Karunaratne, A. C. Willis, *Aust. J. Chem.* **2010**, *63*, 1437; e) J. H. Chang, H.-U. Kang, I.-H. Jung, C.-G. Cho, *Org. Lett.* **2010**, *12*, 2016; f) Y. Feng, Z.-X. Yu, *J. Org. Chem.* **2015**, *80*, 1952.

[5] For recent syntheses of lycoramine, see: a) W. P. Malachowski, T. Paul, S. Phounsavath, *J. Org. Chem.* **2007**, *72*, 6792; b) C.-A. Fan, Y.-Q. Tu, Z.-L. Song, E. Zhang, L. Shi, M. Wang, B. Wang, S.-Y. Zhang, *Org. Lett.* **2004**, *6*, 4691, and references therein.

[6] D. H. R. Barton, G. W. Kirby, *J. Chem. Soc. C* **1962**, 806.

[7] a) K. Tomioka, K. Shimizu, S. Yamada, K. Koga, *Heterocycles* **1977**, *6*, 1752; b) K. Shimizu, K. Tomioka, S. Yamada, K. Koga, *Heterocycles* **1977**, *8*, 277.

[8] a) S. Kodama, Y. Hamashima, K. Nishide, M. Node, *Angew. Chem. Int. Ed.* **2004**, *43*, 2659; *Angew. Chem.* **2004**, *116*, 2713; b) M. Node, S. Kodama, Y. Hamashima, T. Katoh, K. Nishide, T. Kajimoto, *Chem. Pharm. Bull.* **2006**, *54*, 1662.

[9] a) B. M. Trost, F. D. Toste, *J. Am. Chem. Soc.* **2000**, *122*, 11262; b) B. M. Trost, W. Tang, *Angew. Chem. Int. Ed.* **2002**, *41*, 2795; *Angew. Chem.* **2002**, *114*, 2919; c) B. M. Trost, W. Tang, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 14785.

[10] V. Satcharoen, N. J. McLean, S. C. Kemp, N. P. Camp, R. C. D. Brown, *Org. Lett.* **2007**, *9*, 1867.

[11] a) H. Tanimoto, T. Kato, N. Chida, *Tetrahedron Lett.* **2007**, *48*, 6267 (unnatural galanthamine); b) T. Kato, H. Tanimoto, H. Yamada, N. Chida, *Heterocycles* **2010**, *82*, 563.

[12] P. Magnus, N. Sane, B. P. Fauber, V. Lynch, *J. Am. Chem. Soc.* **2009**, *131*, 16045.

[13] P. Chen, X. Bao, L.-F. Zhang, M. Ding, X.-J. Han, J. Li, G.-B. Zhang, Y.-Q. Tu, C.-A. Fan, *Angew. Chem. Int. Ed.* **2011**, *50*, 8161; *Angew. Chem.* **2011**, *123*, 8311.

[14] J.-Q. Chen, J.-H. Xie, D.-H. Bao, S. Liu, Q.-L. Zhou, *Org. Lett.* **2012**, *14*, 2714.

[15] Y. Zang, I. Ojima, *J. Org. Chem.* **2013**, *78*, 4013.

[16] J. Choi, H. Kim, S. Park, J. Tae, *Synlett* **2013**, 379.

[17] For recent reviews, see: a) I. Marek, Y. Minko, M. Pasco, T. Mejuch, N. Gilboa, H. Chechik, J. P. Das, *J. Am. Chem. Soc.* **2014**, *136*, 2682; b) A. Y. Hong, B. M. Stoltz, *Eur. J. Org. Chem.* **2013**, 2745; c) J. P. Das, I. Marek, *Chem. Commun.* **2011**, 47, 4593; d) C. Hawner, A. Alexakis, *Chem. Commun.* **2010**, 46, 7295; e) S. Jautze, R. Peters, *Synthesis* **2010**, 365; for selective recent examples on the catalytic asymmetric formation of all-carbon quaternary centers in natural product syntheses, see: f) Z. Li, S. Zhang, S. Wu, X. Shen, L. Zou, F. Wang, X. Li, F.

- Peng, H. Zhang, Z. Shao, *Angew. Chem. Int. Ed.* **2013**, 52, 4117; *Angew. Chem.* **2013**, 125, 4211; g) Z. Xu, Q. Wang, J. Zhu, *J. Am. Chem. Soc.* **2013**, 135, 19127.
- [18] a) R. C. Larock, *J. Organomet. Chem.* **1999**, 576, 111; b) R. C. Larock, E. K. Yum, M. J. Doty, K. C. Sham, *J. Org. Chem.* **1995**, 60, 3270.
- [19] For the only example of the construction of 3,4-fused furan through intramolecular Larock-type annulation, see: a) A. Kojima, T. Takemoto, M. Sodeoka, M. Shibasaki, *J. Org. Chem.* **1996**, 61, 4876; b) A. Kojima, S. Shimizu, M. Shibasaki, *J. Heterocycl. Chem.* **1998**, 35, 1057.
- [20] T. Konno, J. Chae, T. Ishihara, H. Yamanaka, *Tetrahedron* **2004**, 60, 11695.
- [21] a) X. Li, Z. Xi, S. Luo, J.-P. Cheng, *Adv. Synth. Catal.* **2010**, 352, 1097; b) X. Li, S. Hu, Z. Xi, L. Zhang, S. Luo, J.-P. Cheng, *J. Org. Chem.* **2010**, 75, 8697; c) F. Pesciaoli, X. Tian, G. Bencivenni, G. Bartoli, P. Melchiorre, *Synlett* **2010**, 1704.
- [22] a) X. H. Liu, L. L. Lin, X. M. Feng, *Acc. Chem. Res.* **2011**, 44, 574; b) X. H. Liu, L. L. Lin, X. M. Feng, *Org. Chem. Front.* **2014**, 1, 298.
- [23] a) G. A. Kraus, K. A. Frazier, B. D. Roth, M. J. Taschner, K. Neuenschwander, *J. Org. Chem.* **1981**, 46, 2417; b) G. A. Kraus, J. Shi, D. Reynolds, *J. Org. Chem.* **1990**, 55, 1105.
- [24] H. T. Clarke, H. B. Gillespie, S. Z. Weisshaus, *J. Am. Chem. Soc.* **1933**, 55, 4571.

Received: November 29, 2014

Revised: March 3, 2015

Published online: April 2, 2015