



Hydrophosphination of CO₂ and Subsequent Formate Transfer in the 1,3,2-Diazaphospholene-Catalyzed *N*-Formylation of Amines

Che Chang Chong and Rei Kinjo*

Abstract: Hydrophosphination of CO₂ with 1,3,2-Diazaphospholene (NHP-H; **1**) afforded phosphorus formate (NHP-OCOH; **2**) through the formation of a bond between the electrophilic phosphorus atom in **1** and the oxygen atom from CO₂, along with hydride transfer to the carbon atom of CO₂. Transfer of the formate from **2** to Ph₂SiH₂ produced Ph₂Si(OCHO)₂ (**3**) in a reaction that could be carried out in a catalytic manner by using 5 mol % of **1**. These elementary reactions were applied to the metal-free catalytic *N*-formylation of amine derivatives with CO₂ in one pot under ambient conditions.

The valorization of carbon dioxide (CO₂) has attracted much attention in synthetic chemistry because of the potential use of CO₂ as a nontoxic, abundant, and inexpensive C1 source for the production of useful chemicals.^[1,2] However, the considerable thermodynamic stability of CO₂ hampers facile activation. To address this issue, several metal catalysts have been employed for the transformation of CO₂, which has afforded various commodity chemicals such as formic acid derivatives, methane, methanol, CO, and formaldehyde.^[3]

Metal-free catalytic systems for CO₂ valorization present an alternative method to the prevalent use of conventional metal catalysts.^[4,5] The seminal work by Grimme, Stephan, Erker, et al. in 2009 demonstrated the potential of frustrated Lewis pairs (FLPs), consisting of a Lewis base (LB) and Lewis acid (LA), for CO₂ capture.^[6,7] Since then, a variety of systems based on main-group compounds have been developed for the activation of CO₂.^[8] Depending on the binding modes to CO₂, these systems are mainly classified into three types: one LB adduct (**I**), one LB and one LA adduct (**II**), and one LB and two LA adducts (**III**; Figure 1a).^[9] These are also recognized as key structures of initial intermediates in the functionalization of CO₂ by main-group catalysts. For instance, hydroboration of CO₂ catalyzed by main-group compounds is inferred to proceed via the activation of CO₂ to give insertion of the B–H bond into the activated C=O bond.^[8g–h,10a] However, recent studies by Fontaine and Bourissou suggest that such a mechanistic pathway often involves a high activation barrier.^[10b–d] Meanwhile, it has been reported that the direct reduction of CO₂ by main-group p-




Figure 1. a) Three coordination modes in the activation of CO₂ by main-group compounds. b) Direct insertion of [E]–H bonds into CO₂. c) Examples of products and an intermediate from direct insertion of the P^V–N bond into CO₂. d) Generic structure of 2-*H*-1,3,2-diazaphospholene (**1**).

block hydrides [E]–H (**IV**) gives rise to formate derivatives (**V**) via insertion of the E–H bond (Figure 1b). In this process, the main-group fragment [E] and the H atom of **IV** bond to an oxygen atom and the carbon atom of CO₂, respectively. The E–H bond thus has to be polarized as E(δ⁺)–H(δ[–]) so that it concomitantly displays hydride-donor ability and electrophilic nature at the [E] moiety. As such, extant examples are limited to group 13–14 elements such as B,^[11] Al,^[12] Ga,^[13] Ge,^[14] and Sn.^[15]

With group 15 elements (Figure 1c), Cavell et al. reported the insertion of CO₂ into the P–N bond of Me(F₃C)₃P^V–NMe₂ over two days, which afforded the hexacoordinate phosphorus derivative **VI**.^[16] Stephan et al. demonstrated that amidophosphoranes containing strained four-membered rings readily reacted with CO₂ to produce P^V heterocycles (**VIIa–b**).^[17] Recently, Fontaine et al. described the activation of CO₂ with phosphazenes, in which an intermediate (**VIII**) involving CO₂ insertion into the P^V–N bond was proposed.^[18] In those reactions, the P centre in the + V oxidation state acted as an electrophile to form a bond with an O atom of CO₂. Relevant sequestration of CO₂ with phosphines (R₃P^{III}) is extremely rare,^[19,20] and to the best of our knowledge, formation of a phosphorus(III) formate (R₂P^{III}–OCOH) via P^{III}–H bond insertion into CO₂, that is, hydrophosphination, has not thus far been achieved. This is presumably because phosphines are typically nucleophilic, and therefore the interaction with CO₂


[*] C.-C. Chong, Prof. Dr. R. Kinjo
Division of Chemistry and Biological Chemistry
School of Physical and Mathematical Sciences
Nanyang Technological University
Nanyang Link 21, Singapore 637371 (Singapore)
E-mail: rkinjo@ntu.edu.sg

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201505244>.

should predominantly generate adduct **I** (Figure 1a) rather than the P^{III}–H bond insertion product.^[2d,8p–q] Given the significance of CO₂ valorization, it is highly desirable to develop diverse strategies for the capture of CO₂ and its transformation into value-added compounds.

Recently, we reported that 1,3,2-diazaphospholene (**1**), which exhibits zwitterionic properties (**1'**; Figure 1d),^[21] effectively promotes the hydroboration of carbonyl compounds.^[22] The reaction mechanism involved a facile insertion of the exocyclic P–H bond of **1** into the C=O bond at the initial step. This result prompted us to investigate the reactivity of **1** towards CO₂ since it possesses consecutive C=O bonds. Herein, we report a facile CO₂ insertion into the P–H bond of **1** to form the phosphorus formate **2**. Transfer of the formate from **2** to silane, and its application in the catalytic N-formylation of amines are also described.

A CD₃CN solution of **1** was exposed to 1 atm of CO₂ at ambient temperature. Instantaneously, the yellow solution darkened to a deep brown color, showing a singlet at 111.5 ppm in the ³¹P NMR spectrum, which is shifted downfield compared to that (57.8 ppm) of **1**. After workup, product **2** was isolated as a brown solid in 93 % yield (Scheme 1a). The




Scheme 1. a) Reaction of **1** with CO₂ (¹³CO₂). b) Solid-state structure of **2**. Hydrogen atoms, except for H11, are omitted for clarity. c) Formate transfer from **2** to Ph₂SiH₂ under CO₂ atmosphere.

¹H NMR spectrum of **2** shows a characteristic peak at 8.10 ppm while the ¹³C{¹H} NMR spectrum shows a singlet at 164.2 ppm. We also carried out a ¹³C-labelling study with ¹³CO₂, which produced 2-¹³C (90 % yield). The ¹H NMR spectrum of 2-¹³C displays a doublet at 8.13 ppm that is due to coupling (¹J_{H-C} = 210.8 Hz) with the ¹³C atom arising from ¹³CO₂ (see the Supporting Information). In the solid-state IR spectrum of **2**, a peak corresponding to the C=O stretching vibration was detected at 1720 cm⁻¹. These results indicate the presence of a formate group (OCHO) in **2**, which was decisively confirmed by X-ray diffractometry (Scheme 1b). The P1–O1 distance (1.808(1) Å) is 11 % longer than a typical P–O single bond (1.63 Å). The formation of phosphorus formate **2** rather than adduct **I** (Figure 1a) demonstrates that one of the O atoms in CO₂ attacks the electrophilic P atom, whereas the C atom in CO₂ accepts the H atom as a hydride from **1**. Note that this result presents the first example of hydrophosphination of CO₂.

We postulated that the downfield ³¹P NMR chemical shift and the long P–O bond in **2** indicate an ionic property of the bond that induces zwitterionic character (**2'**; Scheme 1a), from which the formate group should be readily transferred to the appropriate acceptor. To bear out our hypothesis, we examined the reaction of **2** with a hydrosilane because the expected formation of a strong Si–O bond could be a driving force for the reaction. Under CO₂ atmosphere, treatment of **2** with a half equivalent of Ph₂SiH₂ afforded Ph₂Si(OCHO)₂ (**3**) as the major product, concomitant with the formation of the siloxane **4** (3/4 = 2.3:1 after 15 min; Scheme 1c). We also observed that **3** was converted into **4** after 18 h under the reaction conditions (Figures S1, S2 in the Supporting Information). The formation of **3** demonstrates the transfer of formate from **2** to Ph₂SiH₂, along with regeneration of **1**. During the reaction, however, only a peak for **2** was detected in the ³¹P NMR spectrum, probably owing to a rapid reaction between the in situ regenerated **1** and CO₂. Therefore, we examined the subsequent addition insertion of the P–H bond of **1** into CO₂ followed by formate transfer from **2** to silane in a catalytic process.


CO₂ was introduced into a J-young NMR tube filled with a CD₃CN solution of Ph₂SiH₂ and 5 mol % of **1**, and the reaction was monitored by NMR spectroscopy. Within 1 h at room temperature, the conversion of Ph₂SiH₂ into **3** (> 95 %) was confirmed, and only small amounts of siloxane **4** (< 5 %) were detected (Scheme 2, Figure S3-1). We also performed



Scheme 2. Catalytic hydrosilylation of CO₂ with 5 mol % of **1**.

a ¹³C-labeling experiment for the catalytic reaction under ¹³CO₂ atmosphere and confirmed the clean formation of Ph₂Si(O¹³CHO)₂ (**3**-¹³C; ¹J_{H-C} = 232.5 Hz; Figures S3, S4).

Recently, catalytic N-formylation and N-methylation of amines using CO₂ as the carbon source were reported with various organocatalysts.^[23,24] In these reactions, formate derivatives are considered to be key intermediates for the formation of formamides.^[23–25] Therefore, we attempted the one-pot N-formylation of amines with **1** as a catalyst. To investigate the reaction, *N*-methylaniline was utilized as a test substrate. After a brief screening of the reaction conditions (Table S3–1), we obtained the optimized conditions as shown in Scheme 3. Significantly, when ¹³CO₂ was used, ¹³C-labelled *N*-methyl-*N*-phenylformamide was obtained (see the Supporting Information).



Scheme 3. N-formylation of PhMeNH with CO₂ catalyzed by **1**.

Table 1: Scope of catalytic N-formylation of secondary amines with CO₂.

$\text{R}'\text{-N(R)H} \xrightarrow[\text{CD}_3\text{CN, RT}]{\text{cat. 1 (5 mol\%), 3 Ph}_2\text{SiH}_2, \text{CO}_2} \text{R}'\text{-N(R)C(=O)H} \text{ or } \text{R}'\text{-N(Me)C(=O)H}$					
5	t [h]	Product	Yield [%] ^[a]	TON	TOF [hr ⁻¹]
5a	0.5	6a	95	19	38
5b	2	7b	61	12	6
5c	0.5	6c	94	19	38
5d	7	7d	53	11	2
5e	2	6e	94	19	10
5f	5	6f	87	17	4
5g	5	6g	93	19	4
5h	6	6h	78	16	3
5i	18	6i	71	14	1
5j	6	6j	94	19	3
5k	23	6k	64	13	0.6

[a] Yields were determined by ¹H NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.


With the optimized conditions in hand, we examined the scope of the catalytic reaction with various secondary amines (Table 1). Sterically less-hindered aliphatic amines (**5a**, **5c**, **5e**) afforded the corresponding N-formylamines in excellent yields (**6a** 95 %, **6c** 94 %, **6e** 94 %). Meanwhile, reactions with sterically hindered and highly basic amines such as diisopropylamine (**5b**) and 2,2,4,4-tetramethylpiperidine (**5d**), afforded N-methylated amines (**7b**: 61 %, **7d**: 53 %), which were probably formed by subsequent reduction of the corresponding N-formylamines (**6b**, **6d**).^[24,26] Secondary amines containing aryl substituents (**5f–k**) were well tolerated and the corresponding formamides (**6f–k**) were produced in good to excellent yields (94–64 %). To expand the scope of the reaction, we also tested primary amines (Table 2). All of the amines employed (**8a–j**) were well tolerated and the corresponding formamides (**9a–j**) were produced in 72–99 % yields.

To gain insight into the reaction mechanism, we performed control reactions. In the absence of silane, a mixture of *N*-methylaniline and **1** under CO₂ atmosphere only afforded **2**, and no further reaction between **2** and *N*-methylaniline was observed. We also confirmed that *N*-methylaniline does not react with Ph₂SiH₂. Similarly, no apparent reaction was observed between **1** and silane. Mean-

Table 2: Scope of the catalytic N-formylation of primary amines with CO₂.

$\text{R-NH}_2 \xrightarrow[\text{CD}_3\text{CN, RT}]{\text{cat. 1 (5 mol\%), 3 Ph}_2\text{SiH}_2, \text{CO}_2} \text{R-NH-C(=O)H}$					
8	t [h]	Product	Yield [%] ^[a]	TON	TOF [hr ⁻¹]
8a	2	9a	84	17	9
8b	18	9b	97	19	1
8c	24	9c	85	17	0.7
8d	10	9d	> 99	20	2
8e	10	9e	72	14	1
8f	4	9f	97	19	5
8g	2	9g	91	18	9
8h	2	9h	99	20	10
8i	2	9i	82	16	8
8j	2	9j	90	18	9

[a] Yields were determined by ¹H NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.



Scheme 4. Proposed mechanism for the N-formylation of amines with Ph₂SiH₂ and CO₂ catalyzed by **1**.

while, addition of *N*-methylaniline to Ph₂Si(OCO)₂ (**3**) gave *N*-formyl-*N*-methylaniline after 2 h at room temperature in the absence of **1** and **2**. Therefore, we propose a mechanism (Scheme 4) for the N-formylation of amines that involves the following three steps: 1) insertion of the P–H bond of **1** into CO₂ to generate **2**, 2) transfer of a formate from **2** to Ph₂SiH₂ to form Ph₂Si(OCO)₂ (**3**) along with the regeneration of **1**, 3) formation of formamides and also silane byproducts such as silanols and (Ph₂SiO)₃ through the reaction of **3** with amines. Catalyst **1** thus acts as a reducing agent for CO₂ to form **2**, which subsequently transfers its formate moiety to silane. This mechanism differs from that proposed for the NHC–silane system in the N-formylation of amines,^[23,24] where an initial activation of silane by NHC is energetically more favorable than the mechanism encompassing the formation of an NHC–CO₂ adduct.^[26]

In summary, we have demonstrated the first hydrophosphination of CO₂ with **1**, subsequent transfer of formate from

2 to Ph_2SiH_2 , and catalytic N-formylation of amines under metal-free and ambient conditions.

Acknowledgements

We are grateful to the Singapore Ministry of Education (MOE2013-T2-1-005) and Nanyang Technological University (NTU) for financial support. We thank to Dr. Rakesh Ganguly (NTU) for assistance with X-ray crystallographic analysis.

Keywords: CO_2 activation · hydrophosphination · hydrosilylation · metal-free catalysis · N-formylation

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 12116–12120
Angew. Chem. **2015**, *127*, 12284–12288

- [1] a) D. M. D'Alessandro, B. Smit, J. R. Long, *Angew. Chem. Int. Ed.* **2010**, *49*, 6058–6082; *Angew. Chem.* **2010**, *122*, 6194–6219; b) M. Z. Jacobson, *Energy Environ. Sci.* **2009**, *2*, 148–173; c) G. Olah, A. Goepfert, G. K. Surya Prakash, *J. Org. Chem.* **2009**, *74*, 487–498; d) *Carbon Dioxide Capture and Storage* (Eds.: B. Metz, O. Davidson, H. de Coninck, M. Loos, L. Meyer), Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, New York, **2005**.
- [2] a) Q. Liu, L. Wu, R. Jackstell, M. Beller, *Nat. Commun.* **2015**, *6*, 5933; b) A. Tlili, E. Blondiaux, X. Frogneux, T. Cantat, *Green Chem.* **2015**, *17*, 157–168; c) C. S. Yeung, V. M. Dong, *Top. Catal.* **2014**, *57*, 1342–1350; d) C. Maeda, Y. Miyazaki, T. Ema, *Catal. Sci. Technol.* **2014**, *4*, 1482–1497; e) A. M. Appel, J. E. Bercau, A. B. Bocarsly, H. Dobbek, D. L. DuBois, M. Dupuis, J. G. Ferry, E. Fujita, R. Hille, P. J. A. Kenis, C. A. Kerfeld, R. H. Morris, C. H. F. Peden, A. R. Portis, S. W. Ragsdale, T. B. Rauchfuss, J. N. H. Reek, L. C. Seefeldt, R. K. Thauer, G. L. Waldrop, *Chem. Rev.* **2013**, *113*, 6621–6658; f) S. N. Riduan, Y. Zhang, *Dalton Trans.* **2010**, *39*, 3347–3357; g) T. Sakakura, J. Choi, H. Yasuda, *Chem. Rev.* **2007**, *107*, 2365–2387.
- [3] Selected recent examples and references therein: a) N. M. Rezayee, C. A. Huff, M. S. Sanford, *J. Am. Chem. Soc.* **2015**, *137*, 1028–1031; b) H. Suh, L. M. Guard, N. Hazari, *Chem. Sci.* **2014**, *5*, 3859–3872; c) S. Moret, P. J. Dyson, G. Laurenczy, *Nat. Commun.* **2014**, *5*, 4017; d) S. Bontemps, L. Vendier, S. Sabo-Etienne, *J. Am. Chem. Soc.* **2014**, *136*, 4419–4425; e) A. Berkefeld, W. E. Piers, M. Parvez, L. Castro, L. Maron, O. Eisenstein, *Chem. Sci.* **2013**, *4*, 2152–2162; f) S. Bontemps, S. Sabo-Etienne, *Angew. Chem. Int. Ed.* **2013**, *52*, 10253–10255; *Angew. Chem.* **2013**, *125*, 10443–10445; g) R. Dobrovetsky, D. W. Stephan, *Angew. Chem. Int. Ed.* **2013**, *52*, 2516–2519; *Angew. Chem.* **2013**, *125*, 2576–2579; See also: h) C. Lescot, D. U. Nielsen, I. S. Makarov, A. T. Lindhardt, K. Daasbjerg, T. Skrydstrup, *J. Am. Chem. Soc.* **2014**, *136*, 6142–6147.
- [4] F. Fontaine, M. Courtemanche, M. Légaré, *Chem. Eur. J.* **2014**, *20*, 2990–2996.
- [5] a) D. Martin, M. Soleilhavoup, G. Bertrand, *Chem. Sci.* **2011**, *2*, 389–399; b) P. P. Power, *Nature* **2010**, *463*, 171–177; c) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2010**, *49*, 46–76; *Angew. Chem.* **2010**, *122*, 50–81.
- [6] C. M. Mömming, E. Otten, G. Kehr, R. Fröhlich, S. Grimme, D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2009**, *48*, 6643–6646; *Angew. Chem.* **2009**, *121*, 6770–6773.
- [7] D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2015**, *54*, 6400–6441; *Angew. Chem.* **2015**, *127*, 6498–6541.
- [8] Selected recent examples: a) M. Courtemanche, A. P. Pulis, É. Rochette, M. Légaré, D. W. Stephan, F. Fontaine, *Chem. Commun.* **2015**, *51*, 9797–9800; b) G. Ménard, T. M. Gilbert, J. A. Hatnean, A. Kraft, I. Krossing, D. W. Stephan, *Organometallics* **2013**, *32*, 4416–4422; c) G. Ménard, D. W. Stephan, *Angew. Chem. Int. Ed.* **2011**, *50*, 8396–8399; *Angew. Chem.* **2011**, *123*, 8546–8549; d) X. Zhao, D. W. Stephan, *Chem. Commun.* **2011**, *47*, 1833–1835; e) D. A. Dickie, E. N. Coker, R. A. Kemp, *Inorg. Chem.* **2011**, *50*, 11288–11290; f) G. Ménard, D. W. Stephan, *J. Am. Chem. Soc.* **2010**, *132*, 1796–1797; Selected examples of the catalytic transformation of CO_2 : g) T. Wang, D. W. Stephan, *Chem. Eur. J.* **2014**, *20*, 3036–3039; h) T. Wang, D. W. Stephan, *Chem. Commun.* **2014**, *50*, 7007–7010; i) C. Das Neves Gomes, E. Blondiaux, P. Thuéry, T. Cantat, *Chem. Eur. J.* **2014**, *20*, 7098–7106; j) M. Courtemanche, M. Légaré, L. Maron, F. Fontaine, *J. Am. Chem. Soc.* **2013**, *135*, 9326–9329; k) Y. Wang, Y. Wang, W. Zhang, X. Lu, *J. Am. Chem. Soc.* **2013**, *135*, 11996–12003; l) R. Wehmschulte, M. Saleh, D. R. Powell, *Organometallics* **2013**, *32*, 6812–6819; m) M. Courtemanche, J. Larouche, M. Légaré, W. Bi, L. Maron, F. Fontaine, *Organometallics* **2013**, *32*, 6804–6811; n) A. Schäfer, W. Saak, D. Haase, T. Müller, *Angew. Chem. Int. Ed.* **2012**, *51*, 2981–2984; *Angew. Chem.* **2012**, *124*, 3035–3038; o) M. Khandelwal, R. J. Wehmschulte, *Angew. Chem. Int. Ed.* **2012**, *51*, 7323–7326; *Angew. Chem.* **2012**, *124*, 7435–7439; p) S. N. Riduan, Y. Zhang, J. Y. Ying, *Angew. Chem. Int. Ed.* **2009**, *48*, 3322–3325; *Angew. Chem.* **2009**, *121*, 3372–3375; q) Y. Kayaki, M. Yamamoto, T. Ikariya, *Angew. Chem. Int. Ed.* **2009**, *48*, 4194–4197; *Angew. Chem.* **2009**, *121*, 4258–4261.
- [9] a) L. J. Murphy, K. N. Robertson, R. A. Kemp, H. M. Tuononen, J. A. C. Clyburne, *Chem. Commun.* **2015**, *51*, 3942–3956; b) I. Purushothaman, S. De, P. Parameswaran, *RSC Adv.* **2014**, *4*, 60421–60428.
- [10] a) C.-C. Chong, R. Kinjo, *ACS Catal.* **2015**, *5*, 3238–3259; b) R. Declercq, G. Bouhadir, D. Bourissou, M. Légaré, M. Courtemanche, K. S. Nahi, N. Bouchard, F. Fontaine, L. Maron, *ACS Catal.* **2015**, *5*, 2513–2520; c) M. A. Courtemanche, M. A. Légaré, L. Maron, F. Fontaine, *J. Am. Chem. Soc.* **2014**, *136*, 10708–10717; See also: d) C. Lim, A. M. Holder, J. T. Hynes, C. B. Musgrave, *Inorg. Chem.* **2013**, *52*, 10062–10066.
- [11] a) I. Knopf, C. C. Cummins, *Organometallics* **2015**, *34*, 1601–1603; b) Z. Lu, H. Hausmann, S. Becker, H. A. Wagner, *J. Am. Chem. Soc.* **2015**, *137*, 5332–5335; c) K. Fujiwara, S. Yasuda, T. Mizuta, *Organometallics* **2014**, *33*, 6692–6695; d) I. Peuser, R. C. Neu, X. Zhao, M. Ulrich, B. Schirmer, J. A. Tannert, G. Kehr, R. Fröhlich, S. Grimme, G. Erker, D. W. Stephan, *Chem. Eur. J.* **2011**, *17*, 9640–9650; e) A. Berkefeld, W. E. Piers, M. Parvez, *J. Am. Chem. Soc.* **2010**, *132*, 10660–10661; f) A. E. Ashley, A. L. Thompson, D. O'Hare, *Angew. Chem. Int. Ed.* **2009**, *48*, 9839–9843; *Angew. Chem.* **2009**, *121*, 10023–10027; g) T. Wartik, R. K. Pearson, *J. Inorg. Nucl. Chem.* **1958**, *7*, 404–411; h) T. Wartik, R. K. Pearson, *J. Am. Chem. Soc.* **1955**, *77*, 1075.
- [12] a) G. Ménard, D. W. Stephan, *Dalton Trans.* **2013**, *42*, 5447–5453; b) P. Kuo, I. Chen, J. Chang, M. Lee, C. Hu, C. Hung, H. M. Lee, J. Huang, *Eur. J. Inorg. Chem.* **2004**, 4898–4906.
- [13] J. A. B. Abdalla, I. M. Riddlestone, R. Tirfoin, S. Aldridge, *Angew. Chem. Int. Ed.* **2015**, *54*, 5098–5102; *Angew. Chem.* **2015**, *127*, 5187–5191.
- [14] a) A. Jana, G. Tavčar, H. W. Roesky, M. John, *Dalton Trans.* **2010**, *39*, 9487–9489; b) A. Jana, D. Ghoshal, H. W. Roesky, I. Objartel, G. Schwab, D. Stalke, *J. Am. Chem. Soc.* **2009**, *131*, 1288–1293; For Si, see: c) P. Arya, J. Boyer, R. J. P. Corriu, G. F. Lanneau, M. Perrot, *J. Organomet. Chem.* **1988**, *346*, C11–C14; d) M. Mehta, M. H. Holthausen, I. Mallov, M. Pérez, Z. Qu, S. Grimme, D. W. Stephan, *Angew. Chem. Int. Ed.* **2015**, *54*, 8250–8254; *Angew. Chem.* **2015**, *127*, 8368–8372.
- [15] a) A. Jana, H. W. Roesky, C. Schulzke, A. Döring, *Angew. Chem. Int. Ed.* **2009**, *48*, 1106–1109; *Angew. Chem.* **2009**, *121*, 1126–

- 1129; b) G. Albertin, S. Antoniutti, J. Castro, S. García-Fontán, G. Zanardo, *Organometallics* **2007**, 26, 2918–2930.
- [16] a) R. G. Cavell, K. I. The, L. V. Griend, *Inorg. Chem.* **1981**, 20, 3813–3818; b) K. I. The, L. V. Griend, W. A. Whitla, R. G. Cavell, *J. Am. Chem. Soc.* **1977**, 99, 7379–7380.
- [17] L. J. Hounjet, C. B. Caputo, D. W. Stephan, *Angew. Chem. Int. Ed.* **2012**, 51, 4714–4717; *Angew. Chem.* **2012**, 124, 4792–4795.
- [18] M. Courtemanche, M. Légaré, É. Rochette, F. Fontaine, *Chem. Commun.* **2015**, 51, 6858–6861.
- [19] a) B. Chatelet, L. Joucla, J. Dutasta, A. Martinez, K. C. Szeto, V. Dufaud, *J. Am. Chem. Soc.* **2013**, 135, 5348–5351; b) R. W. Light, L. D. Hutchins, R. T. Paine, C. F. Campana, *Inorg. Chem.* **1980**, 19, 3597–3604; c) G. Oertel, H. Malz, H. Holtschmidt, *Chem. Ber.* **1964**, 97, 891–902.
- [20] For relevant examples, see: a) D. R. Kindra, I. J. Casely, M. E. Fieser, J. W. Ziller, F. Furche, W. J. Evans, *J. Am. Chem. Soc.* **2013**, 135, 7777–7787; b) S. Yin, J. Maruyama, T. Yamashita, S. Shimada, *Angew. Chem. Int. Ed.* **2008**, 47, 6590–6593; *Angew. Chem.* **2008**, 120, 6692–6695.
- [21] a) D. Gudat, *Acc. Chem. Res.* **2010**, 43, 1307–1316; b) S. Burck, D. Gudat, M. Nieger, W. W. D. Mont, *J. Am. Chem. Soc.* **2006**, 128, 3946–3955; c) D. Gudat, A. Haghverdi, M. Nieger, *Angew. Chem. Int. Ed.* **2000**, 39, 3084–3086; *Angew. Chem.* **2000**, 112, 3211–3214.
- [22] a) C.-C. Chong, H. Hirao, R. Kinjo, *Angew. Chem. Int. Ed.* **2015**, 54, 190–194; *Angew. Chem.* **2015**, 127, 192–196; b) C.-C. Chong, H. Hirao, R. Kinjo, *Angew. Chem. Int. Ed.* **2014**, 53, 3342–3346; *Angew. Chem.* **2014**, 126, 3410–3414.
- [23] a) C. Das Neves Gomes, O. Jacquet, C. Villiers, P. Thuéry, M. Ephritikhine, T. Cantat, *Angew. Chem. Int. Ed.* **2012**, 51, 187–190; *Angew. Chem.* **2012**, 124, 191–194; b) O. Jacquet, C. D. N. Gomes, M. Ephritikhine, T. Cantat, *J. Am. Chem. Soc.* **2012**, 134, 2934–2937; For a recent example with metal catalyst, see: c) L. Zhang, Z. Han, X. Zhao, Z. Wang, K. Ding, *Angew. Chem. Int. Ed.* **2015**, 54, 6186–6189; *Angew. Chem.* **2015**, 127, 6284–6287.
- [24] a) E. Blondiaux, J. Pouessel, T. Cantat, *Angew. Chem. Int. Ed.* **2014**, 53, 12186–12190; *Angew. Chem.* **2014**, 126, 12382–12386; b) S. Das, F. D. Bobbink, G. Laurenczy, P. J. Dyson, *Angew. Chem. Int. Ed.* **2014**, 53, 12876–12879; *Angew. Chem.* **2014**, 126, 13090–13093.
- [25] a) B. Wang, Z. Cao, *RSC Adv.* **2013**, 3, 14007–14015; See also: b) L. González-Sebastián, M. Flores-Alamo, J. J. García, *Organometallics* **2013**, 32, 7186–7194; c) S. Itagaki, K. Yamaguchi, N. Mizuno, *J. Mol. Catal. A* **2013**, 366, 347–352; d) R. Shintani, K. Nozaki, *Organometallics* **2013**, 32, 2459–2462; e) W. Sattler, G. Parkin, *J. Am. Chem. Soc.* **2012**, 134, 17462–17465.
- [26] a) O. Jacquet, X. Frogneux, C. D. N. Gomes, T. Cantat, *Chem. Sci.* **2013**, 4, 2127–2131; b) I. Sorribes, K. Junge, M. Beller, *Chem. Eur. J.* **2014**, 20, 7878–7883; See also: c) L. González-Sebastián, M. Flores-Alamo, J. J. García, *Organometallics* **2015**, 34, 763–769.

Received: June 10, 2015

Revised: July 5, 2015

Published online: August 14, 2015