


Copper-Catalyzed Formation of α -Alkoxy cycloalkenones from *N*-Tosylhydrazones

Naijing Su, Juliana A. Theorell, Donald J. Wink, and Tom G. Driver*

Abstract: The combination of 20 mol % of copper iodide and lithium *tert*-butoxide triggers the formation of a broad range of substituted, functionalized α -alkoxy 2*H*-naphthalenones from readily available *N*-tosylhydrazones. The data suggests that this transformation occurs through cycloaddition of a copper carbenoid with an ester, followed by a Lewis acid-catalyzed [1,2] alkyl shift of the in situ generated alkoxyepoxide intermediate.

The reactivity of transition-metal carbene complexes has spurred significant innovation which dramatically simplifies the synthesis of complex, functionalized molecules.^[1,2] These divalent reactive intermediates have been widely exploited in cyclopropanation,^[3] cycloaddition,^[4] annulation,^[5] metathesis,^[6] and C–H bond functionalization processes.^[7] In contrast, the use of these reactive intermediates to trigger cascade processes has lagged behind the established transformations.^[8] Our group has pursued the synthesis of *N*-heterocycles from aryl azides by using transition-metal catalysts to generate and control metal *N*-aryl nitrenes in C–H bond-amination reactions.^[9] Our mechanistic experiments suggested that C(sp²)–H functionalization occurred through a tandem reaction in which C–N bond formation preceded C–H bond cleavage,^[10] and we exploited this mechanism by substituting the hydrogen with other migrating groups to achieve the formation of functionalized 3*H*-indoles, such as **4**, from trisubstituted styryl azides (**1**) through an electrocyclozation-[1,2] carboxylate rearrangement (Scheme 1).^[11] During the course of our investigations, we were curious if this reactivity pattern was unique to rhodium *N*-aryl nitrenes or if it was general to other divalent catalytic intermediates, such as metal aryl carbenes. While metal carbenes are established to react with ketones and aldehydes to afford epoxides,^[12,13] esters coordinate with the electrophilic carbenes to produce a dipole which can be captured through a dipolar cycloaddition reaction.^[4] In the absence of a dipolarophile, we anticipated that an electrocyclozation/migration sequence might occur to afford the indene **6**. Alternatively, a [2+1] cycloaddition, unprecedented with an ester, could occur to produce the alkoxy-substituted epoxide **7**, which might rearrange under the reaction conditions because it is well-established that enol-ester-derived epoxides undergo acyloxy




Scheme 1. Divergent reactivity of metal divalent catalytic intermediates. esp = $\alpha,\alpha,\alpha',\alpha'$ -tetramethyl-1,3-benzene-dipropionate.

migrations to produce α -acyloxy ketones.^[14,15] Herein, we report the copper-catalyzed transformation of *N*-tosylhydrazones into α -alkoxy ketones through a [1,2] alkyl-shift of an in situ generated alkoxy-substituted epoxide.^[16]

To determine if related reactivity patterns are embedded in metal carbenoids, the reactivity of the *N*-tosylhydrazone **11a** was investigated (Table 1). *N*-Tosylhydrazones were chosen as the carbene precursor because they are well-established as a safe and nontoxic source of alkyl diazo compounds,^[17,18] and the requisite substrate is rapidly accessible from a Suzuki cross-coupling reaction between the aryl boronate **10** and vinyl triflate **9**. To our delight, exposure of

Table 1: Development of the optimized reaction conditions for 2*H*-naphthalenone.



Entry	Catalyst	mol %	T [°C]	Yield [%] ^[a]
1	CuI	20	90	98
2	CuBr	20	90	68
3	CuCl	20	90	30
4	(CuOTf) ₂ ·PhMe	20	90	43
5	Cu(OTf) ₂	20	90	20
6	CuBr ₂	20	90	36
7	[Rh ₂ (O ₂ CMe) ₄]	5	90	dec.
8	AlCl ₃	5	90	dec.
9	none	n.a.	90	10
10	CuI	10	90	76
11	CuI	20	50	20

[a] Determined using ¹H NMR spectroscopy with CH₂Br₂ as an internal standard. DME = dimethoxyethane, OTf = trifluoromethylsulfonate, Ts = 4-methyltoluenesulfonate.

* N. Su, J. A. Theorell, Prof. Dr. D. J. Wink, Prof. Dr. T. G. Driver
Department of Chemistry, University of Illinois at Chicago
845 West Taylor Street, Chicago, IL 60607 (USA)
E-mail: tgdriver@uic.edu
Homepage: <http://www2.chem.uic.edu/driver>

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201505993>.


Table 3: (Continued)

Entry	13	N-tosylhydrazone	2H-naphthalenone	Yield [%] ^[a]
14	13n			quant. d.r. 55:45 ^[d]
15	13o			40 ^[f]

[a] Yield of **14** isolated after neutral alumina chromatography, only product obtained. [b] Reaction performed in PhH. [c] Determined using ¹H NMR spectroscopy. [d] Determined using SFC-MS. [e] Identity of the major diastereomer determined by nOe spectroscopic experiments. [f] Styrene elimination product formed in 23 %. Ac = acetyl, Boc = *tert*-butoxycarbonyl.

the hydrazone R¹ group. When the aldehyde-derived **13c** was submitted to the reaction conditions, the naphthalenol **14c** was produced (Table 3, entry 3). Next, the composition of the *ortho*-alkenyl substituent was modified (entries 4–12). While shrinking the ring-size led to an attenuated yield (entry 4), substrates containing medium-sized *o*-cycloalkenyl rings were smoothly converted into the α -methoxy-2H-naphthalenones **14e** and **14f** (entries 5 and 6). The reaction also tolerated *o*-heterocyclic substituents to afford **14g–i** (entries 7–9). This reaction does tolerate acyclic *o*-alkenyl groups: the hydrazone **13j** was efficiently converted into the 2H-naphthalenone **14j** albeit with a reduced yield (entry 10). The diastereoselectivity of the reaction was next probed (entries 11–14). While exposure of the hydrazone **13k** to the reaction conditions provided a 3:2 mixture of diastereomers, the chiral, non-racemic *o*-norbornyl **13l** afforded the 2H-naphthalenone product as a single diastereomer. Next, the effect of moving the one of the bridge attachments from the allylic to the homoallylic position was examined with the β -pinene-derived **13m** (entry 13). Alleviating the steric pressure around the carboxylate substituent improved the yield of the transformation without the loss of stereoselectivity to produce **14m** in 71 % as a single diastereomer. In contrast, no diastereoselectivity was observed with the (–)-menthol-derived carboxylate **13n** (entry 14). This reaction could also be extended to the hydrazone **13o**, which contained an *o*-aryl group, to produce **14o** in 40 % (entry 15). Elimination, however, also occurred to form the 2-substituted styrene in 23 %.^[19]


The conversion of the *N*-tosylhydrazones into α -methoxy-2H-naphthalenones is proposed to occur through the tandem cycloaddition/[1,2] rearrangement of a copper carbene (Scheme 2). The reaction of *N*-tosylhydrazone with *tert*-butoxide generates the diazo **15**,^[20] which is decomposed by the copper(I) salt to produce the copper carbene **16**.^[21] The epoxide **18** could be produced through either a stepwise reaction with the proximal methyl carboxylate via the ylide **17** or directly through a [2+1] cycloaddition reaction.^[12] Alter-



Scheme 2. Possible mechanisms for 2H-naphthalenone formation.

natively, **18** could be formed through a noncarbene mechanism.^[22] In this mechanism, the copper catalyst activates the ester for attack by **15** to produce **22**.^[23] Loss of N₂ forms the benzyl cation **23**, which is attacked by the copper alkoxide to produce **18**. Ring-opening of the epoxide generates the oxocarbenium ion **19**, which triggers a [1,2] methyl shift to afford **20**.^[16] Dissociation of the copper catalyst produces the product **12a**.

Several experiments were performed in an attempt to gain insight into the reactivity of the proposed catalytic intermediates.^[19] First, the addition of TEMPO (2,2,6,6-tetramethylpiperidine 1-oxyl) to the reaction mixture did not produce any new products to indicate that radical intermediates, if formed, did not escape the solvent sheath. Attempts to trap either **17** or **19** with a dipolarophile (DMAD; dimethylacetylenedicarboxylate)^[4] or an external nucleophile (Et₃SiH) did not change the reaction outcome: only the 2H-naphthalenone **12** was formed.^[19] Next, we surveyed the reactivity of the *N*-tosylhydrazones **24** to determine if the *o*-alkenyl group was required for a successful reaction (Scheme 3). Irrespective of the potential ring formed, only β -hydride elimination was observed. Finally, to examine the potential for an alkoxy-substituted epoxide to rearrange, **27** was synthesized and its reactivity toward Lewis and Brønsted acids was tested. We found that while it remained inert toward mixtures of CuI and



Scheme 3. Experiments to elucidate the mechanism of the tandem reaction.

tert-butanol,^[22] exposure of **27** to 5 mol % of a stronger Lewis acid, AlCl₃, at room temperature triggered the formation of the α -methoxymethyl ketone **28**. We interpret this result to mean that epoxides, such as **18**, can rearrange provided that a strong enough Lewis acid is formed during the reaction.

In conclusion, we have discovered a new reactivity pattern for metal aryl carbenes toward pendant carboxylates to transform readily accessible aryl *N*-tosylhydrazones into functionalized α -alkoxy 2*H*-naphthalenones. Currently, we are working towards understanding the mechanism of this reaction to exploit the reactivity of these metal aryl carbenes and trigger additional tandem reactions to facilitate access to complex functionalized carbocycles.

Acknowledgements

We are grateful to the Office of the Vice Chancellor of Research at the University of Illinois at Chicago for their generous support.

Keywords: carbenes · copper · cyclizations · hydrazones · synthetic methods

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 12942–12946
Angew. Chem. **2015**, *127*, 13134–13138

- [1] a) M. P. Doyle, M. A. McKerver, T. Ye, *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides*, Wiley, New York, **1998**; b) F. Zaragoza-Dorwald, *Metal Carbenes in Organic Synthesis*, Wiley-VCH, Weinheim, **1998**; c) “Metal Carbenes in Organic Synthesis”: *Topics in Organometallic Chemistry*, Vol. 13 (Ed.: K. H. Dötz), Springer, Heidelberg, **2004**; d) “Handbook for Metathesis” (Eds.: R. H. Grubbs, D. J. O’Leary), Wiley-VCH, Weinheim, **2014**.
- [2] a) M. P. Doyle, D. C. Forbes, *Chem. Rev.* **1998**, *98*, 911; b) H. Lebel, J.-F. Marcoux, C. Molinaro, A. B. Charette, *Chem. Rev.* **2003**, *103*, 977; c) P. Müller, *Acc. Chem. Res.* **2004**, *37*, 243; d) Z. Li, C. He, *Eur. J. Org. Chem.* **2006**, 4313; e) H. M. L. Davies, J. R. Manning, *Nature* **2008**, *451*, 417; f) H. M. L. Davies, J. R. Denton, *Chem. Soc. Rev.* **2009**, *38*, 3061; g) M. P. Doyle, R. Duffy, M. Ratnikov, L. Zhou, *Chem. Rev.* **2010**, *110*, 704.
- [3] T. Katsuki, *Adv. Synth. Catal.* **2002**, *344*, 131.
- [4] a) A. Padwa, S. K. Bur, *Tetrahedron* **2007**, *63*, 5341; b) A. Padwa, *Prog. Heterocycl. Chem.* **2009**, *20*, 20; c) A. Padwa, *J. Org. Chem.* **2009**, *74*, 6421.
- [5] a) T. Horneff, S. Chuprakov, N. Chernyak, V. Gevorgyan, V. V. Fokin, *J. Am. Chem. Soc.* **2008**, *130*, 14972; b) S. Chuprakov, S. W. Kwok, L. Zhang, L. Lercher, V. V. Fokin, *J. Am. Chem. Soc.* **2009**, *131*, 18034; c) N. Grimster, L. Zhang, V. V. Fokin, *J. Am. Chem. Soc.* **2010**, *132*, 2510; d) B. Chattopadhyay, V. Gevorgyan, *Angew. Chem. Int. Ed.* **2012**, *51*, 862; *Angew. Chem.* **2012**, *124*, 886.
- [6] a) R. H. Grubbs, S. J. Miller, G. C. Fu, *Acc. Chem. Res.* **1995**, *28*, 446; b) R. R. Schrock, A. H. Hoveyda, *Angew. Chem. Int. Ed.* **2003**, *42*, 4592; *Angew. Chem.* **2003**, *115*, 4740; c) A. Deiters, S. F. Martin, *Chem. Rev.* **2004**, *104*, 2199; d) G. C. Vougioukalakis, R. H. Grubbs, *Chem. Rev.* **2010**, *110*, 1746; e) C. Samojłowicz, M. Bieniek, K. Grela, *Chem. Rev.* **2009**, *109*, 3708.
- [7] a) H. M. L. Davies, R. E. J. Beckwith, *Chem. Rev.* **2003**, *103*, 2861; b) H. M. L. Davies, J. R. Manning, *Nature* **2008**, *451*, 417; c) H. M. L. Davies, J. Du Bois, J.-Q. Yu, *Chem. Soc. Rev.* **2011**, *40*, 1855; d) S. Chuprakov, F. W. Hwang, V. Gevorgyan, *Angew. Chem. Int. Ed.* **2007**, *46*, 4757; *Angew. Chem.* **2007**, *119*, 4841; e) H. M. L. Davies, Y. Lian, *Acc. Chem. Res.* **2012**, *45*, 923.
- [8] a) P. F. Korkowski, T. R. Hoye, D. B. Rydberg, *J. Am. Chem. Soc.* **1988**, *110*, 2676; b) A. Padwa, D. C. Dean, L. Zhi, *J. Am. Chem. Soc.* **1989**, *111*, 6451; c) T. R. Hoye, C. J. Dinsmore, D. S. Johnson, P. F. Korkowski, *J. Org. Chem.* **1990**, *55*, 4518; d) T. R. Hoye, C. J. Dinsmore, *J. Am. Chem. Soc.* **1991**, *113*, 4343; e) A. Padwa, R. L. Chinn, S. F. Hornbuckle, Z. J. Zhang, *J. Org. Chem.* **1991**, *56*, 3271; f) A. Padwa, D. J. Austin, Y. Gareau, J. M. Kassir, S. L. Xu, *J. Am. Chem. Soc.* **1993**, *115*, 2637; g) Y. Ni, J. Montgomery, *J. Am. Chem. Soc.* **2004**, *126*, 11162; h) N. Majumdar, K. A. Korthals, W. D. Wulff, *J. Am. Chem. Soc.* **2012**, *134*, 1357; i) S. Jansone-Popova, J. A. May, *J. Am. Chem. Soc.* **2012**, *134*, 17877; j) S. Jansone-Popova, P. Q. Le, J. A. May, *Tetrahedron* **2014**, *70*, 4118.
- [9] a) B. J. Stokes, H. Dong, B. E. Leslie, A. L. Pumphrey, T. G. Driver, *J. Am. Chem. Soc.* **2007**, *129*, 7500; b) M. Shen, B. E. Leslie, T. G. Driver, *Angew. Chem. Int. Ed.* **2008**, *47*, 5056; *Angew. Chem.* **2008**, *120*, 5134; c) K. Sun, R. Sachwani, K. J. Richert, T. G. Driver, *Org. Lett.* **2009**, *11*, 3598; d) A. L. Pumphrey, H. Dong, T. G. Driver, *Angew. Chem. Int. Ed.* **2012**, *51*, 5920; *Angew. Chem.* **2012**, *124*, 6022; e) Q. Nguyen, K. Sun, T. G. Driver, *J. Am. Chem. Soc.* **2012**, *134*, 7262.
- [10] B. J. Stokes, K. J. Richert, T. G. Driver, *J. Org. Chem.* **2009**, *74*, 6442.
- [11] a) K. Sun, S. Liu, P. M. Bec, T. G. Driver, *Angew. Chem. Int. Ed.* **2011**, *50*, 1702; *Angew. Chem.* **2011**, *123*, 1740; b) B. J. Stokes, S. Liu, T. G. Driver, *J. Am. Chem. Soc.* **2011**, *133*, 4702; c) C. Kong, N. Jana, T. G. Driver, *Org. Lett.* **2013**, *15*, 824; d) C. Jones, Q. Nguyen, T. G. Driver, *Angew. Chem. Int. Ed.* **2014**, *53*, 785; *Angew. Chem.* **2014**, *126*, 804.
- [12] a) P. de March, R. Huisgen, *J. Am. Chem. Soc.* **1982**, *104*, 4952; b) M. Alt, G. Maas, *Tetrahedron* **1994**, *50*, 7435; c) M. P. Doyle, W. Hu, D. J. Timmons, *Org. Lett.* **2001**, *3*, 933; d) H. M. L. Davies, J. DeMeese, *Tetrahedron Lett.* **2001**, *42*, 6803; e) S. Muthusamy, C. Gunanathan, M. Nethaji, *Synlett* **2004**, 639; f) Z.-C. Li, J. Zhang, W.-H. Hu, Z.-Y. Chen, X.-Q. Yu, *Synlett* **2005**, 1711.
- [13] More common is for the metal ylide to transfer the carbene to a sulfide, which reacts with an aldehyde or ketone to form an epoxide. For leading examples, see: a) A. W. Johnson, R. B. LaCount, *J. Am. Chem. Soc.* **1961**, *83*, 417; b) E. J. Corey, M. Chaykovsky, *J. Am. Chem. Soc.* **1965**, *87*, 1353; c) V. K. Aggarwal, H. Abdel-Rahman, R. V. H. Jones, H. Y. Lee, B. D. Reid, *J. Am. Chem. Soc.* **1994**, *116*, 5973; d) V. K. Aggarwal, J. G. Ford, A. Thompson, R. V. H. Jones, M. C. H. Standen, *J. Am. Chem. Soc.* **1996**, *118*, 7004; e) V. K. Aggarwal, A. Ali, M. P. Coogan, *J. Org. Chem.* **1997**, *62*, 8628; f) A.-H. Li, L.-X. Dai, V. K. Aggarwal, *Chem. Rev.* **1997**, *97*, 2341; g) V. K. Aggarwal, E. Alonso, G. Hynd, K. M. Lydon, M. J. Palmer, M. Porcelloni, J. R. Studley, *Angew. Chem. Int. Ed.* **2001**, *40*, 1430; *Angew. Chem.* **2001**, *113*, 1479; h) O. Illa, M. Namutebi, C. Saha, M. Ostovar, C. C. Chen, M. F. Haddow, S. Nocquet-Thibault, M. Lusi, E. M. McGarrigle, V. K. Aggarwal, *J. Am. Chem. Soc.* **2013**, *135*, 11951.
- [14] For seminal references on acid-catalyzed rearrangement of acetoxy-substituted epoxides, see: a) A. H. Soloway, W. J. Conside, D. K. Fukushima, T. F. Gallagher, *J. Am. Chem. Soc.* **1954**, *76*, 2941; b) N. S. Leeds, D. K. Fukushima, T. F. Gallagher, *J. Am. Chem. Soc.* **1954**, *76*, 2943; c) P. D. Gardner, *J. Am. Chem. Soc.* **1956**, *78*, 3421; d) W. S. Johnson, B. Gastambide, R. Pappo, *J. Am. Chem. Soc.* **1957**, *79*, 1991; e) H. J. Shine, G. E. Hunt, *J. Am. Chem. Soc.* **1958**, *80*, 2434; f) K. L. Williamson, W. S. Johnson, *J. Org. Chem.* **1961**, *26*, 4563.
- [15] For more recent examples, see: a) X. Feng, L. Shu, Y. Shi, *J. Am. Chem. Soc.* **1999**, *121*, 11002; b) Y. Zhu, K. J. Manske, Y. Shi, *J. Am. Chem. Soc.* **1999**, *121*, 4080; c) S. Tranchimand, B. Faure, J.-

- V. Naubron, V. Alphand, A. Archelas, G. Iacazio, *Eur. J. Org. Chem.* **2012**, 4365.
- [16] In contrast, [1,2] alkyl rearrangements of epoxyethers are significantly more scarce. For leading reports, see: a) C. L. Stevens, S. J. Dykstra, *J. Am. Chem. Soc.* **1954**, 76, 4402; b) T. I. Temnikova, T. Gissel, B. A. Gontarev, *Zh. Obshch. Khim.* **1960**, 30, 776; c) W. J. Xia, D. R. Li, L. Shi, Y. Q. Tu, *Tetrahedron: Asymmetry* **2001**, 12, 1459; d) D. R. Li, W. J. Xia, L. Shi, Y. Q. Tu, *Synthesis* **2003**, 41.
- [17] a) S. R. Angle, M. L. Neitzel, *J. Org. Chem.* **2000**, 65, 6458; b) V. K. Aggarwal, E. Alonso, G. Fang, M. Ferrara, G. Hynd, M. Porcelloni, *Angew. Chem. Int. Ed.* **2001**, 40, 1433; *Angew. Chem.* **2001**, 113, 1482; c) V. K. Aggarwal, J. de Vicente, R. V. Bonnert, *Org. Lett.* **2001**, 3, 2785; d) V. K. Aggarwal, J. R. Fulton, C. G. Sheldon, J. de Vicente, *J. Am. Chem. Soc.* **2003**, 125, 6034; e) J. Barluenga, P. Moriel, C. Valdés, F. Aznar, *Angew. Chem. Int. Ed.* **2007**, 46, 5587; *Angew. Chem.* **2007**, 119, 5683; f) Q. Xiao, J. Ma, Y. Yang, Y. Zhang, J. Wang, *Org. Lett.* **2009**, 11, 4732; g) J. Barluenga, L. Riesgo, L. A. Lóez, E. Rubio, M. Tomás, *Angew. Chem. Int. Ed.* **2009**, 48, 7569; *Angew. Chem.* **2009**, 121, 7705; h) V. K. Aggarwal, *Nat. Chem.* **2009**, 1, 433; i) L. Zhou, F. Ye, Y. Zhang, J. Wang, *J. Am. Chem. Soc.* **2010**, 132, 13590; j) J. Barluenga, M. Tomás-Gamasa, F. Aznar, C. Valdés, *Angew. Chem. Int. Ed.* **2010**, 49, 4993; *Angew. Chem.* **2010**, 122, 5113.
- [18] For reviews, see: a) V. K. Aggarwal, C. L. Winn, *Acc. Chem. Res.* **2004**, 37, 611; b) J. R. Fulton, V. K. Aggarwal, J. de Vicente, *Eur. J. Org. Chem.* **2005**, 1479; c) J. Barluenga, C. Valdés, *Angew. Chem. Int. Ed.* **2011**, 50, 7486; *Angew. Chem.* **2011**, 123, 7626; d) X. Zhao, Y. Zhang, J. Wang, *Chem. Commun.* **2012**, 48, 10162; e) Q. Xiao, Y. Zhang, J. Wang, *Acc. Chem. Res.* **2013**, 46, 236.
- [19] Refer to the Supporting Information for more details.
- [20] a) W. R. Bamford, T. S. Stevens, *J. Chem. Soc.* **1952**, 4735; b) H. W. Davies, M. Schwarz, *J. Org. Chem.* **1965**, 30, 1242.
- [21] For mechanism studies on the formation of metal carbenes from diazo compounds, see: a) R. Cohen, B. Rybtchinski, M. Gandelman, H. Rozenberg, J. M. L. Martin, D. Milstein, *J. Am. Chem. Soc.* **2003**, 125, 6532; b) H. Lu, W. I. Dzik, X. Xu, L. Wojtas, B. de Bruin, X. P. Zhang, *J. Am. Chem. Soc.* **2011**, 133, 8518.
- [22] We thank the reviewer for this mechanistic suggestion.
- [23] For seminal reports of nucleophilic attack of diazo compounds onto carbonyls, see: a) E. Büchner, T. Curtius, *Ber. Dtsch. Chem. Ges.* **1885**, 18, 2371; b) F. Schlotterbeck, *Ber. Dtsch. Chem. Ges.* **1907**, 40, 479.
- [24] Exposure of the epoxide **27** to stoichiometric amounts of LiCl and *tert*-butanol produced only the hydrolyzed α -hydroxy phenyl ketone. See the Supporting Information for more details.

Received: June 30, 2015

Revised: July 22, 2015

Published online: September 7, 2015