

NHC Catalysis

International Edition: DOI: 10.1002/anie.201510106
German Edition: DOI: 10.1002/ange.201510106Enantioselective N-Heterocyclic Carbene Catalysis by the Umpolung of α,β -Unsaturated Ketones

Yuji Nakano and David W. Lupton*

Abstract: N-Heterocyclic carbene-catalyzed formation of β -anionic intermediates from enones has been employed in the enantioselective synthesis of 2-aryl propionates. The reaction was achievable using a homochiral 4-MeOC₆H₄ morpholinone catalyst allowing the first example of enantioselective catalysis by umpolung of α,β -unsaturated ketones. The reaction is high yielding, and shows robustness with reasonable generality. A mechanism is proposed in which the enantiodetermining protonation is achieved using either hexafluoroisopropanol or the formed naphthol product.

Central to many N-heterocyclic carbene (NHC)-catalyzed^[1] reactions are the homoenolate (**1**)^[2] or α,β -unsaturated acyl azolium (**2**; Figure 1).^[3] These form through initial 1,2-addition to an α,β -unsaturated carbonyl, and have allowed a wealth of chemical transformations to be discovered. In contrast, β -azolium ylide **3**, a species afforded through 1,4-addition to α,β -unsaturated carbonyls followed by tautomerization,^[4] as reported by Fu in 2006,^[5] has been largely overlooked in synthesis. While many nucleophilic catalysts undergo 1,4-addition to conjugate acceptors, this usually

triggers reactions by the resultant enolate (**4**).^[6] Tautomerization to reveal a β -anion is rare and, to our knowledge, unique to NHC catalysis. Although β -anion **3** was introduced nearly a decade ago, its utility remains unclear with limited studies reported to date. Notably Matsuoka^[7a,c-g] and Glorius^[7b] have introduced tail-to-tail dimerization of enolates, while Chen has examined polymerization catalysis via intermediates **3** and **4**.^[8] Presumably, the paucity of studies on this intermediate, compared to the plethora on the aforementioned intermediates **1** and **2**, is related to challenges in achieving reaction discovery without triggering reactions of enolate **4**, which are known with NHCs.^[6a-d]

The underdeveloped chemistry of β -azolium ylide **3**, specifically as it relates to enantioselective catalysis, piqued our interest in this area. To this end, we envisaged a cycloisomerization of *bis*-conjugate acceptors (**5**) to 2-aryl propionates (**6**; Eq. 1) as a method to expand knowledge of these neglected intermediates. Herein, we report the outcome of these studies, which have allowed the discovery of the first enantioselective reaction involving the β -anion **3** intermediate. The reaction provides a method to access 2-aryl propionates in up to 91:9 e.r. and good yields. Mechanistic studies, and a discussion of unsuitable substrates, provides a picture of a reaction balanced between competing side reactions, and with sensitivity to alcohol-containing additives. In addition to providing information regarding β -anion **3**, the products contain the 2-aryl propionate motifs found extensively in non-steroidal anti-inflammatory drugs (NSAIDs).^[9]

Studies commenced with the preparation of diene **5a** using procedures modified from Gravel.^[10] When exposed to the Enders TPT^[11] catalyst, we were pleased to isolate 2-aryl propionate **6a** in 92 % yield (Table 1, entry 1). The discovery of an enantioselective variant proved significantly more challenging, and commenced with a survey of new and known chiral catalysts. Based on previous success with morpholinone-based NHCs,^[12] a screen of six catalysts bearing various N-substituents was undertaken (Table 1, entries 2–7). This approach also reflected observations of Chen who found that polymerization of acrylate derivatives was sensitive to the nature of the azolium, and the N-substitution pattern.^[8,13] In this case, N-*t*-butyl (**A1**) and N-C₆F₅ (**A2**) were not viable, while N-Mes (**A3**), N-Ph (**A4**), N-4-MeOC₆H₄ (**A5**), and N-2,6-(MeO)₂C₆H₃ (**A6**) all gave the expected product with moderate enantioselectivity. The highest selectivity of 74:26 e.r. was obtained with N-4-MeOC₆H₄ (**A5**). Both the yield and the enantioselectivity were depressed using alternate bases (Table 1, entries 8 and 9). The decrease with KHMDS was striking, and speaks to the role of the conjugate acid (hexamethyldisilazane) in the enantiodetermining protonation (see below). Introduction of





Figure 1. Overview of NHC reactions.

[*] Y. Nakano, Prof. Dr. D. W. Lupton
School of Chemistry, Monash University
Clayton 3800, Victoria (Australia)
E-mail: david.lupton@monash.edu


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201510106>.

example bis-Michael acceptor **7**, rather than undergoing cyclization, provided dimer **9**, presumably through a reaction analogous to those of Glorius and Matsuoka (Scheme 1, eq. 3).^[7] In addition, increasing the electrophilicity of the second Michael acceptor, as in ketone **10**, diverted the reaction, with the competing Rauhaut–Currier product **11** being formed (eq. 4).

The sensitivity of the reaction to catalyst loading prompted an examination of non-linear effects (NLE). In this case, no NLE was observed (Scheme 2A), implicating a single equivalent of catalyst **A5** in the enantiodetermining event. An alternate explanation for the sensitivity to catalyst loading would involve the phenolic product **6** participating in the enantiodetermining step, and thus the catalyst to substrate ratio would impact the enantioselectivity. Support for this scenario can be gleaned from the enhanced enantioselectivity (84:16 vs. 74:16 e.r.) observed in the formation of **6h** using naphthol **6a** as an additive (Table 2, footnote c). Further-



Scheme 1. Competing reactions.




Scheme 2. Mechanistic studies.

more, in the absence of HFIP, the *ee* of naphthol **6a** progressively increased from 28% after two minutes, to 54% at reaction completion (Scheme 2B). Reintroduction of HFIP gave a reaction that proceeded at 70% *ee* for its duration. Finally, support for this scenario was obtained when the cycloisomerization was performed with deuterated substrate **D-5a**. In addition to deuteration of the phenolic hydrogen, around 25% deuterium incorporation was observed at the 2-position (eq. 6). To determine whether proton transfer between these positions occurs following reaction completion, racemic **6a** and enantioenriched **6a** were subjected to catalyst **A5** and **TPT** respectively. In both cases, no change in the optical purity was observed.^[16] Thus, a plausible mechanism (Scheme 3) for this reaction commences with NHC addition to the terminal olefin to afford enolate **I** which, following tautomerization, provides the key β -azolium ylide **II**. Conjugate addition proceeds through intermediate **II**, with a *trans* arrangement between the azolium and the acceptor, and the azolium positioned to minimize steric interactions between the benzyl group and the phenyl linker, to afford enolate **III**. Diastereoselective protonation of **III** then affords **IV**, which, following E1cB reaction and tautomerization, affords propionate **6a**. An alternate explanation would involve E1cB prior to protonation, a mechanistic scenario difficult to distinguish. When the reaction was monitored by NMR, the only azolium-type structure was the free NHC, indicating that this is likely the resting state of the catalyst.

The kinetics of the reaction are finely balanced, with the desired cyclization (**II**→**III**) moderately faster than reaction with a second equivalent of substrate (that is, **7**→**9**, eq. 3). Furthermore, proton transfer to afford β -azolium ylide **II** is slightly faster than Rauhaut–Currier cyclization (that is, **10**→**11**, eq. 4).

The umpolung of Michael acceptors by 1,4-addition of NHCs is an unusual activation mode, in comparison to the



Scheme 3. Plausible mechanism.

more commonly observed 1,2-addition chemistry. Herein, a cycloisomerization reaction that exploits this reaction pathway is reported. This study has allowed the first enantioselective reaction through the β -azolium ylide to be discovered. For this transformation, electron-rich triazolium-derived NHCs were ideal, while an alcohol additive improved the enantioselectivity. Although the enantioselectivity is not exceptional, these studies provide a strong framework for catalyst and reaction design likely to inform future studies exploiting the β -azolium ylide **3**.

Acknowledgements

The authors thank Dr. Joel Hooper, Dr. Craig Forsyth and Mr. Adam Ametovski (Monash University) for supporting studies, and the Australian Research Council through the Discovery (DP120101315) and Future Fellowship (FT110100319) programs for financial support.

Keywords: 1,4-addition · 2-aryl propionates · enantioselective catalysis · N-heterocyclic carbenes · β -azolium ylides

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 3135–3139
Angew. Chem. **2016**, *128*, 3187–3191

- [1] For a selection of recent reviews on NHC catalysis, see: a) D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.* **2007**, *107*, 5606; For homoenolate chemistry see: b) V. Nair, R. S. Menon, A. T. Biju, C. R. Sinu, R. R. Paul, A. Jose, V. Sreekumar, *Chem. Soc. Rev.* **2011**, *40*, 5336; For acyl azolium enolates see: c) H. U. Vora, P. Wheeler, T. Rovis, *Adv. Synth. Catal.* **2012**, *354*, 1617; d) J. Douglas, G. Churchill, A. D. Smith, *Synthesis* **2012**, *44*, 2295; For cascade catalysis, see: e) A. Grossmann, D. Enders, *Angew. Chem. Int. Ed.* **2012**, *51*, 314; *Angew. Chem.* **2012**, *124*, 320; For acyl anion chemistry, see: f) X. Bugaut, F. Glorius, *Chem. Soc. Rev.* **2012**, *41*, 3511; For applications in total synthesis, see: g) J. Izquierdo, G. E. Hutson, D. T. Cohen, K. A. Scheidt, *Angew. Chem. Int. Ed.* **2012**, *51*, 11686; *Angew. Chem.* **2012**, *124*, 11854; For acyl anion-free catalysis, see: h) S. J. Ryan, L. Candish, D. W. Lupton, *Chem. Soc. Rev.* **2013**, *42*, 4906; For catalysis under oxidative conditions, see: i) S. De Sarkar, A. Biswap, R. C. Samanta, A. Studer, *Chem. Eur. J.* **2013**, *19*, 4664; For acyl azoliums and enol azoliums, see: j) J. Mahatthananchai, J. W. Bode, *Acc. Chem. Res.* **2014**, *47*, 696; For reactions with esters, see: k) P. Chauhan, D. Enders, *Angew. Chem. Int. Ed.* **2014**, *53*, 1485; *Angew. Chem.* **2014**, *126*, 1509; For an introduction to NHCs, see: l) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, *Nature* **2014**, *510*, 485; D. M. Flanagan, F. Romanov-Michailidis, N. A. White, T. Rovis, *Chem. Rev.* **2015**, *115*, 9307.
- [2] Early examples of homolenolate **1** chemistry: a) K. Y.-K. Chow, J. W. Bode, *J. Am. Chem. Soc.* **2004**, *126*, 8126; b) S. S. Sohn, E. L. Rosen, J. W. Bode, *J. Am. Chem. Soc.* **2004**, *126*, 14370; c) C. Burstein, F. Glorius, *Angew. Chem. Int. Ed.* **2004**, *43*, 6205; *Angew. Chem.* **2004**, *116*, 6331.
- [3] Early examples of unsaturated acyl azolium **2** chemistry: a) K. Zeitler, *Org. Lett.* **2006**, *8*, 637; b) B. E. Maki, A. Chean, E. M. Phillips, K. A. Scheidt, *Org. Lett.* **2007**, *9*, 371; c) S. J. Ryan, L. Candish, D. W. Lupton, *J. Am. Chem. Soc.* **2009**, *131*, 14176; d) S. J. Ryan, L. Candish, D. W. Lupton, *J. Am. Chem. Soc.* **2011**, *133*, 4694.
- [4] For studies on related azolium dienamines intermediates formed by substitution chemistry, see: a) C. E. I. Knappke, J.-M. Neudörfl, A. J. von Wangelin, *Org. Biomol. Chem.* **2010**, *8*, 1695; b) C. E. I. Knappke, A. J. Arduengo III, J. Haijun, J.-M. Neudörfl, A. J. von Wangelin, *Synthesis* **2011**, 3784; c) R. N. Reddi, P. K. Prasad, A. Sudalai, *Angew. Chem. Int. Ed.* **2015**, *54*, 14150; *Angew. Chem.* **2015**, *127*, 14356; For formation from addition to styrene, see: d) M. Schedler, N. E. Wurz, C. G. Danilius, F. Glorius, *Org. Lett.* **2014**, *16*, 3134; For studies into the properties of such intermediates, see: e) A. Berkessel, S. Elfert, *Adv. Synth. Catal.* **2014**, *356*, 571; f) B. Maji, M. Horn, H. Mayr, *Angew. Chem. Int. Ed.* **2012**, *51*, 6231; *Angew. Chem.* **2012**, *124*, 6335; g) B. Maji, H. Mayr, *Angew. Chem. Int. Ed.* **2012**, *51*, 10408; *Angew. Chem.* **2012**, *124*, 10554.
- [5] C. Fischer, S. W. Smith, D. A. Powell, G. C. Fu, *J. Am. Chem. Soc.* **2006**, *128*, 1472.
- [6] For NHC reactions through enolate **4**, see: a) D. Enders, K. Breuer, U. Kallfass, T. Balensiefer, *Synthesis* **2003**, 35, 1292; b) L. He, T.-Y. Jian, S. Ye, *J. Org. Chem.* **2007**, *72*, 7466; c) X.-Y. Chen, S. Ye, *Org. Biomol. Chem.* **2013**, *11*, 7991; d) A. Bhunia, S. Thorat, R. G. Gonnade, A. T. Biju, *Chem. Commun.* **2015**, 13690; For a review on MBH chemistry, see: e) Y. Wei, M. Shi, *Comprehensive Enantioselective Organocatalysis: Catalysts, Reactions and Applications* (Ed.: P. I. Dalko), Wiley-VCH, Weinheim, **2013**; For a comparison of nucleophilic catalysts in MBH reactions, see: f) L. Candish, Y. Nakano, D. W. Lupton, *Synthesis* **2014**, *46*, 1823.
- [7] a) S.-i. Matsuoka, Y. Ota, A. Washio, A. Katada, K. Ichioka, K. Takagi, M. Suzuki, *Org. Lett.* **2011**, *13*, 3722; b) A. T. Biju, M. Padmanaban, N. E. Wurz, F. Glorius, *Angew. Chem. Int. Ed.* **2011**, *50*, 8412; *Angew. Chem.* **2011**, *123*, 8562; c) S.-i. Matsuoka, Y. Tochigi, K. Takagi, M. Suzuki, *Tetrahedron* **2012**, *68*, 9836; d) S.-i. Matsuoka, S. Namera, A. Washio, K. Takagi, M. Suzuki, *Org. Lett.* **2013**, *15*, 5916; e) T. Kato, Y. Ota, S.-i. Matsuoka, K. Takagi, M. Suzuki, *J. Org. Chem.* **2013**, *78*, 8739; f) T. Kato, S.-i. Matsuoka, M. Suzuki, *J. Org. Chem.* **2014**, *79*, 4484; g) S.-i. Matsuoka, M. Nakazawa, M. Suzuki, *Bull. Chem. Soc. Jpn.* **2015**, *88*, 1093; h) O.-a. Rajachan, M. Paul, V. R. Yatham, J.-M. Neudörfl, K. Kanokmedhakul, S. Kanokmedhakul, A. Brekesel, *Tetrahedron Lett.* **2015**, *56*, 6537; For 1,4-addition to unsaturated sulfones, see: i) R. L. Aienza, H. S. Ruth, K. A. Scheidt, *Chem. Sci.* **2011**, *2*, 1772.
- [8] a) Y. Zhang, E. Y.-X. Chen, *Angew. Chem. Int. Ed.* **2012**, *51*, 2465; *Angew. Chem.* **2012**, *124*, 2515; b) M. Hong, E. Y.-X. Chen, *Angew. Chem. Int. Ed.* **2014**, *53*, 11900; *Angew. Chem.* **2014**, *126*, 12094.
- [9] For selected approaches to the synthesis of enantioenriched aryl propiolates, see: D. E. Griswold, J. L. Adams, *Med. Res. Rev.* **1996**, *16*, 181; b) C. V. Rao, B. S. Reddy, *Curr. Cancer Drug Targets* **2004**, *4*, 29.
- [10] E. Sanchez-Larios, J. M. Halmes, C. L. Daschner, M. Gravel, *Synthesis* **2011**, *12*, 1896.
- [11] a) D. Enders, K. Breuer, J. H. Teles, *Helv. Chim. Acta* **1996**, *79*, 1217; For the properties and stoichiometric reactions of this carbene, see: b) D. Enders, K. Breuer, G. Raabe, J. Runsink, J. H. Teles, J.-P. Melder, K. Ebel, S. Brode, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1021; *Angew. Chem.* **1995**, *107*, 1119.
- [12] a) L. Candish, C. M. Forsyth, D. W. Lupton, *Angew. Chem. Int. Ed.* **2013**, *52*, 9149; *Angew. Chem.* **2013**, *125*, 9319; b) L. Candish, A. Levens, D. W. Lupton, *J. Am. Chem. Soc.* **2014**, *136*, 14397; c) A. Levens, C. Zhang, L. Candish, C. M. Forsyth, D. W. Lupton, *Org. Lett.* **2015**, *17*, 5332.
- [13] For discussions on the impact of NHC electronics on reaction outcome, see: a) T. Rovis, *Chem. Lett.* **2008**, *37*, 2; b) J. Mahatthananchai, J. W. Bode, *Chem. Sci.* **2012**, *3*, 192; c) C. J. Collett, R. S. Massey, O. R. Maguire, A. S. Batsanov, A. C. O'Donoghue, A. D. Smith, *Chem. Sci.* **2013**, *4*, 1514; d) C. J. Collett, R. S. Massey, J. E. Taylor, O. R. Maguire, A. C. O'Do-

- noghue, A. D. Smith, *Angew. Chem. Int. Ed.* **2015**, *54*, 6887; *Angew. Chem.* **2015**, *127*, 6991.
- [14] For use of carboxylic acid additives in NHC catalysis, see: a) X. D. Zhao, D. A. DiRocco, T. Rovis, *J. Am. Chem. Soc.* **2011**, *133*, 12466; b) G. Liu, P. D. Wilkerson, C. A. Toth, H. Xu, *Org. Lett.* **2012**, *14*, 858; c) Z. Fu, H. Sun, S. Chen, B. Tiwari, G. Li, Y. R. Chi, *Chem. Commun.* **2013**, *49*, 261; d) S. Kuwano, S. Harada, B. Kang, R. Oriez, Y. Yamaoka, K. Takasu, K. Yamada, *J. Am. Chem. Soc.* **2013**, *135*, 11485; e) E. O. McCusker, K. A. Scheidt, *Angew. Chem. Int. Ed.* **2013**, *52*, 13616; *Angew. Chem.* **2013**, *125*, 13861; f) Y. Lin, L. Yang, Y. Deng, G. Zhong, *Chem. Commun.* **2015**, *51*, 8330; For use of catechol additives in NHC catalysis, see: g) D. A. DiRocco, T. Rovis, *J. Am. Chem. Soc.* **2011**, *133*, 10402; For use of HFIP and other alcohols as additives in NHC catalysis, see: h) J. Chen, Y. Huang, *Nat. Commun.* **2014**, *5*, 3437; and reference [7f].
- [15] Single crystal X-ray analysis of anthracene **6k** was performed using CuK α radiation for assignment of absolute stereochemistry. CCDC 1443409 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [16] This result also precludes Brønsted base-mediated chemistry as being the origin of enantioselectivity.

Received: October 30, 2015

Revised: December 24, 2015

Published online: January 28, 2016