

Homogeneous Catalysis


International Edition: DOI: 10.1002/anie.201602155
German Edition: DOI: 10.1002/ange.201602155

Rhodium-Catalyzed Cross-Cyclotrimerization and Dimerization of Allenes with Alkynes

Kazuki Sakashita, Yu Shibata, and Ken Tanaka*

Abstract: It has been established that a cationic rhodium(I)/binap complex catalyzes the cross-cyclotrimerization of two molecules of a monosubstituted allene with one molecule of a functionalized alkyne to give 3,6-dialkylidenecyclohex-1-enes. In contrast, the reactions involving di- or trisubstituted allenenes and/or unfunctionalized alkynes afforded cross-dimerization products, substituted dendralenes, through β -hydrogen elimination from the corresponding rhodacycles.

Transition-metal-catalyzed intermolecular cross-cyclotrimerization and cross-dimerization reactions of alkynes with alkenes are efficient and atom-economical methods for the synthesis of substituted six-membered carbocycles and dienes, respectively.^[1] In contrast, similar transformations using allenenes in place of alkenes have been reported in a limited number of examples.^[2] As examples of the cross-cyclotrimerization of alkynes with allenenes,^[3–5] the nickel-catalyzed reactions of one molecule of a monosubstituted allene with two molecules of an electron-deficient internal alkyne followed by aromatization through olefin isomerization, which affords the corresponding pentasubstituted benzenes, has been reported (Scheme 1a).^[3a] However, the usable alkynes are limited to electron-deficient alkynoates, and electron-rich alkynes cannot be employed.^[3a] Furthermore, a selective and general method for the cross-cyclotrimerization of two molecules of an allene and one molecule of an alkyne has not been reported to date.^[5] As examples of the cross-dimerization of allenenes with alkynes, thermal reactions giving substituted dendralenes have been reported (Scheme 1b).^[6–8] However, the usable alkynes are limited to electron-deficient dimethyl acetylenedicarboxylate and (trifluoromethyl)acetylene.^[6] Herein, we disclose the cross-cyclotrimerization of two molecules of a monosubstituted allene with functionalized alkynes, and cross-dimerization of di- or trisubstituted allenenes with both functionalized and unfunctionalized alkynes.



Scheme 1. Previously reported cross-cyclotrimerization and cross-dimerization of allenenes with alkynes. dppe = 1,2-bis(diphenylphosphino)ethane.


Recently, our research group reported the rhodium-catalyzed enantioselective [2+2+2] cycloaddition of tosylamide-linked 5-allenals and 5-allenones with internal alkynes.^[9] In this study, we found that the reaction of the tosylamide-linked 5-allenone with di-*tert*-butylacetylenedicarboxylate in the presence of a cationic rhodium(I)/binap catalyst does not afford the expected [2+2+2] cycloaddition product, but instead delivers an unexpected cross-cyclotrimerization product, 3,6-dialkylidenecyclohex-1-ene derivative, in good yield (Scheme 2).

Thus, we examined the reaction of the monosubstituted allene **1a** with di-*tert*-butylacetylenedicarboxylate (**2a**) in the presence of the same rhodium catalyst (Table 1). Pleasingly, the cross-cyclotrimerization proceeded at room temperature to give the corresponding 3,6-dialkylidenecyclohex-1-ene **3aa**, although the regioisomer **4aa** was generated as a minor product (entry 1). The effect of biaryl bis(phosphine) ligands was then examined (entries 1–5). Increasing the dihedral angle of ligands improved the ratio of **3aa** to **4aa** (dihedral angle:^[10] H₈-binap > binap > segphos, ratio of **3aa** to **4aa**: H₈-binap > binap > segphos). Increasing the steric


[*] K. Sakashita, Dr. Y. Shibata, Prof. Dr. K. Tanaka
Department of Applied Chemistry
Graduate School of Science and Engineering
Tokyo Institute of Technology
O-okayama, Meguro-ku, Tokyo 152-8550 (Japan)
E-mail: ktanaka@apc.titech.ac.jp
Homepage: <http://www.apc.titech.ac.jp/~ktanaka/>

K. Sakashita, Prof. Dr. K. Tanaka
Department of Applied Chemistry
Graduate School of Engineering
Tokyo University of Agriculture and Technology
Koganei, Tokyo 184-8588 (Japan)

Supporting information for this article can be found under:
<http://dx.doi.org/10.1002/anie.201602155>.



Scheme 2. Unexpected discovery of rhodium-catalyzed cross-cyclotrimerization of an allenone with di-*tert*-butylacetylenedicarboxylate. cod = 1,5-cyclooctadiene.


Table 1: Optimization of reaction conditions for rhodium-catalyzed cross-cyclotrimerization of **1a** with **2a**.^[a]


Entry	Ligand	Catalyst (mol %)	1a / 2a	Conv [%]	Yield [%] ^[b] (3aa / 4aa)
1	binap	10	1:1	> 99	68 (74:26)
2	segphos	10	1:1	> 99	69 (71:29)
3	H ₈ -binap	10	1:1	> 99	52 (83:17)
4	tol-binap	10	1:1	> 99	51 (79:21)
5	xyl-binap	10	1:1	46	18 (85:15)
6 ^[c]	binap	10	1:2	> 99	57 (74:26)
7 ^[d]	binap	10	2:1	> 99	47 (74:26)
8 ^[e]	binap	10	4:1	> 99	45 (74:26)
9 ^[f]	binap	5	1:1	> 99	70 (74:26)

[a] [Rh(cod)₂]BF₄ (0.010 mmol), ligand (0.010 mmol), **1a** (0.10 mmol), **2a** (0.10 mmol), and CH₂Cl₂ (2.0 mL) were used. [b] Yield of isolated product. [c] **1a** (0.10 mmol) and **2a** (0.20 mmol) were used. [d] **1a** (0.10 mmol) and **2a** (0.05 mmol) were used. [e] **1a** (0.20 mmol) and **2a** (0.050 mmol) were used. [f] [Rh(cod)₂]BF₄ (0.010 mmol), ligand (0.010 mmol), **1a** (0.20 mmol), **2a** (0.20 mmol), and CH₂Cl₂ (2.0 mL) were used. Binap = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, H₈-binap = 2,2'-bis(diphenylphosphino)-5,5',6,6',7,7',8,8'-octahydro-1,1'-binaphthyl, segphos = 5,5'-bis(diphenylphosphino)-4,4'-bi-1,3-benzodioxole, tol-binap = 2,2'-bis[di(2,2'-p-tolyl)phosphino]-1,1'-binaphthyl, xyl-binap = 2,2'-bis[di(3,5-xylyl)phosphino]-1,1'-binaphthyl.


bulk of aryl groups on the phosphorus improved the ratio of **3aa** to **4aa** (steric bulk: xyl-binap > tol-binap > binap, ratio of **3aa** to **4aa**: xyl-binap > tol-binap > binap). The use of binap afforded **3aa** in the highest yield, although the ratio of **3aa** to **4aa** was moderate (entry 1). The ratio of **1a** to **2a** was then examined (entries 1 and 6–8), and revealed that a 1:1 ratio afforded **3aa/4aa** in the highest yield (entry 1). In this reaction, the homo-cyclotrimerization of **2a** and the cross-cyclotrimerization of one molecule of **1a** with two molecules of **2a** were not observed. Finally, the catalyst loading could be reduced to 5 mol% without erosion of the product yield (entry 9).

The substrate scope of the rhodium-catalyzed cross-cyclotrimerization is shown in Scheme 3. With respect to alkynes, both electron-deficient and electron-rich symmetrical alkynes (**2a–c**) participated in this reaction to give the corresponding cyclohexenes **3aa–ac/4aa–ac** in good yields. Not only symmetrical alkynes but also the unsymmetrical alkyne **2d** reacted with **1a** to give the corresponding cyclohexenes **3ad/4ad** in good yield. Unfortunately, terminal alkynes could not be employed because of their rapid homo-cyclotrimerization. With respect to allenes, not only the aliphatic allene **1a** but also functionalized and aromatic ones (**1b,c**) reacted with **2a** to give the corresponding cyclohexenes **3ba** and **3ca** in moderate yields. Although the

**Scheme 3.** Rhodium-catalyzed cross-cyclotrimerization of allenes (**1**) with functionalized alkynes (**2**). [Rh(cod)₂]BF₄ (0.010 mmol), binap (0.010 mmol), **1** (0.20 mmol), **2** (0.20 mmol), and CH₂Cl₂ (2.0 mL) were used. The cited yields are of the isolated products.


reason is not clear, the reactions employing allenes **1b,c** did not afford the regioisomers **4** at all.

Interestingly, the reaction of the disubstituted allene **1d** with **2a** in the presence of the same rhodium catalyst did not afford a cyclotrimerization product, but instead the dimerization product, the tetrasubstituted dendralene^[11] **5da**, in good yield (Scheme 4). The use of the trisubstituted allene **1e** significantly increased the yield of the corresponding **5ea**. The substrate scope of this cross-dimerization was then examined. With respect to allenes, not only **1e** but also the functionalized **1f** reacted with **2a** to give the corresponding dendralene **5fa** in good yield. Not only dimethyl but also the cyclohexyl derivative **1g** could be employed. With respect to alkynes, both electron-deficient and electron-rich symmetrical alkynes (**2a–c** and **2e**) participated in this reaction to give the corresponding dendralenes in high yields. Unsymmetrical alkynes (**2d** and **2f**) could also be employed, although regioselectivity was moderate. It is worthy of note that the unfunctionalized alkynes **2g** and **2h** smoothly react with both functionalized and unfunctionalized trisubstituted allenes (**1e–g**) to give the corresponding dendralenes in moderate to good yields. Furthermore, the di- and monosubstituted unfunctionalized allenes (**1d,a**) were capable of reacting with **2g**, although the product yields were low. Not only internal alkynes but also terminal alkynes could be employed for this cross-dimerization with allenes. 1-Dodecyne (**2i**) reacted with **1e–g** to give the corresponding dendralenes **5ei–gi** in high yields. Not only 1-dodecyne (**2i**) but also a variety of alkyl and aryl acetylenes (**2j–o**) could be employed, although olefin isomerization products (**5'**) were generated in small amounts when employing primary-alkyl-substituted alkynes (**1i–k**). In addition, the silyl acetylenes **2p,q** and ethyl propiolate (**2r**) were capable of reacting with **1e** to give the corresponding dendralenes **5ep–er**.



Scheme 4. Rhodium-catalyzed cross-dimerization of allenes (**1**) with internal alkynes (**2**). [Rh(cod)₂]₂BF₄ (0.010 mmol), binap (0.010 mmol), **1** (0.20 mmol), **2** (0.20 mmol), and CH₂Cl₂ (2.0 mL) were used. The cited yields are of the isolated products. [a] d.r. = diastereomer ratio. See the Supporting Information for details. [b] rs = regioselectivity. [c] Major regioisomer is shown. [d] **2** (0.60 mmol) was used. [e] **1** (0.10 mmol) and **2** (0.10 mmol) were used. [f] **2** (0.30 mmol) was used.

Possible reaction mechanisms for the formation of the 3,6-dialkylidenecyclohex-1-enes **3** and **4**, and dendralene **5** from **1** and **2** are shown in Scheme 5. The reaction of **1** and **2** with




Scheme 5. Possible mechanisms for cross-cyclotrimerization and dimerization.

rhodium generates the rhodacyclopentene **A**. Insertion of another molecule of **1** into **A** generates the intermediate **B**, and then reductive elimination furnishes **3**. The minor regioisomer **4** may be generated from the intermediate **C** through insertion of the internal double bond of **1** into **A**. As **4ad** was obtained as a single regioisomer, the formation of **A** may be regioselective for the unsymmetrical alkyne **2d** (R⁴ = CO₂Et, R⁵ = Me). In contrast, the reaction of a di- or trisubstituted allene (**1**) and **2** with rhodium generates the rhodacyclopentene **D**. β-Hydrogen elimination affords the intermediate **E** and subsequent reductive elimination furnishes **5**. Avoiding steric repulsion between the allene substituent and ligand on rhodium might account for the observed regioselective formation of **A**, **B**, and **D**.


It was anticipated that the reaction of the phenyl-substituted allene **1h** and **2a** with rhodium would afford rhodacyclopentene **E'**, in which no β-hydrogen atoms exist (Scheme 6). However, the reaction of **1h** with **2a** afforded the dendralene **5ha** with a different substitution pattern, presumably through β-hydrogen elimination from the rhodacyclopentene **F**. Similarly, **1h** reacted with **2i** to give the regioisomeric dendralenes **5hi** and **5hi'** in high yield.

As dendralenes serve as good cycloaddition partners in Diels–Alder reactions,^[12] the reactions of **5** with tetracyanoethylene (**6**) were examined (Scheme 7). The reaction of an *E/Z* mixture of the pentasubstituted dendralene **5ee** with **6** afforded the corresponding vinylcyclohexene **7** in high yield, and *cis*-**5ee** was recovered. The tetrasubstituted dendralene **5ei** also reacted with **6** to give the corresponding vinylcyclohexene **8** in high yield.

In conclusion, we have established that a cationic rhodium(I)/binap complex catalyzes the cross-cyclotrimerization of two molecules of monosubstituted allenes with one molecule of functionalized alkynes to give 3,6-dialkylidenecyclohex-1-enes. In contrast, the reactions involving di- or trisubstituted allenes and/or unfunctionalized alkynes afforded the cross-dimerization products, substituted dendralenes, through β-hydrogen elimination from the correspond-



Scheme 6. Rhodium-catalyzed cross-dimerization of **1h** with **2a,i**.



Scheme 7. Diels–Alder reactions of **5ee** and **5ei** with **6**.

ing rhodacycles. Future work will focus on further exploration of the cationic rhodium(I) complex-catalyzed C–C bond-forming reactions involving allenes.

Acknowledgments

This work was supported partly by ACT-C from Japan Science and Technology Agency (JST, Japan) and a Grant-in-Aid for Scientific Research (No. 25105714) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT, Japan). We thank Takasago International Corporation for the gift of segphos, H₈-binap, tol-binap, and xyl-binap, and Umicore for generous support in supplying the rhodium complex.

Keywords: alkynes · allenes · cyclization · dimerization · rhodium

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 6753–6757
Angew. Chem. **2016**, *128*, 6865–6869

- [1] For selected recent reviews, see: a) Y. Satoh, Y. Obora, *Eur. J. Org. Chem.* **2015**, 5041; b) M. Amatore, C. Aubert, *Eur. J. Org. Chem.* **2015**, 265; c) G. Domínguez, J. Pérez-Castells in *Comprehensive Organic Synthesis II*, Vol. 5, 2nd ed. (Ed.: P. Knochel), Elsevier, Amsterdam, **2014**, pp. 1537–1581; d) *Transition-Metal-Mediated Aromatic Ring Construction* (Ed.: K. Tanaka), Wiley, Hoboken, **2013**, Chap. 1–11; e) D. L. J. Broere, E. Ruijter, *Synthesis* **2012**, 44, 2639; f) N. Weding, M. Hapke, *Chem. Soc. Rev.* **2011**, 40, 4525; g) R. Hua, M. V. A. Abrenica, P. Wang, *Curr. Org. Chem.* **2011**, 15, 712; h) G. Domínguez, J. Pérez-Castells, *Chem. Soc. Rev.* **2011**, 40, 3430; i) N. Saito, D. Tanaka, M. Mori, Y. Sato, *Chem. Rec.* **2011**, 11, 186; j) A. Pla-Quintana, A. Roglans, *Molecules* **2010**, 15, 9230; k) D. Leboeuf, V. Gandon, M. Malacria in *Handbook of Cyclization Reactions*, Vol. 1 (Ed.: S. Ma), Wiley-VCH, Weinheim, **2010**, p. 367; l) J. A. Varela, C. Saá, *J. Organomet. Chem.* **2009**, 694, 143; m) T. Shibata, K. Tsuchikama, *Org. Biomol. Chem.* **2008**, 6, 1317; n) P. R. Chopade, J. Louie, *Adv. Synth. Catal.* **2006**, 348, 2307; o) Y. Shibata, K. Tanaka, *Synthesis* **2012**, 44, 323; p) M. Fujiwara, I. Ojima in *Modern Rhodium-Catalyzed Organic Reactions* (Ed.: P. A. Evans), Wiley-VCH, Weinheim, **2005**, p. 129.
- [2] For selected recent reviews of the transition-metal-catalyzed oligomerization reactions involving allenes, see: a) A. Lledo, A. Pla-Quintana, A. Roglans, *Chem. Soc. Rev.* **2016**, 45, DOI: 10.1039/C5CS00535C; b) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria, A. Simonneau, *Chem. Rev.* **2011**, 111, 1954; c) S. Yu, S. Ma, *Chem. Commun.* **2011**, 47, 5384; d) S. Kotha, E. Brahmachary, K. Lahiri, *Eur. J. Org. Chem.* **2005**, 4741; e) *Modern Allene Chemistry* (Eds.: N. Krause, A. S. K. Hashmi), Wiley-VCH, Weinheim, **2004**.
- [3] For the transition-metal-catalyzed intermolecular cross-cyclotrimerization of one molecule of an allene with two molecules of an alkyne, see: a) M. Shanmugasundaram, M.-S. Wu, C.-H. Cheng, *Org. Lett.* **2001**, 3, 4233; for the transition-metal-catalyzed intermolecular cyclotrimerization of one molecule of allene with two molecules of benzyne, see: b) J.-C. Hsieh, D. K. Rayabarapu, C.-H. Cheng, *Chem. Commun.* **2004**, 532; c) Y.-L. Liu, Y. Liang, S.-F. Pi, X.-C. Huang, J.-H. Li, *J. Org. Chem.* **2009**, 74, 3199; for the transition-metal-catalyzed intermolecular cyclotrimerization of one molecule of 1,2-cyclohexadiene with two molecules of an alkyne, see: d) I. Quintana, D. Peña, D. Pérez, E. Guitián, *Eur. J. Org. Chem.* **2009**, 5519.
- [4] The nickel-catalyzed intermolecular cross-cyclotrimerization of one molecule of allene with two molecules of an isocyanate has been reported. See: T. Miura, M. Morimoto, M. Murakami, *J. Am. Chem. Soc.* **2010**, 132, 15836.
- [5] The nickel-catalyzed intermolecular cross-cyclotrimerization of two molecules of propadiene with one molecule of acetylene, propyne, phenylacetylene, and vinylacetylene has been reported, while both regioselectivity and product yields were low to moderate. See: a) R. E. Benson, R. V. Lindsey, Jr., *J. Am. Chem. Soc.* **1959**, 81, 4250; b) J. Cermak, V. Blechta, V. Chvalovsky, *Collect. Czech. Chem. Commun.* **1988**, 53, 1274.
- [6] For the thermal cross-dimerization of allenes and alkynes, see: a) J. C. Martin, P. J. Carter, J. L. Chitwood, *J. Org. Chem.* **1971**, 36, 2225; b) H.-A. Chia, B. E. Kirk, D. R. Taylor, *J. Chem. Soc. D* **1971**, 1144; c) H.-A. Chia, B. E. Kirk, D. R. Taylor, *J. Chem. Soc. Perkin Trans. 1* **1974**, 1209; d) C. B. Lee, R. J. J. Newman, D. R. Taylor, *J. Chem. Soc. Perkin Trans. 1* **1978**, 1161; e) B. B. Snider, D. J. Rodini, R. S. E. Conn, S. Sealfon, *J. Am. Chem. Soc.* **1979**, 101, 5283; f) D. J. Pasto, W. Kong, *J. Org. Chem.* **1988**, 53, 4807; g) K. V. Sajna, R. Kotikalapudi, M. Chakravarty, N. N. B. Kumar, K. C. K. Swamy, *J. Org. Chem.* **2011**, 76, 920.
- [7] The transition-metal-mediated reductive cross-coupling reactions of allenes with alkynes have been reported. See: a) K. Suzuki, T. Imai, S. Yamanoi, M. Chino, T. Matsumoto, *Angew.*

- Chem. Int. Ed. Engl.* **1997**, 36, 2469; *Angew. Chem.* **1997**, 109, 2578; b) S. Yatnanoi, T. Imai, T. Matsumoto, K. Suzuki, *Tetrahedron Lett.* **1997**, 38, 3031; c) D. Hideura, H. Urabe, F. Sato, *Chem. Commun.* **1998**, 271; d) S. Yamanoi, T. Matsumoto, K. Suzuki, *Tetrahedron Lett.* **1998**, 39, 9727; e) R. Tanaka, M. Sasaki, F. Sato, H. Urabe, *Tetrahedron Lett.* **2005**, 46, 329; f) H. L. Shimp, G. C. Micalizio, *Chem. Commun.* **2007**, 4531; g) H. L. Shimp, A. Hare, M. McLaughlin, G. C. Micalizio, *Tetrahedron* **2008**, 64, 3437; the transition-metal-catalyzed oxidative cross-coupling reactions of allenes with alkynes have been reported. See: h) G. Takahashi, E. Shirakawa, T. Tsuchimoto, Y. Kawakami, *Adv. Synth. Catal.* **2006**, 348, 837; i) D. Chen, X. Chen, Z. Lu, H. Cai, J. Shen, G. Zhu, *Adv. Synth. Catal.* **2011**, 353, 1474; j) S. Arai, Y. Amako, X. Yang, A. Nishida, *Angew. Chem. Int. Ed.* **2013**, 52, 8147; *Angew. Chem.* **2013**, 125, 8305.
- [8] The thermal intermolecular [2+2] cycloaddition of allenes with alkynes has been reported. See: a) W. Klop, L. Brandsma, *J. Chem. Soc. Chem. Commun.* **1983**, 988; b) H. Sakurai, M. Kudo, K. Sakamoto, Y. Nakadaira, M. Kira, A. Sekiguchi, *Chem. Lett.* **1988**, 1441; c) D. J. Pasto, W. Kong, *J. Org. Chem.* **1989**, 54, 3215; d) D. J. Pasto, K. D. Sugi, D. E. Alonso, *J. Org. Chem.* **1992**, 57, 1146; See also Ref. [6a,c–f]. The Lewis acid catalyzed intermolecular [2+2] cycloaddition of allenes with alkynes has been reported. See: e) M. Hojo, C. Murakami, S. Nakamura, A. Hosomi, *Chem. Lett.* **1998**, 331. The Zr/Cu-mediated intermolecular [2+2] cycloaddition of allenes with alkynes has been reported. See: f) W. Zheng, Y. Wu, F. Zheng, L. Hu, Y. Hong, *Tetrahedron Lett.* **2010**, 51, 4702.
- [9] K. Sakashita, K. Masutomi, K. Noguchi, K. Tanaka, *Chem. Lett.* **2014**, 43, 1260.
- [10] H. Shimizu, I. Nagasaki, T. Saito, *Tetrahedron* **2005**, 61, 5405.
- [11] For the synthesis of dendralenes by the transition-metal-catalyzed C–C bond-forming reactions, see: a) M. Arisawa, T. Sugihara, M. Yamaguchi, *Chem. Commun.* **1998**, 2615; b) T. Miura, T. Biyajima, T. Toyoshima, M. Murakami, *Beilstein J. Org. Chem.* **2011**, 7, 578; c) H. Wang, B. Beiring, D.-G. Yu, K. D. Collins, F. Glorius, *Angew. Chem. Int. Ed.* **2013**, 52, 12430; *Angew. Chem.* **2013**, 125, 12657.
- [12] For examples, see: T. Shibata, Y. Takesue, S. Kadowaki, K. Takagi, *Synlett* **2003**, 0268. See also Ref. [11b,c].

Received: March 3, 2016

Published online: April 25, 2016