

Fused-Ring Systems

International Edition: DOI: 10.1002/anie.201604765
German Edition: DOI: 10.1002/ange.201604765

Total Syntheses of Juglorescein and Juglocombins A and B

Shogo Kamo, Kai Yoshioka, Kouji Kuramochi,* and Kazunori Tsubaki

Abstract: Total syntheses of juglorescein and juglocombins A and B are reported. The highly oxygenated 6/6/5/6/6-fused pentacyclic ring system of these natural products was constructed through a bioinspired dimerization of 1,4-naphthoquinone. Notably, five new stereogenic centers were constructed in a single step by the dimerization reaction. The epoxide intermediate obtained from the dimerization was successfully converted into juglocombins A and B through photoinduced reduction of the epoxide, dehydration, and conversion of the resultant quinone into a hydroquinone derivative. The same epoxide intermediate was also converted into juglorescein through intramolecular lactonization, hydrolysis of the resulting lactone, and removal of the protecting groups. Furthermore, the relative and absolute configurations of juglorescein and juglocombins A and B were determined.

Juglorescein (**1**), and juglocombins A (**2**) and B (**3**) have been isolated from *Streptomyces* sp. 815 and GW4184 (Figure 1a).^[1] Because **2** and **3** are unstable and highly polar, they were converted into dimethyl esters (**2'** and **3'**) and 1'-O-methyljuglocombin B dimethyl ester (**4**) to confirm their structures. Although the structures were confirmed by detailed NMR studies, the relative and absolute configurations of the compounds remained undetermined. These compounds have a highly oxygenated, 6/6/5/6/6-fused A/B/C/D/E ring system. The related naphthoquinone dimers zeylanone,^[2a,b] zeylanone epoxide,^[2c] and two shikometabolins,^[2d] are shown in Figure 1b. Zeylanone shows antibacterial and antifungal activity.^[2b] This compound also exhibits cytotoxic activity against cancer cell lines.^[2b] Shikometabolins E and F show significant neuraminidase inhibitory activity.^[2d,e] Total synthesis of racemic zeylanone and zeylanone epoxide was achieved by our group.^[3] However, no total synthesis of **1–3** has been reported to date, because construction of this carbon framework with multiple contiguous stereogenic centers has proven challenging. Herein, we report the first total syntheses and determination of the absolute configurations of **1–3**. Construction of the ring system,




Figure 1. a) Structures of juglorescein (**1**), juglocombins A (**2**), and B (**3**), and their derivatives (**2'**, **3'**, and **4**). b) Related naphthoquinone dimers zeylanone, zeylanone epoxide, and shikometabolins E and F.


including the five stereogenic centers, was achieved through a bioinspired dimerization of a naphthoquinone monomer.

The proposed biosynthetic pathway to **1–3** involves the dimerization of juglomycin C (**5**; Scheme 1).^[1,4] Sequential intermolecular and intramolecular Michael addition of **5**, followed by oxidation of the resulting hydroquinone, gives **2** and **3**. Compound **3** is then enzymatically oxidized to yield **1**.


Our retrosynthetic analysis of **1–3** is illustrated in Scheme 2. We propose that these compounds can be synthesized from common intermediate **6**, which in turn can be prepared through a bioinspired dimerization reaction of the juglomycin C derivative **7**.^[1] We envisioned that this dimerization would proceed in a highly regio- and stereoselective manner via a five-membered cyclic transition state (**1**). Epoxidation of the resultant hydroquinone monoanion with oxygen^[5] would occur preferentially from the opposite side of the side chain at C-2'.^[3] Compound **7** can readily be prepared

[*] S. Kamo, K. Yoshioka, Prof. Dr. K. Kuramochi, Prof. Dr. K. Tsubaki
Graduate School for Life and Environmental Sciences
Kyoto Prefectural University
1-5 Shimogamo Hanki-cho, Sakyo-ku, Kyoto 606-8522 (Japan)
E-mail: kuramochi@kpu.ac.jp
Prof. Dr. K. Kuramochi
Present address: Department of Applied Biological Science
Faculty of Science and Technology, Tokyo University of Science
2641 Yamazaki, Noda, Chiba 278-8510 (Japan)
E-mail: kuramochi@rs.tus.ac.jp

Supporting information for this article can be found under:
<http://dx.doi.org/10.1002/anie.201604765>.




Scheme 1. Proposed biosynthesis of 1–3.

Scheme 2. Retrosynthetic analysis of 1–3. TBS = *tert*-butyldimethylsilyl, MOM = methoxymethyl.

from the known optically active epoxide (**8**)^[6] and 2-bromonaphthalene (**9**).^[7]

Our synthesis began with the generation of a Grignard reagent from aryl bromide **9**, which was treated with optically active epoxide **8** to give **10** in 57% yield (Scheme 3). The hydroxy group in **10** was protected as a methoxymethyl (MOM) ether to give **11** in 74% yield. Oxidation of **11** with iodobenzene diacetate (PIDA) gave juglomycin C derivative **7** in 90% yield.^[8] The key dimerization reaction of **7** under basic conditions using 1,8-diazabicyclo[5.4.0]undec-7-ene




a solution of **6**, sodium carbonate, sodium hydrosulfite, and **12**^[11] was irradiated under white light to give alcohol **14**.^[12] Thermal dehydration of alcohol **14** proceeded easily, but the resultant naphthoquinone was too unstable to isolate. Therefore, **14** was treated with dihydronicotinamide **13** under heating at 50 °C to give hydroquinone **15**. Without purification, methylation of the two phenolic hydroxy groups in **15**, followed by removal of the *tert*-butyldimethylsilyl (TBS) protecting groups with tetra-*n*-butylammonium fluoride (TBAF)/AcOH gave **16** in 30 % yield over the four steps from **6**. The relative and absolute configurations of **16** were determined by rotating frame nuclear Overhauser effect spectroscopy (ROESY) correlations and comparison of the calculated and experimental circular dichroism (CD) spectra (see Figure S1 in the Supporting Information).

The synthesis of juglocombins A and B is shown in Scheme 4. Oxidation of the primary alcohols in **16** through Dess–Martin oxidation^[13] and Pinnick oxidation^[14] gave dicarboxylic acid **17** in 94 % yield over two steps. Oxidation of the 1,4-dihydroxynaphthalene moiety in **17** with [bis(trifluoroacetoxy)iodo]benzene (PIFA),^[8] followed by removal of the four MOM protecting groups with trifluoroacetic acid (TFA) in CH₂Cl₂ gave juglocombins A (**2**) and B (**3**) as an unstable tautomeric equilibrium mixture. Therefore, **2** and **3** were converted into dimethyl esters **2'** and **3'** through methylation with trimethylsilyldiazomethane. The ¹H and ¹³C NMR spectroscopic data for synthetic **2'** and **3'** were in agreement with those reported for the same compounds derived from natural **2** and **3**. Furthermore, treatment of dimethyl esters **2'** and **3'** with *p*-toluenesulfonic acid in methanol gave 1'-*O*-methyljuglocombin B dimethyl ester (**4**) in 68 % yield ([α]_D²⁴ –133.0 (*c* 0.20, acetone); lit. [α]_D²⁰ –133.3 (*c* 0.2, acetone)). The spectroscopic data and optical rotation of synthetic **4** were identical to those of the same compound derived from natural juglocombins A and B, thus indicating that their absolute configurations were 1'*R*,2'*R*,3'*R*,9*R*,10*R*,10'*S*.


The synthesis of juglorescein (**1**) is shown in Scheme 5. Removal of the two TBS protecting groups in epoxide intermediate **6** gave diol **18** in 78 % yield. Oxidation of the primary alcohols in **18** through Dess–Martin oxidation and Pinnick oxidation gave dicarboxylic acid **19**. Intramolecular lactonization of **19** using a carboxylic acid functionalized silica gel (Chromatorex-COOH) easily afforded six-membered lactone **20** in 81 % yield over the three steps.^[15] Because the carboxylic acid attacks from the opposite side of the epoxide, the *anti* relationship between the epoxide and the side chain at C-9 in **6** was validated. Finally, removal of the MOM protecting groups and hydrolysis of the lactone under acidic conditions gave juglorescein (**1**) in 62 % yield ([α]_D²⁴ –99.1 (*c* 0.58, MeOH); lit. [α]_D²⁰ –107.4 (*c* 0.29, MeOH)). Because the spectroscopic data and optical rotation of synthetic **1** agreed with those reported for natural **1**, the absolute configuration of natural **1** was determined to be 1'*S*,2*R*,3*R*,2'*R*,3'*R*,9*R*,10*R*,10'*S*.

In conclusion, the first total syntheses of juglorescein (**1**) and juglocombins A (**2**) and B (**3**) have been accomplished. Juglorescein (**1**) was synthesized in nine steps and in 11 % overall yield from known compounds **8** and **9**. 1'-*O*-Methyl-



Scheme 4. Reaction conditions for the synthesis of 1'-*O*-methyljuglocombin B dimethyl ester (**4**): a) Dess–Martin periodinane (4.0 equiv), CH₂Cl₂, RT, 25 min; b) NaClO₂ (5.8 equiv), 2-methyl-2-butene (21.6 equiv), NaH₂PO₄·2 H₂O (10.0 equiv), *tert*-BuOH/THF/H₂O (1:1:1), 0 °C, 15 min, 94 % yield over two steps; c) PIFA (1.2 equiv), TFE/H₂O (5:3), 0 °C to RT, 1 h; d) TFA/CH₂Cl₂ (4:1), 0 °C to reflux, 30 min; e) trimethylsilyldiazomethane (8.1 equiv), CH₂Cl₂/MeOH (1:2), RT, 15 min, 56 % over three steps; f) TsOH·H₂O (2.5 equiv), MeOH, RT, 24 h, 68 %. PIFA = [bis(trifluoroacetoxy)iodo]benzene, TFA = trifluoroacetic acid, TsOH = *p*-toluenesulfonic acid.

juglocombin B dimethyl ester (**4**), a common derivative of juglocombins A and B, was synthesized in 14 steps and 3 % overall yield from compounds **8** and **9**. The key feature of this synthesis is a bioinspired dimerization of a juglomycin C derivative, which efficiently provides access to the 6/6/5/6/6-fused core skeleton. A photoinduced reduction of the epoxide in dimer **6**, followed by further transformations, gave juglocombins A and B. Conversion of dimer **6** into dicarboxylic acid **19**, and an acid-promoted intramolecular lactonization followed by hydrolysis of the resultant lactone, afforded juglorescein. Notably, our total syntheses of **1–4** establish the relative and absolute configurations of these compounds. The synthetic route reported herein will support synthetic and biological studies of these natural products and their derivatives.



Scheme 5. Reaction conditions for the synthesis of juglorescein (**1**): a) TBAF (3.0 equiv), AcOH (2.6 equiv), THF, RT, 2 d, 78%; b) Dess–Martin periodinane (3.9 equiv), CH₂Cl₂, RT, 30 min; c) NaClO₂ (5.9 equiv), 2-methyl-2-butene (19.4 equiv), NaH₂PO₄·2 H₂O (9.8 equiv), *tert*-BuOH/THF/H₂O (1:1:1), 0°C, 10 min; d) carboxylic acid functionalized silica gel, CH₂Cl₂, RT, 3 h, 81% yield over three steps; e) Amberlyst 15, 1,4-dioxane/H₂O (4:1), 75–80°C, 4 d, 62% yield.

Acknowledgements

This study was partly supported by a Grant-in-Aid from the Japan Society for the Promotion of Science (JSPS) Fellows (No. 16J00542) to S.K. and a Grant-in-Aid for Scientific Research (C) (KAKENHI No. 15K07416) to K.K.. This study was carried out using the Fourier transform ion cyclotron resonance mass spectrometer at the Joint Usage/Research Center, Kyoto University.

Keywords: 1,4-naphthoquinone · biomimetic synthesis · dimerization · fused-ring systems · total synthesis

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 10317–10320
Angew. Chem. **2016**, *128*, 10473–10476

- [1] H. Lessmann, R. P. Maskey, S. Fotso, H. Lackner, H. Laatsch, *Z. Naturforsch. B* **2005**, *60*, 189–199.
- [2] a) A. V. B. Sankaram, A. S. Rao, J. N. Shoolery, *Tetrahedron* **1979**, *35*, 1777–1782; b) J.-Q. Gu, T. N. Graf, D. Lee, H.-B. Chai, Q. Mi, L. B. S. Kardono, F. M. Setyowati, R. Ismail, S. Riswan, N. R. Farnsworth, G. A. Cordell, J. M. Pezzuto, S. M. Swanson, D. J. Kroll, J. O. Falkinham III, M. E. Wall, M. C. Wani, A. D. Kinghorn, N. H. Oberlies, *J. Nat. Prod.* **2004**, *67*, 1156–1161; c) A. H. Uc-Cachón, G. M. Molina-Salinas, S. Said-Fernández, M. Méndez-González, M. Cáceres-Farfán, R. Borges-Argáez, *Nat. Prod. Res.* **2013**, *27*, 1174–1178; d) Y. Yang, D. Zhao, K. Yuan, G. Zhou, Y. Wang, Y. Xiao, C. Wang, J. Xu, W. Yang, *Nat. Prod. Res.* **2015**, *29*, 908–913; e) Isolation of other shikometabolins, see; M. R. Meselhy, S. Kadota, K. Tsubono, M. Hattori, T. Namba, *Tetrahedron* **1994**, *50*, 3081–3098.
- [3] S. Maruo, K. Nishio, T. Sasamori, N. Tokitoh, K. Kuramochi, K. Tsubaki, *Org. Lett.* **2013**, *15*, 1556–1559.
- [4] H. Lessmann, J. Krupa, H. Lackner, P. G. Jones, *Z. Naturforsch. B* **1989**, *44*, 353–363.
- [5] a) P. Dowd, S. W. Ham, S. J. Geib, *J. Am. Chem. Soc.* **1991**, *113*, 7734–7743; b) P. Dowd, S. W. Ham, R. Hershline, *J. Am. Chem. Soc.* **1992**, *114*, 7613–7617; c) S. W. Ham, J. S. Yoo, *Chem. Commun.* **1997**, 929–930.
- [6] a) R. A. Volkmann, P. R. Kelbaugh, D. M. Nason, V. J. Jasys, *J. Org. Chem.* **1992**, *57*, 4352–4361; b) M. Hwang, S.-J. Han, D.-H. Lee, *Org. Lett.* **2013**, *15*, 3318–3321.
- [7] a) S. S. Scully, J. A. Porco, Jr., *Angew. Chem. Int. Ed.* **2011**, *50*, 9722–9726; *Angew. Chem.* **2011**, *123*, 9896–9900; b) M. E. Jung, J. A. Hagenah, *J. Org. Chem.* **1987**, *52*, 1889–1902.
- [8] H. Tohma, H. Morioka, Y. Harayama, M. Hashizume, Y. Kita, *Tetrahedron Lett.* **2001**, *42*, 6899–6902.
- [9] Two diastereomers of **6** were also obtained as an inseparable 7:1 mixture in 13% yield. Isolation and structural characterization of the minor diastereomers were not achieved.
- [10] a) A. Patra, M. Bandyopadhyay, D. Mal, *Tetrahedron Lett.* **2003**, *44*, 2355–2357; b) C. Hardouin, F. Chevallier, B. Rousseau, E. Doris, *J. Org. Chem.* **2001**, *66*, 1046–1048; c) J. W. Cornforth, R. H. Cornforth, K. K. Mathew, *J. Chem. Soc.* **1959**, 2539–2547.
- [11] H.-J. Xu, Y.-C. Liu, Y. Fu, Y.-D. Wu, *Org. Lett.* **2006**, *8*, 3449–3451.
- [12] Continuous in situ generation of dihydronicotinamide **13** through reduction of **12** with sodium hydrosulfite in the presence of sodium carbonate is necessary to complete the reaction. When **6** was treated with **13** in EtOAc, reduction of the epoxide in **6** did not occur.
- [13] a) D. B. Dess, J. C. Martin, *J. Org. Chem.* **1983**, *48*, 4155–4156; b) D. B. Dess, J. C. Martin, *J. Am. Chem. Soc.* **1991**, *113*, 7277–7287.
- [14] B. S. Bal, W. E. Childers, Jr., H. W. Pinnick, *Tetrahedron* **1981**, *37*, 2091–2096.
- [15] D. A. Henderson, P. N. Collier, G. Pavé, P. Rzepa, A. J. P. White, J. N. Burrows, A. G. M. Barrett, *J. Org. Chem.* **2006**, *71*, 2434–2444.

Received: May 16, 2016

Published online: July 27, 2016