

Asymmetric Synthesis

International Edition: DOI: 10.1002/anie.201607053
German Edition: DOI: 10.1002/ange.201607053


Iridium-Catalyzed Regio- and Enantioselective Allylic Substitution of Trisubstituted Allylic Electrophiles

Ming Chen and John F. Hartwig*


Abstract: The first Ir-catalyzed enantioselective allylation of trisubstituted allylic electrophiles has been developed. Through modification of the leaving group of allylic electrophiles, we found that trisubstituted allylic phosphates are suitable electrophiles for asymmetric allylation. The reaction of allylic phosphates with enol silanes derived from dioxinones gave allylated products in good yields with high enantioselectivities.

Iridium-catalyzed, enantioselective allylic substitution has become a powerful method to access highly enantioenriched molecules.^[1,2] This class of reaction began to occur in high yield and *ee* when a metalacyclic catalyst derived from a phosphoramidite ligand was developed.^[1d] Over the past decade, research with these catalysts has focused on assessing and increasing the scope of nucleophile. In this body of work, cinnamyl carbonates and, in some cases, aliphatic allylic carbonates have been used as electrophiles. With few exceptions,^[3] these carbonates contain one aryl, alkyl or vinyl substituent and one leaving group to generate products in which the terminal alkene unit contains a single substituent (Figure 1).^[4–7]

Well studied reactions:



Unknown reaction:



This work:




Figure 1. Ir-catalyzed enantioselective allylic substitution of trisubstituted electrophiles **1** with enol silane **2**.

[*] M. Chen, Prof. J. F. Hartwig
Departments of Chemistry, University of California
Berkeley, CA 94720 (USA)
E-mail: jhartwig@berkeley.edu


Supporting information for this article can be found under:
<http://dx.doi.org/10.1002/anie.201607053>.

The inability to use allylic electrophiles containing more than one substituent, besides the leaving group, in asymmetric allylic substitutions is a major limitation on the eventual application of such reactions to the syntheses of complex molecules. In the current work, we have sought to develop systems that allow asymmetric allylic substitutions to occur with allylic electrophiles that are more substituted than those used previously, while maintaining the high regioselectivity for formation of branched products that characterizes iridium-catalyzed allylic substitution. In particular, we have sought to develop systems for asymmetric reactions of allylic esters containing 1,2-disubstituted alkene units to form products containing 1,1'-disubstituted olefins. Under the standard conditions comprising a carbonate leaving group and cyclometalated catalyst derived from a phosphoramidite, such reactions do not occur. This lack of reactivity is consistent with the absence of such reactions in the prior publications and highlights the need to find combinations of catalyst, leaving group, and nucleophile that will enable such processes.

Here, we report the first Ir-catalyzed, enantioselective allylic substitution of a trisubstituted allylic electrophile. By using a phosphate as the leaving group, and enol silanes derived from dioxinones^[8] as nucleophiles, the established cyclometalated iridium complexes catalyze allylic substitution with trisubstituted allylic phosphates to form products containing a 1,1'-disubstituted alkenes in good yields with high enantioselectivities and branched-to-linear ratios (Figure 1). The 1,1'-disubstituted olefin unit and the dioxanone moiety can be transformed to a variety of functional groups by established methods.^[9]

We began our studies to achieve allylic substitution of trisubstituted allylic esters by examining the reactions of a series of nucleophiles with the methyl carbonate derived from 2-methyl cinnamyl alcohol in the presence of [Ir(cod)Cl]₂ (cod = cyclooctadiene) and the phosphoramidite ligand (*S_a*, *S_c*, *S_c*)-**L**. We found that the reaction of this methyl carbonate with enol silane **2a** derived from the dioxanone we reported recently to undergo Ir-catalyzed substitution reactions to form products from formal allylation at the less acidic position of a dicarbonyl compound^[6c] gave product **3a** in 23 % yield. Reactions with enol silanes derived from ketones and α,β -unsaturated ketones did not form any detectable amount of product.

To improve the yield of product **3a** formed from the reaction, we surveyed reactions of enol silane **2a** with several allylic esters. As shown in Table 1, treatment of cinnamyl trifluoroacetate (1 equiv) and enol silane **2a** (2 equiv) in the presence of 3 mol % [Ir(cod)Cl]₂ and 6 mol % (*S*)-**L** with KF and 18-crown-6 as additives at 60 °C for 12 hours did not form

Table 1: Evaluation of the reaction conditions for the Ir-catalyzed asymmetric allylic substitution of esters **1** with dienolate **2a**.^[a]



| Entry | LG | b:l ^[b] | Yield (3a) ^[c] | % ee ^[d] |
|-------|---|--------------------|------------------------------------|---------------------|
| 1 | OCOCF ₃ | N.D. | N.R. | N.D. |
| 2 | OCO ₂ Me | 4:1 | 23 % | N.D. |
| 3 | OCO ₂ CH ₂ CCl ₃ | 3:1 | 31 % | N.D. |
| 4 | OCO ₂ <i>t</i> -Bu | N.D. | N.R. | N.D. |
| 5 | OP(O)(OEt) ₂ | 20:1 | 77 % | 98 % |

[a] Reaction conditions: allylic ester **1** (0.1 mmol, 1.0 equiv), dienolate **2a** (0.2 mmol, 2.0 equiv), [Ir(cod)Cl]₂ (3 mol %), (*S,S,S,S*)-**L** (6 mol %), KF (1.0 equiv), 18-crown-6 (1.0 equiv), THF (0.2 mL), 60 °C, 12 h. [b] Ratios of branched (**3a**) to linear product (**4a**) were determined by ¹H NMR analysis of the crude reaction mixtures. [c] Yield of isolated product. [d] *Ee* % of **3a** was determined by HPLC analysis using a chiral stationary phase. N.D. = not determined. N.R. = no reaction.

a detectable amount of product **3a** (entry 1, Table 1). As noted above, the reaction of allyl methyl carbonate with silane **2a** under the same conditions gave a 4:1 ratio of branched product **3a** to the linear adduct **4a** in 23 % yield (entry 2, Table 1). The reaction of cinnamyl trichloroethyl carbonate gave a lower 3:1 branched-to-linear (b/l) ratio in 31 % yield (entry 3, Table 1), and the reaction of the analogous *t*-Bu-carbonate gave no product (entry 4, Table 1). However, allylic substitution of the allyl ethyl phosphate with **2a** provided the branched product **3a** in 77 % yield, 98 % *ee*, and a 20:1 b/l ratio (entry 5, Table 1).

The scope of the allylic phosphate that underwent the Ir-catalyzed enantioselective allylic substitution with enol silane **2a** is summarized in Table 2. Reactions with a variety of trisubstituted allylic phosphates containing a series of substitution patterns on the arene were examined. Asymmetric allylation of substrates bearing substituents at the *meta*-position of the arene gave products **3b,c** in 77–80 % yield with 89–96 % *ee* and between 12:1 and 18:1 b/l ratios (entries 2 and 3, Table 2). Allylic phosphates bearing *para*-substituents also reacted to afford products **3d–f** in 74–90 % yield with 93–97 % *ee* and 13:1 to 20:1 b/l ratios (entries 4–6, Table 2). Allylic substitution of *ortho*-fluoro substituted allylic phosphate provided product **3g** in 86 % yield with 96 % *ee* and a 14:1 b/l ratio (entry 7, Table 2). Similar results were obtained with a naphthyl-substituted allylic phosphate (entry 8, Table 2). A dienyl phosphate reacted to give branched product **3i** selectively (18:1 b/l ratio) in 84 % yield with 94 % *ee*. Heterocycles are tolerated under the developed reaction conditions. Reactions with allylic phosphates containing quinolinyl and furyl groups provided products **3j,k** in 70–76 % yield with 92–98 % *ee* and a 12:1 branched-to-linear ratio (entries 10 and 11, Table 2).

Aliphatic allylic phosphates also reacted to give the substitution products in good yields, stereoselectivity and

Table 2: Scope of the Ir-catalyzed asymmetric allylic substitution of allylic phosphates **1** with dienolate **2a**.^[a–d]


| Product | Yield (%) | % ee | b:l |
|-----------|-----------|-----------|------------|
| 3a | 77% | 98% | 20:1 |
| 3b | 80% | 89% | 12:1 |
| 3c | 77% | 96% | 18:1 |
| 3d | 76% | 97% | 15:1 |
| 3e | 90% | 94% | 20:1 |
| 3f | 74% | 93% | 13:1 |
| 3g | 86% | 96% | 14:1 |
| 3h | 83% | 96% | 15:1 |
| 3i | 84% | 94% | 18:1 |
| 3j | 70% | 98% | 12:1 |
| 3k | 76% | 92% | 12:1 |
| 3l | 68% | 99% | 15:1 |
| 3m | 72% | dr > 20:1 | b:l > 20:1 |
| 3n | 36% | dr = 2:1 | [e] |

[a] Reaction conditions: allylic phosphate **1** (0.2 mmol, 1.0 equiv), dienolate **2a** (0.4 mmol, 2.0 equiv), [Ir(cod)Cl]₂ (3 mol %), (*S,S,S,S*)-**L** (6 mol %), KF (1.0 equiv), 18-crown-6 (1.0 equiv), THF (0.4 mL), 60 °C, 12 h. [b] Branched to linear ratios were determined by ¹H NMR analysis of the crude reaction mixtures. [c] Yields of isolated products are listed (the average of at least two runs). [d] Enantioselectivities were determined by HPLC analysis using a chiral stationary phase. [e] (*R,R,R,R*)-**L** was used.

regioselectivity. In particular, the reaction occurred with cyclic allylic phosphates derived from allylic alcohols. The aliphatic allylic phosphate from cyclohexenyl methanol reacted to give **3l** with high branched selectivity and enantioselectivity (68 % yield, 99 % *ee*, b/l = 15:1; entry 12, Table 2).

The reactions of a related chiral, non-racemic aliphatic allylic phosphate were also studied. The phosphate derived from (*R*)-perillyl alcohol converted under the standard reaction conditions (with (*S,S,S,S*)-**L** as the ligand) to the

allylated product **3m** in 72 % yield with a >20:1 branched selectivity and a >20:1 diastereoselectivity. The reaction with the antipode of the phosphoramidite ligand, (*R_aR_cR_c*)-**L**, under otherwise identical reaction conditions formed a mixture of products **3n** and **3m** in a combined 36 % yield and low diastereoselectivity (2:1), indicating that the reactions of these trisubstituted allylic esters occur with a matched and mismatched pairing of the catalyst with the enantioenriched allylic phosphate.

In contrast to nucleophilic additions of enol silane **2a**, nucleophilic additions of enol silane **2b**, which lacks a substituent at the α -position, can occur at both the α - and γ -positions. It is challenging to achieve high regioselectivity for the reaction of nucleophile **2b** at the γ -position.^[10] To explore whether the allylation could occur regioselectively at the γ -position with enol silane **2b**, allylic substitutions of several allylic electrophiles were examined (Table 3). The reaction of

diethylphosphates were used to evaluate the scope of the reaction of trisubstituted allylic esters with enol silane **2b**.

The results of asymmetric allylic substitution of ethyl phosphates **1** with enol silane **2b** are summarized in Table 4. Reactions of 2-methyl cinnamyl phosphates and alkenyl substituted allylic phosphates with silane **2b** gave allylated products **5a–g** in 62–73 % yields with 92–97 % *ee* and between 5:1 and 8:1 γ/α -selectivities (Table 4). In contrast, the reaction of alkyl substituted allylic electrophiles occurred with low γ/α -selectivities.

Table 4: Scope of the Ir-catalyzed asymmetric allylic substitution of allylic phosphates **1** with dienolate **2b**.^[a–d]





Table 3: Evaluation of the reaction conditions for the Ir-catalyzed asymmetric allylic substitution of allylic esters **1** with dienolate **2b**.^[a]




| Entry | LG | $\gamma:\alpha$ ^[b] | Yield (5a) ^[c] | % <i>ee</i> ^[d] |
|-------|---|--------------------------------|------------------------------------|----------------------------|
| 1 | OCO ₂ Me | 3:1 | 17 % | N.D. |
| 2 | OCOCF ₃ | N.D. | N.R. | N.D. |
| 3 | OCO ₂ <i>t</i> -Bu | N.D. | N.R. | N.D. |
| 4 | OCO ₂ CH ₂ CCl ₃ | N.D. | 11 % | N.D. |
| 5 | OP(O)(OEt) ₂ | 6:1 | 70 % | 92 % |
| 6 | OP(O)(OMe) ₂ | 6:1 | 62 % | 92 % |
| 7 | OP(O)(<i>Oi</i> -Pr) ₂ | 3:1 | 49 % | N.D. |
| 8 | OP(O)(OEH) ₂ | 3:1 | 41 % | N.D. |

[a] Reaction conditions: allylic ester **1** (0.1 mmol, 1.0 equiv), dienolate **2b** (0.2 mmol, 2.0 equiv), [Ir(cod)Cl]₂ (3 mol %), (*S_aS_cS_c*)-**L** (6 mol %), KF (1.0 equiv), 18-crown-6 (1.0 equiv), THF (0.2 mL), 60 °C, 12 h.


[b] Ratios of γ - to α -substitution were determined by ¹H NMR analysis of the crude reaction mixtures. [c] Yield of isolated product (**5a**). [d] *Ee* % of **5a** was determined by HPLC analysis using a chiral stationary phase. N.D. = not determined. N.R. = no reaction. EH = 2-ethyl-hexyl.

the allylic methyl carbonate in Table 3 formed a 3:1 mixture of the product **5a** from reaction at the γ -position of **2a** to the product **6a** from reaction at the α -position of **2a** in a total yield of 17 % (entry 1, Table 3). Reactions with the analogous allylic trifluoroacetate, *t*-Bu-carbonate and trichloroethyl carbonate failed to give product **5a** in a substantial yield (entries 2–4, Table 3). However, the reaction of this nucleophile with the allylic ethyl phosphate as the electrophile occurred with a high ratio of products from addition at the γ -position to addition at the α -position, and the yield was significantly higher (entry 5, Table 3). The ratio of γ -substituted product **5a** to α -substituted product **6a** was 6:1, and the γ -adduct **5a** was isolated in 70 % yield with 92 % *ee* (entry 5, Table 3). Allylic esters bearing dimethyl, diisopropyl, di-2-ethylhexyl phosphates (entries 6–8, Table 3) reacted in the same or lower yield and γ - to α -selectivity. Therefore,



[a] Reaction conditions: allylic phosphate **1** (0.2 mmol, 1.0 equiv), dienolate **2b** (0.4 mmol, 2.0 equiv), [Ir(cod)Cl]₂ (3 mol %), (*S_aS_cS_c*)-**L** (6 mol %), KF (1.0 equiv), 18-crown-6 (1.0 equiv), THF (0.4 mL), 60 °C, 12 h. [b] Ratios of γ - to α -substitution were determined by ¹H NMR analysis of the crude reaction mixtures. [c] Yields of isolated products **5** are listed (the average of at least two runs). [d] Enantioselectivities were determined by HPLC analysis using a chiral stationary phase.


As demonstrated in Tables 1–4, allylic substitution of the trisubstituted allylic esters occurred in high regioselectivity and branched-to-linear selectivity only when conducted with the allylic phosphate. To reveal the relative influences of the leaving group, the ligand, and additives on the selectivity of the reaction, the experiments summarized in Scheme 1 were conducted. In the absence of [Ir(cod)Cl]₂ and the ligand (*S_aS_cS_c*)-**L**, no allylation occurred from the reaction of phosphate **1c** with enol silane **2a** (panel a, Scheme 1). No allylated product **3c** was detected from reactions in the absence of KF (panel b, Scheme 1). However, [Ir(cod)Cl]₂ catalyzes the allylic substitution of phosphate **1c** with enol



Scheme 1. Influence of the ligand and additives on the selectivity.

silane **2a** without the phosphoramidite ligand. As shown in the panel c of Scheme 1, the allylation of **1c** with enol silane **2a** at 60 °C for 12 h gave a 1.5:3:1 mixture of (±)-**3c**, linear product **4c** and the α -adduct (±)-**7** in 52% combined yield in the presence of 3 mol % [Ir(cod)Cl]₂, KF (1 equiv) and 18-crown-6 (1 equiv). The same reaction conducted with the catalyst generated from [Ir(cod)Cl]₂ and the phosphoramidite ligand (*S_a*,*S_c*,*S_c*)-**L** occurred with much higher regioselectivity and branched selectivity (panel d, Scheme 1). These data indicate that the identity of the leaving group of the allylic electrophile alone does not control the site and branched-to-linear selectivities of these reactions. Instead, the combination of the phosphoramidite ligand **L** and the leaving group of the allylic electrophile is crucial to obtaining high regioselectivity at the nucleophile and high branched-to-linear selectivity at the electrophile.

Finally, Scheme 2 shows that the reaction can be conducted on multi-mmol scale and that the product can be converted to the corresponding β -keto ester without eroding enantiomeric excess. The allylic substitution with 2 mmol allylic phosphate proceeded smoothly to give allylated product **3a** in 71% yield with 98% *ee* and a 19:1 branched-to-linear ratio under the same conditions as the reactions run on a smaller scale. The dioxinone moiety of **5a** was converted to the β -keto ester by treatment of **5a** (92% *ee*) with K₂CO₃ and MeOH at ambient temperature to give β -keto ester **8** in 91% yield with the same 92% *ee* as the reactant (Scheme 2).



Scheme 2. A large-scale reaction and derivatization of compound **5a**.

In conclusion, we have developed the first Ir-catalyzed enantioselective allylic alkylation of trisubstituted allylic electrophiles. By utilizing allylic phosphates as electrophiles, asymmetric allylic substitution of enol silanes derived from dioxinones gave allylated products in good yields with high enantioselectivities and branched-to-linear selectivities. The data presented here suggest that the combination of the phosphoramidite ligand and the phosphate leaving group is critical to obtain high regioselectivity and branched-to-linear selectivity.

The origins of these effects await detailed mechanistic studies, but we attribute the lack of reactions of linear, trisubstituted allylic esters to unfavorable thermodynamics for the oxidative addition of trisubstituted allylic esters to the Ir catalyst. Our previous studies have shown that the oxidative addition of a 1,2-disubstituted allyl methyl carbonate is endothermic and reversible.^[1] The oxidative addition of a 1,1,2-trisubstituted allylic ester should be even less favored thermodynamically because trisubstituted allylic electrophiles are sterically more hindered and would bind more weakly to the iridium than would a 1,2-disubstituted allylic electrophile. At the same time, it is known that the leaving group in the allyl electrophile influences the rate, and presumably thermodynamics, for oxidative addition.^[1c,e,11] Thus, we hypothesize that the identity of the leaving group and the nucleophile are crucial to observing these first enantioselective reactions of disubstituted allylic esters because the better leaving group makes the reversible oxidative addition step more favored thermodynamically and the high nucleophilicity of the dioxinone effectively traps the small concentration of the allyliridium intermediate. Further studies to gain insight into the origin of the selectivity in this reaction and to expand the scope of nucleophile are currently underway in this laboratory.

Acknowledgements

Financial support provided by the National Institutes of Health (GM-58108) is gratefully acknowledged.

Keywords: 1,1'-substituted olefins · allylic asymmetric substitution · allylic phosphates · dioxinones · iridium catalysis

How to cite: *Angew. Chem. Int. Ed.* **2016**, 55, 11651–11655
Angew. Chem. **2016**, 128, 11823–11827

- [1] For seminal contributions, see: a) R. Takeuchi, M. Kashio, *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 263; *Angew. Chem.* **1997**, 109, 268; b) J. P. Janssen, G. Helmchen, *Tetrahedron Lett.* **1997**, 38, 8025; c) T. Ohmura, J. F. Hartwig, *J. Am. Chem. Soc.* **2002**, 124, 15164; d) C. A. Kiener, C. Shu, C. Incarvito, J. F. Hartwig, *J. Am. Chem. Soc.* **2003**, 125, 14272; e) D. Marković, J. F. Hartwig, *J. Am. Chem. Soc.* **2007**, 129, 11680; f) S. T. Madrahimov, D. Marković, J. F. Hartwig, *J. Am. Chem. Soc.* **2009**, 131, 7228; g) S. T. Madrahimov, J. F. Hartwig, *J. Am. Chem. Soc.* **2012**, 134, 8136; h) S. T. Madrahimov, Q. Li, A. Sharma, J. F. Hartwig, *J. Am. Chem. Soc.* **2015**, 137, 14968.
- [2] For recent reviews of Ir-catalyzed asymmetric allylation, see: a) G. Helmchen, A. Dahnz, P. Dubon, M. Schelwies, R. Weihofen, *Chem. Commun.* **2007**, 675; b) G. Helmchen in *Iridium Complexes in Organic Synthesis* (Eds.: L. A. Oro, C. Claver), Wiley-VCH, Weinheim, **2009**, p. 211; c) J. F. Hartwig, M. J. Pouy, *Top. Organomet. Chem.* **2011**, 34, 169; d) W.-B. Liu, J.-B. Xia, S.-L. You, *Top. Organomet. Chem.* **2012**, 38, 155; e) P. Tosatti, A. Nelson, S. P. Marsden, *Org. Biomol. Chem.* **2012**, 10, 3147; For a review of the use of phosphoramidite ligands in asymmetric catalysis, see: f) J. F. Teichert, B. L. Feringa, *Angew. Chem. Int. Ed.* **2010**, 49, 2486; *Angew. Chem.* **2010**, 122, 2538.
- [3] For examples of Cu-catalyzed enantioselective allylic substitution of 1,1,2-trisubstituted allyl electrophiles, see: a) Y. Lee, K. Akiyama, D. G. Gillingham, M. K. Brown, A. H. Hoveyda, *J. Am. Chem. Soc.* **2008**, 130, 446; b) F. Gao, J. L. Carr, A. H. Hoveyda, *J. Am. Chem. Soc.* **2014**, 136, 2149.
- [4] a) M. A. Schafroth, D. Sarlah, S. Krautwald, E. M. Carreira, *J. Am. Chem. Soc.* **2012**, 134, 20276; b) J. Y. Hamilton, D. Sarlah, E. M. Carreira, *J. Am. Chem. Soc.* **2013**, 135, 994; c) J. Y. Hamilton, D. Sarlah, E. M. Carreira, *Angew. Chem. Int. Ed.* **2013**, 52, 7532; *Angew. Chem.* **2013**, 125, 7680; d) S. Krautwald, D. Sarlah, M. A. Schafroth, E. M. Carreira, *Science* **2013**, 340, 1065; e) J. Y. Hamilton, D. Sarlah, E. M. Carreira, *J. Am. Chem. Soc.* **2014**, 136, 3006; f) S. Krautwald, M. A. Schafroth, D. Sarlah, E. M. Carreira, *J. Am. Chem. Soc.* **2014**, 136, 3020; g) J. Y. Hamilton, N. Hauser, D. Sarlah, E. M. Carreira, *Angew. Chem. Int. Ed.* **2014**, 53, 10759; *Angew. Chem.* **2014**, 126, 10935; h) S. Breitler, E. M. Carreira, *J. Am. Chem. Soc.* **2015**, 137, 5296; i) J. Y. Hamilton, D. Sarlah, E. M. Carreira, *Angew. Chem. Int. Ed.* **2015**, 54, 7644; *Angew. Chem.* **2015**, 127, 7754; j) T. Sandmeier, S. Krautwald, H. F. Zipfel, E. M. Carreira, *Angew. Chem. Int. Ed.* **2015**, 54, 14363; *Angew. Chem.* **2015**, 127, 14571.
- [5] a) Q.-F. Wu, H. He, W.-B. Liu, S.-L. You, *J. Am. Chem. Soc.* **2010**, 132, 11418; b) K.-Y. Ye, H. He, W.-B. Liu, L.-X. Dai, G. Helmchen, S.-L. You, *J. Am. Chem. Soc.* **2011**, 133, 19006; c) Q.-F. Wu, W.-B. Liu, C.-X. Zhuo, Z.-Q. Rong, K.-Y. Ye, S.-L. You, *Angew. Chem. Int. Ed.* **2011**, 50, 4455; *Angew. Chem.* **2011**, 123, 4547; d) Q.-F. Wu, C. Zheng, S.-L. You, *Angew. Chem. Int. Ed.* **2012**, 51, 1680; *Angew. Chem.* **2012**, 124, 1712; e) W.-B. Liu, C. Zheng, C.-X. Zhuo, L.-X. Dai, S.-L. You, *J. Am. Chem. Soc.* **2012**, 134, 4812; f) C.-X. Zhuo, Q.-F. Wu, Q. Zhao, Q.-L. Xu, S.-L. You, *J. Am. Chem. Soc.* **2013**, 135, 8169; g) Z.-P. Yang, Q.-F. Wu, S.-L. You, *Angew. Chem. Int. Ed.* **2014**, 53, 6986; *Angew. Chem.* **2014**, 126, 7106; h) C. Zheng, C.-X. Zhuo, S.-L. You, *J. Am. Chem. Soc.* **2014**, 136, 16251; i) Z.-P. Yang, Q.-F. Wu, W. Shao, S.-L. You, *J. Am. Chem. Soc.* **2015**, 137, 15899; j) X. Zhang, Z.-P. Yang, L. Hang, S.-L. You, *Angew. Chem. Int. Ed.* **2015**, 54, 1873; *Angew. Chem.* **2015**, 127, 1893; k) C.-X. Zhuo, Q. Chen, W.-B. Liu, Q. Zhao, S.-L. You, *Angew. Chem. Int. Ed.* **2015**, 54, 8475; *Angew. Chem.* **2015**, 127, 8595; l) Q. Cheng, Y. Wang, S.-L. You, *Angew. Chem. Int. Ed.* **2016**, 55, 3496; *Angew. Chem.* **2016**, 128, 3557; m) L. Huang, L.-X. Dai, S.-L. You, *J. Am. Chem. Soc.* **2016**, 138, 5793.
- [6] a) W. Chen, J. F. Hartwig, *J. Am. Chem. Soc.* **2012**, 134, 15249; b) W. Chen, J. F. Hartwig, *J. Am. Chem. Soc.* **2013**, 135, 2068; c) W. Chen, J. F. Hartwig, *J. Am. Chem. Soc.* **2014**, 136, 377; d) M. Chen, J. F. Hartwig, *Angew. Chem. Int. Ed.* **2014**, 53, 8691; *Angew. Chem.* **2014**, 126, 8835; e) M. Chen, J. F. Hartwig, *Angew. Chem. Int. Ed.* **2014**, 53, 12172; *Angew. Chem.* **2014**, 126, 12368; f) W. Chen, M. Chen, J. F. Hartwig, *J. Am. Chem. Soc.* **2014**, 136, 15825; g) M. Chen, J. F. Hartwig, *J. Am. Chem. Soc.* **2015**, 137, 13972; h) X. Jiang, W. Chen, J. F. Hartwig, *Angew. Chem. Int. Ed.* **2016**, 55, 5819; *Angew. Chem.* **2016**, 128, 5913.
- [7] a) T. Kanayama, K. Yoshida, H. Miyabe, Y. Takemoto, *Angew. Chem. Int. Ed.* **2003**, 42, 2054; *Angew. Chem.* **2003**, 115, 2100; b) D. Polet, X. Rathgeb, C. A. Falcioni, J.-B. Langlois, H. S. El, A. Alexakis, *Chem. Eur. J.* **2009**, 15, 1205; c) W.-B. Liu, C. M. Reeves, S. C. Virgil, B. M. Stoltz, *J. Am. Chem. Soc.* **2013**, 135, 10626; d) W.-B. Liu, C. M. Reeves, S. C. Virgil, B. M. Stoltz, *J. Am. Chem. Soc.* **2013**, 135, 17298; e) J. Qu, L. Roßberg, G. Helmchen, *J. Am. Chem. Soc.* **2014**, 136, 1272; f) K. Seehafer, C. Malakar, M. Bender, J. Qu, C. Liang, G. Helmchen, *Eur. J. Org. Chem.* **2016**, 493; For recent applications of Ir-catalyzed asymmetric allylic substitution in natural product syntheses, see: g) J. Deng, S. Zhou, W. Zhang, J. Li, R. Li, A. Li, *J. Am. Chem. Soc.* **2014**, 136, 8185; h) S.-Z. Jiang, X.-Y. Zeng, X. Liang, T. Lei, K. Wei, Y.-R. Yang, *Angew. Chem. Int. Ed.* **2016**, 55, 4044; *Angew. Chem.* **2016**, 128, 4112.
- [8] a) M. F. Carroll, A. R. Bader, *J. Am. Chem. Soc.* **1953**, 75, 5400; b) J. A. Hyatt, P. L. Feldman, R. J. Clemens, *J. Org. Chem.* **1984**, 49, 5105; c) R. J. Clemens, J. A. Hyatt, *J. Org. Chem.* **1985**, 50, 2431; d) M. Sato, S. Sunami, Y. Sugita, C. Kaneko, *Chem. Pharm. Bull.* **1994**, 42, 839; e) M. Sato, S. Sunami, Y. Sugita, C. Kaneko, *Heterocycles* **1995**, 41, 1435; f) R. A. Singer, E. M. Carreira, *J. Am. Chem. Soc.* **1995**, 117, 12360; g) D. A. Evans, J. A. Murry, M. C. Kozlowski, *J. Am. Chem. Soc.* **1996**, 118, 5814; h) J. Krüger, E. M. Carreira, *J. Am. Chem. Soc.* **1998**, 120, 837.
- [9] a) M. Sato, N. Kanuma, T. Kato, *Chem. Pharm. Bull.* **1984**, 32, 106; b) M. Sato, H. Ogasawara, T. Kato, *Chem. Pharm. Bull.* **1984**, 32, 2602; c) F. G. West, P. V. Fisher, G. U. Gunawardena, S. Mitchell, *Tetrahedron Lett.* **1993**, 34, 4583; d) A. C. Hart, A. J. Phillips, *J. Am. Chem. Soc.* **2006**, 128, 1094; e) M. T. Crimmins, A. C. Smith, *Org. Lett.* **2006**, 8, 1003; f) D. W. Custer, T. P. Zabawa, K. A. Scheidt, *J. Am. Chem. Soc.* **2008**, 130, 804; g) T. Hoyer, M. E. Danielson, A. E. May, H. Zhao, *J. Org. Chem.* **2010**, 75, 7052; h) I. Yavari, M. J. Bayat, S. Skoulia, *Synlett* **2013**, 2591.
- [10] a) C.-L. Gu, L. Liu, D. Wang, Y.-J. Chen, *J. Org. Chem.* **2009**, 74, 5754; b) B. Lei, A. Fallis, *Can. J. Chem.* **1991**, 69, 1450; c) R. W. Dugger, C. H. Heathcock, *J. Org. Chem.* **1980**, 45, 1181.
- [11] W. Kinouchi, R. Saeki, H. Kawashima, Y. Kobayashi, *Tetrahedron Lett.* **2015**, 56, 2265.

Received: July 20, 2016

Published online: August 17, 2016