

QT prolongation and Torsades de Pointes in patients previously treated with Anthracyclines

Yaron Arbel, Michael Swartzon and Dan Justo

Anthracyclines reduce myocardial repolarization reserve and might increase the risk for Torsades de Pointes a long time after treatment. We studied all the publications concerning Torsades de Pointes in patients previously treated with anthracyclines to investigate the clinical circumstances leading to this rare life-threatening complication. Our literature search yielded nine reports of 11 patients who had developed Torsades de Pointes anywhere from weeks to years following treatment with anthracyclines. One of the patients was hospitalized in our medical center. Risk factors and triggers for Torsades de Pointes, among other clinical aspects, were analyzed in each report. Most patients ($n=10$; 90.9%) were previously treated with anthracyclines owing to acute leukemias: acute myelogenous leukemia ($n=5$), acute lymphocytic leukemia ($n=3$) and acute promyelocytic leukemia ($n=2$). One patient was previously treated with anthracyclines owing to Hodgkin's lymphoma. Most patients were women ($n=9$; 81.8%). The most prevalent triggers for Torsades de Pointes were the administration of a QT-prolonging agent ($n=10$; 90.9%) and hypokalemia ($n=9$; 81.8%). Azole derivatives were the most prevalent of the QT-prolonging agents that triggered Torsades de Pointes ($n=5$; 45.5%).

Although four patients suffered from anthracycline-induced left ventricular dysfunction and five other patients had only one or two questionable triggers for Torsades de Pointes, in only two of these cases the authors considered previous treatment with anthracyclines as a risk factor for Torsades de Pointes. Previous treatment with anthracycline is an underestimated risk factor for Torsades de Pointes. Possible triggers includes azole derivatives, other QT-prolonging agents and hypokalemia. Women patients are particularly at risk. *Anti-Cancer Drugs* 18:493–498
© 2007 Lippincott Williams & Wilkins.

Anti-Cancer Drugs 2007, 18:493–498

Keywords: anthracycline, QT interval, Torsades de Pointes

Department of Internal Medicine D, Tel-Aviv Sourasky Medical Center, Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel

Correspondence to Dr Dan Justo, MD, Department of Internal Medicine D, Tel-Aviv Sourasky Medical Center, 6 Weizman Street, Tel-Aviv 64239, Israel
Tel: +972 50 6437395; fax: +972 9 7408575;
e-mail: danjusto@post.tau.ac.il

Received 8 September 2006 Revised form accepted 1 November 2006

Introduction

A 61-year-old woman was hospitalized in our medical center because of recent syncope and palpitations. She had a history of essential hypertension, hypothyroidism and diabetes mellitus. Three years earlier, she underwent a total abdominal hysterectomy and was treated with chemotherapy owing to endometrial carcinoma. Three months before her current admission she was treated with mitoxantrone (cumulative dose of 36 mg/m^2) for recently diagnosed acute myelogenous leukemia – probably chemotherapy related. Her medications included glibenclamide, levothyroxine and candesartan. She was also taking oral fluconazole 200 mg/day for the last 3 months owing to systemic candidiasis. The patient had no history of coronary heart disease. Previous electrocardiogram (ECG) was normal; with corrected QT (QTc) of 420. Echo performed 1 month earlier was normal and showed no systolic dysfunction. Initial ECG in the emergency department showed Torsades de Pointes (TdP) (Fig. 1). Intravenous lidocaine, magnesium and potassium were immediately administered, and fluconazole was discontinued. The arrhythmia resolved promptly with no need for electrical defibrillation. The measured QTc interval was 466 and several hours later it was 502 (Fig. 2). Initial laboratory

tests before potassium administration revealed hypokalemia (potassium 2.9 mmol/l). No other electrolyte disturbances were seen. The QTc interval returned to normal the following day. Since then, TdP did not recur.

TdP is a rare ventricular tachycardia which might result in ventricular fibrillation and death. It is associated with congenital long-QT syndrome and drug administration. Patients with cardiomyopathy and congestive heart failure are at risk for TdP [1,2]. These conditions are associated with down-regulation of myocardial potassium channels which can lead to repolarization prolongation [3]. Anthracyclines are associated with late cardiomyopathy and heart failure in lymphoma patients [4] and children with solid tumors or leukemias [5,6]; furthermore, they might reduce myocardial repolarization reserve [7]. Hence, anthracyclines might increase the risk for late TdP – weeks–years after treatment.

Aims of the study

We studied all the publications concerning TdP in patients previously treated with anthracyclines in order to determine the prevalence of any known risk factors for TdP during episodes of TdP and to investigate the

clinical circumstances leading to this life-threatening arrhythmia.

Material and methods

Retrieval of case reports

We performed a literature search in PubMed for all published reports, in all languages, concerning TdP in patients previously treated with anthracyclines, from 1966 until September 2006, using the following keywords: 'TdP', 'Torsade de Pointes', 'QT interval', 'arrhythmia', 'syncope', 'sudden death' and 'anthracyclines'. The references from each report were reviewed for additional publications, as were relevant letters to the editor. We also contacted several authors for additional information and clarification. Dr Bagatell [8] and Dr Sisakova [9] were kind enough to respond.

Exclusion of case reports

The following reports were excluded: reports of sudden death with no documented TdP [10,11], reports of ventricular arrhythmias other than TdP [12–14] and

reports of sudden death or arrhythmias during or directly after administration of anthracyclines [15–17].

Risk factors and triggers for Torsades de Pointes

Each case report was analyzed for the presence of risk factors or triggers for TdP: female sex [18]; cardiomyopathy and congestive heart failure [2]; hypokalemia, defined as potassium serum levels < 3.5 mmol/l [19]; hypocalcaemia, defined as calcium serum levels < 8.5 mg/dl [20]; hypomagnesaemia, defined as magnesium serum levels < 1.8 mg/dl [1]; use of an agent that might trigger TdP [1]; impaired QT-prolonging agent clearance because of liver cirrhosis or renal failure [1]; HIV infection [21]; congenital long-QT syndrome, family history of long-QT syndrome, history of previous TdP and prolonged QT interval in the baseline ECG before drug initiation [22] defined as corrected QTc > 450 and calculated by the Bazett's formula [23].


Statistical analysis

Continuous variables, like age, Anthracyclines cumulative dose, and QTc interval, were summarized by mean, standard deviation, median, and range. Statistical analysis was performed using the SPSS (SPSS Inc., Chicago, USA) system for Windows, version 13.0.

Results

Including the patient who was hospitalized in our medical center, we reviewed nine reports of 11 patients who developed TdP starting at 15 days to over 3 years after treatment with anthracyclines [8,9,24–29]. Table 1 summarizes the clinical characteristics of the above-mentioned patients. The mean age of the reported patients was

Fig. 1



Electrocardiogram showing Torsades de Pointes.

Fig. 2



Electrocardiogram showing prolonged QT interval several hours following Torsades de Pointes.

Table 1 Clinical characteristics of patients with TdP previously treated with Anthracyclines

Number	Author (reference)	Age (years), sex (male/female)	Anthracyclines (cumulative dose)	Indication for anthracyclines	Timing of TdP following anthracyclines administration	Apparent triggers for TdP	Baseline QTc	Long QTc	Echo findings
1	Otsuka <i>et al.</i> [28]	33, female	Daunorubicin (NA)	AML	15 days	Pentamidine, hypokalemia	415	470	NA
2	Tatetsu <i>et al.</i> [26]	55, female	Mitoxantrone (NA)	AML	28 days	Fluconazole, hypokalemia	375	528	NA
3	Unnikrishnan <i>et al.</i> [27]	29, male	Idarubicin (NA)	AML and relapsing AML	NA	Arsenic trioxide, hypokalemia	NA	440	Normal EF 57%
4		43, male	Daunorubicin (72 mg/m ²)						
5	Naito <i>et al.</i> [24]	23, female	Mitoxantrone (36 mg/m ²)	APL and relapsing APL	NA	Arsenic trioxide	435	450	Normal
6		51, female	Idarubicin (92 mg/m ²)						
7	Alkan <i>et al.</i> [29]	15, female	Daunorubicin (720 mg/m ²)	APL	>3 years	Arsenic trioxide, hypokalemia, hypomagnesemia	429	478	Hypokinesia EF 51%
8	Kishi <i>et al.</i> [25]	16, female	Mitoxantrone (24 mg/m ²)	ALL	22 months	Voriconazole, ciprofloxacin, cotrimoxazole, hypokalemia	442	499	Hypokinesia EF 26%
9	Sisakove <i>et al.</i> [9]	33, female	Anthracyclines (NA)	ALL	Few months	Hypokalemia	412	570	Normal
10	Bagatell <i>et al.</i> [8]	16, female	Doxorubicin (125 mg/m ²)	Hodgkin's lymphoma	2 months	Itraconazole, Terfenadine, hypokalemia	NA	500	EF 35%
11		61, female	Aclarubicin (60 mg/m ²)						
			Mitoxantrone (18 mg/m ²)	ALL	Few weeks	Clarithromycin, renal failure, hypokalemia, hypomagnesemia	388	566	NA
11	Present case	61, female	Anthracyclines (NA)	AML	3 months	Fluconazole, hypokalemia	420	466	Normal

ALL, acute lymphocytic leukemia; AML, acute myeloblastic leukemia; APL, acute promyelocytic leukemia; EF, ejection fraction; NA, not applicable; QTc, corrected QT; TdP, Torsade de Pointes.

34.1 ± 16.4 years (median: 33 years, range: 15–61 years). Most of the patients were women ($n = 9$; 81.8%). One patient died as a result of recurrent TdP [27].

Indications for Anthracyclines

Most patients ($n = 10$; 90.9%) were previously treated with anthracyclines owing to acute leukemias: acute myelogenous leukemia ($n = 5$), acute lymphocytic leukemia ($n = 3$) and acute promyelocytic leukemia ($n = 2$). One patient was previously treated with anthracyclines owing to Hodgkin's lymphoma [9].

Anthracycline cumulative dose

Anthracycline cumulative dose was reported in six patients and was within the recommended safe range [30] in all of these patients, except one who received a total of 720 mg/m^2 daunorubicin for acute promyelocytic leukemia [24].

QTc interval

Baseline QTc interval before administration of QT-prolonging agents was reported in eight patients and their mean QTc interval was 415 ± 23 (median: 417, range: 375–442). The QTc interval right after TdP was reported in all patients and their mean QTc interval was 497 ± 45 (median: 489, range: 440–500).

The triggers for Torsades de Pointes

The most prevalent triggers for TdP were the administration of QT-prolonging agents ($n = 10$; 90.9%) and hypokalemia ($n = 9$; 81.8%). Azole derivatives were the most common QT-prolonging agents to trigger TdP ($n = 5$; 45.5%). Arsenic trioxide use was also documented ($n = 4$; 36.3%). Three patients were treated with two or more QT-prolonging agents concurrently [9,24,29]. At the time of the arrhythmia, the patients had 2.3 ± 0.9 triggers for TdP and nine patients (81.8%) had ≥ 2 triggers for TdP.

Anthracyclines-associated Torsades de Pointes

Four patients had obvious Anthracyclines-induced left ventricular dysfunction as per echo [9,24,25]. Five more patients had only one or two questionable triggers for TdP – including the patient we presented here [26–28]. Yet, in seven of these cases, the authors (or the physicians that first treated the patient we presented) did not consider previous treatment with anthracyclines as a risk factor for TdP. Moreover, in three patients, echo was not performed or mentioned [8,26,28]. Four patients had a normal echo, but none of those authors (or the physicians that first treated the patient we presented) discussed the possibility of subclinical cardiotoxicity in their patients [27,29].

Discussion

Anthracyclines are the first-line therapy for many neoplastic processes. Unfortunately, anthracyclines are associated with early cardiotoxicity during the treatment

or late cardiotoxicity – months–years after the treatment. Early cardiotoxicity includes supraventricular tachycardia, ventricular ectopy, myopericarditis, cardiomyopathy and death [31]. Late cardiotoxicity includes cardiomyopathy and heart failure [4–6]. Sarcoplasmic reticulum reduction, vacuole formation and necrosis in the myocardium are all histological features of anthracycline-associated cardiotoxicity. They may be explained by the following mechanisms: adenosine triphosphate depletion owing to mitochondrial dysfunction, iron–doxorubicin complex mediating free radical lipid peroxidation or a decrease in glutathione peroxidase [32].

The risk of fatal arrhythmia from weeks to years after supposedly life-saving treatment is quite troubling. Indeed, certain guidelines for cardiac monitoring and the usage of cardioprotective drugs during anthracycline administration have been established to prevent cardiotoxicity [31–34]. Withdrawal of anthracyclines from the market after years of successful clinical experience would probably harm more patients overall than it would help to prevent TdP. As our findings show that the role of previous treatment with anthracyclines in the pathophysiology of late TdP is underestimated, it is important to study the clinical circumstances by which TdP occurs following anthracycline administration.

According to Isner *et al.* [35], 67.1% of patients treated with anthracyclines have clinical heart failure and/or histological features of cardiotoxicity *post mortem*. Interestingly, 35.9% of patients with histological features of cardiotoxicity, *post mortem* did not have any signs or symptoms of clinical heart failure during their lifetime [35]. Although four of the patients in our study had a normal echo, subclinical cardiotoxicity was still possible, i.e. histological features of cardiotoxicity. This may explain some of our findings. It is also possible that recent administration of anthracycline itself increased the risk for TdP by increasing the dispersion of repolarization [36]. According to Milberg *et al.* [7], reduced repolarization reserve even in the early stages of anthracycline therapy facilitates TdP induced by hypokalemia and potassium currents blockers.

Risk factors for anthracycline-induced cardiotoxicity include female sex, cumulative dose greater than 550 mg/m^2 , hypertension, preexisting cardiac disease, advanced age and prior mediastinal irradiation [35,37]. Although a high cumulative dose of anthracyclines is associated with increased risk for cardiotoxicity [30], some patients do not develop signs of anthracycline-induced cardiotoxicity following very high (above 5000 mg/m^2) cumulative doses of anthracyclines, whereas other patients develop anthracycline-induced cardiotoxicity following very low cumulative doses (below 40 mg/m^2) of anthracycline [38]. Our results are consistent with those observations: nine out of 11 patients were women

and three out of four patients with anthracyclines-induced left ventricular dysfunction received recommended doses of anthracyclines.

Study limitations

The patient we described had two triggers for TdP (i.e. hypokalemia and fluconazole) and one risk factor for TdP (female sex). It is possible that anthracycline itself or subclinical cardiotoxicity following anthracycline administration increased the risk for TdP further. Although this is only a possibility, it should be highlighted as it is underestimated.

This study was based entirely on published case reports. In all of these reports, except one, the patients survived the potentially fatal arrhythmia. We believe that there are many more incidents of anthracycline-associated TdP that remain unpublished, particularly when a similar incidence involving an anthracycline had already appeared in print, when a patient had died or when a report had been rejected from publication because of apparent risk factors for TdP other than the anthracyclines treatment.

Clinical implications and conclusions

Patients with risk factors for anthracycline-induced cardiotoxicity or patients receiving a high cumulative dose of anthracyclines should have a baseline evaluation of cardiac function by echo or radionuclear ventriculography. Some physicians perform serial assessments of left ventricular function after every cycle of chemotherapy. Several simple aspects of the patient's history, however, should alert physicians to a potentially increased risk for TdP. According to our findings, female patients with hematological malignancies previously treated with anthracyclines, who are currently receiving azole derivatives or other QT-prolonging agents, are at increased risk for Anthracyclines-associated TdP. These patients should be advised to promptly report any symptoms, such as palpitations or syncope. Other conditions or therapies that can cause hypokalemia, such as gastroenteritis, or the addition of a diuretic to the patient's regimen can also increase the risk. Obtaining electrocardiograms repeatedly to detect asymptomatic prolongation of the QT interval may be warranted in such high-risk patients during administration of QT-prolonging agents; although, it is unknown whether these precautions would reduce the risk for TdP.

Finally, TdP might have been the first sign for anthracycline-induced cardiotoxicity in our patient. Hence, physicians should consider repeating the echo in patients whom have already had TdP to exclude late cardiotoxicity.

References

- Roden DM. Drug-induced prolongation of the QT interval. *N Engl J Med* 2004; **350**:1013–1022.
- Torp-Pedersen C, Moller M, Bloch-Thomsen PE, Kober L, Sandoe E, Egstrup K, *et al.* Dofetilide in patients with congestive heart failure and left ventricular dysfunction. Danish Investigations of Arrhythmia and Mortality on Dofetilide Study Group. *N Engl J Med* 1999; **341**:857–865.
- Tsuji Y, Opthof T, Kamiya K, Yasui K, Liu W, Lu Z, *et al.* Pacing-induced heart failure causes a reduction of delayed rectifier potassium currents along with decreases in calcium and transient outward currents in rabbit ventricle. *Cardiovasc Res* 2000; **48**:300–309.
- Hequet O, Le QH, Moullet I, Pauli E, Salles G, Espinouse D, *et al.* Subclinical late cardiomyopathy after doxorubicin therapy for lymphoma in adults. *J Clin Oncol* 2004; **22**:1864–1871.
- Pinarli FG, Oguz A, Tunaoglu FS, Karadeniz C, Gokcora N, Elbeg S. Late cardiac evaluation of children with solid tumors after anthracycline chemotherapy. *Pediatr Blood Cancer* 2005; **44**:370–377.
- Sorensen K, Levitt GA, Bull C, Dorup I, Sullivan ID. Late anthracycline cardiotoxicity after childhood cancer: a prospective longitudinal study. *Cancer* 2003; **97**:1991–1998.
- Milberg P, Fleischer D, Stypmann J, Osada N, Monnig G, Engelen MA, *et al.* Reduced repolarization reserve due to anthracycline therapy facilitates torsade de pointes induced by I(Kr) blockers. *Basic Res Cardiol* 2007; **102**:42–51.
- Bagatell R, Hainstock M, Lowe MC, Barber BJ, Samson RA. The perfect storm: torsades de pointes in a child with leukemia. *Pediatr Blood Cancer* 2005; (in press).
- Sisakova M, Toman O, Florianova A, Kadlecova J, Chroust K, Papousek I, *et al.* Extension of QT interval as a consequence of risk factor accumulation – case study. *Vnitř Lek* 2006; **52**:271–273.
- Tsuda N, Oka R, Kajino H, Kajino M, Okuno A. Sudden death of a patient in complete remission after anthracycline therapy for acute lymphoblastic leukemia. *Pediatr Int* 2000; **42**:319–321.
- Westervelt P, Brown RA, Adkins DR, Khoury H, Curtin P, Hurd D, *et al.* Sudden death among patients with acute promyelocytic leukemia treated with arsenic trioxide. *Blood* 2001; **98**:266–271.
- Ohnishi K, Yoshida H, Shigeno K, Nakamura S, Fujisawa S, Naito K, *et al.* Prolongation of the QT interval and ventricular tachycardia in patients treated with arsenic trioxide for acute promyelocytic leukemia. *Ann Intern Med* 2000; **133**:881–885.
- Steinherz LJ, Steinherz PG, Tan C. Cardiac failure and dysrhythmias 6–19 years after anthracycline therapy: a series of 15 patients. *Med Pediatr Oncol* 1995; **24**:352–361.
- Patel SP, Garcia-Manero G, Ferrajoli A, Faderl S, Verstovsek S, Kantarjian H, *et al.* Cardiotoxicity in African-American patients treated with arsenic trioxide for acute promyelocytic leukemia. *Leuk Res* 2006; **30**:362–363.
- Wortman JE, Lucas VS Jr, Schuster E, Thiele D, Logue GL. Sudden death during doxorubicin administration. *Cancer* 1979; **44**:1588–1591.
- Morioka M, Kakinoki Y, Katagiri M, Iwasaki H, Gotohda Y, Kobayashi T, *et al.* Torsade de pointes ventricular tachycardia in a patient with acute myelocytic leukemia. *Rinsho Ketsueki* 1991; **32**:1009–1011.
- Stupp R, Bauer J, Pagani O, Gerard B, Cerny T, Sessa C, *et al.* Ventricular arrhythmia and torsade de pointe: dose limiting toxicities of the MDR-modulator S9788 in a phase I trial. *Ann Oncol* 1998; **9**:1233–1242.
- Makkar RR, Fromm BS, Steinman RT, Meissner MD, Lehmann MH. Female gender as a risk factor for torsades de pointes associated with cardiovascular drugs. *JAMA* 1993; **270**:2590–2597.
- Kay GN, Plumb VJ, Arciniegas JG, Henthorn RW, Waldo AL. Torsade de pointes: the long-short initiating sequence and other clinical features: observations in 32 patients. *J Am Coll Cardiol* 1983; **2**:806–817.
- Napolitano C, Priori SG, Schwartz PJ. Torsade de pointes. Mechanisms and management. *Drugs* 1994; **47**:51–65.
- Nanavati KA, Fisher SD, Miller TL, Lipshultz SE. HIV-related cardiovascular disease and drug interactions. *Am J Cardiovasc Drugs* 2004; **4**:315–324.
- Houltz B, Darpo B, Edvardsson N, Blomstrom P, Brachmann J, Crijns HJ, *et al.* Electrocardiographic and clinical predictors of torsades de pointes induced by almokalant infusion in patients with chronic atrial fibrillation or flutter: a prospective study. *Pacing Clin Electrophysiol* 1998; **21**: 1044–1057.
- Bazzet HC. An analysis of time relations of echocardiograms. *Heart* 1920; **7**:353–367.
- Naito K, Kobayashi M, Sahara N, Shigeno K, Nakamura S, Shinjo K, *et al.* Two cases of acute promyelocytic leukemia complicated by torsade de pointes during arsenic trioxide therapy. *Int J Hematol* 2006; **83**:318–323.
- Kishi S, Yoshida A, Yamauchi T, Tsutani H, Lee JD, Nakamura T, *et al.* Torsade de pointes associated with hypokalemia after anthracycline treatment in a patient with acute lymphocytic leukemia. *Int J Hematol* 2000; **71**:172–179.

- 26 Tatetsu H, Asou N, Nakamura M, Hanaoka N, Matsuno F, Horikawa K, *et al.* Torsades de pointes upon fluconazole administration in a patient with acute myeloblastic leukemia. *Am J Hematol* 2006; **81**:366–369.
- 27 Unnikrishnan D, Dutcher JP, Varshneya N, Lucariello R, Api M, Garl S, *et al.* Torsades de pointes in 3 patients with leukemia treated with arsenic trioxide. *Blood* 2001; **97**:1514–1516.
- 28 Otsuka M, Kanamori H, Sasaki S, Taguchi J, Harano H, Ogawa K, *et al.* Torsades de pointes complicating pentamidine therapy of pneumocystis carinii pneumonia in acute myelogenous leukemia. *Intern Med* 1997; **36**:705–708.
- 29 Alkan Y, Haefeli WE, Burhenne J, Stein J, Yaniv I, Shalit I. Voriconazole-induced QT interval prolongation and ventricular tachycardia: a non-concentration-dependent adverse effect. *Clin Infect Dis* 2004; **39**:e49–e52.
- 30 Hensley ML, Schuchter LM, Lindley C, Meropol NJ, Cohen GI, Broder G, *et al.* American Society of Clinical Oncology clinical practice guidelines for the use of chemotherapy and radiotherapy protectants. *J Clin Oncol* 1999; **17**:3333–3355.
- 31 Bristow MR, Thompson PD, Martin RP, Mason JW, Billingham ME, Harrison DC. Early anthracycline cardiotoxicity. *Am J Med* 1978; **65**:823–832.
- 32 Floyd JD, Nguyen DT, Lobins RL, Bashir Q, Doll DC, Perry MC. Cardiotoxicity of cancer therapy. *J Clin Oncol* 2005; **23**:7685–7696.
- 33 Steinhert LJ, Graham T, Hurwitz R, Sondheimer HM, Schwartz RG, Shaffer EM, *et al.* Guidelines for cardiac monitoring of children during and after anthracycline therapy: report of the Cardiology Committee of the Childrens Cancer Study Group. *Pediatrics* 1992; **89**:942–949.
- 34 Seymour L, Bramwell V, Moran LA. Use of dexrazoxane as a cardioprotectant in patients receiving doxorubicin or epirubicin chemotherapy for the treatment of cancer. The Provincial Systemic Treatment Disease Site Group. *Cancer Prev Control* 1999; **3**:145–159.
- 35 Isner JM, Ferrans VJ, Cohen SR, Witkind BG, Virmani R, Gottdiener JS, *et al.* Clinical and morphologic cardiac findings after anthracycline chemotherapy. Analysis of 64 patients studied at necropsy. *Am J Cardiol* 1983; **51**:1167–1174.
- 36 Kelso EJ, Geraghty RF, McDermott BJ, Cameron CH, Nicholls DP, Silke B. Characterisation of a cellular model of cardiomyopathy, in the rabbit, produced by chronic administration of the anthracycline, epirubicin. *J Mol Cell Cardiol* 1997; **29**:3385–3397.
- 37 Lipshultz SE, Lipsitz SR, Mone SM, Goorin AM, Sallan SE, Sanders SP, *et al.* Female sex and drug dose as risk factors for late cardiotoxic effects of doxorubicin therapy for childhood cancer. *N Engl J Med* 1995; **332**:1738–1743.
- 38 Von Hoff DD, Layard MW, Basa P, Davis HL Jr, Von Hoff AL, Rozenzweig M, *et al.* Risk factors for doxorubicin-induced congestive heart failure. *Ann Intern Med* 1979; **91**:710–717.