

Redox Control of Prion and Disease Pathogenesis

Neena Singh, Ajay Singh, Dola Das, and Maradumane L. Mohan

Abstract

Imbalance of brain metal homeostasis and associated oxidative stress by redox-active metals like iron and copper is an important trigger of neurotoxicity in several neurodegenerative conditions, including prion disorders. Whereas some reports attribute this to end-stage disease, others provide evidence for specific mechanisms leading to brain metal dyshomeostasis during disease progression. In prion disorders, imbalance of brain-iron homeostasis is observed before end-stage disease and worsens with disease progression, implicating iron-induced oxidative stress in disease pathogenesis. This is an unexpected observation, because the underlying cause of brain pathology in all prion disorders is PrP-scrapie (PrP^{Sc}), a β -sheet-rich conformation of a normal glycoprotein, the prion protein (PrP^C). Whether brain-iron dyshomeostasis occurs because of gain of toxic function by PrP^{Sc} or loss of normal function of PrP^C remains unclear. In this review, we summarize available evidence suggesting the involvement of oxidative stress in prion-disease pathogenesis. Subsequently, we review the biology of PrP^C to highlight its possible role in maintaining brain metal homeostasis during health and the contribution of PrP^{Sc} in inducing brain metal imbalance with disease progression. Finally, we discuss possible therapeutic avenues directed at restoring brain metal homeostasis and alleviating metal-induced oxidative stress in prion disorders. *Antioxid. Redox Signal.* 12, 1271–1294.

I. Introduction	1271
II. Prion Disorders: An Overview	1272
A. The unique nature of prion disorders	1272
1. The “prion” agent	1273
2. Mechanism(s) underlying prion-associated neurotoxicity	1274
B. Redox metals, oxidative stress, and prion-disease pathogenesis	1275
III. Redox Control of Prion Protein: Physiological Implications	1277
A. PrP and copper interaction	1277
B. PrP and other metals, in particular iron	1278
C. The antioxidant activity of PrP	1282
IV. Redox Homeostasis and Prion-Disease Pathogenesis	1283
A. PrP and metal interaction: the ironic connection	1283
B. Prion disorders and brain-iron homeostasis	1284
V. Therapeutic Options	1284
A. Antioxidants	1285
B. Metal chelators	1285
VI. Summary and Perspective	1286

I. Introduction

NEURODEGENERATIVE CONDITIONS such as prion disorders, Alzheimer’s disease (AD), amyotrophic lateral sclerosis (ALS), Parkinson’s disease (PD), and Huntington’s disease (HD) share a common pathogenic event involving the

accumulation of β -sheet-rich aggregates of a specific protein in the brain parenchyma of affected individuals. Such protein aggregates are believed to be the proximate cause of neurotoxicity in some conditions and are considered to be an end-stage product of a cascade of events in others (181). The underlying mechanism leading to the development of protein

Reviewing Editors: Andrew F. Hill, Sophie Mouillet-Richard, Jesús R. Requena, and Jason Shearer

Department of Pathology, Case Western Reserve University, Cleveland, Ohio.

aggregates and accompanying neurotoxicity is relatively clear for some of these conditions, whereas in others, it is still debated. Recent evidence suggests the imbalance of brain metal homeostasis as a common cause of neuronal death in several of these disorders (9, 45, 48, 60, 61, 70, 111, 148, 171, 203, 242). It is believed that a redox-active metal interacts with a specific protein and is reduced in its presence, resulting in the generation of reactive oxygen species (ROS), hydrogen peroxide (H_2O_2), and hydroxyl radicals (OH^\bullet) that cause aggregation of the involved protein (16, 206, 215, 224, 225). Prominent redox-active metals that undergo this type of reaction include copper and iron, metals that are often detected in association with protein aggregates specific to AD, PD, and prion disorders (9, 150, 171, 206, 252). Whether the accumulation of these metals is a cause or consequence of the disease process is a subject of much dispute. Although significant information on the contribution of redox-active metals in the pathogenesis of AD, PD, and HD is available, similar information on prion disease-associated neurotoxicity is limited.

Studies aimed at understanding the role of metals in prion-disease pathogenesis have been hindered because of the insoluble nature of PrP^{Sc} , the principal agent involved in the pathogenesis of prion disorders (37, 49, 176). Although the biology and pathophysiology of metal- PrP^{Sc} interaction has been difficult to elucidate, the interaction of metals with PrP^C , the normal counterpart of and the substrate for PrP^{Sc} , has been studied extensively by using both *in vitro* and *in vivo* techniques. Collectively, the available evidence points to a functional role for PrP^C in the uptake and metabolism of copper and iron (117, 142, 143, 204, 205), leading to the hypothesis that dysfunction of PrP^C due to conversion to the PrP^{Sc} form may contribute to the imbalance of brain metal homeostasis observed in prion disorders (203). Other mechanisms leading to altered metal metabolism and disease pathogenesis have been proposed, and consensus on this issue is still lacking (40, 44).

At the center of this debate is PrP^{Sc} , the principal molecule involved in the pathogenesis of all prion disorders through mechanisms that are still unclear (1–3). Several pathways of neurotoxicity by PrP^{Sc} have been suggested, the most prominent being the induction of toxic signals through PrP^C at the neuronal plasma membrane (136). However, neuronal death is often seen in the absence of PrP^{Sc} and *vice versa*, suggesting the presence of alternative pathways of neurotoxicity in addition to PrP^{Sc} (119). One of these factors that is often ignored is the depletion of PrP^C due to conversion to the PrP^{Sc} form. Because PrP^C plays a prominent role in protecting neurons against oxidative stress, a significant reduction in PrP^C levels may increase the susceptibility of neurons to toxic insults (31, 46, 129, 135, 189). Keeping these facts in mind, both gain of toxic function by PrP^{Sc} and loss of normal function of PrP^C have been proposed as possible mechanisms of neurotoxicity, and information on both fronts is still emerging. The contribution of these factors individually to disease pathogenesis is difficult to assess because the normal function of PrP^C itself is not entirely clear. Diverse functions have been attributed to PrP^C based on the model and the experimental design used for evaluation (126, 129, 239). However, because PrP^C is involved in copper and iron uptake, it is possible that loss of normal function of PrP^C in copper and iron metabolism, combined with gain of toxic function by redox-active PrP^{Sc} -ferritin aggregates, induce a state of brain metal im-

balance, resulting in metal-induced oxidative stress and neurotoxicity (12, 166, 203–205, 239). The accumulated redox-active iron and copper would further react with dioxygen species abundant in the metabolically active environment of the brain, further aggravating the oxidative damage (169).

In this review, we describe the unique nature of prion disorders and the prevailing hypotheses on the mechanism of neurotoxicity in these conditions. Emphasis is placed on reports that support or contradict the involvement of redox-active metals and associated oxidative stress in the pathogenesis of these disorders. Evidence from other neurodegenerative conditions in which the pathogenic role of redox-active metals has been established is discussed with relevance to the available information on prion disorders. Specific information on the pathophysiology of redox-active metals in prion-disease pathogenesis is divided into three sections: (a) physiologic functions of PrP^C as an antioxidant and in copper and iron uptake, (b) pathologic implications of PrP^C -metal interaction, and (c) the origin and functional significance of brain-iron dyshomeostasis in prion disorders. We conclude by summarizing possible therapeutic strategies based on the premise that redox-active metal-mediated oxidative stress is a significant cause of neurotoxicity in prion disorders.


II. Prion Disorders: An Overview

A. The unique nature of prion disorders

Prion disorders affect humans and animals and are unique among neurodegenerative conditions of protein aggregation because of their infectious nature. Human prion disorders include sporadic Creutzfeldt-Jakob disease (sCJD), which comprises 80% of all cases, the less-common inherited forms classified as familial CJD, Gerstmann-Straussler-Scheinker (GSS) disease, and fatal familial insomnia (FFI), based primarily on their phenotypic presentation, and the relatively rare infectious forms such as variant CJD. Animal prion disorders include bovine spongiform encephalopathy (BSE) of cattle, scrapie of sheep, chronic wasting disease (CWD) of deer and elk, and prion diseases of other animals. The principal event in all prion disorders is the change in conformation of PrP^C from a mainly α -helical to a β -sheet-rich PrP^{Sc} form. Deposits of PrP^{Sc} in the brain parenchyma are considered to be the principal cause of neurotoxicity and infectivity in most prion disorders, although prion disease-associated neurotoxicity is sometimes seen in the absence of PrP^{Sc} and *vice versa* (1–3, 40, 79, 115).

PrP^C is a normal cell-surface glycoprotein linked to the plasma membrane by a glycosylphosphatidylinositol (GPI) membrane anchor. Like other glycoproteins, PrP^C is synthesized as a precursor form with N- and C-terminal signal peptides and is translocated into the endoplasmic reticulum co-translationally (Fig. 1). The N-signal peptide is cleaved soon after translocation, and the C-terminal signal peptide is replaced by a preassembled GPI-anchor in a transamidation reaction within 5 min of synthesis. Subsequently, high-mannose glycans and a disulfide bond are added, and the glycans are trimmed and modified as the protein traverses the Golgi apparatus on its way to the plasma membrane (Fig. 1) (41). The change in conformation of PrP^C to the PrP^{Sc} form renders it resistant to limited digestion by proteinase-K (PK), sparing the C-terminal fragment comprising of amino acids 90 to 231 and the GPI anchor (Fig. 1). In most instances, the

FIG. 1. Precursor and post-translationally processed PrP^C. Precursor PrP^C is translocated co-translationally into the ER where the N-signal peptide is cleaved by a signal peptidase. After translocation, the C-terminal GPI-signal peptide is replaced by a preassembled GPI anchor in a transamidation reaction within 5 min of synthesis. Subsequently, the disulfide bond and mannose-rich glycans are added in the ER, and the protein is transported to the Golgi complex where immature glycans are processed to their complex form. The mature protein is transported to the plasma membrane through secretory vesicles. The change in conformation of PrP^C to PrP^{Sc} renders the protein partially resistant to digestion by limited amounts of PK that cleaves PrP^{Sc} at the indicated sites. The C-terminal PK-resistant fragment is sufficient by itself for prion disease-associated neurotoxicity and infectivity. CHO, carbohydrates; S-S, disulfide bond; PK, proteinase K; N-SP, N-terminal signal peptide; GPI-SP, C-terminal signal peptide.



PK-resistant C-terminal fragment is sufficient to transmit prion infection and induce neurotoxicity in the recipient animal. Both full-length and the C-terminal fragment of PrP^{Sc} aggregate easily, are insoluble in non-ionic detergents, and resist degradation when exposed to a variety of denaturing agents (Table 1) (1-3). Often referred to as the prion agent (176), PrP^{Sc} derives its infectious nature from its unique characteristic of transforming additional PrP^C molecules to its own β -sheet-rich PrP^{Sc} conformation (Fig. 2). However, the mechanistic details of the conversion process are not completely clear. In sporadic prion disorders, the conversion of PrP^C to PrP^{Sc} is believed to occur as a random event. In inherited disorders, certain point mutations in the prion protein-coding region induce the transformation of PrP^C to PrP^{Sc}, and in infectious disorders, PrP^{Sc} from an exogenous source induces the conversion of host PrP^C to its own conformation (1-3, 40, 176). Figure 3 demonstrates aggregates of PrP^{Sc} in the brain parenchyma of a sCJD case immunostained with PrP-specific antibody and spongiform change in the surrounding cells.

Infectious prion disorders were first recognized as Kuru in the Fore tribe of Papua New Guinea, where cannibalism was a ritualistic practice, and later as variant CJD that was most likely transmitted to humans through the consumption of BSE-infected meat (36, 50). Currently, CWD is spreading in

certain parts of the United States, and the possibility of horizontal spread of this agent and transmission to cattle and humans is uncertain (56, 237). In addition to the oral route, the prion agent can be transmitted through the bloodstream or peripheral nerves, either accidentally, iatrogenically, or through skin abrasions (128, 221). Although much publicized, infectious prion disorders of humans are relatively rare and comprise only 1% of all reported cases of prion disorders (Table 2). It is important to note that although infectivity of PrP^{Sc} has been demonstrated for most cases, some forms are relatively resistant to transmission. The prion disorders of humans and animals and the underlying cause of disease are described in Table 2 (2, 40, 53, 176). For discussion, we do not distinguish between PrP^{Sc} of sporadic, inherited, or infectious origin, and consider it as the common pathogenic agent for all prion disorders.

1. The “prion” agent. Mammalian prions are considered synonymous with PrP^{Sc}, and the word “prion” is derived from proteinaceous infectious particle to symbolize the unique characteristic of a protein as the principal infectious agent (176). Over the years, diverse experimental models have been

TABLE 1. BIOCHEMICAL AND BIOPHYSICAL CHARACTERISTICS OF PrP^C AND PrP^{Sc}

Biochemical characteristic	PrP ^C	PrP ^{Sc}
Secondary structure	43% α -helix	30% α -helix 43% β -sheet
Solubility in non-ionic detergents	Soluble	Insoluble
Sensitivity to limited digestion with PK	PK sensitive	PK resistant
Aggregation	Not aggregated	Aggregated
Infection	Not infectious	Infectious




FIG. 2. Structural differences between PrP^C and PrP^{Sc}. A model of the α -helical structure of PrP^C and the β -sheet-rich PrP^{Sc} form implicated in prion-disease pathogenesis.




FIG. 3. Deposits of PrP^{Sc} in the brain. Brain section from a case of sCJD immunostained with anti-PrP antibody 3F4 shows immunoreactive deposits and spongiform change in the surrounding tissue.

used to understand the mechanism(s) of conversion of PrP^C to PrP^{Sc} and the cause of the associated neurotoxicity. Several important facts have emerged from these studies. For example, it is now clear that for prion transmission, a certain degree of homology between the exogenous PrP^{Sc} and host PrP^C is required, the absence of which accounts for the species barrier in prion transmission and the presence of prion-resistant genotypes (2, 79, 115). In some cases, however, heterologous PrP^{Sc} has been demonstrated to induce subclinical disease or

adapt to an apparently resistant host and to cause disease on subsequent passage (92, 178). In familial disorders, point mutations are believed to render mutant PrP forms susceptible to a change in conformation to PrP^{Sc} at an advanced age, although the contributing factors that precipitate this change remain unidentified (2, 176). The fact that PrP^{Sc} can be taken up by neuronal cells fairly nonspecifically, but only a few cell lines can sustain persistent PrP^{Sc} infection, points to cell-specific factors that are necessary for successful propagation (19, 59, 134, 229). Several experimental paradigms have been tried to understand the generation of PrP^{Sc} *in vitro* and *in vivo*. The most prominent *in vitro* method involves conversion of cellular, brain-derived, or recombinant PrP^C to a form similar to PrP^{Sc} by the protein-misfolding cyclic-amplification reaction (PMCA) by using brain-derived PrP^{Sc} as the inoculum (39). The conversion process is enhanced by RNA molecules, redox-active metals like copper, certain denaturants, sulfated glycans, solvents, pH, temperature, reducing agents, inhibition of glycosylation, and a combination of buffer conditions and metal ions (156, 197, 246). However, PrP^{Sc} generated by only some of these reactions can change additional PrP^C to PrP^{Sc} *in vitro* and cause disease when introduced to recipient animals (39, 54). Other forms resemble PrP^{Sc} in certain biochemical characteristics and either are not infectious, or have not been assessed in bioassays to evaluate infectivity (8, 156).

2. Mechanism(s) underlying prion-associated neurotoxicity. Despite evidence suggesting PrP^{Sc} as the principal pathogenic and transmissible agent responsible for all prion disorders, it is important to note that prion-specific neurotoxicity is sometimes observed in the absence of detectable PrP^{Sc}, and PrP^{Sc} deposits may occur in the absence of neurotoxicity (43, 67, 119, 173). Thus, other players besides PrP^{Sc}

TABLE 2. HUMAN AND ANIMAL PRION DISORDERS

<i>Prion disease</i>	<i>Underlying cause of disease</i>
Human	
Sporadic	Spontaneous conversion of PrP ^C to PrP ^{Sc}
Creutzfeldt-Jakob disease (sCJD), Fatal	
Familial insomnia (sFFI)	
Familial	Point mutations in the coding region of <i>PRNP</i> gene leading to conversion of mutant PrP to the PrP ^{Sc} form
fCJD, FFI	
Gerstmann-Sträussler-Scheinker syndrome (GSS)	
Infectious	Acquired from an exogenous source of PrP ^{Sc} through food products or iatrogenically
Variant CJD	
Iatrogenic CJD	
Kuru	
Animal	
Sporadic	Spontaneous conversion of PrP ^C to PrP ^{Sc}
Scrapie of sheep	
Chronic wasting disease (CWD) of deer and elk	
Infectious	Horizontal and vertical transmission of PrP ^{Sc} from feed and through poorly understood mechanisms
Bovine encephalopathy of cattle (BSE)	
Exotic ungulate encephalopathy	
Feline spongiform encephalopathy	
Scrapie and CWD	
Transmissible mink encephalopathy	

that influence prion transmission and neurotoxicity exist and are gradually being discovered. Several important facts have emerged from these studies. For example, it is clear that regardless of PrP^{Sc} load, expression of PrP^C on the host neuronal plasma membrane is essential for neurotoxicity. Transgenic mice expressing anchorless PrP^C accumulate infectious PrP^{Sc} in the extracellular milieu but do not demonstrate neuronal death, suggesting that the neurotoxic signal is transmitted through plasma membrane-associated PrP^C (43, 79, 136). However, astrocyte-specific expression of PrP^C is sufficient to support the replication and neurotoxicity of infectious PrP^{Sc}, even in the absence of neuronal PrP^C expression, indicating the involvement of other neurotoxic molecules or mechanisms in this process (98). Controversies about the neurotoxic nature of PrP^{Sc} have led to the identification of other PrP conformers that may be involved in the pathology of prion disorders (44, 81). Prominent among these are the C-transmembrane and cytosolic forms of PrP that induce neurotoxicity by poorly understood mechanisms (72, 82–84, 130, 131, 145). Investigations of mutant PrP forms by using cell models of familial prion disorders indicate that a combination of direct and indirect effects of the mutation alter posttranslational processing, transport, and the cellular site of accumulation of specific PrP forms, resulting in toxicity due to aggregation or conversion of PrP^C to a PrP^{Sc}-like form at that site (71–75, 101, 145). In addition, PrP amyloid fibrils (157), certain structurally misfolded intermediates of PrP^C, and small oligomers in the pathway of PrP^{Sc} formation (79, 202) also are considered neurotoxic, possibly because of characteristics unique to their structure. Cross-linking of PrP^C on the plasma membrane has been noted to induce neurotoxicity (209), perhaps by activating certain apoptotic pathways or by compromising the normal function of PrP^C (174, 182–184).


A compelling set of observations suggest that subversion or loss of normal function of PrP^C is an equally significant cause of prion disease-associated neurotoxicity (34, 66, 79, 155). The specific function of PrP^C involved in this process, however, has been difficult to identify because of the variety of functional activities attributed to this protein, based on the experimental design and the method used. Among these, the neuroprotective function of PrP^C demonstrated in cell lines,

primary neurons, and *in vivo* models is noteworthy (174, 183). Studies of primary human neurons suggest that PrP^C protects cells against Bax-induced cell death by inhibiting the conformational change of Bax, supporting the loss-of-function hypothesis of neuronal death (184, 185). Other studies suggest a protective role of PrP^C by virtue of its property to function as a Cu/Zn superoxide dismutase (SOD), although several other studies refute this claim (26, 32, 96). It is likely that compromised function of this enzyme in prion disease-affected brains (245) increases the susceptibility of neurons to toxic signals that are generated as a by-product of normal cellular metabolic processes. It has also been proposed that PrP^C is a free radical-scavenging protein, and loss of this activity in mutant PrP forms increases the susceptibility of neurons to toxicity (236). The protective function of PrP^C is further exemplified by the fact that expression of a single copy of PrP^C can rescue the ataxic phenotype of transgenic mice expressing a deletion construct of PrP (PrP Δ 32-121/134) (199), slow neurodegeneration in transgenic mice expressing a pathogenic mutation of PrP (217), rescue neurons from Doppel-induced toxicity (149), and protect brain tissue from ischemia- and trauma-induced insult (210, 238). Moreover, PrP^C is believed to function as a copper- and iron-uptake protein, suggesting a role in the maintenance of cellular copper and iron homeostasis (33, 126, 205, 226). Collectively, these reports suggest that loss of normal function of PrP^C can be as significant as accumulation of PrP^{Sc} in inducing prion disease-associated neurotoxicity (Fig. 4). Although it is difficult to distinguish between these possibilities because PrP^C is the substrate for PrP^{Sc}, these observations underscore the importance of the physiologic function of PrP^C in disease-associated neurotoxicity.

B. Redox metals, oxidative stress, and prion-disease pathogenesis

Among the redox-active metals, iron and copper are most relevant because PrP^C is involved in their metabolism, thereby increasing the likelihood of their involvement in prion-disease pathogenesis. A significant amount of information is already available on the neurotoxic role of iron in disorders

FIG. 4. Loss of PrP^C function as a contributing factor in prion disease-associated neurotoxicity. PrP^C has been ascribed several physiologic functions, including a role in protecting cells from oxidative stress. The change in conformation of PrP^C to the PrP^{Sc} form is likely to compromise the normal protective function of PrP^C, resulting in neurotoxicity.



such as AD, multiple sclerosis, PD, tardive dyskinesia, Pick's disease, HD, Hallervorden-Spatz disease, Friedreich's ataxia, and aceruloplasminemia (10, 11, 68, 250–253). In addition to iron excess, a deficiency of this metal is equally harmful and can result in compromised motor and cognitive function, in addition to other manifestations of anemia (15, 133). Likewise, an increase in brain copper content is neurotoxic (213, 218), and copper deficiency is known to compromise the function of Cu/Zn-SOD, an important antioxidant enzyme essential for cell survival (13). Zinc is redox inactive but is important for neurotransmission, and inadequate supplies can result in compromised neuronal function (235). The role of other metals in brain function has been reviewed elsewhere (148).

Observations from prion disease-affected human and mouse brains suggest that iron- and copper-induced oxidative damage is a significant factor affecting disease pathogenesis (107, 109–111, 142, 171, 201). Although the underlying cause of metal imbalance is not clear, recent reports indicating a facilitative role of PrP^C in iron and copper uptake and interference with manganese uptake suggest that loss of this function as a result of aggregation to the PrP^{Sc} form may contribute to brain metal dyshomeostasis, resulting in metal-induced oxidative damage (147, 166, 204, 205). Furthermore, iron and copper may remain associated with PrP^{Sc}, thus rendering the complex redox active and neurotoxic (12, 203). Observations reporting the generation of H₂O₂ by certain fragments of PrP, when exposed to iron or copper, lend credence to this hypothesis (106, 224, 225). Whether these fragments exist *in vivo* and undergo such a reaction is not clear, but it suggests that abnormal forms of PrP or its fragments are likely to contribute to the generation of free radicals. Similar observations have been reported for amyloid-beta (A β), the principal component of neurotoxicity in AD. The interaction of A β with iron and copper is believed to mediate its protease resistance and oxygen-dependent production of free radicals (4, 207). Normally, antioxidants such as glutathione transferase and catalase detoxify the free radicals produced through these reactions to protect the cells from the highly neurotoxic OH \cdot radical generated by the Fenton reaction. However, increased production combined with compromised capability to detoxify such radicals is likely to result in the neurotoxicity associated with AD.

A similar process of oxidative stress and associated neurotoxicity is likely to occur during prion-disease progression. Markers of metal-induced oxidative stress such as products of lipid peroxidation, including free malondialdehyde (MDA) and increased levels of Fe²⁺ and Fe³⁺ ions, have been identified in the cerebral cortex, striatum, and brainstem of prion disease-affected human and mouse brains, supporting this assumption (5, 108, 109, 111, 171, 203). The state of increased oxidative stress could result from direct toxicity by PrP-metal complexes, increased susceptibility of neuronal cells to free radicals due to the compromised function of PrP^C, or a combination of both processes (22, 140, 141, 182). It is more likely that both processes operate simultaneously because PrP knockout (PrP^{-/-}) mice that lack PrP expression exhibit increased vulnerability to superoxide, H₂O₂, and copper ions, and show evidence of oxidative stress, such as elevated levels of protein carbonyls and ubiquitin-protein conjugates in their brains (20, 112, 187, 244, 245). In scrapie-infected mouse brains and cells replicating mouse prions, activities of antioxidant enzymes glutathione-peroxidase, reductase, and SOD are re-

duced, whereas markers of oxidative stress such as MDA, 4-hydroxyalkenals (HAE), and reactive aldehyde products of lipid peroxidation are increased (25, 141, 249). In addition, levels of 3-nitrotyrosine, heme oxygenase-1, carbon monoxide, and iron are increased, further supporting these observations (47, 77, 203, 249). These changes cause oxidative damage to proteins, nucleic acids, and lipids, prominent features of scrapie-infected mouse and CJD-affected human brains (64, 76, 162).

Evidence of oxidative stress is also noted in the cerebrospinal fluid (CSF) and plasma of sCJD patients that show lipid peroxidation and reduced levels of polyunsaturated fatty acids, ascorbate, and α -tocopherol (6). In certain cases of GSS and vCJD, deposits of iron and its storage protein, ferritin, are seen in the brain parenchyma in association with and in the tissue surrounding PrP^{Sc} plaques (171), suggesting a causal relation. In addition, sCJD- and scrapie-infected Syrian hamster brains show a combination of oxidative, glycoxidative, lipoxidative, and nitrative protein damage, accompanied by increased oxidative response, as indicated by elevated levels of the oxidized nucleic acid base 8-hydroxyguanosine (8OHG) and HNE-modified proteins, neuronal, endothelial, and inducible nitric oxide synthase (nNos, eNos and iNos), SOD 1, and SOD 2, glutamic and aminoadipic semialdehydes (products of metal-catalyzed oxidation), malondialdehydelysine (product of lipoxidation), *N*- ϵ -carboxyethyl lysine (product of glycoxidation), and *N*- ϵ -carboxymethyl lysine (generated by lipoxidation and glycoxidation) (64, 76, 162, 249). An *in vivo* study of scrapie-infected mice links peroxidative damage directly to neuronal loss and *de novo* PrP^{Sc} propagation, establishing a causal link between PrP^{Sc} accumulation, oxidative damage, and disease pathogenesis (25). PrP^{Sc} itself is a target of increased oxidative modifications (162), suggesting the presence of chronic oxidative stress in diseased brains. Interestingly, metal-induced oxidative stress has been reported to cause the aggregation and insolubility of PrP^C to a form similar to PrP^{Sc} in cell models (12) and to decrease infectivity by causing the hydroxylation and degradation of PrP^{Sc} (164), indicating a role in both the generation and degradation of PrP^{Sc}. Besides, it has been suggested that the conversion of PrP^C to PrP^{Sc} involves the formation of a redox-active copper complex within the plasma membrane (125, 142, 154), supporting the idea that the generation of PrP^{Sc} is facilitated by redox ions. Although apparently contradictory, these observations probably reflect a continuum of the same process; redox-iron induced aggregation of PrP^C to PrP^{Sc}, followed by degradation of oxidatively modified PrP^{Sc} by scavenger cells using a similar metal-catalyzed reaction.

In addition to the direct damage to proteins and lipids noted earlier, reactive oxygen species (ROS) also sensitize the cells to endoplasmic reticulum (ER) stress (88). This is an important observation because several mutant forms of PrP associated with familial prion disorders induce ER stress due to misfolding or aggregation or both. This would aggravate the toxicity due to ER and oxidative stress individually, providing a common pathway of neuronal death for sporadic and familial prion disorders (88, 89). Furthermore, activity of NF- κ B, a transcriptional activator of proinflammatory cytokines, and proinflammatory cytokines such as TNF- α and interleukins 1 α and 1 β are elevated in scrapie-infected mouse brains, a situation that can further induce the upregulation of iron-

uptake proteins and increase the content of brain iron, thus creating an ongoing state of oxidative stress (108, 114, 150).

Collectively, this information leaves little doubt that oxidative stress plays a prominent role in prion disease-associated neurotoxicity. However, it is difficult to conclude whether this is a cause or consequence of the disease process. Subsequently, we review available information on PrP-metal interaction, the physiologic and pathologic implications of this association, and the likely cause(s) of brain metal dys-homeostasis in prion disorders.

III. Redox Control of Prion Protein: Physiological Implications


The physiologic role of PrP^C in cellular redox control is poorly understood. However, enough evidence exists to implicate PrP^C in redox reactions that serve the general purpose of intracellular signaling (174, 193), protection against oxidative stress (120, 219), and modulation of synaptic activity (85, 117, 228). The redox potential of PrP^C is mainly because of its association with redox-active metals such as copper and iron, although PrP is known to interact with other metals, including zinc, manganese, and nickel (23, 29, 97, 233). Copper and zinc interact with the octapeptide repeat region of PrP^C at the N-terminus and influence its functional characteristics in distinct ways (142, 233). It has been suggested that this interaction facilitates the folding of the largely unstructured N-terminus, providing stability to the protein (121). Because prion disease-affected brains contain altered concentrations of Cu and other divalent cations (87, 219, 243), it is likely that a deficiency or excess of these metals further contributes to the instability of PrP^C and an increased propensity to convert to the PrP^{Sc} form (125). Manganese also interacts with PrP^C and alters its function (28, 48). Relatively less is known about the interaction of PrP^C with iron, a metal with significantly higher redox potential than copper. Because both copper and iron are essential for cellular metabolism and are toxic if mismanaged, the interaction of PrP^C with these metals has both physiologic and pathologic implications. We first summarize information on PrP^C-copper interaction, because this is better understood, followed by available information on the interaction of PrP^C with other metals, including iron.

A. PrP and copper interaction

A growing body of evidence suggests that PrP^C is involved in copper metabolism. This phenomenon has been studied extensively by using recombinant PrP^C, peptide fragments of PrP^C, cell models, and mouse models (29, 52, 102, 113, 226). Collectively, these studies indicate that human PrP^C binds five Cu²⁺ ions under physiologic conditions. Four copper-binding sites are within the octapeptide repeat sequence Pro-His-Gly-Gly-Gly-Trp-Gly-Gln, between residues 61 and 91 of PrP (Fig. 5). This region binds copper ions with femto- and nanomolar affinity, and other metal ions like Ni²⁺, Zn²⁺, and Mn²⁺ with affinities lower by at least three orders of magnitude. The fifth copper-binding domain is between residues 91 and 111 and is coordinated by histidine residues at positions 96 and 111 of the PrP sequence (Fig. 5). This site has a lower affinity for copper than the octapeptide repeat region (7, 93–95, 103, 195). The binding is optimal at physiologic pH 7.4, and decreases precipitately at acidic pH (241). However, the *in vitro* results should be interpreted with caution because these studies use recombinant PrP^C, and the affinity of proteins for metals is known to differ by several orders of magnitude when they are in their native conformation and environment. It is notable that the octapeptide repeat region is conserved among mammalian species, and insertion of one or more octapeptide units in the PrP sequence is associated with familial CJD in humans and prion disease in experimental mice (211). The mechanism by which additional repeats induce disease, however, is not entirely clear. It is probable that altered copper concentrations in the brain due to increased binding sites play a role (116). Further investigations are required to understand this phenomenon fully.

Although the binding sites and affinity of PrP^C for copper are known, the redox state of bound copper is not clear. Some reports suggest that PrP^C-associated copper is in its reduced form because of the presence of tryptophan residues within the PrP sequence that function as redox acceptors (186). Contradictory reports indicate that copper is bound in the nonreduced form and sequesters potentially harmful free radicals, thereby serving a protective function (198). Exposure of PrP^C-bound copper to H₂O₂ oxidizes Cu¹⁺ to Cu²⁺ with the generation of hydroxyl radicals that are highly toxic and are

FIG. 5. Copper- and iron-binding sites on PrP^C. A model of PrP^C with known copper-binding sites in the octapeptide repeat region and the histidine residue 111. The putative iron-binding sites are based on data from recombinant full-length PrP^C and its fragments. Denaturation of PrP^C releases all bound iron, suggesting that the interaction of PrP^C with iron is sensitive to its secondary or tertiary structure.






FIG. 6. Regulated and ROS-mediated cleavage of PrP^C. PrP^C undergoes a regulated cleavage at amino acids 110/111 mediated by the ADAM family of proteases in an endocytic compartment, termed α -cleavage. The resulting C-terminal fragment is transported back to the plas-

ma membrane. In addition, PrP^C undergoes ROS-mediated cleavage near amino acid 90, termed β -cleavage. This event is believed to protect PrP^C-expressing cells from free radical damage.

believed to cause the aggregation of PrP^C. The redox-active form of copper reacts with free radicals and induces the generation of ROS, resulting in β -cleavage of PrP^C near the octapeptide repeat region, a reaction facilitated by tryptophan residues that act as one-electron donors to mediate the reduction of the Cu²⁺ center to Cu¹⁺ (Fig. 6) (234). Based on these observations, PrP^C has been proposed as a “sacrificial quencher” of free radicals (154). Similar aggregation of PrP^C is noted when the cells are exposed to a source of redox-iron (12), suggesting that the PrP-metal interaction may serve a protective role by sequestering extracellular redox-active ions, and aggregation of PrP^C may in part be a consequence of this process. The affinity of PrP^C for copper is optimal to quench excess copper ions and to function as a copper-uptake protein, supporting these observations (125, 126, 142).

The functional role of PrP^C in copper uptake from the extracellular milieu has been studied in cultured cells and mouse models. In mouse neuroblastoma cells, extracellular copper ions stimulate the endocytosis of PrP^C, supporting the idea that PrP^C may bind and deliver extracellular copper ions to endocytic compartments. Deletion of the octapeptide repeat region or mutation of histidine residues within this region abolishes copper uptake by PrP^C, emphasizing the role of this region in copper binding (35, 166, 167, 198, 212, 216). A similar mechanism of copper uptake has been demonstrated in PrP^C-GFP-expressing SN56 cells (122). In addition to serving as a copper-uptake protein, it is believed that the octapeptide repeat region of PrP^C reduces captured Cu²⁺ ions before their delivery to Cu¹⁺-specific intracellular copper-carrying proteins (147), serving an important role in copper transport across the endosomal membrane. Studies using radioactive copper indicate that nonchelated ⁶⁷Cu is taken up by PrP^C-deficient neurons as efficiently as neurons expressing PrP^C, whereas ⁶⁷Cu chelated with histidine is taken up in proportion to neuronal PrP^C expression. Interestingly, ⁶⁷Cu levels in SOD 1 are in direct proportion to PrP^C-expression levels, suggesting that decreased SOD 1 activity in PrP^C-deficient cells may in part be due to deficient delivery of copper to the enzyme (33). Similar observations have been reported for a rat kidney cell line RK13A, in which expression of PrP^C increases copper binding and the activities of antioxidant enzymes without altering copper delivery (180). Although these observations implicate PrP^C as a major copper-uptake and -delivery protein, this is unlikely, because the brain copper and zinc content of mice expressing wild-type levels or 10-fold higher levels of PrP^C are similar to those in mice lacking PrP (PrP^{-/-}) (231), leaving the matter unsettled. A plausible explanation for these disparate findings could be that PrP^C alters the distribution of copper rather than its overall content in the brain, as observed for zinc (179). This

assumption is supported by the fact that the copper content of synaptosomal membranes that express high levels of PrP^C is twofold higher in wild-type mice compared with PrP^{-/-} mice, an observation that has prompted the suggestion that PrP^C regulates copper concentration at the synapse by serving as a sink for excess copper ions released during synaptic vesicle fusion (117, 228). Such a function would maintain physiologically safe levels of copper in presynaptic cytosol, a site that is exposed to high concentrations of copper because of release of copper ions from nerve endings after depolarization. In addition, such a function would prevent the potentially harmful participation of unbound copper in Fenton- and Heber-Weiss-type redox reactions that generate reactive oxygen species and redistribute the released copper back to presynaptic cytosol for recycling (228). Figure 7 is a diagrammatic representation of the putative mechanism(s) of copper uptake and transport by PrP^C.

Interestingly, copper ions have been demonstrated to upregulate PrP^C expression, and conversely, PrP^C regulates the copper content of cells. The former function is achieved by activating a metal-responsive element in the promoter region of PrP^C (17, 227), and the latter, by shedding PrP^C into the culture medium when exposed to copper or to a zinc metalloprotease (165, 223). Upregulation of PrP^C by copper is reversed by copper chelators, confirming the specificity of the reaction. A similar upregulation of PrP^C is noted when cells are exposed to iron or cadmium, but not to zinc or manganese (12, 226). In this regard, the effect of metals on PrP^C expression differs from metallothionins and Cu/Zn-SOD genes that are induced by all three metals: copper, cadmium, and zinc (153, 179, 247).

Together, these studies suggest a role for PrP^C in cellular copper uptake and transport, maintenance of physiologically safe copper concentrations at the synapse, upregulation of PrP^C expression in response to copper, and cleavage, shedding, and aggregation of copper-bound PrP^C when exposed to free radicals. Other less-defined outcomes of PrP^C-copper interaction have also been reported, leading to the conclusion that the role of PrP^C in copper metabolism is complex and is likely to involve presently unidentified pathways that influence cellular copper homeostasis.

B. PrP and other metals, in particular iron

Physiological implications of the interaction of PrP^C with other metals, such as manganese, iron, zinc, and nickel, are poorly understood (86, 87). The majority of these metals induce the aggregation of purified or recombinant PrP^C to a form resembling PrP^{Sc} in certain characteristics (156). How-




FIG. 7. Mechanism of copper uptake by PrP^C. PrP^C is a GPI-linked protein that normally resides in cholesterol- and sphingolipid-rich rafts on the plasma membrane. On exposure to copper, PrP^C binds copper in the octapeptide repeat region and relocates to the vicinity of a transmembrane protein, and the complex is internalized in clathrin-coated pits. Bound copper is released in an endosomal compartment and reduced by the octapeptide repeat region of PrP^C before transfer to cytosolic copper-carrier proteins. The N-terminal region of PrP^C is cleaved proteolytically in an endocytic compartment (α -cleavage), and the C-terminal 18-kDa fragment is transported back to the plasma membrane.

ever, these evaluations involve *in vitro* experiments, and the influence of these metals in the *in vivo* situation is unclear, except for zinc, which induces rapid turnover of PrP^C (35). Limited studies suggest that manganese can displace copper and bind to PrP^C, which then mediates its uptake into cells and is rendered protease resistant by this association (28).

The interaction of PrP^C with iron deserves a special note, because iron is required for optimal neuronal growth (14), and, like copper, is considered a toxin because of its ability to exist in two oxidation states, ferric (Fe³⁺) and ferrous (Fe²⁺) (104, 220). Free iron can catalyze the conversion of hydrogen peroxide to reactive hydroxyl radicals by the Fenton reaction, resulting in oxidative damage. Furthermore, iron-dependent lipid peroxidation generates potentially toxic peroxy/alkoxy radicals (78), and iron is known to convert neutral catechols to neurotoxic intermediates, compounding the neurotoxicity (192). Figure 8 provides a partial list of normal functions of iron in the brain and possible mechanisms of iron-mediated toxicity, and Fig. 9 illustrates the possible contribution of PrP^C and PrP^{Sc} in this process.

It has been demonstrated that brain iron increases with age in humans, rats, and mice, increasing the chances of metal-

induced free radical injury detected most often in the basal ganglia, hippocampus, and cerebellar nuclei (51, 137, 208). Levels of ferritin also increase with aging in human brains, suggesting a precarious balance of iron metabolism and an increased propensity for dysregulation with the slightest insult (251). Because imbalance of cellular iron homeostasis can result in the generation of ROS, the transport of iron in and out of the cells is tightly regulated. Within cells, iron is present in ferritin, which serves the general function of intracellular iron sequestration, detoxification, and storage. Under conditions in which capacity of ferritin to detoxify iron is exceeded, Fe²⁺ can participate in one-electron transfer reactions, resulting in the formation of reactive intermediates, including OH[•] radicals that can catalyze the oxidation of proteins, lipids, carbohydrates, and nucleic acids, ultimately leading to cell death by apoptosis or necrosis. Cells have therefore developed sophisticated mechanisms to regulate iron metabolism through coordinated control of transferrin (Tf) and transferrin receptor (TfR)-mediated uptake, and ferritin-mediated sequestration in the cytosol. Ferritin regulates the labile iron pool within cells and is itself regulated by iron-regulatory proteins (IRP) 1 and 2. When cells are iron deplete, IRP 1 and 2 bind to




FIG. 8. Brain-iron dyshomeostasis as a possible cause of prion disease-associated neurotoxicity. Maintenance of normal brain-iron homeostasis is essential for several vital metabolic processes, and iron deficiency is likely to compromise essential brain functions. Excess iron can induce neurotoxicity because of its redox-active nature, underscoring the necessity for maintaining brain-iron levels within normal limits.

iron-responsive elements (IREs) in the TfR and ferritin mRNAs, blocking degradation of the former and decreasing translation of the latter. The net result is an increase of a labile iron pool for metabolic use. The opposite scenario takes effect when cells are iron replete (132, 150, 163).

Within this tightly orchestrated mechanism of homeostasis by well-characterized proteins, it is surprising that PrP^{C} influences iron uptake and transport (204, 205). Efforts aimed at understanding the stoichiometry of PrP^{C} -iron interaction and specific binding sites have yielded mixed results. Recombinant full-length PrP (PrP^{23-231}) can be radiolabeled with iron (^{59}Fe) and fractionated on nondenaturing gels, indicating a sufficiently strong interaction (12). Cell-associated PrP can also be radiolabeled, and the PrP-iron complex can be isolated by immunoprecipitation, indicating a biologically relevant interaction (12). Surprisingly, ^{59}Fe bound to PrP^{23-231} is not transferred to apo-transferrin, a protein with a very high affinity for iron (Fig. 10A, lanes 1–6) except in the presence of citrate, which forms a ^{59}Fe -citrate complex for transfer to transferrin. This is evident from the appearance of a faster-migrating band with increasing concentrations of apo-Tf that

co-migrates with ^{59}Fe -charged Tf (Fig. 10A, lanes 1–11) (172). These results indicate that the association of ^{59}Fe with PrP^{23-231} is not due to nonspecific sticking or an artifact of the experimental procedure. Interestingly, denaturation of recombinant PrP^{23-231} with guanidinium hydrochloride releases the bound iron completely, whereas reduction of disulfide bonds before or after the addition of the ^{59}Fe -citrate complex has no measurable effect, suggesting that the interaction of PrP^{C} with iron is sensitive to its conformation (Fig. 10B, lanes 1 to 4). However, these observations are of limited relevance, because cell- and brain-associated PrP^{C} does not show a similar affinity for iron (12, 205). It is therefore likely that under physiological conditions, the interaction of PrP^{C} with iron is transient or of low affinity, such that the bound iron is released during experimental manipulation (205).

Despite uncertainties about the affinity and binding site(s) of PrP^{C} for iron, recent studies leave little doubt that PrP^{C} plays a significant role in cellular iron metabolism (204, 205). Observations on human neuroblastoma cells show that overexpression of PrP^{C} increases the cellular labile iron pool (LIP) and iron saturation of ferritin, whereas pathogenic and





FIG. 9. Possible mechanisms of neurotoxicity in prion disorders. Several triggers induce the conversion of PrP^{C} to the disease-causing PrP^{Sc} form, including metal-induced oxidative stress. Once formed, PrP^{Sc} deposits increase the content of redox-active iron in the brain, thus feeding into the generation of additional PrP^{Sc} and worsening the state of brain-metal homeostasis. Neurotoxicity results from the direct effect of PK-resistant or certain species of PK-sensitive PrP^{Sc} , or both, and because of the redox-active nature of PrP^{Sc} complexes through poorly understood pathways.

FIG. 10. Recombinant full-length PrP binds iron with relatively high affinity. (A) PrP²³⁻²³¹ (1 μ g) was radiolabeled with ⁵⁹Fe-citrate complex for 10 min, dialyzed extensively, and incubated with increasing concentrations of apo-Tf in the absence (lanes 1–6) or presence (lanes 7–10) of citrate, followed by fractionation on a native gel (172, 203–205). Autoradiography reveals a prominent ⁵⁹Fe-labeled band of PrP²³⁻²³¹ (lanes 1–6) that releases bound ⁵⁹Fe to a faster-migrating band that comigrates with ⁵⁹Fe-charged Tf only in the presence of citrate (lanes 7–11, *open arrow-heads*). (B) Denaturation of ⁵⁹Fe-PrP²³⁻²³¹ with guanidine hydrochloride (GdnHCl) releases ⁵⁹Fe completely (lanes 1 and 2), whereas reduction of disulfides with DTT either after or before exposure to ⁵⁹Fe has no effect on PrP²³⁻²³¹-iron interaction (lanes 3 and 4).



nonpathogenic mutations of PrP overexpressed to the same extent as PrP^C alter cellular LIP and ferritin iron levels in a manner that is specific to each mutation (205). The difference in the iron content of these cells is maintained when they are exposed to excess extracellular iron, suggesting a dominant role for PrP^C in cellular iron uptake and transport (205). Whether PrP^C mediates iron uptake by a novel pathway or modulates the conventional pathway(s) of iron uptake through Tf/TfR-mediated endocytosis is unclear from published reports. However, stimulation of endocytosis by a PrP^C-specific antibody increases intracellular iron stores, whereas expression of anchorless PrP^C that is not expressed on the plasma membrane abolishes this effect, suggesting an active role for PrP^C in iron uptake at the plasma membrane (205). It is clear that PrP^C is not involved in iron efflux from cells, reinforcing the idea that it influences iron uptake, not export (100, 205). The possible mechanism by which PrP^C modulates cellular iron levels is unknown at present. It is also possible that extracellular iron induces the movement of PrP^C from detergent-insoluble membrane domains, where it normally resides, to the proximity of TfR, as suggested for copper (147). Here it may enhance the binding of iron to Tf, the binding of iron-loaded Tf to its receptor, or stimulate the en-

docytosis of Tf/TfR complex by direct or indirect mechanisms (Fig. 11). Conflicting reports suggest that PrP^C undergoes clathrin-mediated endocytosis after associating with a transmembrane protein through its N-terminal domain (200) or through a caveolae-mediated endosomal pathway (170). It is interesting to note that PrP co-localizes with Tf and TfR within the endosomes, an observation suggestive of a functional association rather than co-residence due to a common mode of endocytosis (170). Other proteins known to modulate iron uptake by regulating the interaction of the Tf/TfR complex include hereditary hemochromatosis protein (HFE), although HFE decreases iron uptake in contrast to PrP^C, which has the opposite effect (232). It also is possible that PrP^C functions as a ferric reductase to facilitate the transport of Fe³⁺ iron released from Tf to cytosolic ferritin, as described for copper, with which PrP^C is believed to reduce Cu²⁺ before transfer to Cu¹⁺-specific trafficking proteins in the cytosol (147). Figure 11 represents possible pathways of uptake and transport of iron by PrP^C.

The functional role of PrP^C in iron uptake is further exemplified by PrP^{-/-} mice that contain a targeted deletion of *PRNP*, the gene encoding PrP, and do not express PrP^C. As noted in cell models, lack of PrP^C expression induces a




FIG. 11. Possible mechanisms of iron uptake by PrP^C. Cell-surface PrP^C may bind iron from the extracellular milieu and mediate its uptake directly by endocytosis, or modulate the uptake of iron by the Tf/TfR complex. Alternately, PrP^C may mediate the transport of iron across the endosomal membrane by functioning as a ferric reductase. Iron transported to the cytosol enters the labile iron pool for metabolic processes, and excess is stored within ferritin in a relatively inert form.

phenotype of iron deficiency in PrP^{-/-} mice relative to matched wild-type controls (204). The levels of iron in the plasma, liver, spleen, and brain of PrP^{-/-} mice are significantly lower than those in wild-type controls, and neuronal TfR levels are upregulated, demonstrating a state of neuronal iron deficiency. It is noteworthy that the absence of PrP^C in PrP^{-/-} mice decreases the transport of iron from the intestinal epithelium to the bloodstream and hampers subsequent uptake by cells of all major organs, including hematopoietic progenitor cells (204). Analysis of other hematologic parameters shows minimal differences in the number of red cells and hematocrit, but a significant increase in reticulocytes in the peripheral blood of PrP^{-/-} mice relative to wild-type controls. Likewise, a proliferation of red cell precursors occurs in the bone marrow of PrP^{-/-} animals, indicating an attempt by the iron homeostatic machinery to compensate for the iron deficiency in these animals. The iron-deficient phenotype of PrP^{-/-} mice is reversed by expressing wild-type PrP^C in the PrP^{-/-} background, reinforcing the idea that PrP^C plays a functional role in iron uptake and transport (204). Because the iron deficiency of PrP^{-/-} mice is largely compensated and the animals live normally except for specific deficiencies mainly restricted to the central nervous system (127, 187, 191, 222), we believe that PrP^C modulates the function of other iron-uptake proteins or is involved in a parallel pathway of iron uptake that compensates for its absence. Because distinct iron-modulating proteins are involved in mediating iron transport from the intestine and uptake by hematopoietic cells, the hypothesis that PrP^C acts at a point downstream from these pathways deserves further consideration. Further investigations are necessary to understand the underlying mechanism

of iron transport and the identity of iron-modulating proteins that interact with PrP^C to influence iron levels in different cell types.

C. The antioxidant activity of PrP

PrP^C is believed to protect cells against oxidative stress, thus functioning as an antioxidant. The underlying mechanism of this activity, however, has remained elusive. It has been reported that neurons devoid of PrP^C expression show increased sensitivity to superoxide anions (31, 188), hydrogen peroxide (240), manganese (46), and copper toxicity (30) compared with wild-type controls. In PrP^{-/-} mice that lack PrP expression, markers of oxidative stress are evident in the brain (244). Likewise, loss of PrP^C function due to conversion to the PrP^{Sc} form may account for the increase in protein damage and markers of oxidative stress observed in sCJD and scrapie-infected mouse brains (64, 245). Continued oxidative stress is likely to deplete glutathione, a potent free-radical scavenger in the brain, thus intensifying toxicity (57).

Conflicting reports on the mechanism of the antioxidant function of PrP^C suggest that it influences the activity of cytosolic Cu/Zn SOD, or functions as an SOD enzyme itself (32, 96, 190). It remains plausible, though, that PrP^C modulates the activity of Cu/Zn-SOD indirectly by functioning as a copper-uptake and -transport protein, as observed in cultured cells incubated with radioactive copper, in which PrP^C modulates copper incorporation into the active site of this enzyme (32). Other studies suggest that association of PrP^C with copper in the octapeptide repeat region induces a conformational change in the C-terminal region that is considered important

for its antioxidant property (27, 95, 146). Furthermore, PrP^C is believed to quench extracellular free radicals, an activity that may account for its function as an antioxidant (154).

Collectively, these observations suggest that PrP^C protects cells against oxidative stress, although the mechanisms sustaining this action are still elusive.

IV. Redox Homeostasis and Prion-Disease Pathogenesis

A. PrP and metal interaction: the ionic connection

In addition to serving as physiologic ligands for PrP^C, copper and iron have significant pathologic implications because of their redox-active nature (150, 158). Recent studies demonstrate that copper-bound PrP^C is capable of accepting and donating electrons cyclically, although the significance of these observations is not clear (196). Likewise, the influence of the PrP^C-copper interaction on the generation of PrP^{Sc} is controversial. In some experimental paradigms, copper induces PrP^{Sc} formation, whereas in others, it has the opposite effect. For example, exposure of purified PrP^C to copper induces its conversion to a form similar to PrP^{Sc} (177), whereas addition of copper to synthetic prions slows the formation of amyloid, indicating an inhibitory role (21, 159). When added to scrapie-infected cells, copper reduces the accumulation of PrP^{Sc} (91), whereas in scrapie-infected mouse brains, copper is associated with PrP^{Sc} deposits, and chelation of copper delays the onset of prion disease (201). In addition, copper induces the interconversion of PrP^{Sc} strains *in vitro* from clinically distinct subtypes of CJD, most likely by altering their metal-ion occupancy and secondary structure (230). Furthermore, humans carrying an expansion of the copper-binding octapeptide repeat region are more prone to a familial form of CJD, whereas deletion of these repeats has no effect (62, 211), suggesting a direct role of copper levels with prion-disease pathogenesis.

Some of the contradictory results pertaining to beneficial or harmful effects of copper noted earlier can be explained by observations from *in vitro* studies in which copper inhibits amplification of PrP^{Sc} from purified brain-derived and recombinant PrP^C by stabilizing the structure of the latter (93, 143). At the same time, copper is known to enhance the β -sheet structure when added to preformed PrP fibrils, increasing their overall PrP^{Sc} content (21). Thus, copper could delay or augment disease pathogenesis, based on the time within the incubation period when it is introduced. Alternately, these disparate findings could be explained by the redox-active nature of copper and the metal content of PrP^C. Purified brain-derived and recombinant PrP^C are associated with little copper or iron and are not likely to generate free radicals in response to added copper. Preformed fibrils, conversely, are associated with both copper and iron and are likely to aggregate further when exposed to added copper because of the initiation of the Fenton reaction (124).

Cell-associated PrP^C responds to redox-active metals in a similar manner. Exposure of cells expressing PrP^C to a source of redox-iron causes aggregation of PrP^C in association with cellular ferritin, and the complex mimics PrP^{Sc} in several biochemical characteristics (12). It is likely that interaction of metal-associated PrP^C on the cell surface with a source of redox-iron initiates the Fenton reaction, resulting in its aggregation. These aggregates are diverted to lysosomes for

degradation where they come into contact with ferritin and assume a redox-active nature because of the associated iron and copper. The aggregated PrP-ferritin complexes initiate the generation of additional PrP-ferritin aggregates, propagating PrP^{Sc}-like conformation within the cells (12). Cells exposed to hemin, an iron-containing compound, show a similar aggregation, internalization, and degradation of PrP^C, demonstrating the sensitivity of PrP^C to free radicals and the significance of metal-induced oxidative stress in prion-disease pathogenesis (123). It is interesting to note that a marginal increase in the concentration of redox-active copper or iron in the culture medium of PrP^C-expressing cells results in accumulation of PrP^{Sc}-like aggregates with the redox activity of the metal preserved, suggesting that PrP co-aggregates with the metal and becomes redox active (12, 118, 154). A similar aggregation of α -synuclein is noted when it is exposed to ferrous chloride *in vitro* or when expressed in cells (69, 80, 160), suggesting that protein aggregation by redox-active metals is not specific to PrP^C. A partial list of neurodegenerative conditions associated with metal-ion-induced protein aggregation is provided in Table 3.

The association of PrP^{Sc} with ferritin was also observed in sCJD-affected brain homogenates and mouse scrapie-infected cell lines ScN2a and SMB (144, 203). Interestingly, ferritin is not degraded by exposure to proteinase-K, and the interaction of ferritin with PrP^{Sc} is maintained in the presence of digestive enzymes (144, 203), raising the possibility that PrP^{Sc} is co-internalized by intestinal epithelial cells along with ferritin. *In vitro* experiments with the epithelial cell line Caco-2 suggest that the PrP^{Sc}-ferritin complex is indeed transported intact across a monolayer of these cells, suggesting an important role for ferritin in this process (144). Whether the interaction of PrP^{Sc} with ferritin is essential for its PK-resistant nature is not entirely clear. However, chelation of iron from prion disease-affected brain homogenates decreases the total amount of PK-resistant PrP^{Sc}, suggesting that iron is somehow involved in the stability of PrP^{Sc} (12).

A similar role for iron has been reported in scrapie-infected mouse neuroblastoma cells (ScN2a) that actively replicate mouse prions in culture. These cells show alteration of cellular iron homeostasis and increased susceptibility to iron-induced oxidative stress (60, 61). The total iron content and calcein chelatable iron pool in ScN2a cells is twofold lower, and unexpectedly, the activity of iron regulatory proteins IRE1 and 2 also is lower by 40% and 50%, indicating mismanagement of cellular iron homeostasis (61). Moreover, the levels of redox-active Fe²⁺ iron are higher in ScN2a cells, increasing their susceptibility to free radical-induced toxicity (61). Exposure of ScN2a cells to exogenous iron results in increased production of ROS, further supporting the idea that accumulation of PrP^{Sc} causes mismanagement of cellular iron homeostasis

TABLE 3. NEURODEGENERATIVE DISEASES INVOLVING METAL-INDUCED PROTEIN AGGREGATION

Protein involved	Metal	Disease
PrP ^C or PrP ^{Sc}	Cu, Fe, Mn	Prion disorders
A β	Cu, Fe, Zn	Alzheimer's disease
SOD1	Cu	ALS
α -Synuclein	Fe, Cu	Parkinson's disease




FIG. 12. A model of possible mechanisms of iron-mediated neurotoxicity in prion disorders. Free radicals generated from normal metabolic processes in the brain are normally detoxified by antioxidant enzymes and proteins, including PrP^C. When the antioxidant capability of the brain is exceeded, free radicals are likely to react with PrP^C-bound copper and iron and induce their aggregation to the PrP^{Sc} form that co-aggregates with ferritin and becomes redox active (12, 203). This causes further production of free radicals, resulting in a state of brain-iron imbalance. This increases the production of ROS, resulting in further aggregation of PrP^C and neurotoxicity.

(60). GT-1 cells infected with scrapie show a similar response to exogenous iron, ruling out artifactual effects due to clonal selection of either cell line (60).

These studies suggest that the association of PrP^C with copper and iron can have deleterious consequences under certain circumstances because of their redox-active nature. Perhaps the site and nature of PrP-metal interaction and the structure of PrP^C itself are important underlying factors in this process, because the formation of β -sheet-rich aggregates on exposure to free radicals has been reported only for a few proteins such as PrP^C and α -synuclein (215). Other major iron- and copper-binding proteins do not show this response. It is likely that sequestration of iron in PrP^{Sc} aggregates renders them redox active, thus accentuating the associated toxicity. Future investigations are necessary to understand this phenomenon fully.

B. Prion disorders and brain-iron homeostasis

The contribution of redox-active PrP^{Sc} aggregates and altered brain-iron homeostasis to prion-disease pathogenesis is further supported by observations in diseased human and animal brains. In human brains affected with vCJD, redox-active iron is detected in association with PrP^{Sc} plaques, along with iron- and ferritin-rich microglia in the region surrounding the plaque (171). Although the underlying mechanism leading to iron deposits is not clear, redox-active iron is likely to induce oxidative damage in the surrounding neuronal population (99, 132). Other reports indicate alteration of the activity of iron-regulatory proteins 1 and 2 and expression of iron-storage protein ferritin in the hippocampus and cerebral cortex of prion-infected mouse brains, suggesting a state of iron imbalance (107). This assumption is supported by ob-

servations indicating a consistent increase in total and redox-active Fe²⁺ iron, and paradoxically, a phenotype of iron deficiency in sCJD-affected brain tissue (203). As a consequence, major iron-uptake proteins Tf and TfR are upregulated to compensate for the deficiency, worsening the state of iron imbalance. The deficiency of iron is evident in Purkinje cells of sCJD brains, suggesting that the neuronal population is especially affected by this metabolic alteration. This phenotype of brain-iron deficiency develops during the incubation period and shows a direct correlation with PrP^{Sc} levels, supporting the hypothesis that PrP^{Sc} itself or PrP^{Sc}-ferritin aggregates sequester iron in a biologically unavailable form, resulting in a state of iron deficiency in the presence of excess iron (203). Because PrP^{Sc}-ferritin aggregates are themselves redox active, it is likely that once initiated, this complex propagates PrP^{Sc} accumulation and iron deficiency, worsening the state of iron dyshomeostasis and associated neurotoxicity. Figure 12 summarizes the possible mechanisms of iron-induced neurotoxicity in prion disease-affected brains.

Although studies supporting imbalance of iron homeostasis as a significant contributing factor of prion-disease pathogenesis are limited, the observations noted here provide a viable means of developing a prophylactic and therapeutic strategy, both of which are lacking at this time. Equally strong evidence supporting other mechanisms of prion replication and disease-associated neurotoxicity exists, but has not been included in this review in an attempt to keep the focus limited to the redox activity of prion-disease pathogenesis.

V. Therapeutic Options

Investigations directed at possible therapeutic options for prion disorders are still at a formative stage and demand a

clear objective and in-depth evaluation to design and test successfully possible therapies. Because all prion disorders except vCJD have a relatively long incubation period, effective treatment can be initiated much before clinical symptoms appear. Given the focus of this review, two possible avenues hold promise: (a) antioxidants and free radical scavengers (63, 139, 214), and (b) metal chelators (38, 65, 248). Limited studies conducted in this direction have been quite promising and are reviewed here.

A. Antioxidants

Studies on prion-infected cell and mouse models have provided useful information on the therapeutic potential of antioxidants. In cell models, both direct application of cell-permeable antioxidants and indirect methods to restore endogenous antioxidant levels have been tried (24). Flupirtine, a triaminopyridine compound, has become quite popular among researchers because it can act as an *N*-methyl-D-aspartate (NMDA) antagonist without binding to NMDA receptors. This drug has the exceptional ability to normalize intracellular glutathione levels and restore oxidative balance within the cell, thereby combating accumulation of ROS and other free radicals. The associated upregulation of anti-apoptotic protein Bcl-2 and the relatively favorable pharmacokinetic profile of flupirtine make it a promising therapeutic agent to treat prion disorders (151, 161, 168, 194). An equally promising agent is the nonpsychoactive cannabis constituent cannabidiol, by virtue of its antioxidant property, NMDA antagonism, reduction in glutamate release, and blockade of microglial migration and activation, all of which are detrimental factors that aggravate PrP^{Sc}-mediated neurotoxicity (55, 138). Similarly, the disaccharide trehalose, known for its ability to reduce A β -mediated toxicity by inhibiting its aggregation, also protects prion-infected cells from oxidative damage (18). Another effective agent is EUK-189, a potent Mn SOD/catalase mimetic, that reduces oxidative damage in prion-infected mouse models, as evidenced by reduction in nitritative damage to vital cellular proteins, prolongation of incubation time, and decreased spongiform change in the brains of terminally ill mice (24). Although a clinically viable antioxidant that can alleviate prion disease-associated neurotoxicity is lacking, these observations argue that counteracting oxidative stress may have therapeutic benefit in prion disease and provide the basis for future investigations in this area.

B. Metal chelators

Most of the strategies aimed at metal chelation are targeted toward copper because the association of iron with prion-disease pathogenesis is a relatively new observation (203). Contradictory observations have been reported for copper, in which both increased and reduced levels of brain copper have been implicated in disease-associated neurotoxicity. In an experimental paradigm in which the loss of neurons and astrogliosis was induced by introduction of copper into the dorsal hippocampus of rats, co-injection of a synthetic peptide corresponding to the octapeptide repeat domain of PrP (PrP⁵⁹⁻⁹¹) that binds copper reduced neuronal death (42). With a similar premise, chelation of copper with D-penicillamine, a drug used routinely for treating Wilson disease, decreased brain-copper content of prion-infected mice by 30% and in-

creased the incubation period, supporting the idea that increased levels of brain copper promote diseases (58, 201). However, contradictory reports suggest a protective role for copper in prion disorders. It was observed that neuroblastoma cells cultured in the presence of copper ions lost the ability to bind and internalize PrP^{Sc}, thereby evading infection and toxicity. A significant delay in the onset of clinical disease also was observed in scrapie-infected hamsters given a dietary supplement of copper (91), supporting these observations. It is likely that the protective effect of copper reflects internalization and degradation of PrP^C on exposure to copper, the substrate for PrP^{Sc} generation, although a direct effect on the generation of PrP^{Sc} cannot be ruled out because inhibition of PrP^{Sc} accumulation is observed after the addition of copper *in vitro* to PMCA reactions, a procedure used to amplify PrP^{Sc} (39, 159).

The involvement of redox iron and imbalance of brain-iron homeostasis in prion disease-associated neurotoxicity is a relatively new observation, and the effect of iron chelation on disease pathogenesis has not been tried in cell or mouse models. However, chelation of iron from prion disease-affected human and mouse brain homogenates *in vitro* reduces the amount of disease-associated PrP^{Sc}, suggesting that this method may be used prophylactically to decrease prion infectivity in consumable products (12). A similar reduction of PrP^{Sc} levels *in vivo* may prove useful in decreasing PrP^{Sc} load, although optimal iron chelators that are nontoxic at therapeutic doses and can cross the blood-brain barrier effectively have not been developed. Studies in MPTP mouse models of Parkinson disease report significant benefit from the concomitant administration of blood-brain barrier-permeable iron chelator VK-28 [5-(4-(2-hydroxyethyl) piperazin-1-yl) (methyl)-8-hydroxyquinoline] and its derivative M30 [5-(*N*-methyl-*N*-propargylaminomethyl)-8-hydroxyquinoline], providing direct evidence for the involvement of iron in disease pathogenesis (253). However, the applicability of these compounds in prion disease-associated neurotoxicity is yet to be investigated. Although apparently encouraging, the reduction of brain iron may aggravate the disease by increasing iron uptake by surviving cells, warranting caution in using such compounds. Restoring brain-iron homeostasis in diseased brains is therefore a daunting task, because complex biochemical pathways are involved in iron metabolism.

However, iron chelation as a means to reduce the toxicity associated with its redox-active nature has been pursued actively in diseases like AD and PD. Several chelators have been tried, the most prominent ones being desferrioxamine (DFO) and 5-chloro-7-iodo-quinolin-8-ol (Clioquinol) (90). Although DFO showed some success in studies with AD patients, it does not cross the blood-brain barrier effectively and is toxic in therapeutic doses, making it an unsuitable drug for the treatment of AD or prion disorders. Clioquinol is an antibiotic that binds to Zn, Cu, and iron, and crosses the blood-brain barrier effectively (90). Structurally, Clioquinol is related to quinacrine analogues that have been used effectively in other studies on prion disease-affected experimental models and in human trials and could be a safe drug for *in vivo* use (152). The use of Clioquinol in scrapie-infected hamsters increases the incubation time modestly, suggesting a future potential for the use of this drug in humans (175). Likewise, promising results were observed when Clioquinol was administered orally to mouse models of AD and PD (38, 105).

The use of antioxidants and metal ion chelators are the major approach toward developing therapies for prion disorders; however, most of the drugs are effective when administered at a very early stage of the disease. The imposed problem is that prion disorders in most cases go unnoticed at early stages and can be diagnosed much later when most of the damage and neurodegeneration have occurred. Hence, proper treatment of prion disorders also demands a better diagnostic tool to allow detection at an early stage that can complement the use of reported drugs with greater efficiency.

VI. Summary and Perspective

The complexity and multiplicity of factors involved in the pathogenesis of prion disorders has hampered our progress toward the development of a therapeutic strategy, although each report brings us closer to an answer. Collective evidence from different models suggests a key role for PrP^{Sc}, a β -sheet-rich form of the cell-surface glycoprotein PrP^C, as the principal neurotoxic element. Equally strong data support loss of protective function of PrP^C as a significant contributing factor in disease-associated neuronal death. Several mechanisms of neurotoxicity have been suggested for PrP^{Sc}, and an equally diverse set of functions has been proposed for PrP^C, the loss of which could induce neuronal death. Consensus on either of these issues is still lacking. Recent reports on the association of PrP^C with redox-active metals such as copper and iron provide a new perspective on the mechanisms underlying prion disease-associated neurotoxicity. Human PrP^C binds to five copper ions with a relatively high affinity and mediates the uptake of copper ions from the extracellular milieu. The octapeptide repeat region of PrP^C also functions as a reductase, facilitating the transport of copper from the endosomes to cytosolic carrier proteins. The binding site and affinity of PrP^C for iron is less well defined, although it is clear that PrP^C mediates the uptake and transport of iron, and lack of PrP^C induces a phenotype of iron deficiency in PrP^{-/-} mice. Thus, the association of PrP^C with copper and iron has significant physiologic implications. Conversely, the inherently redox-active nature of these metals increases the susceptibility of PrP^C to free radicals that causes its aggregation to a form resembling PrP^{Sc} in several characteristics. The PrP^{Sc} thus formed co-aggregates with ferritin and associated iron, rendering it redox active. This complex induces the aggregation of additional PrP^C to PrP^{Sc}, and the process continues. The lack of bioavailability of sequestered iron results in a state of iron deficiency, resulting in brain-iron imbalance and associated neurotoxicity. These observations suggest a significant role for metal-induced oxidative stress in prion disease-associated neurotoxicity, and the prospect of using antioxidants and iron chelators such as Clotrimazole that can cross the blood-brain barrier effectively as useful therapeutic agents. However, given the complex nature of brain-iron metabolism, chelation *per se* may provide only limited help or even worsen the situation. Future investigations are necessary to resolve this issue, keeping in mind other aspects of prion-disease pathogenesis that are not discussed here due to the specific focus of this review.

Acknowledgment

N.S. is supported by grants from the National Institutes of Health.

References

1. Abid K and Soto C. The intriguing prion disorders. *Cell Mol Life Sci* 63: 2342–2351, 2006.
2. Aguzzi A and Polymenidou M. Mammalian prion biology: one century of evolving concepts. *Cell* 116: 313–327, 2004.
3. Aguzzi A, Sigurdson C, and Heikenwaelder M. Molecular mechanisms of prion pathogenesis. *Annu Rev Pathol Mech Dis* 3: 11–40, 2008.
4. Allsop D, Mayes J, Moore S, Masad A, and Tabner BJ. Metal-dependent generation of reactive oxygen species from amyloid proteins implicated in neurodegenerative disease. *Biochem Soc Trans* 36: 1293–1298, 2008.
5. Andreoletti O, Levavasseur E, Uro-Coste E, Tabouret G, Sarradin P, Delisle MB, Berthon P, Salvayre R, Schelcher F, and Negre-Salvayre A. Astrocytes accumulate 4-hydroxynonenal adducts in murine scrapie and human Creutzfeldt-Jakob disease. *Neurobiol Dis* 11: 386–393, 2002.
6. Arlt S, Kontush A, Zerr I, Buhmann C, Jacobi C, Schröter A, Poser S, and Beisiegel U. Increased lipid peroxidation in cerebrospinal fluid and plasma from patients with Creutzfeldt-Jakob disease. *Neurobiol Dis* 10: 150–156, 2002.
7. Aronoff-Spencer E, Burns CS, Avdievich NI, Gerfen GJ, Peisach J, Antholine WE, Ball HL, Cohen FE, Prusiner SB, and Millhauser GL. Identification of the Cu²⁺ binding sites in the N-terminal domain of the prion protein by EPR and CD spectroscopy. *Biochemistry* 39: 13760–13771, 2000.
8. Atarashi R, Moore RA, Sim VL, Hughson AG, Downward DW, Onwubiko HA, Priola SA, and Caughey B. Ultra-sensitive detection of scrapie prion protein using seeded conversion of recombinant prion protein. *Nat Methods* 4: 645–650, 2007.
9. Barnham KJ and Bush AI. Metals in Alzheimer's and Parkinson's diseases. *Curr Opin Chem Biol* 12: 222–228, 2008.
10. Barnham KJ, Cappai R, Beyreuther K, Masters CL, and Hill AF. Delineating common molecular mechanisms in Alzheimer's and prion diseases. *Trends Biochem Sci* 31: 465–472, 2006.
11. Bartzokis G. In vivo evaluation of brain iron in Alzheimer's disease and normal subjects using MRI. *Biol Psychiatry* 35: 480–487, 1994.
12. Basu S, Mohan ML, Luo X, Kundu B, Kong Q, and Singh N. Modulation of PK-resistant PrP^{Sc} in cells and infectious brain homogenate by redox-iron: implications for prion replication and disease pathogenesis. *Mol Biol Cell* 18: 3302–3312, 2007.
13. Bayer TA, Schäfer S, Breyhan H, Wirths O, Treiber C, and Multhaup G. A vicious circle: role of oxidative stress, intraneuronal A β and Cu in Alzheimer's disease. *Clin Neuropathol* 25: 163–171, 2006.
14. Beard JL and Connor JR. Iron status and neural functioning. *Annu Rev Nutr* 23: 41–58, 2003.
15. Beard JL. Why iron deficiency is important in infant development. *J Nutr* 138: 2534–2536, 2008.
16. Behl C, Davis JB, Lesley R, and Schubert D. Hydrogen peroxide mediates amyloid beta protein toxicity. *Cell* 77: 817–827, 1994.
17. Bellingham SA, Coleman LA, Masters CL, Camakaris J, and Hill AF. Regulation of prion gene expression by transcription factors SP1 and metal transcription factor-1. *J Biol Chem* 284: 1291–1301, 2009.
18. Beranger F, Crozet C, Goldsborough A, and Lehman S. Trehalose impairs aggregation of PrP^{Sc} molecules and protects prion-infected cells against oxidative damage. *Biochem Biophys Res Commun* 374: 44–48, 2008.

19. Bieschke J, Weber P, Sarafoff N, Beekes M, Giese A, and Kretschmar H. Autocatalytic self-propagation of misfolded prion protein. *Proc Natl Acad Sci U S A* 101: 12207–12211, 2004.
20. Bio KF, Dal Pizzol F, Conte da Frota ML, Walz R, Andrades ME, da Silva EG, Brentani RR, Izquierdo I, and Fonseca Moreira JC. Imbalance of antioxidant defense in mice lacking cellular prion protein. *Free Radic Biol Med* 30: 1137–1144, 2001.
21. Bocharova OV, Breydo L, Salnikov VV, and Baskakov IV. Copper(II) inhibits in vitro conversion of prion protein into amyloid fibrils. *Biochemistry* 44: 6776–6787, 2005.
22. Bounhar Y, Zhang Y, Goodyer CG, and LeBlanc A. Prion protein protects human neurons against Bax-mediated apoptosis. *J Biol Chem* 276: 39145–39149, 2001.
23. Brazier MW, Davies P, Player E, Marken F, Viles JH, and Brown DR. Manganese binding to the prion protein. *J Biol Chem* 283: 12831–12839, 2008.
24. Brazier MW, Doctrow SR, Masters CL, and Collins SJ. A manganese-superoxide dismutase/catalase mimetic extends survival in a mouse model of human prion disease. *Free Radic Biol Med* 45: 184–192, 2008.
25. Brazier MW, Lewis V, Cicotosto GD, Klug GM, Lawson VA, Cappai R, Ironside JW, Masters CL, Hill AF, White AR, and Collins S. Correlative studies support lipid peroxidation is linked to PrP(res) propagation as an early primary pathogenic event in prion disease. *Brain Res Bull* 68: 346–354, 2006.
26. Brown DR and Besinger A. Prion protein expression and superoxide dismutase activity. *Biochem J* 334: 423–429, 1998.
27. Brown DR, Clive C, and Haswell SJ. Antioxidant activity related to copper binding of native prion protein. *J Neurochem* 76: 69–76, 2001.
28. Brown DR, Hafiz F, Glassmith LL, Wong BS, Jones IM, Clive C, and Haswell SJ. Consequences of manganese replacement of copper for prion protein function and proteinase resistance. *EMBO J* 19: 1180–1186, 2000.
29. Brown DR, Qin K, Herms JW, Madlung A, Manson J, Strome R, Fraser PE, Kruck T, von Bohlen A, Schulz-Schaeffer W, Giese A, Westaway D, and Kretschmar H. The cellular prion protein binds copper in vivo. *Nature* 390: 684–687, 1997.
30. Brown DR, Schmidt B, and Kretschmar HA. Effects of copper on survival of prion protein knockout neurons and glia. *J Neurochem* 70: 1686–1693, 1998.
31. Brown DR, Schultz-Schaeffer WJ, Schmidt B, and Kretschmar HA. Prion protein deficient cells show altered response to oxidative stress due to decreased SOD-1 activity. *Exp Neurol* 146: 104–112, 1997.
32. Brown DR, Wong BS, Hafiz F, Clive C, Haswell SJ, and Jones M. Normal prion protein has an activity like that of super-oxide dismutase. *Biochem J* 344: 1–5, 1999.
33. Brown DR. Prion protein expression aids cellular uptake and veratridine-induced release of copper. *J Neurosci Res* 58: 717–725, 1999.
34. Brown DR. PrP^{Sc}-like prion protein peptide inhibits the function of cellular prion protein. *Biochem J* 352: 511–518, 2000.
35. Brown LR and Harris DA. Copper and zinc cause delivery of the prion protein from the plasma membrane to a subset of early endosomes and the Golgi. *J Neurochem* 87: 353–363, 2003.
36. Bruce ME, Will RG, Ironside JW, McConnell I, Drummond D, Suttie A, McCardle L, Chree A, Hope J, Birkett C, Cousens S, Fraser H, and Bostock CJ. Transmissions to mice indicate that “new variant” CJD is caused by the BSE agent. *Nature* 389: 498–501, 1997.
37. Büeler H, Aguzzi A, Sailer A, Greiner RA, Autenried P, Auget M, and Weissmann C. Mice devoid of PrP are resistant to scrapie. *Cell* 73: 1339–1347, 1993.
38. Bush AI and Tanzi RE. Therapeutics for Alzheimer’s disease based on the metal hypothesis. *Neurotherapeutics* 5: 421–432, 2008.
39. Castilla J, Saá P, Hetz C, and Soto C. In vitro generation of infectious scrapie prions. *Cell* 121: 195–206, 2005.
40. Caughey B and Baron GS. Prions and their partners in crime. *Nature* 443: 803–810, 2006.
41. Caughey B, Race RE, Ernst D, Buchmeier MJ, and Chesebro B. Prion protein biosynthesis in scrapie-infected and uninfected neuroblastoma cells. *J Virol* 63: 175–181, 1989.
42. Chacón MA, Barriá MI, Lorca R, Huidobro-Toro JP, and Inestrosa NC. A human prion protein peptide [PrP(59–91)] protects against copper neurotoxicity. *Mol Psychiatry* 8: 853–862, 2003.
43. Chesebro B, Trifilo M, Race R, Meade-White K, Teng C, LaCasse R, Raymond L, Favara C, Baron G, Priola S, Caughey B, Masliah E, and Oldstone M. Anchorless prion protein results in infectious amyloid disease without clinical scrapie. *Science* 308: 1435–1439, 2005.
44. Chiesa R and Harris DA. Prion diseases: what is the neurotoxic molecule? *Neurobiol Dis* 8: 743–763, 2001.
45. Chinta SJ and Andersen JK. Redox imbalance in Parkinson’s disease. *Biochim Biophys Acta* 1780: 1362–1367, 2008.
46. Choi CJ, Anantharam V, Saetveit NJ, Houk RS, Kanthasamy A, and Kanthasamy AG. Normal cellular prion protein protects against manganese-induced oxidative stress and apoptotic cell death. *Toxicol Sci* 98: 495–509, 2007.
47. Choi YG, Kim JI, Lee HP, Jin JK, Choi EK, Carp RI, and Kim YS. Induction of heme oxygenase-1 in the brains of scrapie-infected mice. *Neurosci Lett* 289: 173–176, 2000.
48. Choi CJ, Kanthasamy A, Anantharam V, Kanthasamy AG. Interaction of metals with prion protein: possible role of divalent cations in the pathogenesis of prion diseases. *Neurotoxicology* 27: 777–787, 2006.
49. Cohen FE, Pan KM, Huang Z, Baldwin M, Fletterick RJ, and Prusiner SB. Structural clues to prion replication. *Science* 264: 530–531, 1994.
50. Collinge J, Sidle KCL, Meads J, Ironside J, and Hill AF. Molecular analysis of prion strain variation and the aetiology of “new variant” CJD. *Nature* 383: 685–690, 1996.
51. Cook CI and Yu BP. Iron accumulation in aging: modulation by dietary restriction. *Mech Ageing Dev* 102: 1–13, 1998.
52. Davies P and Brown DR. The chemistry of copper binding to PrP: is there sufficient evidence to elucidate a role for copper in protein function? *Biochem J* 410: 237–244, 2008.
53. DeArmond SJ and Ironside JW. Neuropathology of prion diseases. In: *Prion biology and diseases*, edited by Prusiner SB. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 1999, pp. 585–652.
54. Deleault NR, Harris BT, Rees JR, and Supattapone S. Formation of native prions from minimal components in vitro. *Proc Natl Acad Sci U S A* 104: 9741–9746, 2007.
55. Dirikoc S, Priola SA, Marella M, Zsuzsger N, and Chabry J. Nonpsychoactive Cannabidiol prevents prion accumulation and protects neurons against prion toxicity. *J Neurosci* 27: 9537–9544, 2007.
56. Doherr MG. Brief review on the epidemiology of transmissible spongiform encephalopathies (TSE). *Vaccine* 25: 5619–5624, 2007.

57. Dukhande VV, Malthankar-Phatak GH, Hugus JJ, Daniels CK, and Lai JC. Manganese-induced neurotoxicity is differentially enhanced by glutathione depletion in astrocytoma and neuroblastoma cells. *Neurochem Res* 11: 1349–1357, 2006.
58. El-Youssef M. Wilson disease. *Mayo Clin Proc* 78: 1126–1136, 2003.
59. Fasano C, Campana V, and Zurzolo C. Prions: protein only or something more? Overview of potential prion cofactors. *J Mol Neurosci* 29: 195–214, 2006.
60. Fernaues S and Land T. Increased iron-induced oxidative stress and toxicity in scrapie-infected neuroblastoma cells. *Neurosci Lett* 382: 133–136, 2005.
61. Fernaues S, Halldin J, Bedecs K, and Land T. Changed iron regulation in scrapie-infected neuroblastoma cells. *Brain Res Mol Brain Res* 133: 266–273, 2005.
62. Flechsig E, Shmerling D, Hegyi I, Raeber AJ, Fischer M, Cozzio A, von Mering C, Aguzzi A, and Weissmann C. Prion protein devoid of the octapeptide repeat region restores susceptibility to scrapie in PrP knockout mice. *Neuron* 27: 399–408, 2000.
63. Floyd RA and Hensley K. Oxidative stress in brain aging: implications for therapeutics of neurodegenerative diseases. *Neurobiol Aging* 23: 795–807, 2002.
64. Freixes M, Rodriguez A, Dalfó E, and Ferrer I. Oxidation, glycoxidation, lipoxidation, nitration, and responses to oxidative stress in the cerebral cortex in Creutzfeldt-Jakob disease. *Neurobiol Aging* 27: 1807–1815, 2006.
65. Gaeta A and Hider RC. The crucial role of metal ions in neurodegeneration: the basis for a promising therapeutic strategy. *Br J Pharmacol* 146: 1041–1059, 2005.
66. Gains MJ and LeBlanc AC. Canadian Association of Neurosciences Review: prion protein and prion diseases: the good and the bad. *Can J Neurol Sci* 34: 126–145, 2007.
67. Gambetti P, Dong Z, Yuan J, Xiao X, Zheng M, Alshekhlee A, Castellani R, Cohen M, Barria MA, Gonzalez-Romero D, Belay ED, Schonberger LB, Marder K, Harris C, Burke JR, Montine T, Wisniewski T, Dickson DW, Soto C, Hulette CM, Mastrianni JA, Kong Q, and Zou WQ. A novel human disease with abnormal prion protein sensitive to protease. *Ann Neurol* 63: 697–708, 2008.
68. Gerlach M, Ben-Shachar D, Riederer P, and Youdim MB. Altered brain metabolism of iron as a cause of neurodegenerative diseases. *J Neurochem* 63: 793–807, 1994.
69. Goldberg MS and Lansbury PT Jr. Is there a cause-and-effect relationship between alpha-synuclein fibrillization and Parkinson's disease? *Nat Cell Biol* 2: E115–E119, 2000.
70. Gregory A and Hayflick SJ. Neurodegeneration with brain iron accumulation. *Folia Neuropathol* 43: 286–296, 2005.
71. Gu Y, Hinnerwisch J, Fredricks R, Kalepu S, Mishra RS, and Singh N. Identification of cryptic nuclear localization signals in the prion protein: implications for prion disease pathogenesis. *Neurobiol Dis* 12: 133–149, 2003.
72. Gu Y, Luo X, Basu S, Fujioka H, and Singh N. Cell-specific metabolism and pathogenesis of transmembrane prion protein. *Mol Cell Biol* 26: 2697–2715, 2006.
73. Gu Y, Singh A, Bose S, and Singh N. Pathogenic mutations in the glycosylphosphatidylinositol signal peptide of PrP modulate its membrane topology in neuroblastoma cells. *Mol Cell Neurosci* 37: 647–656, 2007.
74. Gu Y, Verghese S, Bose S, Mohan ML, and Singh N. Mutant prion protein D202N associated with familial prion disease forms “curly” intracellular aggregates and sensitizes cells to oxidative stress. *J Mol Neurosci* 32: 90–96, 2007.
75. Gu Y, Verghese S, Mishra RS, Xu X, Shi Y, and Singh N. Mutant prion protein-mediated aggregation of normal prion protein in the endoplasmic reticulum: implications for prion propagation and neurotoxicity. *J Neurochem* 84: 10–22, 2003.
76. Guentchev M, Siedlak SL, Jarius C, Tagliavini F, Castellani RJ, Perry G, Smith MA, and Budka H. Oxidative damage to nucleic acids in human prion disease. *Neurobiol Dis* 9: 275–281, 2002.
77. Guentchev M, Voigtlander T, Haberler C, Groschup MH, and Budka H. Evidence for oxidative stress in experimental prion disease. *Neurobiol Dis* 7: 270–273, 2000.
78. Gutteridge JM. The role of superoxide and hydroxyl radicals in phospholipid peroxidation catalysed by iron salts. *FEBS Lett* 150: 454–458, 1982.
79. Harris DA and True HL. New insights into prion structure and toxicity. *Neuron* 50: 353–357, 2006.
80. Hashimoto M, Hsu LJ, Xia Y, Takeda A, Sisk A, Sundsmo M, and Masliah E. Oxidative stress induces amyloid-like aggregate formation of NACP/alpha-synuclein in vitro. *Neuroreport* 10: 717–721, 1999.
81. Hegde RS and Rane NS. Prion protein trafficking and the development of neurodegeneration. *Trends Neurosci* 26: 337–339, 2003.
82. Hegde RS, Mastrianni JA, Scott MR, Defea KA, Tremblay P, Torchia M, DeArmond SJ, Prusiner SB, and Lingappa VR. A transmembrane form of the prion protein in neurodegenerative disease. *Science* 279: 827–834, 1998.
83. Hegde RS, Tremblay P, Groth D, DeArmond SJ, Prusiner SB, and Lingappa VR. Transmissible and genetic prion diseases share a common pathway of neurodegeneration. *Nature* 402: 822–826, 1999.
84. Heller U, Winklhofer KF, Heske J, Reintjes A, and Tatzelt J. Post-translational import of the prion protein into the endoplasmic reticulum interferes with cell viability: a critical role for the putative transmembrane domain. *J Biol Chem* 278: 36139–36147, 2003.
85. Herms J, Tings T, Gall S, Madlung A, Giese A, Siebert H, Schürmann P, Windl O, Brose N, and Kretzschmar H. Evidence of presynaptic location and function of the prion protein. *J Neurosci* 19: 8866–8875, 1999.
86. Hesketh S, Sassoon J, Knight R, and Brown DR. Elevated manganese levels in blood and CNS in human prion disease. *Mol Cell Neurosci* 37: 590–598, 2008.
87. Hesketh S, Sassoon J, Knight R, Hopkins J, and Brown DR. Elevated manganese levels in blood and central nervous system occur before onset of clinical signs in scrapie and bovine spongiform encephalopathy. *J Anim Sci* 85: 1596–1609, 2007.
88. Hetz C, Castilla J, and Soto C. Perturbation of endoplasmic reticulum homeostasis facilitates prion replication. *J Biol Chem* 282: 12725–12733, 2007.
89. Hetz CA and Soto C. Stressing out the ER: a role of the unfolded protein response in prion-related disorders. *Curr Mol Med* 6: 37–43, 2006.
90. Hider RC, Ma Y, Molina-Holgado F, Gaeta A, and Roy S. Iron chelation as a potential therapy for neurodegenerative disease. *Biochem Soc Trans* 36: 1304–1308, 2008.
91. Hijazi N, Shaked Y, Rosenmann H, Ben-Hur T, and Gabizon R. Copper binding to PrP^C may inhibit prion disease propagation. *Brain Res* 12: 192–200, 2003.
92. Hill AF, Joiner S, Linehan J, Desbruslais M, Lantos PL, and Collinge J. Species-barrier-independent prion replication in apparently resistant species. *Proc Natl Acad Sci U S A* 97: 10248–10253, 2000.

93. Hodak M, Chisnell R, Lu W, and Bernholc J. Functional implications of multistage copper binding to the prion protein. *Proc Natl Acad Sci U S A* 106: 11576–11581, 2009.
94. Hornshaw MP, McDermott JR, and Candy JM. Copper binding to the N-terminal tandem repeat regions of mammalian and avian prion protein. *Biochem Biophys Res Commun* 207: 621–629, 1995.
95. Hornshaw MP, McDermott JR, Candy JM, and Lakey JH. Copper binding to the N-terminal tandem repeat region of mammalian and avian prion protein: structural studies using synthetic peptides. *Biochem Biophys Res Commun* 214: 993–999, 1995.
96. Hutter G, Heppner FL, and Aguzzi A. No superoxide dismutase activity of cellular prion protein in vivo. *Biol Chem* 384: 1279–1285, 2003.
97. Jackson GS, Murray I, Hosszu LL, Gibbs N, Waltho JP, Clarke AR, and Collinge J. Location and properties of metal-binding sites on the human prion protein. *Proc Natl Acad Sci U S A* 98: 8531–8535, 2001.
98. Jeffrey M, Goodsir CM, Race RE, and Chesebro B. Scrapie-specific neuronal lesions are independent of neuronal PrP expression. *Ann Neurol* 55: 781–792, 2004.
99. Jenner P. Oxidative damage in neurodegenerative disease. *Lancet* 344: 796–798, 1994.
100. Jeong SY and David S. Glycosylphosphatidylinositol-anchored ceruloplasmin is required for iron efflux from cells in the central nervous system. *J Biol Chem* 278: 27144–27148, 2003.
101. Jin T, Gu Y, Zanusso G, Sy M, Kumar A, Cohen M, Gambetti P, and Singh N. The chaperone protein BiP binds to a mutant prion protein and mediates its degradation by the proteasome. *J Biol Chem* 275: 38699–38704, 2000.
102. Jobling MF, Huang X, Stewart LR, Barnham KJ, Curtain C, Volitakis I, Perugini M, White AR, Cherny RA, Masters CL, Barrow CJ, Collins SJ, Bush AI, and Cappai R. Copper and zinc binding modulates the aggregation and neurotoxic properties of the prion peptide PrP106–126. *Biochemistry* 40: 8073–8084, 2001.
103. Jones CE, Abdelraheim SR, Brown DR, and Viles JH. Preferential Cu²⁺ coordination by His96 and His111 induces beta-sheet formation in the unstructured amyloidogenic region of the prion protein. *J Biol Chem* 279: 32018–32027, 2004.
104. Kaplan J. Mechanisms of cellular iron acquisition: another iron in the fire. *Cell* 111: 603–606, 2002.
105. Kaur D, Yantiri F, Rajagopalan S, Kumar J, Mo JQ, Boonplueang R, Viswanath V, Jacobs R, Yang L, Beal MF, DiMonte D, Volitakis I, Ellerby L, Cherny RA, Bush AI, and Andersen JK. Genetic or pharmacological iron chelation prevents MPTP-induced neurotoxicity in vivo: a novel therapy for Parkinson's disease. *Neuron* 37: 889–890, 2003.
106. Kawano T. Prion-derived copper-binding peptide fragments catalyze the generation of superoxide anion in the presence of aromatic monoamines. *Int J Biol Sci* 3: 57–63, 2007.
107. Kim BH, Jun YC, Jin JK, Kim JI, Kim NH, Leibold EA, Connor JR, Choi EK, Carp RI, and Kim YS. Alteration of iron regulatory proteins (IRP1 and IRP2) and ferritin in the brains of scrapie-infected mice. *Neurosci Lett* 422: 158–163, 2007.
108. Kim JI, Ju WK, Choi JH, Choi E, Carp RI, Wisniewski HM, and Kim YS. Expression of cytokine genes and increased nuclear factor-kappa B activity in the brains of scrapie-infected mice. *Brain Res Mol Brain Res* 73: 17–27, 1999.
109. Kim JL, Choi SI, Kim NH, Jin JK, Choi EK, Carp RI, and Kim YS. Oxidative stress and neurodegeneration in prion diseases. *Ann N Y Acad Sci* 928: 182–186, 2001.
110. Kim NH, Choi JK, Jeong BH, Kim JI, Kwon MS, Carp RI, and Kim YS. Effect of transition metals (Mn, Cu, Fe) and deoxycholic acid (DA) on the conversion of PrP^C to PrP^{res}. *FASEB J* 19: 783–785, 2005.
111. Kim NH, Park SJ, Jin JK, Kwon MS, Choi EK, Carp RI, and Kim YS. Increased ferric iron content and iron-induced oxidative stress in the brains of scrapie-infected mice. *Brain Res* 884: 98–103, 2000.
112. Klamt F, Dal-Pizzol F, Conte da Frota ML, Walz R, Andrades ME, da Silva EG, Brentani RR, Izquierdo I, and Fonseca Moreira JC. Imbalance of antioxidant defense in mice lacking cellular prion protein. *Free Radic Biol Med* 30: 1137–1144, 2001.
113. Klewpatinond M, Davies P, Bowen S, Brown DR, and Viles JH. Deconvoluting the Cu²⁺ binding modes of full-length prion protein. *J Biol Chem* 283: 1870–1881, 2008.
114. Kordek R, Nerurkar VR, Liberski PP, Isaacson S, Yanagihara R, and Gajdusek DC. Heightened expression of tumor necrosis factor alpha, interleukin 1 alpha, and glial fibrillary acidic protein in experimental Creutzfeldt-Jakob disease in mice. *Proc Natl Acad Sci U S A* 93: 9754–9758, 1996.
115. Kovacs GG and Budka H. Prion diseases: from protein to cell pathology. *Am J Pathol* 172: 555–565, 2008.
116. Krasemann S, Zerr I, Weber T, Poser S, Kretzschmar H, Hunsmann G, and Bodemer W. Prion disease associated with a novel nine octapeptide repeat insertion in the PRNP gene. *Mol Brain Res* 34: 173–176, 1995.
117. Kretzschmar HA, Tings T, Madlung A, Giese A, and Herms J. Function of PrP(C) as a copper-binding protein at the synapse. *Arch Virol Suppl* 16: 239–249, 2000.
118. Kuczius T, Buschmann A, Zhang W, Karch H, Becker K, Peters G, and Groschup MH. Cellular prion protein acquires resistance to proteolytic degradation following copper ion binding. *Biol Chem* 385: 739–747, 2004.
119. Lasmézas CI, Deslys JP, Robain O, Jaegly A, Beringue V, Peyrin JM, Fournier JG, Hauw JJ, Rossier J, and Dormont D. Transmission of the BSE agent to mice in the absence of detectable abnormal prion protein. *Science* 275: 402–405, 1997.
120. Lasmézas CI. Putative functions of PrP(C). *Br Med Bull* 66: 61–70, 2003.
121. Leclerc E, Serban H, Prusiner SB, Burton DR, and Williamson RA. Copper induces conformational changes in the N-terminal part of cell-surface PrP^C. *Arch Virol* 151: 2103–2109, 2006.
122. Lee KS, Magalhães AC, Zanata SM, Brentani RR, Martins VR, and Prado MA. Internalization of mammalian fluorescent cellular prion protein and N-terminal deletion mutants in living cells. *J Neurochem* 79: 79–87, 2001.
123. Lee KS, Raymond LD, Schoen B, Raymond GJ, Kett L, Moore RA, Johnson LM, Taubner L, Speare JO, Onwubiko HA, Baron GS, Caughey WS, and Caughey B. Hemin interactions and alterations of the subcellular localization of prion protein. *J Biol Chem* 282: 36525–36533, 2007.
124. Lee S and Eisenberg D. Seeded conversion of recombinant prion protein to a disulfide-bonded oligomer by a reduction-oxidation process. *Nat Struct Biol* 10: 725–730, 2003.
125. Lehmann S. Metal ions and prion diseases. *Curr Opin Chem Biol* 6: 187–192, 2002.
126. Linden R, Martins VR, Prado MA, Cammarota M, Izquierdo I, and Brentani RR. Physiology of the prion protein. *Physiol Rev* 88: 673–728, 2008.
127. Lledo PM, Tremblay P, DeArmond SJ, Prusiner SB, and Nicoll RA. Mice deficient for prion protein exhibit normal

- neuronal excitability and synaptic transmission in the hippocampus. *Proc Natl Acad Sci U S A* 93: 2403–2407, 1996.
128. Llewelyn CA, Hewitt PE, Knight RSG, Amar K, Cousens S, Mackenzie J, and Will RG. Possible transmission of variant Creutzfeldt-Jakob disease by blood transfusion. *Lancet* 363: 417–421, 2004.
 129. Lo RY, Shyu WC, Lin SZ, Wang HJ, Chen SS, and Li H. New molecular insights into cellular survival and stress responses: neuroprotective role of cellular prion protein (PrP^C). *Mol Neurobiol* 35: 236–244, 2007.
 130. Ma J and Lindquist S. Conversion of PrP to a self-perpetuating PrP^{Sc}-like conformation in the cytosol. *Science* 298: 1785–1788, 2002.
 131. Ma J, Wollmann R, and Lindquist S. Neurotoxicity and neurodegeneration when PrP accumulates in the cytosol. *Science* 298: 1781–1785, 2002.
 132. MacKenzie EL, Iwasaki K, and Tsuji Y. Intracellular iron transport and storage: from molecular mechanisms to health implications. *Antioxid Redox Signal* 10: 997–1030, 2001.
 133. Madsen E and Gitlin JD. Copper and iron disorders of the brain. *Annu Rev Neurosci* 30: 317–337, 2007.
 134. Magalhães AC, Baron GS, Lee KS, Steele-Mortimer O, Dorward D, Prado MA, and Caughey B. Uptake and neuritic transport of scrapie prion protein coincident with infection of neuronal cells. *J Neurosci* 25: 5207–5216, 2005.
 135. Malaisé M, Schätzl HM, and Bürkle A. The octarepeat region of prion protein, but not the TM1 domain, is important for the antioxidant effect of prion protein. *Free Radic Biol Med* 45: 1622–1630, 2008.
 136. Mallucci G, Dickinson A, Linehan J, Kohn PC, Brandner S, and Collinge J. Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* 302: 871–874, 2003.
 137. Martin WR, Ye FQ, and Allen PS. Increasing striatal iron content associated with normal aging. *Mov Disord* 13: 281–286, 1998.
 138. Mechoulam R, Panikashvili D, and Shohami E. Cannabinoids and brain injury: therapeutic implications. *Trends Mol Med* 8: 58–61, 2002.
 139. Melov S, Doctrow SR, Schneider JA, Haberson J, Patel M, Coskun PE, Huffman K, Wallace DC, and Malfroy B. Lifespan extension and rescue of spongiform encephalopathy in superoxide dismutase 2 nullizygous mice treated with superoxide dismutase-catalase mimetics. *J Neurosci* 21: 8348–8353, 2001.
 140. Milharet O and Lehmann S. Oxidative stress and the prion protein in transmissible spongiform encephalopathies. *Brain Res Brain Res Rev* 38: 328–339, 2002.
 141. Milharet O, McMahon HE, Rachidi W, Nishida N, Katamine S, Mangé A, Arlotto M, Casanova D, Riondel J, Favier A, and Lehmann S. Prion infection impairs the cellular response to oxidative stress. *Proc Natl Acad Sci USA* 97: 13937–13942, 2000.
 142. Millhauser GL. Copper and the prion protein: methods, structures, function, and disease. *Annu Rev Phys Chem* 58: 299–320, 2007.
 143. Millhauser GL. Copper binding in the prion protein. *Acc Chem Res* 37: 79–85, 2004.
 144. Mishra RS, Basu S, Gu Y, Luo X, Zou WQ, Mishra R, Li R, Chen SG, Gambetti P, Fujioka H, and Singh N. Protease-resistant human prion protein and ferritin are co-transported across Caco-2 epithelial cells: implications for species barrier in prion uptake from the intestine. *J Neurosci* 24: 11280–11290, 2004.
 145. Mishra RS, Gu Y, Bose S, Verghese S, Kalepu S, and Singh N. Cell surface accumulation of a truncated transmembrane prion protein in Gerstmann-Straussler-Scheinker disease P102L. *J Biol Chem* 277: 24554–24561, 2002.
 146. Miura T, Hori-I A, and Takeuchi H. Metal-dependent alpha-helix formation promoted by the glycine-rich octapeptide region of prion protein. *FEBS Lett* 396: 248–252, 1996.
 147. Miura T, Sasaki S, Toyama A, and Takeuchi H. Copper reduction by the octapeptide repeat region of prion protein: pH dependence and implications in cellular copper uptake. *Biochemistry* 44: 8712–8720, 2005.
 148. Molina-Holgado F, Hider RC, Gaeta A, Williams R, and Francis P. Metals ions and neurodegeneration. *Biomaterials* 20: 639–654, 2007.
 149. Moore RC, Mastrangelo P, Bouzamondo E, Heinrich C, Legname G, Prusiner SB, Hood L, Westaway D, DeArmond SJ, and Tremblay P. Doppel-induced cerebellar degeneration in transgenic mice. *Proc Natl Acad Sci USA* 98: 15288–15293, 2001.
 150. Moos T and Morgan EH. The metabolism of neuronal iron and its pathogenic role in neurological disease. *Ann N Y Acad Sci* 1012: 14–26, 2004.
 151. Muller WE, Laplane JL, Ushijima H, and Schroder HC. Novel approaches in diagnosis and therapy of Creutzfeldt-Jakob disease. *Mech Ageing Dev* 116: 193–218, 2000.
 152. Murakami-Kubo I, Doh-Ura K, Ishikawa K, Kawatake S, Sasaki K, Kira J, Ohta S, and Iwaki T. Quinoline derivatives are therapeutic candidates for transmissible spongiform encephalopathies. *J Virol* 78: 1281–1288, 2004.
 153. Murata M, Gong P, Suzuki K, and Koizumi S. Differential metal response and regulation of human heavy metal-inducible genes. *J Cell Physiol* 180: 105–113, 1999.
 154. Nadal RC, Abdelraheim SR, Brazier MW, Rigby SE, Brown DR, and Viles JH. Prion protein does not redox-silence Cu²⁺, but is a sacrificial quencher of hydroxyl radicals. *Free Radic Biol Med* 42: 79–89, 2007.
 155. Nicolas O, Gavín R, and Del Río JA. New insights into cellular prion protein (PrP(c)) functions: the “ying and yang” of a relevant protein. *Brain Res Rev* 61: 170–184, 2009.
 156. Nishina K, Jenks S, and Supattapone SJ. Ionic strength and transition metals control PrP^{Sc} protease resistance and conversion-inducing activity. *Biol Chem* 279: 40788–40794, 2004.
 157. Novitskaya V, Bocharova OV, Bronstein I, and Baskakov IV. Amyloid fibrils of mammalian prion protein are highly toxic to cultured cells and primary neurons. *J Biol Chem* 281: 13828–13836, 2006.
 158. Olanow CW and Arendash GW. Metals and free radicals in neurodegeneration. *Curr Opin Neurol* 7: 548–558, 1994.
 159. Orem NR, Geoghegan JC, Deleault NR, Kascak R, and Supattapone S. Copper (II) ions potently inhibit purified PrP^{res} amplification. *J Neurochem* 96: 1409–1415, 2006.
 160. Ostrerova-Golts N, Petrucelli L, Hardy J, Lee JM, Farer M, and Wolozin B. The A53T alpha-synuclein mutation increases iron-dependent aggregation and toxicity. *J Neurosci* 20: 6048–6054, 2000.
 161. Otto M, Copek L, Ratzka P, Doehlinger S, Boekhoff I, Wiltfang J, Irle E, Perganda G, Ellers-Lenz B, Windl O, Ketzscldmar HA, Poser S, and Prange H. Efficacy of flupirtine on cognitive function in patients with CJD: a double blind study. *Neurology* 62: 714–718, 2004.

162. Pamplona R, Naudí A, Gavín R, Pastrana MA, Sajani G, Ilieva EV, Del Río JA, Portero-Otín M, Ferrer I, and Requena JR. Increased oxidation, glycooxidation, and lipoxidation of brain proteins in prion disease. *Free Radic Biol Med* 45: 1159–1166, 2008.
163. Pantopoulos K. Iron metabolism and the IRE/IRP regulatory system: an update. *Ann N Y Acad Sci* 1012: 1–13, 2004.
164. Park SJ, Kim NH, Jeong BH, Jin JK, Choi JK, Park YJ, Kim JI, Carp RI, and Kim YS. The effect of Fenton reaction on protease-resistant prion protein (PrP^{Sc}) degradation and scrapie infectivity. *Brain Res* 1238: 172–180, 2008.
165. Parkin ET, Watt NT, Turner AJ, and Hooper NM. Dual mechanisms for shedding of the cellular prion protein. *J Biol Chem* 279: 11170–11178, 2004.
166. Pauly PC and Harris DA. Copper stimulates endocytosis of the prion protein. *J Biol Chem* 273: 33107–33110, 1998.
167. Perera WS and Hooper NM. Ablation of the metal ion-induced endocytosis of the prion protein by disease-associated mutation of the octarepeat region. *Curr Biol* 11: 519–523, 2001.
168. Perovic S, Schröder HC, Pergande G, Ushijima H, and Müller WE. Effect of flupirtine on Bcl-2 and glutathione level in neuronal cells treated in vitro with the prion protein fragment (PrP 106–126). *Exp Neurol* 147: 518–524, 1997.
169. Perry G, Taddeo MA, Petersen RB, Castellani RJ, Harris PL, Siedlak SL, Cash AD, Liu Q, Nunomura A, Atwood CS, and Smith MA. Adventitiously-bound redox active iron and copper are at the center of oxidative damage in Alzheimer disease. *Biometals* 16: 77–81, 2003.
170. Peters PJ, Mironov A Jr, Peretz D, van Donselaar E, Leclerc E, Erpel S, DeArmond SJ, Burton DR, Williamson RA, Vey M, and Prusiner SB. Trafficking of prion proteins through a caveolae-mediated endosomal pathway. *J Cell Biol* 162: 703–717, 2003.
171. Petersen RB, Siedlak SL, Lee HG, Kim YS, Nunomura A, Tagliavini F, Ghetti B, Cras P, Moreira PI, Castellani RJ, Guentchev M, Budka H, Ironside JW, Gambetti P, Smith MA, and Perry G. Redox metals and oxidative abnormalities in human prion diseases. *Acta Neuropathol (Berl)* 110: 232–238, 2005.
172. Petrak JV and Vyoral D. Detection of iron-containing proteins contributing to the cellular labile iron pool by a native electrophoresis metal blotting technique. *J Inorg Biochem* 86: 669–675, 2001.
173. Piccardo P, Manson JC, King D, Ghetti B, and Barron RM. Accumulation of prion protein in the brain that is not associated with transmissible disease. *Proc Natl Acad Sci U S A* 104: 4712–4717, 2007.
174. Pietri M, Caprini A, Mouillet-Richard S, Pradines E, Ermonval M, Grassi J, Kellermann O, and Schneider B. Overstimulation of PrP^C signaling pathways by prion peptide 106–126 causes oxidative injury of bioaminergic neuronal cells. *J Biol Chem* 281: 28470–28479, 2006.
175. Pollera C, Lucchini B, Formentin E, Bareggi S, Poli G, and Ponti W. Evaluation of anti-prionic activity of clioquinol in an *in vivo* model (*Mesocricetus auratus*). *Vet Res Commun* 29: 253–255, 2005.
176. Prusiner SB. Prions. *Proc Natl Acad Sci USA* 95: 3363–3383, 1998.
177. Quaglio E, Chiesa R, and Harris DA. Copper converts the cellular prion protein into a protease-resistant species that is distinct from the scrapie isoform. *J Biol Chem* 276: 11432–11438, 2001.
178. Race R, Raines A, Raymond GJ, Caughey B, and Chesebro B. Long-term subclinical carrier state precedes scrapie replication and adaptation in a resistant species: analogies to bovine spongiform encephalopathy and variant Creutzfeldt-Jakob disease in humans. *J Virol* 75: 10106–10112, 2001.
179. Rachidi W, Chimienti F, Aouffen M, Senator A, Guiraud P, Seve M, and Favier A. Prion protein protects against zinc-mediated cytotoxicity by modifying intracellular exchangeable zinc and inducing metallothionein expression. *J Trace Elem Med Biol* 23: 214–223, 2009.
180. Rachidi W, Vilette D, Guiraud P, Arlotto M, Riondel J, Laude H, Lehmann S, and Favier A. Expression of prion protein increases cellular copper binding and antioxidant enzyme activities but not copper delivery. *J Biol Chem* 278: 9064–9072, 2003.
181. Ross CA and Poirier MA. Protein aggregation and neurodegenerative disease. *Nat Med* 10(Suppl): S10–S17, 2004.
182. Roucou X and LeBlanc AC. Cellular prion protein neuroprotective function: implications in prion diseases. *J Mol Med* 83: 3–11, 2005.
183. Roucou X, Gains M, and LeBlanc AC. Neuroprotective functions of prion protein. *J Neurosci Res* 75: 153–161, 2004.
184. Roucou X, Giannopoulos PN, Zhang Y, Jodoin J, Goodyer CG, and LeBlanc A. Cellular prion protein inhibits proapoptotic Bax conformational change in human neurons and in breast carcinoma MCF-7 cells. *Cell Death Differ* 12: 783–795, 2005.
185. Roucou X, Guo G, Zhang Y, Goodyer CG, and LeBlanc AC. Cytosolic prion protein is not toxic and protects against Bax mediated cell death in human primary neurons. *J Biol Chem* 278: 40877–40881, 2003.
186. Ruiz FH, Silva E, and Inestrosa NC. The N-terminal tandem repeat region of human prion protein reduces copper: role of tryptophan residues. *Biochem Biophys Res Commun* 269: 491–495, 2000.
187. Sakaguchi S, Katamine S, Nishida N, Moriuchi R, Shigematsu K, Sugimoto T, Nakatani A, Kataoka Y, Houtani T, Shirabe S, Okada H, Hasegawa S, Miyamoto T, and Noda T. Loss of cerebellar Purkinje cells in aged mice homozygous for a disrupted PrP gene. *Nature* 380: 528–531, 1996.
188. Sakudo A, Lee DC, Saeki K, Nakamura Y, Inoue K, Matsumoto Y, Itohara S, and Onodera T. Impairment of superoxide dismutase activation by N-terminally truncated prion protein (PrP) in PrP-deficient neuronal cell line. *Biochem Biophys Res Commun* 308: 660–667, 2003.
189. Sakudo A, Lee DC, Yoshimura E, Nagasaka S, Nitta K, Saeki K, Matsumoto Y, Lehmann S, Itohara S, Sakaguchi S, and Onodera T. Prion protein suppresses perturbation of cellular copper homeostasis under oxidative conditions. *Biochem Biophys Res Commun* 313: 850–855, 2004.
190. Sakudo A, Nakamura I, Tsuji S, and Ikuta K. GPI-anchorless human prion protein is secreted and glycosylated but lacks superoxide dismutase activity. *Int J Mol Med* 2: 217–222, 2008.
191. Sakudo A, Onodera T, Suganuma Y, Kobayashi T, Saeki K, and Ikuta K. Recent advances in clarifying prion protein functions using knockout mice and derived cell lines. *Mini Rev Med Chem* 6: 589–601, 2006.
192. Schipper HM. Brain iron deposition and the free radical-mitochondrial theory of ageing. *Ageing Res Rev* 3: 265–301, 2004. (Erratum in *Ageing Res Rev* 5: 117–118, 2006).
193. Schneider B, Mutel V, Pietri M, Ermonval M, Mouillet-Richard S, and Kellermann O. NADPH oxidase and

- extracellular regulated kinases 1/2 are targets of prion protein signaling in neuronal and nonneuronal cells. *Proc Natl Acad Sci U S A* 100: 13326–13331, 2003.
194. Schroder HC and Muller WE. Neuroprotective effect of flupirtine in prion disease. *Drugs Today (Barc)* 38: 49–55, 2002.
 195. Shearer J, Soh P, and Lentz S. Both Met(109) and Met(112) are utilized for Cu(II) coordination by the amyloidogenic fragment of the human prion protein at physiological pH. *J Inorg Biochem* 102: 2103–2113, 2008.
 196. Shearer J and Soh P. The copper (II) adduct of the unstructured region of the amyloidogenic fragment derived from the human prion protein is redox-active at physiological pH. *Inorg Chem* 46: 710–719, 2007.
 197. Shi S, Dong CF, Tian C, Zhou RM, Xu K, Zhang BY, Gao C, Han J, and Dong XP. The propagation of hamster-adapted scrapie PrP^{Sc} can be enhanced by reduced pyridine nucleotide in vitro. *FEBS J* 276: 1536–1545, 2009.
 198. Shiraishi N, Ohta Y, and Nishikimi M. The octapeptide repeat region of prion protein binds Cu(II) in the redox-inactive state. *Biochem Biophys Res Commun* 267: 398–402, 2000.
 199. Shmerling D, Hegyi I, Fischer M, Blättler T, Brandner S, Götz J, Rüllicke T, Flechsig E, Cozzio A, von Mering C, Hangartner C, Aguzzi A, and Weissmann C. Expression of amino-terminally truncated PrP in the mouse leading to ataxia and specific cerebellar lesions. *Cell* 93: 203–214, 1998.
 200. Shyng SL, Moulder KL, Lesko A, and Harris DA. The N-terminal domain of a glycolipid-anchored prion protein is essential for its endocytosis via clathrin-coated pits. *J Biol Chem* 270: 14793–14800, 1995.
 201. Sigurdsson EM, Brown DR, Alim MA, Scholtzova H, Carp R, Meeker HC, Prelli F, Frangione B, and Wisniewski T. Copper chelation delays the onset of prion disease. *J Biol Chem* 278: 46199–46202, 2003.
 202. Silveira JR, Raymond GJ, Hughson AG, Race RE, Sim VL, Hayes SF, and Caughey B. The most infectious prion protein particles. *Nature* 437: 257–261, 2005.
 203. Singh A, Isaac AO, Luo X, Mohan ML, Cohen ML, Chen F, Kong Q, Bartz J, and Singh N. Abnormal brain iron homeostasis in human and animal prion disorders. *PLoS Pathog* 5: e1000336, 2009.
 204. Singh A, Kong Q, Luo X, Petersen RB, and Meyerson H. Singh N. Prion protein (PrP) knock-out mice show altered iron metabolism: a functional role for PrP in iron uptake and transport. *PLoS One* 4: e6115, 2009.
 205. Singh A, Mohan ML, Isaac AO, Luo X, and Singh N. Prion protein modulates cellular iron metabolism: implications for prion disease pathogenesis. *PLoS One* 4: e4468, 2009.
 206. Smith DG, Cappai R, and Barnham KJ. The redox chemistry of the Alzheimer's disease amyloid beta peptide. *Biochim Biophys Acta* 1768: 1976–1990, 2007.
 207. Smith MA, Harris PL, Sayre LM, and Perry G. Iron accumulation in Alzheimer disease is a source of redox-generated free radicals. *Proc Natl Acad Sci U S A* 94: 9866–9868, 1997.
 208. Sohal RS, Wennberg-Kirch E, Jaiswal K, Kwong LK, and Forster MJ. Effect of age and caloric restriction on bleomycin-chelatable and nonheme iron in different tissues of C57BL/6 mice. *Free Radic Biol Med* 27: 287–293, 1999.
 209. Solfrosi L, Criado JR, McGavern DB, Wirz S, Sánchez-Alavez M, Sugama S, DeGiorgio LA, Volpe BT, Wiseman E, Abalos G, Masliah E, Gilden D, Oldstone MB, Conti B, and Williamson RA. Cross-linking cellular prion protein triggers neuronal apoptosis in vivo. *Science* 303: 1514–1516, 2004.
 210. Spudich A, Frigg R, Kilic E, Kilic U, Oesch B, Raeber A, Bassetti CL, and Hermann DM. Aggravation of ischemic brain injury by prion protein deficiency: role of ERK-1/-2 and STAT-1. *Neurobiol Dis* 20: 442–449, 2005.
 211. Stevens DJ, Walter ED, Rodríguez A, Draper D, Davies P, Brown DR, and Millhauser GL. Early onset prion disease from octarepeat expansion correlates with copper binding properties. *PLoS Pathog* 5: e1000390, 2009.
 212. Stöckel J, Safar J, Wallace AC, Cohen FE, and Prusiner SB. Prion protein selectively binds copper(II) ions. *Biochemistry* 20: 7185–7193, 1998.
 213. Strausak D, Mercer JF, Dieter HH, Stremmel W, and Mülthaup G. Copper in disorders with neurological symptoms: Alzheimer's, Menkes, and Wilson diseases. *Brain Res Bull* 55: 175–185, 2001.
 214. Suh JH, Moreau R, Heath SH, and Hagen TM. Dietary supplementation with (R)-alpha-lipoic acid reverses the age-related accumulation of iron and depletion of antioxidants in the rat cerebral cortex. *Redox Rep* 10: 52–60, 2005.
 215. Tabner BJ, Turnbull S, Fullwood NJ, German M, and Allsop D. The production of hydrogen peroxide during early-stage protein aggregation: a common pathological mechanism in different neurodegenerative diseases? *Biochem Soc Trans* 33: 548–550, 2005.
 216. Taylor DR, Watt NT, Perera WS, and Hooper NM. Assigning functions to distinct regions of the N-terminus of the prion protein that are involved in its copper-stimulated, clathrin-dependent endocytosis. *J Cell Sci* 118: 5141–5153, 2005.
 217. Telling GC, Haga T, Torchia M, Tremblay P, DeArmond SJ, and Prusiner SB. Interactions between wild-type and mutant prion proteins modulate neurodegeneration in transgenic mice. *Genes Dev* 10: 1736–1750, 1996.
 218. Texel SJ, Xu X, and Harris ZL. Ceruloplasmin in neurodegenerative diseases. *Biochem Soc Trans* 36: 1277–1281, 2008.
 219. Thackray AM, Knight R, Haswell SJ, Bujdoso R, and Brown DR. Metal imbalance and compromised antioxidant function are early changes in prion disease. *Biochem J* 362: 253–258, 2002.
 220. Thompson KJ, Shoham S, and Connor JR. Iron and neurodegenerative disorders. *Brain Res Bull* 55: 155–164, 2001.
 221. Thomzig A, Schulz-Schaeffer W, Wrede A, Wemheuer W, Brenig B, Kratzel C, Lemmer K, and Beekes M. Accumulation of pathological prion protein PrP^{Sc} in the skin of animals with experimental and natural scrapie. *PLoS Pathog* 3: e66, 2007.
 222. Tobler I, Gaus SE, Deboer T, Achermann P, Fischer M, Rulicke T, Moser M, Oesch B, McBride PA, and Manson JC. Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* 380: 639–642, 1996.
 223. Toni M, Massimino ML, Griffoni C, Salvato B, Tomasi V, and Spisni E. Extracellular copper ions regulate cellular prion protein (PrP^C) expression and metabolism in neuronal cells. *FEBS Lett* 579: 741–744, 2005.
 224. Turnbull S, Tabner BJ, Brown DR, and Allsop D. Copper-dependent generation of hydrogen peroxide from the toxic prion protein fragment PrP106–126. *Neurosci Lett* 336: 159–162, 2003.
 225. Turnbull S, Tabner BJ, Brown DR, and Allsop D. Generation of hydrogen peroxide from mutant forms of the prion protein fragment PrP121–231. *Biochemistry* 42: 7675–7681, 2003.

226. Varela-Nallar L, Toledo EM, Chacón MA, and Inestrosa NC. The functional links between prion protein and copper. *Biol Res* 39: 39–44, 2006.
227. Varela-Nallar L, Toledo EM, Larrondo LF, Cabral AL, Martins VR, and Inestrosa NC. Induction of cellular prion protein gene expression by copper in neurons. *Am J Physiol Cell Physiol* 290: C271–C281, 2005.
228. Vassallo N and Herms J. Cellular prion protein function in copper homeostasis and redox signalling at the synapse *J Neurochem* 86: 538–544, 2003.
229. Vilette D. Cell models of prion infection. *Vet Res* 39: 10, 2008.
230. Wadsworth JDF, Hill AF, Joiner S, Jackson GS, Clarke AR, and Collinge J. Strain specific prion protein conformation determined by metal ions. *Nat Cell Biol* 1: 55–59, 1999.
231. Waggoner DJ, Drisaldi B, Bartnikas TB, Casareno RL, Prohaska JR, Gitlin JD, and Harris DA. Brain copper content and cuproenzyme activity do not vary with prion protein expression level. *J Biol Chem* 275: 7455–7458, 2000.
232. Waheed A, Grubb JH, Zhou XY, Tomatsu S, Fleming RE, Costaldi ME, Britton RS, Bacon BR, and Sly WS. Regulation of transferrin-mediated iron uptake by HFE, the protein defective in hereditary hemochromatosis. *Proc Natl Acad Sci U S A* 99: 3117–3122, 2002.
233. Walter ED, Stevens DJ, Visconte MP, and Millhauser GL. The prion protein is a combined zinc and copper binding protein: Zn^{2+} alters the distribution of Cu^{2+} coordination modes. *J Am Chem Soc* 129: 15440–15441, 2007.
234. Watt NT and Hooper NM. Reactive oxygen species (ROS)-mediated beta-cleavage of the prion protein in the mechanism of the cellular response to oxidative stress. *Biochem Soc Trans* 33: 1123–1125, 2005.
235. Watt NT and Hooper NM. The prion protein and neuronal zinc homeostasis. *Trends Biochem Sci* 28: 406–410, 2003.
236. Watt NT, Taylor DR, Gillott A, Thomas DA, Perera WS, and Hooper NM. Reactive oxygen species-mediated beta-cleavage of the prion protein in the cellular response to oxidative stress. *J Biol Chem* 280: 35914–35921, 2005.
237. Watts JC, Balachandran A, and Westaway D. The expanding universe of prion diseases. *PLoS Pathog* 2: e26, 2006.
238. Weise J, Crome O, Sandau R, Schulz-Schaeffer W, Bähr M, and Zerr I. Upregulation of cellular prion protein (PrP^C) after focal cerebral ischemia and influence of lesion severity. *Neurosci Lett* 372: 146–150, 2004.
239. Westergaard L, Christensen HM, and Harris DA. The cellular prion protein (PrP^C): its physiological function and role in disease. *Biochim Biophys Acta* 1772: 629–644, 2007.
240. White AR, Collin SJ, Maher F, Jobling MF, Stewart LR, Thyer JM, Beyreuther K, Masters CL, and Cappai R. Prion protein-deficient neurons reveal lower glutathione reductase activity and increased susceptibility to hydrogen peroxide toxicity. *Am J Pathol* 155: 1723–1730, 1999.
241. Whittal RM, Ball HL, Cohen FE, Burlingame AL, Prusiner SB, and Baldwin MA. Copper binding to octarepeat peptides of the prion protein monitored by mass spectrometry. *Protein Sci* 9: 332–343, 2000.
242. Wong BS, Brown DR, Pan T, Whiteman M, Liu T, Bu X, Li R, Gambetti P, Olesik J, Rubenstein R, and Sy MS. Oxidative impairment in scrapie-infected mice is associated with brain metals perturbations and altered antioxidant activities. *J Neurochem* 79: 689–698, 2001.
243. Wong BS, Chen SG, Colucci M, Xie Z, Pan T, Liu T, Li R, Gambetti P, Sy MS, and Brown DR. Aberrant metal binding by prion protein in human prion disease. *J Neurochem* 78: 1400–1408, 2001.
244. Wong BS, Liu T, Li R, Pan T, Petersen RB, Smith MA, Gambetti P, Perry G, Manson JC, Brown DR, and Sy MS. Increased levels of oxidative stress markers detected in the brains of mice devoid of prion protein. *J Neurochem* 76: 565–572, 2001.
245. Wong BS, Pan T, Liu T, Li R, Petersen RB, Jones IM, Gambetti P, Brown DR, and Sy MS. Prion disease: a loss of antioxidant functions? *Biochem Biophys Res Commun* 275: 249–252, 2000.
246. Wong C, Xiong LW, Horiuchi M, Raymond L, Wehrly K, Chesebro B, and Caughey B. Sulfated glycans and elevated temperature stimulate PrP(Sc)-dependent cell-free formation of protease-resistant prion protein. *EMBO J* 20: 377–386, 2001.
247. Yoo HY, Chang MS, and Rho HM. Heavy metal-mediated activation of the rat Cu/Zn superoxide dismutase gene via a metal-responsive element. *Mol Gen Genet* 262: 310–313, 1999.
248. Youdim MB, Stephenson G, and Ben Shachar D. Ironing iron out in Parkinson's disease and other neurodegenerative diseases with iron chelators: a lesson from 6-hydroxydopamine and iron chelators, desferal and VK-28. *Ann N Y Acad Sci* 1012: 306–325, 2004.
249. Yun SW, Gerlach M, Riederer P, and Klein MA. Oxidative stress in the brain at early preclinical stages of mouse scrapie. *Exp Neurol* 201: 90–98, 2006.
250. Zecca L, Stroppolo A, Gatti A, Tampellini D, Toscani M, Gallorini M, Giaveri G, Arosio P, Santambrogio P, Fariello RG, Karatekin E, Kleinman MH, Turro N, Hornykiewicz O, and Zucca FA. The role of iron and copper molecules in the neuronal vulnerability of locus coeruleus and substantia nigra during aging. *Proc Natl Acad Sci U S A* 101: 9843–9848, 2004.
251. Zecca L, Youdim MBH, Riederer P, Connor JR and Crichton RR. Iron, brain ageing and neurodegenerative disorders. *Nat Rev Neurosci* 5: 863–873, 2004.
252. Zhu X, Su B, Wang X, Smith MA, and Perry G. Causes of oxidative stress in Alzheimer disease. *Cell Mol Life Sci* 64: 2202–2210, 2007.
253. Zhu W, Xie W, Pan T, Xu P, Fridkin M, Zheng H, Jankovic J, Youdim MBH, and Le W. Prevention and restoration of lactacystin-induced nigrostriatal dopamine neuron degeneration by novel brain-permeable iron chelators. *FASEB J* 21: 3835–3844, 2007.

Address correspondence to:

Neena Singh

Department of Pathology

5527, Wolstein Research Building

Case Western Reserve University

Cleveland, OH 44106

E-mail: neena.singh@case.edu

Date of first submission to ARS Central, April 14, 2009; date of final revised submission, September 22, 2009; date of acceptance, October 3, 2009.

Abbreviations Used

8OHG = 8-hydroxyguanosine
 $A\beta$ = amyloid- β
 AD = Alzheimer's disease
 ADAM10 and
 ADAM17 = members of the disintegrin
 and metalloproteinase family
 ALS = amyotrophic lateral sclerosis
 Bcl-2 = family of antiapoptotic proteins
 that derive their name from B-cell
 lymphoma 2 protein
 BSE = bovine spongiform encephalopathy
 CJD = Creutzfeldt-Jakob disease
 CSF = cerebrospinal fluid
 DFO = desferrioxamine
 ER = endoplasmic reticulum
 ^{59}Fe = radioactive iron
 Fe^{2+} = ferrous iron
 Fe^{3+} = ferric iron
 FFI = fatal familial insomnia
 GPI = glycosylphosphatidylinositol
 GSS = Gerstmann-Strausler-Scheinker
 disease
 GT-1 = a hypothalamic neuronal cell line
 HAE = 4-hydroxyalkenal
 HD = Huntington's disease
 HFE = hereditary hemochromatosis protein
 H_2O_2 = hydrogen peroxide
 IRE = iron-responsive element
 IRP 1 and 2 = iron-responsive proteins 1 and 2
 LIP = labile iron pool
 MDA = malondialdehyde
 Mn^{2+} = manganese
 N2a = mouse neuroblastoma cell
 $\text{NF-}\kappa\text{B}$ = nuclear factor κB (transcription factor)
 Ni^{2+} = nickel
 NMDA = *N*-methyl-D-aspartate
 NOS = nitric oxide synthase
 OH^\bullet = hydroxyl radical
 PD = Parkinson's disease
 PK = proteinase-K
 PMCA = protein-misfolding cyclic amplification
 PRNP = gene encoding PrP
 $\text{PrP}^{-/-}$ = mice lacking PrP expression
 PrP^{C} = prion protein
 PrP^{C} -GFP = PrP^{C} tagged with green fluorescent
 protein
 PrP^{Sc} = PrP-scrapie
 RK13A = rat kidney cell line
 ROS = reactive oxygen species
 ScN2a = scrapie infected mouse
 neuroblastoma cell
 SDS-PAGE = sodium dodecylsulfate polyacrylamide
 gel electrophoresis
 SMB = mouse brain cell of
 mesenchymal origin
 SOD = Cu/Zn superoxide dismutase
 Tf = transferrin
 TfR = transferrin receptor
 $\text{TNF-}\alpha$ = tumor necrosis factor alpha
 Zn^{2+} = zinc

This article has been cited by:

1. Feimeng Zhou, Glenn L. Millhauser. 2012. The rich electrochemistry and redox reactions of the copper sites in the cellular prion protein. *Coordination Chemistry Reviews* **256**:19-20, 2285-2296. [[CrossRef](#)]
2. Paola D'Angelo, Stefano Della Longa, Alessandro Arcovito, Giordano Mancini, Andrea Zitolo, Giovanni Chillemi, Gabriele Giachin, Giuseppe Legname, Federico Benetti. 2012. Effects of the Pathological Q212P Mutation on Human Prion Protein Non-Octarepeat Copper-Binding Site. *Biochemistry* **51**:31, 6068-6079. [[CrossRef](#)]
3. Sepehr Ehsani, Ashkan Salehzadeh, Hairu Huo, William Reginold, Cosmin L. Pocanschi, Hezhen Ren, Hansen Wang, Kelvin So, Christine Sato, Mohadeseh Mehrabian, Robert Strome, William S. Trimble, Lili-Naz Hazrati, Ekaterina Rogaeva, David Westaway, George A. Carlson, Gerold Schmitt-Ulms. 2012. LIV-1 ZIP Ectodomain Shedding in Prion-Infected Mice Resembles Cellular Response to Transition Metal Starvation. *Journal of Molecular Biology* . [[CrossRef](#)]
4. Ajay Singh, Liuting Qing, Qingzhong Kong, Neena Singh. 2011. Change in the characteristics of ferritin induces iron imbalance in prion disease affected brains. *Neurobiology of Disease* . [[CrossRef](#)]
5. Lin Liu, Dianlu Jiang, Alex McDonald, Yuanqiang Hao, Glenn L. Millhauser, Feimeng Zhou. 2011. Copper Redox Cycling in the Prion Protein Depends Critically on Binding Mode. *Journal of the American Chemical Society* 110718073144042. [[CrossRef](#)]
6. Gianluca Lepore, Sergio Gadau, Antonella Peruffo, Arcadia Mura, Emilio Mura, Antonello Floris, Francesca Balzano, Marco Zedda, Vittorio Farina. 2011. Aromatase expression in cultured fetal sheep astrocytes after nitrosative/oxidative damage. *Cell and Tissue Research* **344**:3, 407-413. [[CrossRef](#)]
7. Salvatore Bongarzone, Hoang Ngoc Ai Tran, Andrea Cavalli, Marinella Roberti, Michela Rosini, Paolo Carloni, Giuseppe Legname, Maria Laura Bolognesi. 2011. Hybrid Lipoic Acid Derivatives to Attack Prion Disease on Multiple Fronts. *ChemMedChem* **6**:4, 601-605. [[CrossRef](#)]
8. Isidro Ferrer, Margarita Carmona, Rosa Blanco, Maria Jesus Rey Recio, R. M. San Segundo. 2011. Gerstmann-Straüssler-Scheinker PRNP P102L-129V mutation. *Translational Neuroscience* **2**:1, 23-32. [[CrossRef](#)]
9. Snehadrinarayan Khatua, Jina Kang, Kibong Kim, Jung Oh Huh, Junseong Lee, Chang Seop Hong, David G. Churchill. 2010. Crystal Structures and Magnetic Properties of Newly Synthesized Mono- and Dinuclear CuII Schiff-Base Complexes. *European Journal of Inorganic Chemistry* **2010**:31, 5018-5026. [[CrossRef](#)]