

Oxidative Stress and Blood–Brain Barrier Dysfunction Under Particular Consideration of Matrix Metalloproteinases

Christine Lehner,^{1,2} Renate Gehwolf,^{1,3} Herbert Tempfer,⁴ Istvan Krizbai,⁵
Bernhard Hennig,⁶ Hans-Christian Bauer,^{1,4} and Hannelore Bauer¹

Abstract

A cell's "redox" (oxidation and reduction) state is determined by the sum of all redox processes yielding reactive oxygen species (ROS), reactive nitrogen species (RNS), and other reactive intermediates. Low amounts of ROS/RNS are generated by different mechanisms in every cell and are important regulatory mediators in many signaling processes (redox signaling). When the physiological balance between the generation and elimination of ROS/RNS is disrupted, oxidative/nitrosative stress with persistent oxidative damage of the organism occurs. Oxidative stress has been suggested to act as initiator and/or mediator of many human diseases. The cerebral vasculature is particularly susceptible to oxidative stress, which is critical since cerebral endothelial cells play a major role in the creation and maintenance of the blood–brain barrier (BBB). This article will only contain a focused introduction on the biochemical background of redox signaling, since this has been reported already in a series of excellent recent reviews. The goal of this work is to increase the understanding of basic mechanisms underlying ROS/RNS-induced BBB disruption, with a focus on the role of matrix metalloproteinases, which, after all, appear to be a key mediator in the initiation and progression of BBB damage elicited by oxidative stress. *Antioxid. Redox Signal.* 15, 1305–1323.

Introduction

Oxidative stress:

Cellular and physiological background

REACTIVE OXYGEN SPECIES (ROS) and reactive nitrogen species (RNS), summarized in this article as ROS, are present at moderate levels in every living cell where they fulfill vitally important functions. Provided that the redox homeostasis is well balanced, redox signaling enables the cells to respond properly to endogenous and exogenous stimuli of various kinds. Any imbalance between generation and elimination of ROS leads to a physiologic phenomenon termed oxidative stress. Oxidative stress processes are involved directly or indirectly in many human pathologies, including cancers, neurodegenerative diseases, diabetes, atherosclerosis, ischemic stroke, and many more (for reviews, see Refs. 34, 164, 168).

ROS are produced by a great variety of mechanisms and originate from many sources. Exogenous sources include

exposure to di-oxygen, ozone (a powerful oxidizing agent that can oxidize biological components directly), ionizing and nonionizing irradiation, UV radiation, air pollutants, cigarette smoke, industrial contaminants, the invasion of pathogens, and food that may contain various kinds of oxidants, including peroxides, aldehydes, oxidized fatty acids, and transition metals (89). Endogenously, two ROS-generating systems exist: a) the leakage of activated oxygen from mitochondria during oxidative phosphorylation; and b) oxygen-metabolizing enzymatic reactions such as xanthine oxidase, cytochromes P450, NADPH oxidases, myeloperoxidase, and nitric oxidase synthase (140).

ROS can basically be subdivided into two groups: a) The radical group containing superoxide ion radical, hydroxyl radical, nitric oxide radical, peroxy and alkoxy radicals, and one form of singlet oxygen; and b) nonradical compounds comprising a large variety of substances such as hypochlorous acid, hydrogen peroxide, organic peroxides, aldehydes, ozone, and oxygen (89). The terms reactive oxygen species,

¹Department of Organismic Biology, Development Biology Group, ²Clinic for Sports Injuries and Trauma Surgery, and ³Christian Doppler Clinic, University Hospital of Salzburg, Salzburg, Austria.

⁴Paracelsus Private Medical University, Salzburg, Austria.

⁵Institute of Biophysics, Biology Research Center, Szeged, Hungary.

⁶Molecular and Cell Nutrition Laboratory, University of Kentucky, Lexington, Kentucky.

reactive nitrogen species, oxygen-derived species, pro-oxidants and oxidants are often used synonymously in the literature. In the following, oxidative stress and nitrosative stress will be referred to as *oxidative stress only*.

Longstanding medical interest has focused on the deleterious effect of oxidative stress on the cerebral microvasculature. The compromised functional integrity of the BBB leads to increased permeability of cerebral microvessels, which contributes to the pathology of a number of neurological diseases with inflammatory characteristics (reviewed in Ref. 140). A list of major prooxidant enzymes in cerebral endothelial cells (cEC) is shown in Figure 1.

Reactive oxygen species

The superoxide anion (O_2^-) is formed by the univalent reduction of triplet-state molecular oxygen (3O_2). Its intracellular generation is mediated by enzymes such as NAD(P)H oxidases and xanthine oxidase or nonenzymatically by redox-reactive compounds such as the semi-ubiquinone compound of the mitochondrial electron transport chain or in auto-oxidation reactions. O_2^- is relatively unreactive with a half-life time of about 10^{-3} sec, compared to the highly reactive hydroxyl radical ($\cdot OH$) (half-life time of about 10^{-10} sec) and has a rather poor ability to cross biological membranes. O_2^- can enzymatically and nonenzymatically be converted to hydrogen peroxide (H_2O_2) which, in turn, may either be converted into water by the enzymes catalase or glutathione peroxidase (GSHPx), or, in the presence of reduced transition metals (ferrous or cuprous ions), to the hydroxyl radical $\cdot OH$ (34).

O_2^- is capable of acting directly on cEC, thereby modulating the permeability of the BBB. It has been suggested that O_2^- is produced during the firm adhesion of monocytes to cEC, which triggers cytoskeletal rearrangements allowing infiltration of monocytes into the brain (170).

Reactive nitrogen species

NO is formed by three different mammalian types of NO synthases: the constitutive isoforms endothelial NO synthase (eNOS/NOS3) and neuronal synthase (nNOS/NOS1), and the inducible isoform (iNOS/NOS2). NO plays an important role in the regulation of cerebral blood flow and neuronal cell viability; however, excessive production of NO by iNOS and nNOS in the brain participates in neurotoxicity (162).

Actually, NO is a weak oxidant and, by itself, is hardly capable of inducing oxidative stress. However, as shown in a

rodent brain perfusion model, various NO redox forms ($NO\cdot$, NO^+ , NO^-) exert different permeability effects on the BBB. While $NO\cdot$ causes only modest BBB disruption, a combination of $NO\cdot/NO^-$ or $NO\cdot/NO^-/NO^+$ elicits severe BBB damage, as reflected by ^{14}C -sucrose uptake (18).

By interaction of $NO\cdot$ and O_2^- , formation of the highly noxious peroxynitrite ($ONOO^-$) occurs. The protonated form of $ONOO^-$ is a powerful oxidizing agent and exerts deleterious effects in biological systems by causing the oxidation of DNA and the nitration of aromatic amino acid residues in proteins (e.g., forming 3-nitrotyrosine) (34, 89). At physiological pH, $ONOO^-$ may decompose through the intermediate formation of its conjugated acid $ONOOH$, which may either form the very reactive radicals $NO_2\cdot$ and $\cdot OH$ or may react in biological conditions with carbon dioxide (154).

BBB dysfunction elicited by NO and NO-derivatives is well documented (129, 156). Transcriptional upregulation of eNOS and iNOS, as well as increased production of 3-nitrotyrosine, appears to be characteristic of BBB breakdown and cerebral edema following cerebral trauma (120). iNOS-deficient mice showed significantly reduced disruption of the BBB following experimental bacterial induction of meningitis compared to wild-type mice, which was associated with a lack of $ONOO^-$ and 3-nitrotyrosine formation (180). The purine metabolite uric acid proved to be a powerful scavenger of $ONOO^-$ in inflammatory brain disease (83). In a mouse model of active experimental autoimmune encephalomyelitis (the mouse correlate of human multiple sclerosis), exogenously administered uric acid was shown to act at two levels. It reduced BBB damage and penetrated the compromised BBB to block $ONOO^-$ -mediated tyrosine nitration and apoptotic cell death in inflamed neuronal tissue (67).

Targets of oxidative damage

Numerous studies have demonstrated that ROS can attack virtually all biological molecules. The major targets are lipids, proteins, and DNA and are discussed in more detail below.

Lipids. Free radicals attack the phospholipid components of cellular membranes, leading to alterations of membrane permeability and lipid fluidity, as well as modified lipid-protein interactions (58, 191). The damage to lipids, termed *lipid peroxidation*, is a chain reaction. It is initiated by any primary free radical that has sufficient reactivity to abstract a hydrogen atom from a reactive methylene group of a polyunsaturated fatty acid. The formation of the initiating complex is followed by a bond rearrangement. The lipid radical then takes up oxygen to form the peroxy radical, which can combine with each other and even attack proteins.

The breakdown of lipid peroxides generates radicals that propagate lipid peroxidation and also nonradical fragmentation products such as malondialdehyde, alkanals, 2-alkenals, and 4-hydroxyalkenal, many of which are biologically active. The highly reactive 4-hydroxynonenal (4-HNE) is the biologically most significant class of peroxidation products. The main aldehydes that have been found in peroxidizing biological samples are malondialdehyde, hexanal, propanal, and 4-HNE (26, 28, 37, 38).

Increased plasma levels of malondialdehyde and nitrates/nitrites were found to be associated with hypoxic/ischemic encephalopathy (97). *In vitro*, H_2O_2 -induced oxidative stress

Prooxidant enzymes in cerebral endothelial cells		
iNOS (NOS2)	Inducible nitric oxidase synthase	$NO\cdot$
eNOS (NOS3)	Endothelial nitric oxide synthase	$NO\cdot$
XDH/XOD	Xanthine dehydrogenase/oxidase	O_2^-
COX-1	Cyclooxygenase-1	O_2^-
COX-2	Cyclooxygenase-2	O_2^-
MAO	Monoaminoxidase	H_2O_2
NAD(P)HO	NAD(P)H oxidase	O_2^-

($NO\cdot$) nitric oxide radical; (O_2^-) superoxide anion; (H_2O_2) hydrogen peroxide; (NADP) Nicotinamide-adenin-dinucleotide-phosphate

FIG. 1. Prooxidant enzymes in cerebral endothelial cells (according to (24), with modifications).

was shown to induce lipid peroxidation and membrane dysfunction in cerebral endothelial cells, which could be reduced by decomposition of free radicals by nitronyl nitroxides (17). Further, in an *in vitro* BBB model, 4-HNE was shown to increase the barrier permeability, obviously by decreasing the content of reduced glutathione in cEC (113). A genotoxic effect of 4-HNE was reported by Karlhuber *et al.* (82), showing that treatment of cultured cEC with 4-HNE resulted in chromosomal aberrations and the formation of micronuclei.

Proteins. ROS-mediated oxidation of proteins can result in protein crosslinking and increased proteolytic degradation, inactivation, or altered cellular functions. Numerous modifications of amino acid side chains may occur, including hydroxylation of aromatic groups and aliphatic amino acid side chains, nitration of aromatic amino acid residues, nitrosylation of sulfhydryl groups, sulfoxidation of methionine residues, chlorination of aromatic groups and primary amino groups, and conversion of some amino acid residues to carbonyl derivatives (26, 145). Methionine residues and cysteine residues of proteins are particularly sensitive to oxidation by ROS but, unlike oxidation of other amino acid residues, the oxidation of these sulfur amino acids is reversible. Protein damage is mainly caused by $\cdot\text{OH}$, $\text{RO}\cdot$, and nitrogen-reactive radicals, yielding protein oxidation products such as aldehydes, keto compounds, and carbonyls. During the last couple of years, increasing attention has been paid to the importance of protein oxidation in aging, supported by the observation that levels of oxidized proteins increase with age (144, 146). This age-related accumulation of oxidized proteins may reflect age-related increases in rates of ROS generation, decreases in antioxidant activities, or losses in the capacity to degrade oxidized proteins. As mentioned above, one of the major adducts during protein oxidation is produced by the interaction between $\text{ONOO}\cdot$ and other nitrogen reactive radicals with the amino acid tyrosine, forming 3-nitrotyrosine. This adduct can easily be detected in tissues and plasma and is used as a marker for oxidative damage of proteins (130). The increased production of 3-nitrotyrosine contributes to BBB disruption. For instance, in A β 42 peptide-injected rat hippocampus, the selective inhibition of iNOS reduced 3-nitrotyrosine levels and diminished BBB leakiness to IgG (139).

DNA. Some ROS produce a multiplicity of modifications in DNA, such as release of free bases from DNA and generation of strand breaks with various sugar modifications and damage to the DNA repair system. Not all ROS can cause damage. $\cdot\text{O}_2^-$ and H_2O_2 are normally not reactive towards DNA. However, upon metal catalyzation, H_2O_2 and $\cdot\text{O}_2$ might be converted to $\cdot\text{OH}$ known to readily attack DNA. A wide spectrum of oxidative modifications, which yield oxidation products such as 8-hydroxydeoxyguanosine (8-OHdG) (reviewed in (89)), or 8 (or 4-, 5-)-hydroxyadenine, thymine peroxide, thymine glycols, and uracil glycol, are elicited by ROS. In addition, peroxynitrite can easily cause DNA damage similar to that obtained when $\cdot\text{OH}$ is involved. The biological consequences of many of the oxidative products are known. For example, thymine glycol interferes with DNA replication and is potentially lethal to cells, while 8-oxoG is readily bypassed by the DNA polymerase and is highly mutagenic. Unrepaired 8-oxoG will mispair with

deoxyadenosine (dA), leading to an increase in G (guanine) to T (thymine) transition mutations (138).

Cytotoxic and genotoxic effects of oxidative stress were demonstrated using cultured cEC (21). Aglycemic culture conditions exacerbated the cytotoxic effects of oxidative insult, as evidenced by increased apoptosis and decreased mitotic activity of cEC (21).

Endothelial and Epithelial Tissue Barriers: Biological Systems Vulnerable to Oxidative Stress

All epithelial and endothelial tissue barriers, such as the vertebrate BBB, the blood–retina barrier, or the blood–cerebrospinal fluid barrier, rely on the establishment of particularly tight intercellular contacts (tight junctions, TJs) between neighboring cells. TJs are the most restrictive intercellular junctions located at the apical border of the lateral membrane where they fulfill two main tasks: They act as a permeability barrier in the paracellular space and as a fence that separates apical and baso-lateral domains within the plasma membrane of epithelial and endothelial cells. In due course, a distinct apical-basal polarity is created, allowing the vectorial transport of ions and macromolecules across the cell. The structural features of TJs resemble all other occluding or adhesive junctions acting between epithelial and endothelial cells (*i.e.*, a single- or multi-pass transmembrane component is connected to cytoplasmic proteins, commonly referred to as peripheral junctional proteins). The extracellular domains of the integral junctional proteins of two opposing cells interact in a homophilic or heterophilic manner to create a paracellular “seal” (25a, 94, 166).

Two groups of proteins are involved in the formation, function and modulation of TJs: a) integral membrane proteins that bridge the intercellular space, and b) peripheral proteins, which assemble at the cytoplasmic surface of the junction site where they interact directly or indirectly with the transmembrane components (6, 35).

Integral membrane proteins found at endothelial TJs include the tetraspan proteins claudins (125a) and occludin (17a), which create two extracellular loops, exposing their N- and C-termini to the cytosol. The junctional adhesion molecules (JAMs) (13a) and the endothelial cell-selective adhesion molecule (ESAM) (66), which belong to the immunoglobulin superfamily due to their Ig-like domains located at the extracellular portion of the proteins, represent the major single span proteins contributing to the endothelial BBB (13, 66, Fig. 2). ESAM mediates homophilic adhesion between endothelial cells and was shown to bind to the scaffolding protein MAGI-1 (membrane-associated guanylate kinase with inverted domain structure 1) (177). ESAM is specifically expressed in endothelial cells and platelets and is involved in platelet aggregation and leukocyte transmigration across the endothelial wall (147,178).

So far, a plethora of peripheral proteins located at TJs have been discovered, the role of which has just partly been unravelled. Proteins localizing at the cytoplasmic surface of TJs comprise adaptor or scaffolding proteins, signaling molecules, and transcriptional regulators (6, 11, 35, 52, 57). The zonula occludens (ZO) proteins, ZO-1 and ZO-2, were shown to localize to the cytoplasm and to the nucleus in epithelial and endothelial cells (165). These proteins exert a dual role, being structural components at the junctional site and regulators of gene expression in epithelial and endothelial cells (52a).




FIG. 2. The mammalian blood-brain barrier (BBB) and structural components of endothelial tight junctions (TJs). **(A)** The BBB in development. Mouse embryo (day 13 after conception), microperfused with Trypan blue and fixed with PFA. The dye (bound to plasma proteins) stains all parts of the body with exception of the brain and the spinal cord. This suggests that at this early stage of development the BBB is functional at least for proteins (10). The arrow indicates a slight staining in the developing choroid plexus. **(B)** Schematic drawing of the neurovascular unit. Cerebral endothelial cells are tightly connected by tight junctions. Pericytes are in close contact with the endothelial cells. A basal lamina surrounds both endothelial cells and pericytes. Capillary walls are ensheathed by astrocytic end feet. Occasionally, neurites are also in direct contact with capillaries and so are microglia. Afp, astrocytic foot process; BL, basal lamina; cEC, cerebral endothelial cell; CL, capillary lumen; MG, microglia; n, nucleus; PC, pericyte, TJ, tight junction. **(C)** Integral proteins locating at endothelial TJs. Peripheral junctional proteins may be subdivided into three classes: i) Adaptors and scaffolding proteins, ii)

regulatory and signaling proteins, and iii) transcriptional and posttranscriptional regulators. AJs, adherens junctions; ESAM (endothelial cell-specific adhesion molecule); JAMs, junctional adhesion molecules; TJs, tight junctions. (To see this illustration in color, the reader is referred to the web version of this article at www.liebertonline.com/ars).

The vertebrate BBB is one of the key interfaces between blood and neural tissue of higher vertebrates, including mammals. TJs of the BBB restrict the paracellular movement of ions and polar solutes between the blood and brain and impede the transport of macromolecules to the central nervous system (CNS) and vice versa (1, 36, 141), thereby TJs contribute to the creation and maintenance of a specific intracerebral homeostatic milieu. cEC are in close contact with other non-neural and neural cell types (pericytes, astrocytes, neurons, and microglia) and the extracellular matrix (ECM), a complex commonly referred to as the "neurovascular unit" (63). Disturbance of TJs by oxidative stress leads to impairment of the BBB, compromising the CNS (95, 104). Apparently, a finely tuned interplay of the cells constituting the BBB is important to cope with oxidative insult. In this context, astrocytes were found to be less susceptible to oxidative stress compared to cECs, which may protect and support the maintenance of the BBB during oxidative insult (21). In

addition, astrocytes are involved in the regulation of neuronal glutathione homeostasis and, in this way, may contribute to neuronal protection during oxidative stress (176). The role of pericytes, which are intimately attached to the cerebral capillary wall, has been studied extensively (for review, see Ref. 96). Although their supportive role in BBB function is generally accepted, pericytes were shown to exhibit sustained contraction upon ischemia-induced oxidative stress, which impairs the restoration of the cerebral microcirculation and negatively affects the survival of neuronal tissue (184).

Selected Pathological Conditions Eliciting Oxidative Stress

Tumors

Alterations in oxygen metabolism appear to be a common feature of tumor cells, although the genetic profile of each type of tumor varies considerably. In normal cells, oxygen is used

to produce energy (ATP) through oxidative phosphorylation. During the last couple of years, intriguing evidence has accumulated suggesting that cancer cells activate glycolysis (in the presence of O_2) to cope with their increased energy demand. As a consequence, part of the oxygen is available for generation of ROS, particularly $\cdot O_2^-$ and H_2O_2 . This metabolic switch from oxidative phosphorylation to aerobic glycolysis has repeatedly been observed in cancer cells (48). Accumulation of ROS and oxidative stress play an important role in carcinogenesis and is subject of many comprehensive reviews (87, 105, 127, 163).

In addition to carcinogenic agents, the activation of several oncogenes and mutations in the tumor suppressor gene p53 were found to increase the cellular production of ROS (91, 109, 167).

The critical role of ROS in tumor biology has prompted the elaboration of novel approaches for antioxidant drug treatments in cancer therapy (redox chemotherapeutics). In addition, induction of oxidative stress by pro-oxidant agents is emerging as an attractive anticancer strategy. Some anticancer agents produce H_2O_2 which is known to be an efficient inducer of cell death in cancer cells (3, 117). Knockout of antioxidant defense enzymes was shown to increase oxidative damage promoting age-related cancer (59).

In recent years, a particular focus has been directed towards the role of ROS in brain tumors, particularly gliomas (astrocytomas/glioblastomas, ependymomas, oligodendrogliomas), the largest group of primary brain tumors. Oxidative DNA damage (8-oxoguanine) is high in some glioblastoma multiforme (GBM) cell lines and many GBM tumors. Also, the incidence of HNE-immunopositive tumor cells in astrocytic and ependymal tumors parallels the increasing grades of malignancy. Again, highest levels of HNE were found in tumor cells of GBM (78).

A significant decrease of antioxidant enzyme activity such as glutathione reductase (GRx) and superoxide dismutase (SOD) activity was observed in most types of brain tumor cases, leading to the assumption that the degree of decline in antioxidant levels may indicate severity of malignancy in brain tumors (132).

Downregulation of endothelial TJ proteins and impairment of the BBB in glioma was reported severalfold (72, 101, 181). Particularly, expression of claudin-1, -3, -5, and occludin appears to be affected in microvessels of human glioblastoma and astrocytoma. Sufficient evidence suggests that VEGF (vascular endothelial growth factor) also contributes to BBB disruption with subsequent edema in brain tumors (106).

Diabetes and diabetic retinopathy

Diabetes in general is characterized by pancreatic beta-cell dysfunction. Oxidative stress produced by islet-infiltrating immune cells has been considered to contribute to the destruction of beta cells. Diabetes is usually accompanied by both, production of free radicals and impaired antioxidant defenses (reviewed in Refs. 39, 108, 179).

Under diabetic conditions, chronic hyperglycemia gradually deteriorates beta-cell function, and exacerbates insulin resistance, a process called "glucose toxicity." Oxidative stress activates the JNK pathway, which is likely to be involved in pancreatic beta cell dysfunction and insulin resistance (81). Increased levels of ROS contribute to insulin

resistance, as demonstrated by Houstis *et al.* using two cellular models (68).

The main source of free radicals is glucose, which is converted into reactive ketoaldehydes and to superoxide anion radicals. In addition, the hyperglycemic environment leads to nonenzymatic glycosylation of numerous proteins forming advanced glycation end products (AGEs) that accumulate in diabetic plasma and tissues (142). A well-characterized receptor for AGEs was shown to be present on monocytes, macrophages, and neutrophils (27, 42). AGEs alone trigger minimal $\cdot O_2^-/H_2O_2$ production by neutrophils, but upon chemoattractant stimulation the production of ROS becomes markedly increased. This process seems to be related directly to glycemic control involving protein kinase C (PKC), NADPH oxidase, and the calcium-independent phospholipase A(2) (iPLA2) (5, 54, 64). However, recent evidence suggests that diabetic cells are capable of developing a mechanism of self-defense by increasing their antioxidant content, which results in high levels of glutathione and overexpression of a large set of genes encoding antioxidant proteins such as hemeoxygenase-1 and the transcription factor Nrf2 (nuclear erythroid-related factor-2/NF-E2-related factor 2) (98).

Diabetes causes severe pathologic aberrations in the retina, collectively termed as "diabetic retinopathy" (RP). Diabetic RP affects vascular and nonvascular cells of the retina. Vascular damage ranges from increased vascular permeability and leukostasis (*i.e.*, abnormal attraction and adhesion of leukocytes to the vascular wall) to capillary occlusion and degeneration. Increased expression of inflammation-related proteins, including iNOS, COX-2, ICAM-1, caspase 1, VEGF, and NF- κ B, and increased production of ROS and inflammatory cytokines is characteristic of early stages of diabetic RP (85). Antioxidants have been found to inhibit the development of inflammatory changes in retinas of diabetic animals (92, 93). For detailed reviews on diabetic RP, see Frey and Antonetti (43a), and for inflammation of cerebral barriers see Coisne and Engelhardt (25a).

Disruption of the BBB is an early event in the development of diabetic RP. Again, VEGF was shown to play a key role in the pathogenesis of diabetic retinopathy by inducing the PKC-dependent phosphorylation and subsequent degradation of occludin (61, 119).

Glucocorticoids reduce retinal endothelial cell permeability and stimulate the synthesis and dephosphorylation of occludin (40). In an *in vitro* model hydrocortisone was shown to increase barrier properties in cEC by inducing enhanced expression of occludin (43). In primary retinal endothelial cells, glucocorticoids increased the transcription of occludin and claudin-5 through a transactivation mechanism that does not involve direct promoter binding by the glucocorticoid receptor (41).

Ischemic stroke and reperfusion injury

The immediate loss of blood supply to the cerebral tissue is commonly referred to as ischemic stroke. Microvascular responses to ischemia are very rapid and highly coordinated, although they vary topographically with the degree of ischemic injury (32). Within 1 to 2 hours following middle carotid artery occlusion, initial loss of the primary microvessel permeability barrier occurs, producing transudation of small

molecules, fibrinogen, and plasma (32). The pathogenesis of ischemic stroke involves various mechanisms, among which oxidative injury and inflammatory processes appear to be the most critical events. ROS are substantially involved in the pathological pathways of cerebral ischemia and reperfusion (99). In an initial phase, ROS production by nNOS and eNOS type I and type III is activated, while at a later stage increased activity of iNOS in glia and neutrophils may be observed (182). Actually, all cells of the neurovascular unit are able to activate one or more enzyme systems (including NOS, enzymes of the mitochondrial respiratory chain, xanthine oxidase, and NADPH-dependent oxidases) to contribute to the intra- and extracellular formation of ROS (153). ROS may potentiate the activation of stress-related transcription factors, including hypoxia inducible factor-1 and NF- κ B (20, 114, 159). In turn, the expression of cytokines and adhesion molecules in cerebral endothelial cells is induced, which further facilitates activation and recruitment of inflammatory cells to the infarcted site (185).

Overexpression of radical-scavenging enzymes was shown to protect against stroke, and animals lacking such enzymes were more susceptible to cerebral ischemic damage (50). Migration of monocytes across a monolayer of rat cEC was reduced by scavenging O_2^- with SOD and by blocking the ROS-producing enzyme xanthine oxidase (XO) with allopurinol (170). The gp91phox containing NADPH oxidase contributes to ROS formation by glia cells, fibroblasts, vascular endothelial cells, and most of all by leukocytes (53, 169). In leukocytes, NADPH oxidase is responsible for the respiratory burst in which the cell generates toxic amounts of superoxide anions. In NADPH oxidase-knockout (gp91phox $^{-/-}$) mice, middle cerebral artery occlusion-induced BBB disruption and lesion volume were shown to be largely diminished compared with those in wild-type mice (80).

A critical role of ROS in cerebral injury accompanying ischemic stroke is also reflected by the fact that even delayed treatment with antioxidants and free radical scavengers could be neuroprotective in such pathological states (50, 168, 173). However, in aged rats, antioxidant treatment (*i.e.*, suppression of NADPH oxidase activity) even increased BBB permeability to sucrose following ischemic stroke, suggesting that age adds a further level of variability to the treatment of oxidative brain damage (84).

Increased expression of metalloproteinases and inflammatory cytokines and reduced levels of occludin and claudin-5 were observed in ischemic rat brains. The loss of TJ proteins was reflected by BBB disruption, as evidenced by increased Evans blue extravasation into the brain (73). A decrease in ZO-1 and occludin protein levels, and an increase in tyrosine phosphorylation of occludin, was suggested to be critical for the disruption of TJs and dysfunction of the BBB after cerebral ischemia (79). Prevention of occludin phosphorylation by the Src tyrosine kinase inhibitor PP2 inhibited BBB leakage and decreased the infarct volume following transient focal ischemia in rats (155). Moreover, selective inhibition of the EP1 subtype of prostaglandin E2 receptors in a mouse model of focal cerebral ischemia, produced by middle cerebral artery occlusion, decreased occludin phosphorylation and reduced BBB disruption, pointing towards an additional mechanism underlying the deleterious effects of prostaglandin E2 in the setting of cerebral ischemia (44).

Mediators Through Which Oxidative Stress Acts on BBB Function (ROS, Cytokines, Leukocytes)

The disruption of epithelial and endothelial barriers seems to be a crucial step in the acquisition of inflammation after oxidative insults (133).

Endothelial cells may be both a source and target of ROS. ROS attack lipids and proteins of their target cells, leading to the release and activation of cytokines and proteases that propagate the vascular damage. The most studied cytokines related to inflammation in acute ischemic stroke are tumor necrosis factor- α (TNF- α), the interleukins (IL), IL-1 β , IL-6, IL-20, IL-10, and IL-17, and transforming growth factor (TGF)- β (reviewed in Ref. 99).


IL-17A was demonstrated to induce NADPH oxidase- or XO-dependent ROS production. The resulting oxidative stress activated the endothelial contractile machinery, which was accompanied by downregulation of occludin. Blocking either ROS formation or myosin light chain phosphorylation prevented IL-17A-induced BBB disruption (70).

Inflammatory cytokines were shown to induce the expression of various cell adhesion molecules (CAMs) on endothelial cells (174, 175). The appearance of P-selectin, intercellular adhesion molecule-1 (ICAM-1), and E-selectin on microvessel endothelium, together with their counter-receptors (*e.g.*, P-selectin glycoprotein-1 (PSGL-1) and the β 2-integrin CD18) on PMN (polymorphonuclear) leukocytes was shown to accompany the initial movement of inflammatory cells into the ischemic region (32). ROS treatment of pulmonary endothelial cells induced the phosphorylation of the focal adhesion kinase pp125^{FAK}, a cytosolic tyrosine kinase implicated in the oxidant-mediated adhesion process (171). In addition, expression of chemokines, such as the monocyte chemoattractant protein-1 (MCP-1) and macrophage inflammatory protein-1 α (MIP-1 α) in cerebral cells, further supports leukocyte entry and increased oxidative damage (115, 116, 175).

TNF- α plays a pivotal role in various types of inflammatory brain diseases (8, 22, 45, 46, 186). O_2^- appears to be a primary mediator of TNF- α -induced oxidative stress in the cerebral vascular endothelium. An important cellular source generating proapoptotic O_2^- in response to TNF- α in cerebral vascular endothelium is the xanthine oxidase/reductase system (126). XO and XDH (xanthine dehydrogenase) are interconvertible forms of the same enzyme (123), known as xanthine oxidoreductase (XOR). Normally this enzyme is present *in vivo* as an NAD-dependent cytosolic dehydrogenase (XDH), incapable of ROS production. Following conversion of XDH to XO by sulfhydryl oxidation or limited proteolysis, the enzyme produces O_2^- and H_2O_2 (122). One of the primary roles of XOR is the conversion of hypoxanthine to xanthine and xanthine to uric acid (Fig. 3).

In hypoxic endothelial cells, XO is phosphorylated, at least in part, via casein kinase II and the stress-activated p38 MAP kinase which seems to be necessary for the hypoxia-induced enzymatic activation. XO was shown to be released into the blood stream in human ischemia-reperfusion injury (157). Interestingly, it has been proposed that circulating XO binds to sulphated glycosaminoglycans on the surface of endothelial cells (2). This endothelial cell-bound XO continues to produce oxidants, which can further trigger endothelial dysfunction and may contribute to organ injury also in remote organs (157).

FIG. 3. Oxidative damage of the BBB occurs through the xanthine-oxidase dependent production of ROS during reperfusion following cerebral ischemia. In endothelial cells xanthine oxidase (XO/XOD) is the key source of $\cdot\text{O}_2^-$ metabolites in acute ischemia/reperfusion conditions. During the course of ischemia, transmembrane ion gradients are dissipated, allowing cytosolic concentrations of Ca^{2+} to rise, which in turn, activates proteolytic conversion of XDH to XO. Simultaneously, cellular ATP is catabolized to hypoxanthine. During the reperfusion, XO uses readmitted oxygen and generates $\cdot\text{O}_2^-$ and H_2O_2 through conversion of hypoxanthine to xanthine and xanthine to uric acid. SOD converts $\cdot\text{O}_2^-$ to H_2O_2 which is further processed to H_2O by catalase (schematic drawing according to (126) with modifications).



ROS selectively activate signaling cascades involving RhoA, PI3 kinase, and protein kinase B (PKB/Akt) to increase monocyte migration across an *in vitro* BBB model. Thereby, rearrangements of the actin cytoskeleton and spatial redistribution and disappearance of occludin and claudin-5 take place (143). Hypoxia-induced cytoskeletal alterations contributing to BBB disruption were also reported by Hicks *et al.* (65). The authors showed that hypoxic stress leads to redistribution of moesin and VASP (vasodilator-stimulated phosphoprotein), two proteins involved in actin polymerization in BBB endothelial cells (65). Oxidative stress also induces the expression of distinct Toll-like receptors in cerebral endothelial cells *in vitro*, which leads to downregulation and delocalization of occludin and claudin-5 and a concomitant increase in permeability (121).

Redox-Sensitive Transcriptional Regulation

During oxidative stress, the proper function and integrity of the BBB depends on various redox-sensitive systems, which are active at different ROS levels (Fig. 4). Redox-sensitive signaling proteins and downstream transcription factors act in concert to maintain the intracellular prooxidant-antioxidant homeostasis (128). The redox-sensitive thioredoxins (Trxs) constitute a family of enzymes which, together with other redox-sensitive systems, play important roles in maintaining the intracellular redox balance. Many recent publications have shown that redox regulation by Trxs is critically involved in the pathogenesis of various oxidative stress-associated disorders (90, 128). In the nucleus, Trxs interact physically with the ubiquitously expressed bifunctional protein Redox-factor-1 (Ref-1/Ape1/HAP1/APEX1) (16, 160). Ref-1 is involved in DNA repair but also acts as transcriptional activator by reducing oxidized cysteine

residues within the DNA-binding domain of a number of transcription factors, such as activator protein-1 (AP-1), nuclear factor κB (NF- κB), p53, early growth response protein-1 (Egr-1), activating transcription factor/cAMP-response element-binding protein (ATF/CREB), hypoxia-inducible factor (HIF)-1 α , paired box-containing proteins (Pax), and many more. The reduced form of Ref-1 is restored by Trx in a redox cycle (reviewed in Refs. 124, 160). In addition, Ref-1 suppresses the intracellular production of ROS by negatively regulating the activity of the Ras-related GTPase Rac1 (131). In phagocytic cells, Rac proteins have been previously shown to regulate $\cdot\text{O}_2^-$ production through the neutrophil NADPH oxidase system (19). Similar to the NADPH oxidase complex in phagocytic cells, an FAD-binding protein regulated by Rac1 may regulate ROS levels in nonphagocytic cells such as fibroblasts (150).

Both *in vivo* and *in vitro* studies demonstrated that the elevation of intracellular ROS induces a rapid and transient upregulation of Ref-1 protein expression, leading to an increase in cell resistance toward oxidative stress and DNA damage (55, 131). Activation of Ref-1 may also be obtained by a process independent of *de novo* protein synthesis. This involves the cytoplasmic-nuclear redistribution of Ref-1 during oxidative stress or upon physiologic increase in intracellular ROS production (131). Transgenic mice overexpressing Trx1 show an extended life span and are relatively resistant to ischemia-mediated brain damage (128). Also, intravenously administered Trx was shown to reduce oxidative cerebral damage in mice following transient focal ischemia (62).

Oxidative Stress and Metalloproteinases

A number of proteases, including neutral proteases (caspases), serine proteases, tissue-type and urokinase-type




FIG. 4. BBB function is modulated by various redox-sensitive systems which are active at different ROS levels. Reactive oxygen species (ROS) are synthesized constitutively in eukaryotic cells. At low levels, all three SOD isoforms dismutate $\cdot\text{O}_2^-$, forming H_2O_2 , which is scavenged by peroxisomal catalase or GPx. At increasing levels of ROS additional systems are activated. The redox-sensitive thioredoxins interact physically with the ubiquitously expressed bi-functional protein Ref-1/Ape1, usually involved in DNA repair. Increased levels of Ref-1 accumulate in the nucleus where they act as transcriptional regulator. Ref-

1 reduces oxidized cysteine residues within the DNA-binding domain of a number of transcription factors (TFs) including AP-1, NF- κ B, HIF-1 α , and others. Thereby, the DNA binding capacity of these TFs is increased and transcription of a number of downstream genes, which are involved in the inflammatory and adaptive response to oxidative stress, is activated. AP-1/NF κ B target genes include MMP-9, COX-2 and iNOS. HIF-1 α target genes include VEGF, iNOS, and HO-1. Ref-1 per se is capable of reducing intracellular ROS production by inhibiting the small GTPase Rac-1. Nrf2 regulates the expression of a set of detoxifying enzymes and antioxidant proteins. Nrf-2 is normally sequestered in the cytoplasm where it is attached to its repressor protein Keap-1. Upon increasing (mediate) levels of ROS, Nrf-2 dissociates from Keap-1 and translocates to the nucleus where it binds to distinct upstream sequences (ARE sequences) of its target genes, leading to increased transcription. ARE-containing target genes include SOD, GPx, catalase, HO-1, and many others (all citations are given in the text). AP-1, Activator protein-1; ARE, antioxidant responsive element; COX-2, cyclooxygenase-2; GPx, glutathione peroxidase; HIF-1 α , hypoxia inducible factor- α ; HO-1, heme oxygenase-1; iNOS, inducible nitric oxidase; Keap-1, (Kelch-like ECH-associated protein-1); MMP-9, matrix metalloproteinase-9; NF- κ B, nuclear factor- κ B; Nrf-2, nuclear erythroid-related factor-2; Ref-1/Ape1, redox factor-1/AP endonuclease-1; SOD, superoxide dismutase.

plasminogen activator (tPA, uPA), two families of metalloproteinases (matrix metalloproteinases, MMPs), and the adamalysins (ADAMs), are involved in the vascular response to oxidative insult. In this main section, we will focus on the role of MMPs which appear to be key mediators in the pathogenesis of cerebral oxidative damage and concomitant BBB dysfunction.

Biology of MMPs

MMPs are proteolytic enzymes (zinc-dependent endopeptidases) which are secreted as zymogens. To be active, these zymogens have to be cleaved by other MMPs or proteases such as tPA or uPA (149, 172). For instance, MMP-3 was shown to activate proMMP-1, -7, -8, -9, and -13, while MMP-2 can activate proMMP-1, -9, and -13 (29). Oxidation and S-nitrosylation are two protease-independent processes that can activate MMPs (23).

A detailed study of the complex regulatory mechanisms of MMP-9 expression and the role of multiple co-activators regulating MMP-9 transcription was reported recently (190). The authors showed that MMP-9 gene expression depends on binding of the sequence specific activators AP-1 and NF- κ B to their corresponding *cis*-elements on the MMP-9 promoter. In addition, members of three classes of coactivators are essential for MMP-9 expression, including the CREB-binding protein/p300 (CBP/p300), the p300/CBP-associated factor (PCAF),

the coactivator-associated arginine methyltransferase 1 (CARM1), and the glucocorticoid receptor interacting protein-1/steroid receptor coactivator-2 (GRIP-1/SRC-2).

Since MMPs are capable of degrading all kinds of ECM proteins, they play a central role in tissue repair and ECM remodeling during development, morphogenesis, and angiogenesis. Besides being responsible for ECM degradation processes, MMPs are involved in the cleavage of cell surface receptors and the release of apoptotic ligands (shedding) and in cytokine inactivation (23).

Based on their substrate specificity, cellular localization and protein domain structure, four to five major classes of MMPs may be defined, including collagenases, stromelysins and matrilysins, gelatinases, and membrane-type metalloproteinases (MT-MPs). Gelatinases, stromelysins, and MT-MPs are the main MMPs found in the brain. Although MMPs show particulate substrate preferences, there is much overlap in substrate specificity between subgroups.

The protein structures of MMPs share a basic common pattern of propeptide and catalytic domains (86). The matrilysin MMP-7 has the simplest domain structure, with the propeptide domain and a zinc-containing catalytic domain, although it degrades most ECM components. MMP-14, -15, and -16 contain an additional transmembrane domain, while the stromelysins (MMP-1, -3, and -10) are characterized by a hemopexin domain. The gelatinases (MMP-2 and -9) have a unique fibronectin binding do-

main, allowing them to bind to basement membranes. MMP-2 and -9 (alternatively referred to as gelatinase-A, and -B) specifically digest type IV collagen, laminin, and fibronectin. The latency state of MMPs is provided by the common pro-peptide.

MMPs are inhibited by four endogenous inhibitors, termed tissue inhibitors to the metalloproteinases (TIMPs). The balance between MMPs and TIMPs is critical for cell survival. TIMP-1 is a 28 kD molecule that forms a complex with MMP-9, whereas TIMP-2 complexes with MMP-2 (at higher concentrations), although TIMPs are able to inhibit all of the MMPs. TIMP-3 is unique in that it is bound to the cell surface where it inhibits preferentially MMP-3 and the tumor necrosis factor- α converting enzyme (TACE) (29).

MMPs and vascular diseases

Expression of MMPs in the adult brain is very low but many MMPs are rapidly upregulated in response to injury. MMP-2, MMP-3, and MMP-9 are the main soluble MMPs produced in the CNS (23).

The role of MMPs in cerebral diseases, including ischemia/reperfusion and stroke, has been a focus of intensive research during the last decade. There is general agreement that MMPs are critically involved in the initiation and progression of ischemic/hypoxic cerebral insults preceding BBB disruption (23, 107). Results derived from numerous *in vivo* and *in vitro* studies concerning the role of MMPs in cerebral oxidative stress have been summarized in a series of comprehensive reviews (23, 29, 33, 47, 75, 86, 107, 136, 187).

Activation of MMPs in response to oxidative stress

The role of MMPs in oxidative stress-induced cerebral diseases is complex, and various proposed mechanisms have been based upon the knowledge derived from ischemic/hypoxic insult and inflammatory processes in other tissues. There is evidence that MMP-2, which is constitutively expressed by astrocytes, becomes activated upon the formation of a ternary complex involving MMP-2, TIMP-2, and MT1-MMP. This provides the basis for a first and local event of proteolysis, leading to basal lamina degradation and BBB damage. In addition, expression and release of pro-MMP-9 is augmented as a response to the increase of ROS and inflammatory molecules released by activated astrocytes and due to infiltrating inflammatory cells (leukocytes, monocytes/neutrophils, macrophages) (23) (Fig. 5).

During hypoxic/ischemic and inflammatory conditions, all major cell types of the neurovascular unit, including neurons, cEC, reactive astrocytes, and microglia, express MMPs, either through induction by inflammatory cytokines (TNF- α , IL-6, IL-1 β , endothelin-1) or growth factors (23, 102, 151).

Evidence that ROS mediate BBB disruption through activation of MMPs

Free radicals can regulate expression and activity of MMPs both directly and indirectly. Direct regulation may occur through oxidation or S-nitrosylation of MMPs, resulting in the unmasking of the propeptide (without cleavage of the pro-domain) and subsequent activation of MMPs (56, 112). In addition, ROS-induced inflammatory cytokines (*e.g.*, IL1- β , IL-6, TNF- α) may activate signaling pathways (including




FIG. 5. The increase in MMP-9 activity is indicative of BBB disruption. A complex pattern of MMP expression occurs in cerebral hypoxia/ischemia and reperfusion. In reperfusion animal models an early (within the first hours of reperfusion) increase of MMP-2 is observed, which correlates with an early reversible opening of the BBB. The initial opening is followed by a second opening of the BBB, within the time range of 24 to 48 hours after the onset of reperfusion. At that stage, a marked increase in MMP-9 activity occurs, which leads to severe damage of the BBB. Impairment of BBB integrity is reflected by the loss of TJ-related proteins such as occludin and claudin-5. ROS levels increase during ischemia/reperfusion due to the production of ROS by cells of the neurovascular unit (cECs, astrocytes, pericytes, microglia) and by infiltrating inflammatory cells (monocytes, neutrophils, leukocytes and macrophages). The suppression of ROS production and the detoxification and scavenging of ROS by endogenous and exogenous antioxidant systems and molecules prevent BBB disruption and attenuate the oxidative damage elicited by ischemic stroke. BL, basal lamina; COX, cyclooxygenase; eTJ, endothelial tight junctions; LMW antioxidants, low molecular weight antioxidants; Nrf2, nuclear erythroid-related factor-2; SOD, superoxide dismutase.

MEK1-Erk, MAPK P38, and the phosphatidylinositol 3 (PI3)-kinase), resulting in the increased binding of transcription factors (NF- κ B and AP-1) to redox-sensitive cis-regulatory elements in the MMP-9 promoter. Thereby, the transcription of MMP-9 is regulated (69) (Fig. 6).

Results from a mouse transient focal cerebral ischemia model revealed that during reperfusion, pro-MMP-9 was increased more in SOD1-deficient mice than in their wild-type counterparts. In addition, active MMPs were found to colocalize with $\cdot\text{O}_2^-$ at the level of capillary walls and astrocytic processes in ischemic brain regions. In contrast, transgenic mice overexpressing SOD1 showed decreased lesion size and edema along with decreased activity of MMP-9, compared with wild-type littermates (118).




FIG. 6. Inflammatory cells contribute to the elevation of MMP-9 levels in neural tissue and cEC constituting the BBB. Monocytes, macrophages and leukocytes play a critical role in many pathological conditions, including oxidative stress. They are a major source of MMP-9 and ROS attacking the BBB. In addition, they release inflammatory cytokines and growth factors, such as interleukin (IL)-1 β , IL-6, TNF α , FGF, EGF, which can stimulate the production and secretion of MMP-9 in all cells constituting the BBB. The signaling pathways involved include activation of MEK1-Erk, MAPK P38, and PI3K-Akt. Transcription of MMP-9 is regulated through activating AP-1 and NF- κ B, which subsequently bind to cis elements on the MMP-9 promoter. AP-1 and NF- κ B act in concert with various chromatin remodeling complexes and co-activators to induce MMP-9 expression. These include CBP/P300, PCAF, CARM1, and GRIP-1/SRC-2. Protease-independent activation of MMPs occurs by S-nitrosylation or oxidation to unmask the catalytic domain without pro-domain cleavage (citations are given in the text). AP-1, activator protein-1; CARM1, coactivator-associated arginine methyltransferase 1; CBP, CREB-binding protein; CREB, cAMP-response-element binding protein; EGF, epidermal growth factor; ERK, extracellular signal activated protein kinase; FGF, fibroblast growth factor; GRIP-1, glucocorticoid receptor interacting protein-1; MAPK, mitogen activated protein kinase; MEK1-ERK, mitogen activated ERK activating kinase; NF- κ B, nuclear factor κ B; PCAF, p300/CBP associated factor; PI3K, phosphatidylinositol 3-kinase; SRC-2, steroid receptor coactivator-2; TNF α , tumor necrosis factor alpha.

ROS were also found to increase BBB permeability and monocyte migration *in vitro* in a protein tyrosine kinase (PTK)-dependent manner (60). The authors showed that ROS not only increased activation of MMP-1, -2, and -9, which paralleled the degree of degradation of basement membrane components, but also enhanced tyrosine phosphorylation of tight junction-related proteins (60).

Evidence regarding the involvement of NO in the induction and activation of MMPs is still controversial. A dose-

dependent, biphasic regulatory effect of NO on the activity of MMPs (MMP-9, -1, and -13) secreted from murine macrophages was described recently (135). However, a detailed study on the mechanism of NO-mediated activation of MMPs has shown that NO is incapable of directly activating proMMP-9, while a modest activation was found through S-nitrosylation by the N-donor S-nitrosocysteine (110).

The role of MMP-9 in BBB/TJ disturbance during oxidative stress

Accumulating evidence suggests that activation of MMP-9 appears to be a key step in the impairment of BBB function in focal cerebral ischemia (23, 29, 33, 86, 99). Following transient and permanent middle cerebral artery closure, the significant increase of proMMP-9 and induction of MMP-9 activity correlated with an increase of BBB permeability, which further confirms the important role for MMP-9 in the early formation of vasogenic edema after transient focal cerebral ischemia. Synthetic MMP inhibitors restore the early integrity of the BBB but are ineffective in the later BBB opening (137).

The infiltration of neutrophils in the infarcted tissue is a major source of MMP-9 (see also Fig. 6). In this context, MMP-9 was shown to be prepackaged in neutrophils and released under neuroinflammatory conditions (4, 49). MMP-9 deficient mice showed diminished BBB disruption and decreased inflammation following ischemic conditions (4, 152).

Early inhibition (day 1) of MMP-9 was shown to reduce infarction of day 14 in an ischemia/stroke model. However, the benefit was lost when the treatment was delayed until day 3 (187), suggesting that the effect of MMP-inhibition strongly depends on the timing of treatment in relation to the stage of brain injury.

It is important to mention that MMPs may have a different role during delayed phases after stroke. Because MMPs modulate vascular ECM, they may mediate beneficial plasticity and remodeling during the stroke recovery phase. If treatment with MMP inhibitors is administered too late (*i.e.*, at a timepoint when neurovascular remodeling in the infarcted area has already started) recovery is impaired and tissue damage is exacerbated.

Compelling evidence has suggested that MMPs are involved in BBB disruption by acting directly on TJ-related proteins (183). Exposure of mice to prolonged hypoxia was demonstrated to increase MMP-9 activity and cerebral vascular permeability. Concomitantly, expression of occludin was decreased, resulting in disrupted continuity of occludin and zonula occludens-1 (ZO-1) staining in endothelial cells. Inhibition of MMP activity reduced vascular leakage and attenuated the disorganization of TJs (9). Treatment with normobaric hyperoxia could attenuate MMP-9-mediated early BBB disruption following ischemic stroke in rats (103). Treatment of brain slices or isolated microvessels with purified MMP-9 resulted in specific degradation of occludin (103). Significant degradation of occludin and claudin-5 was observed in cEC as a response to overexpression of MMP-9 (25). The observed alterations of TJ proteins correlated with increased BBB permeability to fluorescein isothiocyanate-dextran tracers (25). Further, live cell imaging has shown that transendothelial migration of monocytes coincides with the local disappearance of occludin and subsequent endothelial

gap formation, which could be prevented by administration of a broad-spectrum inhibitor of MMPs (134).

Defense Mechanisms of the Brain and the BBB Against Oxidative Stress

Due to the impermeability of the BBB, antioxidative treatment of intraneural tissue is particularly difficult. Antioxidants should protect the endothelial barrier itself and must be able to transmigrate through the BBB to reach the brain parenchyma. A comprehensive survey of neuroprotective agents, supposed to be effective in the treatment of ischemic stroke, has been published previously (51).


Under normal physiological conditions, all cells are equipped with an elaborated antioxidant system for protection against reactive metabolites (for review, see Ref. 124). Basically, the antioxidant system can be classified into two major groups: enzymes and low molecular weight antioxidants. The enzymes include a limited number of proteins, SOD, catalase, GSHPx, GRx, and peroxiredoxins. All of the corresponding genes are under the control of Nrf2 (77, 99). Under normal conditions, Nrf2 is sequestered in the cytoplasm and associated with its repressor Kelch-like protein (Keap1), an actin-binding protein. On exposure of cells to oxidative stress, Nrf2 dissociates from Keap1 and translocates

into the nucleus. Nrf2 binds to a recognition sequence, referred to as antioxidant response element (ARE), located in the upstream promoters of many genes encoding antioxidant proteins (88).

Activation of Nrf2 represents a promising treatment strategy to ameliorate cerebral oxidative insult (31, 76). In a rat model, the naturally occurring alkylating agent, sulforaphane, protected the brain from ischemic oxidative damage. The systemically administered sulforaphane was shown to enter the brain, as determined by increased mRNA and protein levels of the Nrf2-responsive gene heme oxygenase 1 (188). Even delayed administration of sulforaphane efficiently reduced the loss of TJ-related proteins (occludin, claudin-5) and preserved the BBB in a rodent model of brain injury (189). Further, short sulforaphane treatment of astrocytes resulted in long-lasting elevation of endogenous antioxidants and increased resistance to superoxide-induced cell damage (14).

Low molecular weight molecules comprise indirectly acting antioxidants (*e.g.*, chelating agents) and directly acting antioxidants (*e.g.*, scavengers and chain breaking antioxidants). The latter are extremely important in combating against oxidative stress. Common nonenzymatic antioxidants include the vitamins A, C, and E, glutathione, lipoic acid, mixed carotenoids, coenzyme Q10, bioflavonoids, antioxidant minerals (copper, zinc, manganese, and selenium), folic acid,

FIG. 7. TJs, establishing the epithelial and endothelial brain barriers are multifunctional complexes. Tight junctions are based on protein complexes locating at epithelial and endothelial intercellular contact sites. Their major constituents forming the intercellular seal are represented by integral membrane proteins, such as claudins, the MARVEL proteins occludin, tricellulin, and MarvelD3, and the junctional adhesion molecules (JAMs). A great variety of peripheral junctional proteins are located at the cytoplasmic surface of TJs forming a functional network between



the plasma membrane and the cytoskeleton. For review, see Refs. 6, 12, 35, 52, 100, 148. Although junctional components of endothelial and epithelial TJs are largely comparable, several unique features may be attributed to endothelial barriers. For instance, endothelial cells express the vascular specific occludin-5, but not claudin-1. Claudin-5 and -12 appear to be the key claudins expressed at the BBB (30). In addition, occludin, tricellulin (marveld2), and JAM4 are highly enriched in cEC (30). Tricellulin is known to seal TJs at tricellular contacts (71), which may be critical for forming a high electrical resistance barrier. Through their "fence" function, TJs establish a particular apical-basal polarity, enabling the vectorial transport of solutes and molecules across the barrier. Accumulating evidence suggests that TJs are involved in intracellular signaling (52) and regulation of gene expression (12, 111). TJs, tight junctions; TJS, tight junction strands. *Square brackets*: proteins associating exclusively with adherens junctions. Apg-2, ATP and peptide-binding protein in germ cells; aPKC, atypical protein kinase C; ARVCF, armadillo repeat gene deleted in velo-cardio-facial syndrome; CAR, coxsackie and adenovirus receptor; CDK-4, cyclin-dependent kinase-4; C/EBP, CCAAT/enhancer binding protein; CLMP, coxsackie and adenovirus receptor-like membrane protein; CRB1, human crumbs homologue; DEP-1, density-enhanced phosphatase-1; ESAM, endothelial cell-selective adhesion molecule; GEF, GDP/GTP-exchange factor; JAMs, junctional adhesion molecules; MAP-K, mitogen-activated protein kinases; Pals1, protein associated with Lin-7; PAR-3, -6, partitioning defective protein-3, -6; PATJ, Pals1 associated tight junction protein; PK-A, -G, protein kinase-A, -G; PP1, protein phosphatase1; PP2A, phosphatase2A; SAF-B, scaffold attachment factor-B; ZAK, ZO-1 associated kinase; ZO-1, -2, zonula occludens protein-1, -2; ZONAB, ZO-1-associated nucleic acid binding protein.

and vitamins B1, 2, 6, and 12. Only a few of these molecules, such as glutathione, are synthesized by the cell itself. The majority, including ascorbic acid, lipoic acid, polyphenols, and carotenoids, are derived from dietary sources.

As outlined above, a considerable number of compounds are available which are potentially effective in reducing or preventing oxidative damage to neural cells *in vitro*. However, their successful application *in vivo* has been hampered due to the impermeability of the BBB. Therefore, sufficient knowledge of the molecular and functional characteristics of the BBB

is essential for the successful design of systemically applied antioxidants.

So far, much progress has been made concerning the elucidation of the molecular components of the intra- and intercellular network of tight endothelial and epithelial cell-cell contacts (Fig. 7) (see also Ref. 25a). As already outlined above, major integral proteins of TJs (occludin, claudin-5) are affected by oxidative stress (17a, 125a). In addition, peripheral junctional proteins, such as the multi-domain ZO proteins appear to be major targets of oxidative




FIG. 8. Nuclear translocation of zonula occludens (ZO-2) protein-2 induces downregulation of MMP-9 at the transcriptional level and suppresses the migratory activity of cerebral endothelial cells. Cerebral endothelial cells were stably transfected with a ZO-2 mutant carrying a strong nuclear targeting information signal. Cloning of the ZO-2 mutant was done as described previously (165) (A) cECs were shown to express nuclear ZO-2 in addition to the endogenous ZO-2 expressed at cell borders (*upper panel*). Control cells exhibit ZO-2 staining mainly at cell-cell contact sites. Antibodies: polyclonal anti-ZO-2 (Zymed), Alexa-Fluor568 (Invitrogen). Immunostaining was performed as described earlier (165). (B) Quantitative PCR demonstrated that expression of MMP-9 is significantly reduced in cEC overexpressing nuclear ZO-2, whereas expression of MMP-2 appears unaltered. Control levels are represented as relative gene expression 1.0. One representative experiment is shown. Isolation of RNA, cDNA synthesis, and quantitative PCR was done as described by us previously (161). Primers used were: mGAPDH (internal control) forward 5' GTG AAC GGA TTT GGC CGT AT 3'; mGAPDH reverse: 5' CTT CTC CAT GGT GGT GAA GA 3'; rat MMP-9 forward: 5' TTC GAC TTG AAG TCT CAG AAG GT 3'; rat MMP-9 reverse: 5' TAG GTC ACG TAG GCC ACG TGG TC 3'. (C) cEC were grown on coverslips coated with collagen IV. After wounding, the migratory activity of the cells was monitored. cEC overexpressing nuclear ZO-2 showed strongly decreased migration compared to controls. Pictures were taken 4 days after wounding. *Dotted line* indicates the border of the wounded area (day 0). (D) ZO-2 is considered a "dual residency" protein. It is normally located at the cytoplasmic surface of junctional sites and interacts with various types of integral membrane proteins (11). In subconfluent monolayers and in response to environmental stress conditions, ZO-2 translocates to the nucleus (74, 165). Nuclear activities of ZO-2 include its interaction with SAF-B/HET (165), a negative regulator of hsp27 (125), and with transcription factors of the AP-1 complex (15). It remains to be clarified whether one of these already described nuclear activities of ZO-2 is involved in the pathway leading to transcriptional downregulation of MMP-9. AP-1, activator protein-1; hsp27, heat shock protein27; SAF-B/HET, scaffold attachment factor-B/Hsp27-ERE-TATA-binding protein. (To see this illustration in color, the reader is referred to the web version of this article at www.liebertonline.com/ars).

insult (52a). The involvement of ZO proteins in oxidative stress response is of particular interest since ZO-proteins (ZO-1 and ZO-2), are "dual residency" proteins, capable of shuttling between the junctional site and the nucleus (for further details, see Ref. 11). Their role in transcriptional regulation has been shown several-fold (7, 15, 158). *In vitro* studies have demonstrated that nuclear translocation of ZO-2 in epithelial cells is induced upon environmental stress (heat shock, toxic insult) (165). Moreover, increased nuclear accumulation of ZO-2 was found in cEC *in vitro* subjected to oxidative stress (H. Bauer, unpublished observation). In addition, experimentally-induced nuclear overexpression of ZO-2 in cEC led to the downregulation of MMP-9 transcription (H. Bauer, unpublished observation) (Fig. 8). MMP-9 is a major mediator of oxidative damage in the brain and MMP9-deficient mice show reduced BBB disruption following ischemic conditions (4, 112). Although further studies are needed, it may be speculated that the nuclear targeting of ZO proteins represents a novel mechanism that is involved in a signaling cascade, leading to alterations of gene expression and probably BBB protection in response to oxidative stress.

Conclusion

Considerable medical and experimental evidence has suggested a critical role for ROS and MMPs in the pathophysiology of cerebral oxidative damage and BBB disruption. There is evidence that activation of the cellular antioxidant defense, either through enzymatic or nonenzymatic means, can provide some protection against BBB dysfunction.

Much of the information concerning the role of oxidative stress in BBB disruption has emerged from observations and experiments *in vitro* or *in vivo*, using co-culture systems mimicking the BBB or defined animal models. So far, several compounds were shown to protect the BBB *in vitro* and to reduce the loss of TJs at the BBB in brain microvessel endothelial cells. However, in practice, most of the drugs used in these experimental systems failed to provide neuroprotection during and following oxidative insult in clinical trials.

The BBB contributes to oxidative cerebral damage in two ways: First, the BBB itself is a target of ROS-mediated cellular damage (lipid peroxidation, protein/DNA modifications, junctional alterations) and, second, the BBB impedes the transport of many systemically administered antioxidative compounds.

Therefore, any drug design that enables BBB transport will depend on new knowledge of the molecular and functional biology of brain capillary endothelial cells *in vivo* and *in vitro*. Additional work will have to focus on the role of transcriptional regulators inducing endogenous antioxidant gene expression in cerebral tissue. Because cEC are a critical component responsible for normal BBB function, changes in intracellular signaling of BBB endothelial cells in response to an imbalance in the oxidative stress/antioxidant status should be a critical focus of future BBB research. In particular, one of the challenges will be to further expand our understanding of the physiological and molecular mechanisms underlying the interplay of ROS and MMPs in oxidative brain injury. This knowledge should help to develop new ideas for

designing and elaborating effective medical treatment approaches to prevent or reduce the deleterious outcome of oxidative cerebral insults.

Acknowledgments

This work was supported by the Austrian FWF P-18743 and by research grants from the Paracelsus Private Medical University (PMU) Salzburg, Austria. H. Bauer and H.C. Bauer HC are participants of the NEUROBID consortium (EU 7th FP).

References

- Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, and Begley DJ. Structure and function of the blood-brain barrier. *Neurobiol Dis* 37: 13–25, 2010.
- Adachi T, Fukushima T, Usami Y, and Hirano K. Binding of human xanthine oxidase to sulphated glycosaminoglycans on the endothelial-cell surface. *Biochem J* 289: 523–527, 1993.
- Alexandre J, Batteux F, Nicco C, Chereau C, Laurent A, Guillemin L, Weill B, and Goldwasser F. Accumulation of hydrogen peroxide is an early and crucial step for paclitaxel-induced cancer cell death both *in vitro* and *in vivo*. *Int J Cancer* 119: 41–48, 2006.
- Asahi M, Asahi K, Jung JC, del Zoppo GJ, Fini ME, and Lo EH. Role for matrix metalloproteinase 9 after focal cerebral ischemia: Effects of gene knockout and enzyme inhibition with BB-94. *J Cereb Blood Flow Metab* 20: 1681–1689, 2000.
- Ayilavarapu S, Kantarci A, Fredman G, Turkoglu O, Omori K, Liu H, Iwata T, Yagi M, Hasturk H, and Van Dyke TE. Diabetes-induced oxidative stress is mediated by Ca²⁺-independent phospholipase A2 in neutrophils. *J Immunol* 184: 1507–1515, 2010.
- Balda MS and Matter K. Tight junctions at a glance. *J Cell Sci* 121: 3677–3682, 2008.
- Balda MS and Matter K. Tight junctions and the regulation of gene expression. *Biochim Biophys Acta* 1788: 761–767, 2009.
- Barone FC, Arvin B, White RF, Miller A, Webb CL, Willette RN, Lysko PG, and Feuerstein GZ. Tumor necrosis factor- α . A mediator of focal ischemic brain injury. *Stroke* 28: 1233–1244, 1997.
- Bauer AT, Burgers HF, Rabie T, and Marti HH. Matrix metalloproteinase-9 mediates hypoxia-induced vascular leakage in the brain via tight junction rearrangement. *J Cereb Blood Flow Metab* 30: 837–848, 2010.
- Bauer H, Sonnleitner U, Lametschwandtner A, Steiner M, Adam H, and Bauer HC. Ontogenic expression of the erythroid-type glucose transporter (Glut 1) in the telencephalon of the mouse: Correlation to the tightening of the blood-brain barrier. *Brain Res Dev Brain Res* 86: 317–325, 1995.
- Bauer H, Zweimueller-Mayer J, Steinbacher P, Lametschwandtner A, and Bauer HC. The dual role of zonula occludens (ZO) proteins. *J Biomed Biotechnol* 2010: 402593, 2010.
- Bauer HC, Traweger A, Zweimueller-Mayer J, Lehner C, Tempfer H, Krizbai I, Wilhelm I, and Bauer H. New aspects of the molecular constituents of tissue barriers. *J Neural Transm* 118: 7–21, 2011.
- Bazzoni G. The JAM family of junctional adhesion molecules. *Curr Opin Cell Biol* 15: 525–530, 2003.

- 13a. Bazzoni G. Pathobiology of junctional adhesion molecules. *Antioxid Redox Signal* 15: 1221–1234, 2011.
14. Bergstrom P, Andersson HC, Gao Y, Karlsson JO, Nodin C, Anderson MF, Nilsson M, and Hammarsten O. Repeated transient sulforaphane stimulation in astrocytes leads to prolonged Nrf2-mediated gene expression and protection from superoxide-induced damage. *Neuropharmacology* 60: 343–353, 2011.
15. Betanzos A, Huerta M, Lopez-Bayghen E, Azuara E, Amerena J, and Gonzalez-Mariscal L. The tight junction protein ZO-2 associates with Jun, Fos and C/EBP transcription factors in epithelial cells. *Exp Cell Res* 292: 51–66, 2004.
16. Bhakat KK, Mantha AK, and Mitra S. Transcriptional regulatory functions of mammalian AP-endonuclease (APE1/Ref-1), an essential multifunctional protein. *Antioxid Redox Signal* 11: 621–638, 2009.
17. Blasig IE, Mertsch K, and Haseloff RF. Nitronyl nitroxides, a novel group of protective agents against oxidative stress in endothelial cells forming the blood–brain barrier. *Neuropharmacology* 43: 1006–1014, 2002.
- 17a. Blasig IE, Bellmann C, Cording J, del Vecchio G, Zwanziger D, Huber O, and Haseloff RF. Occludin protein family: oxidative stress and reducing conditions. *Antioxid Redox Signal* 15: 1195–1219, 2011.
18. Boje KM and Lakhman SS. Nitric oxide redox species exert differential permeability effects on the blood–brain barrier. *J Pharmacol Exp Ther* 293: 545–550, 2000.
19. Bokoch GM and Zhao T. Regulation of the phagocyte NADPH oxidase by Rac GTPase. *Antioxid Redox Signal* 8: 1533–1548, 2006.
20. Bonello S, Zahringer C, BelAiba RS, Djordjevic T, Hess J, Michiels C, Kietzmann T, and Gorlach A. Reactive oxygen species activate the HIF-1 α promoter via a functional NF κ B site. *Arterioscler Thromb Vasc Biol* 27: 755–761, 2007.
21. Bresgen N, Karlhuber G, Krizbai I, Bauer H, Bauer HC, and Eckl PM. Oxidative stress in cultured cerebral endothelial cells induces chromosomal aberrations, micronuclei, and apoptosis. *J Neurosci Res* 72: 327–333, 2003.
22. Bruce AJ, Boling W, Kindy MS, Peschon J, Kraemer PJ, Carpenter MK, Holtsberg FW, and Mattson MP. Altered neuronal and microglial responses to excitotoxic and ischemic brain injury in mice lacking TNF receptors. *Nat Med* 2: 788–794, 1996.
23. Candelario-Jalil E, Yang Y, and Rosenberg GA. Diverse roles of matrix metalloproteinases and tissue inhibitors of metalloproteinases in neuroinflammation and cerebral ischemia. *Neuroscience* 158: 983–994, 2009.
24. Chan PH. Reactive oxygen radicals in signaling and damage in the ischemic brain. *J Cereb Blood Flow Metab* 21: 2–14, 2001.
25. Chen F, Ohashi N, Li W, Eckman C, and Nguyen JH. Disruptions of occludin and claudin-5 in brain endothelial cells *in vitro* and in brains of mice with acute liver failure. *Hepatology* 50: 1914–1923, 2009.
26. Coleman JD, Prabhu KS, Thompson JT, Reddy PS, Peters JM, Peterson BR, Reddy CC, and Vanden Heuvel JP. The oxidative stress mediator 4-hydroxynonenal is an intracellular agonist of the nuclear receptor peroxisome proliferator-activated receptor-beta/delta (PPARbeta/delta). *Free Radic Biol Med* 42: 1155–1164, 2007.
27. Collison KS, Parhar RS, Saleh SS, Meyer BF, Kwaasi AA, Hammami MM, Schmidt AM, Stern DM, and Al-Mohanna FA. RAGE-mediated neutrophil dysfunction is evoked by advanced glycation end products (AGEs). *J Leukoc Biol* 71: 433–444, 2002.
28. Comporti M. Lipid peroxidation and biogenic aldehydes: From the identification of 4-hydroxynonenal to further achievements in biopathology. *Free Radic Res* 28: 623–635, 1998.
29. Cunningham LA, Wetzel M, and Rosenberg GA. Multiple roles for MMPs and TIMPs in cerebral ischemia. *Glia* 50: 329–339, 2005.
30. Daneman R, Zhou L, Agalliu D, Cahoy JD, Kaushal A, and Barres BA. The mouse blood–brain barrier transcriptome: A new resource for understanding the development and function of brain endothelial cells. *PLoS One* 5: e13741, 2010.
31. de Vries HE, Witte M, Hondius D, Rozemuller AJ, Drukarch B, Hoozemans J, and van Horssen J. Nrf2-induced antioxidant protection: A promising target to counteract ROS-mediated damage in neurodegenerative disease? *Free Radic Biol Med* 45: 1375–1383, 2008.
32. del Zoppo GJ and Mabuchi T. Cerebral microvessel responses to focal ischemia. *J Cereb Blood Flow Metab* 23: 879–894, 2003.
33. Dong X, Song YN, Liu WG, and Guo XL. Mmp-9, a potential target for cerebral ischemic treatment. *Curr Neuropharmacol* 7: 269–275, 2009.
34. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47–95, 2002.
35. Ebnet K. Organization of multiprotein complexes at cell–cell junctions. *Histochem Cell Biol* 130: 1–20, 2008.
36. Engelhardt B and Sorokin L. The blood–brain and the blood–cerebrospinal fluid barriers: Function and dysfunction. *Semin Immunopathol* 31: 497–511, 2009.
37. Esterbauer H, Benedetti A, Lang J, Fulceri R, Fauler G, and Comporti M. Studies on the mechanism of formation of 4-hydroxynonenal during microsomal lipid peroxidation. *Biochim Biophys Acta* 876: 154–166, 1986.
38. Esterbauer H, Cheeseman KH, Dianzani MU, Poli G, and Slater TF. Separation and characterization of the aldehydic products of lipid peroxidation stimulated by ADP-Fe²⁺ in rat liver microsomes. *Biochem J* 208: 129–140, 1982.
39. Evans JL, Goldfine ID, Maddux BA, and Grodsky GM. Oxidative stress and stress-activated signaling pathways: A unifying hypothesis of type 2 diabetes. *Endocr Rev* 23: 599–622, 2002.
40. Felinski EA and Antonetti DA. Glucocorticoid regulation of endothelial cell tight junction gene expression: Novel treatments for diabetic retinopathy. *Curr Eye Res* 30: 949–957, 2005.
41. Felinski EA, Cox AE, Phillips BE, and Antonetti DA. Glucocorticoids induce transactivation of tight junction genes occludin and claudin-5 in retinal endothelial cells via a novel cis-element. *Exp Eye Res* 86: 867–878, 2008.
42. Festa A, Schmolzer B, Schernthaner G, and Menzel EJ. Differential expression of receptors for advanced glycation end products on monocytes in patients with IDDM. *Diabetologia* 41: 674–680, 1998.
43. Forster C, Silwedel C, Golenhofen N, Burek M, Kietz S, Mankertz J, and Drenckhahn D. Occludin as direct target for glucocorticoid-induced improvement of blood–brain barrier properties in a murine *in vitro* system. *J Physiol* 565: 475–486, 2005.
- 43a. Frey T and Antonetti DA. Alterations to the blood–retinal barrier in diabetes: cytokines and reactive oxygen species. *Antioxid Redox Signal* 15: 1271–1284, 2011.
44. Fukumoto K, Takagi N, Yamamoto R, Moriyama Y, Takeo S, and Tanonaka K. Prostanoid EP1 receptor antagonist

- reduces blood-brain barrier leakage after cerebral ischemia. *Eur J Pharmacol* 640: 82–86, 2010.
45. Gao X, Xu X, Belmadani S, Park Y, Tang Z, Feldman AM, Chilian WM, and Zhang C. TNF- α contributes to endothelial dysfunction by upregulating arginase in ischemia/reperfusion injury. *Arterioscler Thromb Vasc Biol* 27: 1269–1275, 2007.
46. Gao X, Zhang H, Belmadani S, Wu J, Xu X, Elford H, Potter BJ, and Zhang C. Role of TNF- α -induced reactive oxygen species in endothelial dysfunction during reperfusion injury. *Am J Physiol Heart Circ Physiol* 295: H2242–2249, 2008.
47. Gasche Y, Soccal PM, Kanemitsu M, and Copin JC. Matrix metalloproteinases and diseases of the central nervous system with a special emphasis on ischemic brain. *Front Biosci* 11: 1289–1301, 2006.
48. Gatenby RA and Gillies RJ. Why do cancers have high aerobic glycolysis? *Nat Rev Cancer* 4: 891–899, 2004.
49. Gidday JM, Gasche YG, Copin JC, Shah AR, Perez RS, Shapiro SD, and Chan PH, Park TS. Leukocyte-derived matrix metalloproteinase-9 mediates blood-brain barrier breakdown and is proinflammatory after transient focal cerebral ischemia. *Am J Physiol Heart Circ Physiol* 289: H558–568, 2005.
50. Gilgun-Sherki Y, Melamed E, and Offen D. Oxidative stress induced-neurodegenerative diseases: The need for antioxidants that penetrate the blood-brain barrier. *Neuropharmacology* 40: 959–975, 2001.
51. Ginsberg MD. Neuroprotection for ischemic stroke: Past, present and future. *Neuropharmacology* 55: 363–389, 2008.
52. Gonzalez-Mariscal L, Tapia R, and Chamorro D. Crosstalk of tight junction components with signaling pathways. *Biochim Biophys Acta* 1778: 729–756, 2008.
- 52a. González-Mariscal L, Quirós M, and Díaz-Coránguez M. ZO proteins and redox-dependent processes. *Antioxid Redox Signal* 15: 1235–1253, 2011.
53. Gorlach A, Brandes RP, Nguyen K, Amidi M, Dehghani F, and Busse R. A gp91phox containing NADPH oxidase selectively expressed in endothelial cells is a major source of oxygen radical generation in the arterial wall. *Circ Res* 87: 26–32, 2000.
54. Groemping Y, Lapouge K, Smerdon SJ, and Rittinger K. Molecular basis of phosphorylation-induced activation of the NADPH oxidase. *Cell* 113: 343–355, 2003.
55. Grosch S, Fritz G, and Kaina B. Apurinic endonuclease (Ref-1) is induced in mammalian cells by oxidative stress and involved in clastogenic adaptation. *Cancer Res* 58: 4410–4416, 1998.
56. Gu Z, Kaul M, Yan B, Kridel SJ, Cui J, Strongin A, Smith JW, Liddington RC, and Lipton SA. S-nitrosylation of matrix metalloproteinases: Signaling pathway to neuronal cell death. *Science* 297: 1186–1190, 2002.
57. Guillemot L, Paschoud S, Pulimeno P, Foglia A, and Citi S. The cytoplasmic plaque of tight junctions: A scaffolding and signalling center. *Biochim Biophys Acta* 1778: 601–613, 2008.
58. Gutierrez J, Ballinger SW, Darley-Usmar VM, and Landar A. Free radicals, mitochondria, and oxidized lipids: The emerging role in signal transduction in vascular cells. *Circ Res* 99: 924–932, 2006.
59. Halliwell B. Oxidative stress and cancer: Have we moved forward? *Biochem J* 401: 1–11, 2007.
60. Haorah J, Ramirez SH, Schall K, Smith D, Pandya R, and Persidsky Y. Oxidative stress activates protein tyrosine kinase and matrix metalloproteinases leading to blood-brain barrier dysfunction. *J Neurochem* 101: 566–576, 2007.
61. Harhaj NS, Felinski EA, Wolpert EB, Sundstrom JM, Gardner TW, and Antonetti DA. VEGF activation of protein kinase C stimulates occludin phosphorylation and contributes to endothelial permeability. *Invest Ophthalmol Vis Sci* 47: 5106–5115, 2006.
62. Hattori I, Takagi Y, Nakamura H, Nozaki K, Bai J, Kondo N, Sugino T, Nishimura M, Hashimoto N, and Yodoi J. Intravenous administration of thioredoxin decreases brain damage following transient focal cerebral ischemia in mice. *Antioxid Redox Signal* 6: 81–87, 2004.
63. Hawkins BT and Davis TP. The blood-brain barrier/neurovascular unit in health and disease. *Pharmacol Rev* 57: 173–185, 2005.
64. Heyworth PG and Badwey JA. Protein phosphorylation associated with the stimulation of neutrophils. Modulation of superoxide production by protein kinase C and calcium. *J Bioenerg Biomembr* 22: 1–26, 1990.
65. Hicks K, O'Neil RG, Dubinsky WS, and Brown RC. TRPC-mediated actin-myosin contraction is critical for BBB disruption following hypoxic stress. *Am J Physiol Cell Physiol* 298: C1583–1593, 2010.
66. Hirata K, Ishida T, Penta K, Rezaee M, Yang E, Wohlge-muth J, and Quertermous T. Cloning of an immunoglobulin family adhesion molecule selectively expressed by endothelial cells. *J Biol Chem* 276: 16223–16231, 2001.
67. Hooper DC, Scott GS, Zborek A, Mikheeva T, Kean RB, Koprowski H, and Spitsin SV. Uric acid, a peroxynitrite scavenger, inhibits CNS inflammation, blood-CNS barrier permeability changes, and tissue damage in a mouse model of multiple sclerosis. *FASEB J* 14: 691–698, 2000.
68. Houstis N, Rosen ED, and Lander ES. Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature* 440: 944–948, 2006.
69. Huang CY, Fujimura M, Noshita N, Chang YY, and Chan PH. SOD1 down-regulates NF- κ B and c-Myc expression in mice after transient focal cerebral ischemia. *J Cereb Blood Flow Metab* 21: 163–173, 2001.
70. Huppert J, Closhen D, Croxford A, White R, Kulig P, Pietrowski E, Bechmann I, Becher B, Luhmann HJ, Waisman A, and Kuhlmann CR. Cellular mechanisms of IL-17-induced blood-brain barrier disruption. *FASEB J* 24: 1023–1034, 2010.
71. Ikenouchi J, Furuse M, Furuse K, Sasaki H, and Tsukita S. Tricellulin constitutes a novel barrier at tricellular contacts of epithelial cells. *J Cell Biol* 171: 939–945, 2005.
72. Ishihara H, Kubota H, Lindberg RL, Leppert D, Gloor SM, Errede M, Virgintino D, Fontana A, Yonekawa Y, and Frei K. Endothelial cell barrier impairment induced by glioblastomas and transforming growth factor beta2 involves matrix metalloproteinases and tight junction proteins. *J Neuropathol Exp Neurol* 67: 435–448, 2008.
73. Ishrat T, Sayeed I, Atif F, Hua F, and Stein DG. Progesterone and allopregnanolone attenuate blood-brain barrier dysfunction following permanent focal ischemia by regulating the expression of matrix metalloproteinases. *Exp Neurol* 226: 183–190, 2010.
74. Islas S, Vega J, Ponce L, and Gonzalez-Mariscal L. Nuclear localization of the tight junction protein ZO-2 in epithelial cells. *Exp Cell Res* 274: 138–148, 2002.

75. Jian Liu K and Rosenberg GA. Matrix metalloproteinases and free radicals in cerebral ischemia. *Free Radic Biol Med* 39: 71–80, 2005.
76. Johnson JA, Johnson DA, Kraft AD, Calkins MJ, Jakel RJ, Vargas MR, and Chen PC. The Nrf2-ARE pathway: An indicator and modulator of oxidative stress in neurodegeneration. *Ann NY Acad Sci* 1147: 61–69, 2008.
77. Jung KA and Kwak MK. The Nrf2 system as a potential target for the development of indirect antioxidants. *Molecules* 15: 7266–7291, 2010.
78. Juric-Sekhar G, Zarkovic K, Waeg G, Cipak A, Zarkovic N. Distribution of 4-hydroxynonenal-protein conjugates as a marker of lipid peroxidation and parameter of malignancy in astrocytic and ependymal tumors of the brain. *Tumori* 95: 762–768, 2009.
79. Kago T, Takagi N, Date I, Takenaga Y, Takagi K, and Takeo S. Cerebral ischemia enhances tyrosine phosphorylation of occludin in brain capillaries. *Biochem Biophys Res Commun* 339: 1197–1203, 2006.
80. Kahles T, Luedike P, Endres M, Galla HJ, Steinmetz H, Busse R, Neumann-Haefelin T, and Brandes RP. NADPH oxidase plays a central role in blood–brain barrier damage in experimental stroke. *Stroke* 38: 3000–3006, 2007.
81. Kaneto H, Matsuoka TA, Katakami N, Kawamori D, Miyatsuka T, Yoshiuchi K, Yasuda T, Sakamoto K, Yamasaki Y, and Matsuhisa M. Oxidative stress and the JNK pathway are involved in the development of type 1 and type 2 diabetes. *Curr Mol Med* 7: 674–686, 2007.
82. Karlhuber GM, Bauer HC, and Eckl PM. Cytotoxic and genotoxic effects of 4-hydroxynonenal in cerebral endothelial cells. *Mutat Res* 381: 209–216, 1997.
83. Kean RB, Spitsin SV, Mikheeva T, Scott GS, and Hooper DC. The peroxynitrite scavenger uric acid prevents inflammatory cell invasion into the central nervous system in experimental allergic encephalomyelitis through maintenance of blood–central nervous system barrier integrity. *J Immunol* 165: 6511–6518, 2000.
84. Kelly KA, Li X, Tan Z, VanGilder RL, Rosen CL, Huber JD. NOX2 inhibition with apocynin worsens stroke outcome in aged rats. *Brain Res* 1292: 165–172, 2009.
85. Kern TS. Contributions of inflammatory processes to the development of the early stages of diabetic retinopathy. *Exp Diabetes Res* 2007: 95103, 2007.
86. Kim HY and Han SH. Matrix metalloproteinases in cerebral ischemia. *J Clin Neurol* 2: 163–170, 2006.
87. Klaunig JE, Kamendulis LM, and Hocevar BA. Oxidative stress and oxidative damage in carcinogenesis. *Toxicol Pathol* 38: 96–109, 2010.
88. Kobayashi M and Yamamoto M. Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation. *Antioxid Redox Signal* 7: 385–394, 2005.
89. Kohen R and Nyska A. Oxidation of biological systems: Oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. *Toxicol Pathol* 30: 620–650, 2002.
90. Kondo N, Nakamura H, Masutani H, and Yodoi J. Redox regulation of human thioredoxin network. *Antioxid Redox Signal* 8: 1881–1890, 2006.
91. Kopnin PB, Agapova LS, Kopnin BP, and Chumakov PM. Repression of sestrin family genes contributes to oncogenic Ras-induced reactive oxygen species up-regulation and genetic instability. *Cancer Res* 67: 4671–4678, 2007.
92. Kowluru RA, Koppolu P, Chakrabarti S, and Chen S. Diabetes-induced activation of nuclear transcriptional factor in the retina, and its inhibition by antioxidants. *Free Radic Res* 37: 1169–1180, 2003.
93. Kowluru RA, Tang J, and Kern TS. Abnormalities of retinal metabolism in diabetes and experimental galactosemia. VII. Effect of long-term administration of antioxidants on the development of retinopathy. *Diabetes* 50: 1938–1942, 2001.
94. Krause G, Winkler L, Mueller SL, Haseloff RF, Piontek J, and Blasig IE. Structure and function of claudins. *Biochim Biophys Acta* 1778: 631–645, 2008.
95. Krizbai IA, Bauer H, Bresgen N, Eckl PM, Farkas A, Szatmari E, Traweger A, Wejksza K, and Bauer HC. Effect of oxidative stress on the junctional proteins of cultured cerebral endothelial cells. *Cell Mol Neurobiol* 25: 129–139, 2005.
96. Krueger M and Bechmann I. CNS pericytes: Concepts, misconceptions, and a way out. *Glia* 58: 1–10, 2010.
97. Kumar A, Mittal R, Khanna HD, and Basu S. Free radical injury and blood–brain barrier permeability in hypoxic-ischemic encephalopathy. *Pediatrics* 122: e722–727, 2008.
98. Lacraz G, Figeac F, Movassat J, Kassis N, Coulaud J, Galinier A, Leloup C, Bailbe D, Homo-Delarche F, and Portha B. Diabetic beta-cells can achieve self-protection against oxidative stress through an adaptive up-regulation of their antioxidant defenses. *PLoS One* 4: e6500, 2009.
99. Lakhani SE, Kirchgessner A, and Hofer M. Inflammatory mechanisms in ischemic stroke: Therapeutic approaches. *J Transl Med* 7: 97, 2009.
100. Liebner S, Cavallaro U, and Dejana E. The multiple languages of endothelial cell-to-cell communication. *Arterioscler Thromb Vasc Biol* 26: 1431–1438, 2006.
101. Liebner S, Fischmann A, Rascher G, Duffner F, Grote EH, Kalbacher H, and Wolburg H. Claudin-1 and claudin-5 expression and tight junction morphology are altered in blood vessels of human glioblastoma multiforme. *Acta Neuropathol* 100: 323–331, 2000.
102. Liu JF, Crepin M, Liu JM, Barritault D, and Ledoux D. FGF-2 and TPA induce matrix metalloproteinase-9 secretion in MCF-7 cells through PKC activation of the Ras/ERK pathway. *Biochem Biophys Res Commun* 293: 1174–1182, 2002.
103. Liu W, Hendren J, Qin XJ, Shen J, and Liu KJ. Normobaric hyperoxia attenuates early blood–brain barrier disruption by inhibiting MMP-9-mediated occludin degradation in focal cerebral ischemia. *J Neurochem* 108: 811–820, 2009.
104. Lochhead JJ, McCaffrey G, Quigley CE, Finch J, DeMarco KM, Nametz N, and Davis TP. Oxidative stress increases blood–brain barrier permeability and induces alterations in occludin during hypoxia-reoxygenation. *J Cereb Blood Flow Metab* 30: 1625–1636, 2010.
105. Lopez-Lazaro M. A new view of carcinogenesis and an alternative approach to cancer therapy. *Mol Med* 16: 144–153, 2010.
106. Machein MR and Plate KH. VEGF in brain tumors. *J Neurooncol* 50: 109–120, 2000.
107. Malemud CJ. Matrix metalloproteinases (MMPs) in health and disease: An overview. *Front Biosci* 11: 1696–1701, 2006.
108. Maritim AC, Sanders RA, and Watkins JB, 3rd. Diabetes, oxidative stress, and antioxidants: A review. *J Biochem Mol Toxicol* 17: 24–38, 2003.
109. Matoba S, Kang JG, Patino WD, Wragg A, Boehm M, Gavrilova O, Hurley PJ, Bunz F, and Hwang PM. p53 regulates mitochondrial respiration. *Science* 312: 1650–1653, 2006.
110. McCarthy SM, Bove PF, Matthews DE, Akaike T, and van der Vliet A. Nitric oxide regulation of MMP-9 activation

- and its relationship to modifications of the cysteine switch. *Biochemistry* 47: 5832–5840, 2008.
111. McCrea PD, Gu D, and Balda MS. Junctional music that the nucleus hears: Cell–cell contact signaling and the modulation of gene activity. *Cold Spring Harb Perspect Biol* 1: a002923, 2009.
112. Meli DN, Christen S, and Leib SL. Matrix metalloproteinase-9 in pneumococcal meningitis: Activation via an oxidative pathway. *J Infect Dis* 187: 1411–1415, 2003.
113. Mertsch K, Blasig I, and Grune T. 4-Hydroxynonenal impairs the permeability of an *in vitro* rat blood–brain barrier. *Neurosci Lett* 314: 135–138, 2001.
114. Michiels C, Minet E, Mottet D, and Raes M. Regulation of gene expression by oxygen: NF-kappaB and HIF-1, two extremes. *Free Radic Biol Med* 33: 1231–1242, 2002.
115. Minami M, Katayama T, and Satoh M. Brain cytokines and chemokines: Roles in ischemic injury and pain. *J Pharmacol Sci* 100: 461–470, 2006.
116. Minami M and Satoh M. Chemokines and their receptors in the brain: Pathophysiological roles in ischemic brain injury. *Life Sci* 74: 321–327, 2003.
117. Mizutani H, Tada-Oikawa S, Hiraku Y, Kojima M, and Kawanishi S. Mechanism of apoptosis induced by doxorubicin through the generation of hydrogen peroxide. *Life Sci* 76: 1439–1453, 2005.
118. Morita-Fujimura Y, Fujimura M, Gasche Y, Copin JC, and Chan PH. Overexpression of copper and zinc superoxide dismutase in transgenic mice prevents the induction and activation of matrix metalloproteinases after cold injury-induced brain trauma. *J Cereb Blood Flow Metab* 20: 130–138, 2000.
119. Murakami T, Felinski EA, and Antonetti DA. Occludin phosphorylation and ubiquitination regulate tight junction trafficking and vascular endothelial growth factor-induced permeability. *J Biol Chem* 284: 21036–21046, 2009.
120. Nag S, Picard P, and Stewart DJ. Expression of nitric oxide synthases and nitrotyrosine during blood–brain barrier breakdown and repair after cold injury. *Lab Invest* 81: 41–49, 2001.
121. Nagyoszi P, Wilhelm I, Farkas AE, Fazakas C, Dung NT, Hasko J, and Krizbai IA. Expression and regulation of toll-like receptors in cerebral endothelial cells. *Neurochem Int* 57: 556–564, 2010.
122. Nishino T. The conversion of xanthine dehydrogenase to xanthine oxidase and the role of the enzyme in reperfusion injury. *J Biochem* 116: 1–6, 1994.
123. Nishino T, Okamoto K, Eger BT, Pai EF, and Nishino T. Mammalian xanthine oxidoreductase: Mechanism of transition from xanthine dehydrogenase to xanthine oxidase. *FEBS J* 275: 3278–3289, 2008.
124. Novo E and Parola M. Redox mechanisms in hepatic chronic wound healing and fibrogenesis. *Fibrogenesis Tissue Repair* 1: 5, 2008.
125. Oesterreich S, Lee AV, Sullivan TM, Samuel SK, Davie JR, and Fuqua SA. Novel nuclear matrix protein HET binds to and influences activity of the HSP27 promoter in human breast cancer cells. *J Cell Biochem* 67: 275–286, 1997.
- 125a. Overgaard CE, Daugherty BL, Mitchell LA, and Koval M. Claudins: control of barrier function and regulation in response to oxidant stress. *Antioxid Redox Signal* 15: 1179–1193, 2011.
126. Pacher P, Nivorozhkin A, and Szabo C. Therapeutic effects of xanthine oxidase inhibitors: Renaissance half a century after the discovery of allopurinol. *Pharmacol Rev* 58: 87–114, 2006.
127. Panayiotidis M. Reactive oxygen species (ROS) in multi-stage carcinogenesis. *Cancer Lett* 266: 3–5, 2008.
128. Patenaude A, Murthy MR, AND Mirault ME. Emerging roles of thioredoxin cycle enzymes in the central nervous system. *Cell Mol Life Sci* 62: 1063–1080, 2005.
129. Phares TW, Fabis MJ, Brimer CM, Kean RB, AND Hooper DC. A peroxynitrite-dependent pathway is responsible for blood–brain barrier permeability changes during a central nervous system inflammatory response: TNF-alpha is neither necessary nor sufficient. *J Immunol* 178: 7334–7343, 2007.
130. Rabbani N and Thornalley PJ. Assay of 3-nitrotyrosine in tissues and body fluids by liquid chromatography with tandem mass spectrometric detection. *Methods Enzymol* 440: 337–359, 2008.
131. Ramana CV, Boldogh I, Izumi T, and Mitra S. Activation of apurinic/aprimidinic endonuclease in human cells by reactive oxygen species and its correlation with their adaptive response to genotoxicity of free radicals. *Proc Natl Acad Sci USA* 95: 5061–5066, 1998.
132. Rao GM, Rao AV, Raja A, Rao S, and Rao A. Role of antioxidant enzymes in brain tumours. *Clin Chim Acta* 296: 203–212, 2000.
133. Rao R. Oxidative stress-induced disruption of epithelial and endothelial tight junctions. *Front Biosci* 13: 7210–7226, 2008.
134. Reijerkerk A, Kooij G, van der Pol SM, Khazen S, Dijkstra CD, and de Vries HE. Diapedesis of monocytes is associated with MMP-mediated occludin disappearance in brain endothelial cells. *FASEB J* 20: 2550–2552, 2006.
135. Ridnour LA, Windhausen AN, Isenberg JS, Yeung N, Thomas DD, Vitek MP, Roberts DD, and Wink DA. Nitric oxide regulates matrix metalloproteinase-9 activity by guanylyl-cyclase-dependent and -independent pathways. *Proc Natl Acad Sci USA* 104: 16898–16903, 2007.
136. Rosenberg GA. Matrix metalloproteinases in neuroinflammation. *Glia* 39: 279–291, 2002.
137. Rosenberg GA and Yang Y. Vasogenic edema due to tight junction disruption by matrix metalloproteinases in cerebral ischemia. *Neurosurg Focus* 22: E4, 2007.
138. Russo MT, De Luca G, Degan P, and Bignami M. Different DNA repair strategies to combat the threat from 8-oxoguanine. *Mutat Res* 614: 69–76, 2007.
139. Ryu JK and McLarnon JG. Minocycline or iNOS inhibition block 3-nitrotyrosine increases and blood–brain barrier leakiness in amyloid beta-peptide-injected rat hippocampus. *Exp Neurol* 198: 552–557, 2006.
140. Sarsour EH, Kumar MG, Chaudhuri L, Kalen AL, and Goswami PC. Redox control of the cell cycle in health and disease. *Antioxid Redox Signal* 11: 2985–3011, 2009.
141. Saunders NR, Ek CJ, Habgood MD, and Dziegielewska KM. Barriers in the brain: A renaissance? *Trends Neurosci* 31: 279–286, 2008.
142. Schmidt AM, Yan SD, Yan SF, and Stern DM. The multi-ligand receptor RAGE as a progression factor amplifying immune and inflammatory responses. *J Clin Invest* 108: 949–955, 2001.
143. Schreibelt G, Kooij G, Reijerkerk A, van Doorn R, Gringhuis SI, van der Pol S, Weksler BB, Romero IA, Couraud PO, Piontek J, Blasig IE, Dijkstra CD, Ronken E, and de Vries HE. Reactive oxygen species alter brain endothelial

- tight junction dynamics via RhoA, PI3 kinase, and PKB signaling. *FASEB J* 21: 3666–3676, 2007.
144. Stadtman ER. Protein oxidation and aging. *Free Radic Res* 40: 1250–1258, 2006.
 145. Stadtman ER and Levine RL. Free radical-mediated oxidation of free amino acids and amino acid residues in proteins. *Amino Acids* 25: 207–18, 2003.
 146. Stadtman ER, Van Remmen H, Richardson A, Wehr NB, and Levine RL. Methionine oxidation and aging. *Biochim Biophys Acta* 1703: 135–140, 2005.
 147. Stalker TJ, Wu J, Morgans A, Traxler EA, Wang L, Chatterjee MS, Lee D, Quertermous T, Hall RA, Hammer DA, Diamond SL, and Brass LF. Endothelial cell specific adhesion molecule (ESAM) localizes to platelet-platelet contacts and regulates thrombus formation *in vivo*. *J Thromb Haemost* 7: 1886–1896, 2009.
 148. Steed E, Balda MS, and Matter K. Dynamics and functions of tight junctions. *Trends Cell Biol* 20: 142–149, 2010.
 149. Sternlicht MD and Werb Z. How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol* 17: 463–516, 2001.
 150. Sundaresan M, Yu ZX, Ferrans VJ, Sulciner DJ, Gutkind JS, Irani K, Goldschmidt-Clermont PJ, and Finkel T. Regulation of reactive-oxygen-species generation in fibroblasts by Rac1. *Biochem J* 318: 379–382, 1996.
 151. Suyama K, Shapiro I, Guttman M, and Hazan RB. A signaling pathway leading to metastasis is controlled by N-cadherin and the FGF receptor. *Cancer Cell* 2: 301–314, 2002.
 152. Svedin P, Hagberg H, Savman K, Zhu C, and Mallard C. Matrix metalloproteinase-9 gene knock-out protects the immature brain after cerebral hypoxia-ischemia. *J Neurosci* 27: 1511–1518, 2007.
 153. Swanson RA, Ying W, and Kauppinen TM. Astrocyte influences on ischemic neuronal death. *Curr Mol Med* 4: 193–205, 2004.
 154. Szabo C, Ischiropoulos H, and Radi R. Peroxynitrite: Biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov* 6: 662–680, 2007.
 155. Takenaga Y, Takagi N, Murotomi K, Tanonaka K, and Takeo S. Inhibition of Src activity decreases tyrosine phosphorylation of occludin in brain capillaries and attenuates increase in permeability of the blood-brain barrier after transient focal cerebral ischemia. *J Cereb Blood Flow Metab* 29: 1099–1108, 2009.
 156. Tan KH, Harrington S, Purcell WM, and Hurst RD. Peroxynitrite mediates nitric oxide-induced blood-brain barrier damage. *Neurochem Res* 29: 579–587, 2004.
 157. Tan S, Gelman S, Wheat JK, and Parks DA. Circulating xanthine oxidase in human ischemia reperfusion. *South Med J* 88: 479–482, 1995.
 158. Tapia R, Huerta M, Islas S, Avila-Flores A, Lopez-Bayghen E, Weiske J, Huber O, and Gonzalez-Mariscal L. Zona occludens-2 inhibits cyclin D1 expression and cell proliferation and exhibits changes in localization along the cell cycle. *Mol Biol Cell* 20: 1102–1117, 2009.
 159. Taylor CT and Cummins EP. The role of NF-kappaB in hypoxia-induced gene expression. *Ann NY Acad Sci* 1177: 178–184, 2009.
 160. Tell G, Quadrioglio F, Tiribelli C, and Kelley MR. The many functions of APE1/Ref-1: Not only a DNA repair enzyme. *Antioxid Redox Signal* 11: 601–620, 2009.
 161. Tempfer H, Wagner A, Gehwolf R, Lehner C, Tauber M, Resch H, and Bauer HC. Perivascular cells of the supraspinatus tendon express both tendon- and stem cell-related markers. *Histochem Cell Biol* 131: 733–741, 2009.
 162. Toda N, Ayajiki K, and Okamura T. Cerebral blood flow regulation by nitric oxide: Recent advances. *Pharmacol Rev* 61: 62–97, 2009.
 163. Toyokuni S. Novel aspects of oxidative stress-associated carcinogenesis. *Antioxid Redox Signal* 8: 1373–1377, 2006.
 164. Trachootham D, Lu W, Ogasawara MA, Nilsa RD, and Huang P. Redox regulation of cell survival. *Antioxid Redox Signal* 10: 1343–1374, 2008.
 165. Traweger A, Fuchs R, Krizbai IA, Weiger TM, Bauer HC, and Bauer H. The tight junction protein ZO-2 localizes to the nucleus and interacts with the heterogeneous nuclear ribonucleoprotein scaffold attachment factor-B. *J Biol Chem* 278: 2692–2700, 2003.
 166. Tsukita S, Furuse M, and Itoh M. Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* 2: 285–293, 2001.
 167. Vafa O, Wade M, Kern S, Beeche M, Pandita TK, Hampton GM, and Wahl GM. c-Myc can induce DNA damage, increase reactive oxygen species, and mitigate p53 function: a mechanism for oncogene-induced genetic instability. *Mol Cell* 9: 1031–1044, 2002.
 168. Valko M, Leibfriz D, Moncol J, Cronin MT, Mazur M, and Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 39: 44–84, 2007.
 169. Vallet P, Charnay Y, Steger K, Ogier-Denis E, Kovari E, Herrmann F, Michel JP, and Szanto I. Neuronal expression of the NADPH oxidase NOX4, and its regulation in mouse experimental brain ischemia. *Neuroscience* 132: 233–238, 2005.
 170. Van der Goes A, Wouters D, Van Der Pol SM, Huizinga R, Ronken E, Adamson P, Greenwood J, Dijkstra CD, and De Vries HE. Reactive oxygen species enhance the migration of monocytes across the blood-brain barrier *in vitro*. *FASEB J* 15: 1852–1854, 2001.
 171. Vepa S, Scribner WM, Parinandi NL, English D, Garcia JG, and Natarajan V. Hydrogen peroxide stimulates tyrosine phosphorylation of focal adhesion kinase in vascular endothelial cells. *Am J Physiol* 277: L150–158, 1999.
 172. Visse R and Nagase H. Matrix metalloproteinases and tissue inhibitors of metalloproteinases: Structure, function, and biochemistry. *Circ Res* 92: 827–839, 2003.
 173. Wang JY, Wen LL, Huang YN, Chen YT, and Ku MC. Dual effects of antioxidants in neurodegeneration: Direct neuroprotection against oxidative stress and indirect protection via suppression of glia-mediated inflammation. *Curr Pharm Des* 12: 3521–3533, 2006.
 174. Wang X and Feuerstein GZ. Induced expression of adhesion molecules following focal brain ischemia. *J Neurotrauma* 12: 825–832, 1995.
 175. Wang X, Yue TL, Barone FC, and Feuerstein GZ. Monocyte chemoattractant protein-1 messenger RNA expression in rat ischemic cortex. *Stroke* 26: 661–665; discussion 665–666, 1995.
 176. Watts LT, Rathinam ML, Schenker S, and Henderson GI. Astrocytes protect neurons from ethanol-induced oxidative stress and apoptotic death. *J Neurosci Res* 80: 655–666, 2005.
 177. Wegmann F, Ebnet K, Du Pasquier L, Vestweber D, and Butz S. Endothelial adhesion molecule ESAM binds directly to the multidomain adaptor MAGI-1 and recruits it to cell contacts. *Exp Cell Res* 300: 121–133, 2004.

178. Wegmann F, Petri B, Khandoga AG, Moser C, Khandoga A, Volkery S, Li H, Nasdala I, Brandau O, Fassler R, Butz S, Krombach F, and Vestweber D. ESAM supports neutrophil extravasation, activation of Rho, and VEGF-induced vascular permeability. *J Exp Med* 203: 1671–1677, 2006.
179. Wiensperger NF. Oxidative stress: The special case of diabetes. *Biofactors* 19: 11–18, 2003.
180. Winkler F, Koedel U, Kastenbauer S, and Pfister HW. Differential expression of nitric oxide synthases in bacterial meningitis: Role of the inducible isoform for blood–brain barrier breakdown. *J Infect Dis* 183: 1749–1759, 2001.
181. Wolburg H, Wolburg-Buchholz K, Kraus J, Rascher-Eggstein G, Liebner S, Hamm S, Duffner F, Grote EH, Risau W, and Engelhardt B. Localization of claudin-3 in tight junctions of the blood-brain barrier is selectively lost during experimental autoimmune encephalomyelitis and human glioblastoma multiforme. *Acta Neuropathol* 105: 586–592, 2003.
182. Wong CH and Crack PJ. Modulation of neuro-inflammation and vascular response by oxidative stress following cerebral ischemia-reperfusion injury. *Curr Med Chem* 15: 1–14, 2008.
183. Yang Y, Estrada EY, Thompson JF, Liu W, and Rosenberg GA. Matrix metalloproteinase-mediated disruption of tight junction proteins in cerebral vessels is reversed by synthetic matrix metalloproteinase inhibitor in focal ischemia in rat. *J Cereb Blood Flow Metab* 27: 697–709, 2007.
184. Yemisci M, Gursoy-Ozdemir Y, Vural A, Can A, Topalkara K, and Dalkara T. Pericyte contraction induced by oxidative-nitrative stress impairs capillary reflow despite successful opening of an occluded cerebral artery. *Nat Med* 15: 1031–1037, 2009.
185. Yilmaz G and Granger DN. Leukocyte recruitment and ischemic brain injury. *Neuromolecular Med* 12: 193–204, 2010.
186. Zhang C, Xu X, Potter BJ, Wang W, Kuo L, Michael L, Bagby GJ, and Chilian WM. TNF- α contributes to endothelial dysfunction in ischemia/reperfusion injury. *Arterioscler Thromb Vasc Biol* 26: 475–480, 2006.
187. Zhao BQ, Wang S, Kim HY, Storrie H, Rosen BR, Mooney DJ, Wang X, and Lo EH. Role of matrix metalloproteinases in delayed cortical responses after stroke. *Nat Med* 12: 441–445, 2006.
188. Zhao J, Kobori N, Aronowski J, and Dash PK. Sulforaphane reduces infarct volume following focal cerebral ischemia in rodents. *Neurosci Lett* 393: 108–112, 2006.
189. Zhao J, Moore AN, Redell JB, and Dash PK. Enhancing expression of Nrf2-driven genes protects the blood brain barrier after brain injury. *J Neurosci* 27: 10240–10248, 2007.
190. Zhao X and Benveniste EN. Transcriptional activation of human matrix metalloproteinase-9 gene expression by multiple co-activators. *J Mol Biol* 383: 945–956, 2008.
191. Zmijewski JW, Landar A, Watanabe N, Dickinson DA, Noguchi N, and Darley-USmar VM. Cell signalling by oxidized lipids and the role of reactive oxygen species in the endothelium. *Biochem Soc Trans* 33: 1385–1389, 2005.

Address correspondence to:

Dr. Hannelore Bauer

Department of Organismic Biology

Developmental Biology Group

Hellbrunner Strasse 34

University of Salzburg

5020 Salzburg

Austria

E-mail: hannelore.bauer@sbg.ac.at

Date of first submission to ARS Central, January 29, 2011;

date of acceptance, February 5, 2011.

Abbreviations Used

ARE = antioxidant response element
BBB = blood–brain barrier
cEC = cerebral endothelial cells
CNS = central nervous system
COX = cyclooxygenase
CSF = cerebrospinal fluid
ECM = extracellular matrix
eNOS = endothelial nitric oxide synthase
ESAM = endothelial cell-specific adhesion molecule
GBM = glioblastoma multiforme
GRx = glutathione reductase
GSHPx = glutathione peroxidase
HNE = hydroxynonenal
iNOS = inducible nitric oxide synthase
JAMs = junctional adhesion molecules
Keap1 = Kelch-like protein 1
nNOS = neuronal nitric oxide synthase
Nrf2 = nuclear erythroid related factor-2
RNS = reactive nitrogen species
ROS = reactive oxygen species
RP = retinopathy
SOD = superoxide dismutase
TJs = tight junctions
VEGF = vascular endothelial cell growth factor
XDH = xanthine dehydrogenase
XO = xanthine oxidase
XOR = xanthine oxidoreductase
ZO = zonula occludens

This article has been cited by:

1. Jin-Seok Lee, Hyung-Geug Kim, Jong-Min Han, Jong-Suk Lee, Seung-Wan Son, Yo-Chan Ahn, Chang-Gue Son. 2012. Myelophil ameliorates brain oxidative stress in mice subjected to restraint stress. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **39**:2, 339-347. [[CrossRef](#)]
2. Chao Liu, Jiliang Wu, Ming-Hui Zou. 2012. Activation of AMP-activated protein kinase alleviates High-glucose-induced dysfunction of brain microvascular endothelial cell tight-junction dynamics. *Free Radical Biology and Medicine* **53**:6, 1213-1221. [[CrossRef](#)]
3. Timo Kahles, Ralf P. Brandes. 2012. NADPH oxidases as therapeutic targets in ischemic stroke. *Cellular and Molecular Life Sciences* **69**:14, 2345-2363. [[CrossRef](#)]
4. Katherine R. Spindler, Tien-Huei Hsu. 2012. Viral disruption of the blood–brain barrier. *Trends in Microbiology* **20**:6, 282-290. [[CrossRef](#)]
5. Marta Skowrońska, Magdalena Zielińska, Luiza Wójcik-Stanaszek, Joanna Ruszkiewicz, Dejan Milatovic, Michael Aschner, Jan Albrecht. 2012. Ammonia increases paracellular permeability of rat brain endothelial cells by a mechanism encompassing oxidative/nitrosative stress and activation of matrix metalloproteinases. *Journal of Neurochemistry* no-no. [[CrossRef](#)]
6. Lorenza González-Mariscal , Miguel Quirós , Monica Díaz-Coránguez . ZO Proteins and Redox-Dependent Processes. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
7. Caroline Coisne , Britta Engelhardt . Tight Junctions in Brain Barriers During Central Nervous System Inflammation. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
8. Ingolf E. Blasig , Reiner F. Haseloff . Tight Junctions and Tissue Barriers. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
9. Christian E. Overgaard , Brandy L. Daugherty , Leslie A. Mitchell , Michael Koval . Claudins: Control of Barrier Function and Regulation in Response to Oxidant Stress. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
10. Gianfranco Bazzoni . Pathobiology of Junctional Adhesion Molecules. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
11. Anna Carrano , Jeroen J.M. Hoozemans , Saskia M. van der Vies , Annemieke J.M. Rozemuller , Jack van Horssen , Helga E. de Vries . Amyloid beta Induces Oxidative Stress-Mediated Blood–Brain Barrier Changes in Capillary Amyloid Angiopathy. *Antioxidants & Redox Signaling*, ahead of print. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]