

## Short communication

## Chemokine scavenging by the human cytomegalovirus chemokine decoy receptor US28 does not inhibit monocyte adherence to activated endothelium

J.M. Boomker<sup>a,b</sup>, E.K. de Jong<sup>a,b</sup>, L.F.M.H. de Leij<sup>a,b</sup>, M.C. Harmsen<sup>a,b,\*</sup><sup>a</sup> Department of Pathology and Laboratory Medicine, Medical Biology Section, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands<sup>b</sup> GUIDE, Groningen University Institute for Drug Exploration, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

Received 8 July 2005; accepted 8 November 2005

### Abstract

The human cytomegalovirus has found smart ways to exploit the chemokine network in order to subvert immune attack. Chemokines trigger the arrest and firm adhesion of inflammatory cells to the vascular wall. Scavenging of chemokines by viral decoy receptors, such as US28, might prevent arrest of leukocytes to the vascular wall and impair an antiviral immune response. We determined the effect of chemokine scavenging by endothelium-expressed signaling mute US28 (US28R129A) on static monocyte adhesion. Despite the chemokine scavenging capacity of US28R129A, expression of this construct by endothelial cells was insufficient to disrupt leukocyte adhesion to cytokine-activated monolayers. Our results suggest that the concentrations of chemokines that trigger firm leukocyte adhesion are too high to be efficiently scavenged by viral chemokine decoy receptors like US28. From the results of this experimental model a role for US28 in viral immune evasion by chemokine scavenging would appear therefore unlikely.

© 2005 Elsevier B.V. All rights reserved.

**Keywords:** Immune evasion; Human cytomegalovirus; Leukocyte adhesion; Chemokine receptor; Chemokine

Chemokines are important mediators of immunity against viral infections. They activate cytotoxic T lymphocytes and enhance the recruitment of inflammatory cells (Mahalingam et al., 2003; Salazar-Mather and Hokeness, 2003). Scavenging of chemokines by virus-encoded decoy receptors is a strategy to impair leukocyte recruitment and subsequent antiviral activity. The human cytomegalovirus gene US28 encodes a chemokine receptor homolog that internalizes  $\beta$ -chemokines with high efficiency in vitro (Billstrom et al., 1999; Bodaghi et al., 1998; Randolph-Habecker et al., 2002). Immobilized chemokines on the apical surface of activated endothelial cells trigger leukocytes to become adherent by inducing rapid integrin avidity and affinity changes (Alon and Feigelson, 2002; Johnston and Butcher, 2002). In monocytes,  $\beta$ -chemokines, such as MCP-1/CCL2, MIP-1 $\alpha$ /CCL3 and RANTES/CCL5, increase the expression of  $\beta$ 2 (CD18) integrins and facilitate adher-

ence (Vaddi and Newton, 1994). A chemokine gradient directs extravasated inflammatory cells towards the site of infection. Although chemokine scavenging was sufficient to reduce monocyte chemotaxis in vitro (Randolph-Habecker et al., 2002), the effect of chemokine scavenging by HCMV-infected endothelial cells on leukocyte arrest is not known.

In addition to chemokine scavenging, US28 also constitutively activates pro-inflammatory transcription factors such as NF- $\kappa$ B and CREB (Casarosa et al., 2001; Waldhoer et al., 2002), which might upregulate the expression of adhesion molecules, cytokines and chemokines that enable leukocyte adhesion. Expression of US28 in vascular smooth muscle cells results in a chemotactic response towards RANTES/CCL5 and MCP-1/CCL2 gradients (Streblow et al., 1999). The role of US28 in the viral life cycle remains unclear. More likely, chemokine scavenging and signaling are two different activities of US28 that occur simultaneously. The significance of signaling and scavenging by US28 may differ between different phases of the viral life cycle. Evasion of antiviral immune responses enables the virus to persist, whereas

\* Corresponding author. Tel.: +31 50 3614776; fax: +31 50 3619911.  
E-mail address: [m.c.harmsen@med.umcg.nl](mailto:m.c.harmsen@med.umcg.nl) (M.C. Harmsen).

for virus replication and dissemination immune activation is indispensable.

We aimed to investigate the effect of chemokine scavenging by US28 on leukocyte arrest. To exclude the influence of US28 signaling, we constructed a mutant receptor US28R129A that did not activate NF- $\kappa$ B (data not shown). Chemokine internalization was measured in confluent monolayers of CHO-k1 cells that were grown in 24 well plates. At 24 h after transfection, 44 nCi of  $^{125}$ I-labeled RANTES/CCL5 was added to each well and incubated for 30, 60, 90 and 120 min at 37 °C. To remove cell surface-bound chemokines, cells were washed with 0.01 M phosphate buffer followed by incubation with 200 mM acetic acid/50 mM NaCl, pH 2.5, for 5 min. Subsequently cells were lysed by addition of 0.5 M NaOH. Total activity in the cell lysate was quantified in an LKB multichannel counter. We found for both constructs that RANTES/CCL5 is rapidly sequestered within the intracellular fraction (Fig. 1A). The US28R129A appears to internalize RANTES/CCL5 more efficiently than the wild type US28. The uptake of RANTES/CCL5 is temperature-

sensitive as no uptake was found after incubation for 90 min at 4 °C. This indicates that chemokine uptake depends on active transport mechanism, and is not due to spontaneous diffusion of the protein (Fig. 1B). The observation that US28 rapidly internalizes chemokines and that mutation of the DRAI motif blocks G protein-mediated signaling is in concordance with published data (Michelson et al., 1997; Waldhoer et al., 2003). The US28R129A construct appeared to be slightly more efficient in chemokine internalization than the wild type receptor. This might reflect an increased recycling rate of the signaling mute receptor (Waldhoer et al., 2003) as transfection efficiencies were comparable for both constructs (~10%, data not shown).

For leukocyte adhesion studies, we selected the human premonocytic cell line MonoMac-6 as this cell line responds to  $\beta$ -chemokines (Cross et al., 1997), such as RANTES/CCL5. MonoMac-6 is also well characterized in a static adhesion assay (Erl et al., 1995). Furthermore, MonoMac-6 cells express the integrin Mac-1 that interacts with ICAM on endothelial cells that is required for sustained adherence (Erl et al., 1995). As a model for endothelial cells, we used primary cell cultures of human umbilical vein endothelial cells (HUVEC) in which we expressed the US28 gene through retroviral transduction. For the construction of retroviruses we used the bicistronic LZRS-linker-IRES-GFP vector in which the gene of interest was linked to a downstream internal ribosomal entry site (IRES) and the green fluorescent protein (GFP) marker gene that allowed independent translation of the products of both genes in the transduced target cells (Heemskerk et al., 1997). Recombinant retroviruses were produced after transfection of the 293T-based  $\Phi$ NX-A amphotropic packaging cell line (kindly provided by Dr. G. Nolan, Stanford University, Palo Alto, CA) with the LZRS-US28R129Amyc-IRES-GFP. HUVECs were grown to subconfluency and incubated for 18 h with undiluted retrovirus-containing supernatant containing 5  $\mu$ g/ml polybrene (Sigma-Aldrich). Efficiency of transduction was determined by measuring the number of GFP-positive cells by flow cytometry. Previously, it was shown that expression of US28 in HUVEC resulted in efficient chemokine internalization (Billstrom et al., 1999). The static adhesion assay was performed according to a protocol adapted from Molema et al. (1998). At 7 days after transduction, subconfluent monolayers of HUVECs were grown in 24 well plates (Corning, Cambridge, MA) and stimulated with 20 ng/ml TNF $\alpha$  (Boehringer Ingelheim GmbH, Basel, Swiss) overnight. For fluorescent staining, MonoMac-6 cells were loaded with 0.8  $\mu$ g/ml of 1,1'-diiodo-3,3',3'-tetramethylindocarbocyanide perchlorate (DiI) (Molecular Probes, OR) at 37 °C for 1 h in 5% CO<sub>2</sub> atmosphere and intensively washed before use. To each well 0.25  $\times 10^6$  MonoMac-6 cells were added. After incubation at 37 °C for 30 min, adherent cells were washed three times using pre-warmed PBS. HUVECs and adherent MonoMac-6 cells were harvested by adding trypsin/EDTA solution (0.5/0.2 mg/ml; Sigma-Aldrich); and short incubation at 37 °C. After adding 25  $\mu$ l of Flow Count Fluorospheres (Coulter Corporation, Miami) samples were analyzed by flow cytometry (Coulter EPICS Elite Flow Cytometer, Coulter Corporation). Based on their fluorescence, the number of




Fig. 1. RANTES binding and internalization by transfected CHO-k1 cells. Per 12 wells approximately 100 pg of  $^{125}$ I-RANTES was added. Activity of the internalized fraction was measured over time. Cells were incubated at 37 °C for 90 min (A). Furthermore, the ligand uptake was measured at 37 °C and 4 °C for 90 min (B). Indicated are the average and S.E.M. of three independent triplicate measurements.

adherent cells relative to the number of fluorospheres could be easily determined. RANTES/CCL5 responsiveness was determined by addition of increasing concentrations (10–100 ng/ml) of RANTES/CCL5 (Peprotech) to TNF $\alpha$ -activated HUVEC monolayers. For chemokine scavenging, RANTES/CCL5 was added 2 h prior to addition of MonoMac-6 cells.

Activated endothelial cells secrete chemokines that are sequestered by proteoglycans and trigger the arrest of circulating monocytes at the endothelial cell surface. Chemokine receptor signaling in monocytes increases affinity and avidity of integrins upon which they can interact with endothelial cell adhesion molecules. Chemokine scavenging at the cell surface may impair monocyte integrin avidity and affinity change and subsequent adhesion. To check whether the expression of US28 molecules was sufficient to reduce monocyte arrest we designed an experiment in which we validated the chemokine-induced adhesion and the impact of US28R129A expression. We found that only high concentrations of exogenously added RANTES/CCL5 triggered a significant increase of monocyte arrest (Fig. 2A). Subsequently, retroviruses were used to inte-

grate the US28R129A gene into the genome of HUVECs. Undiluted tissue culture supernatant from hybridoma 5/3–2.1 (IgG1) was used for the detection of ICAM. PE-conjugated goat-anti-mouse IgG was used as a secondary antibody. Mean fluorescence intensities were determined by flow cytometric analysis (Coulter EPICS Elite Flow Cytometer, Coulter Corporation).

Neither retrovirus transduction alone, nor the expression of US28R129A affected ICAM surface expression level of endothelial cells (Fig. 2B). For chemokine scavenging in the absence of US28 signaling, we transduced HUVECs with US28R129A-retrovirus. US28R129A expression on endothelial cells did not inhibit chemokine-induced MonoMac-6 adhesion (Fig. 2C). The chemokine scavenging potential of US28R129A was not sufficient to significantly reduce leukocyte adhesion to endothelial cells, even when a substantial number of cells was transduced ( $\sim 30\%$ , data not shown). In our static leukocyte adhesion model, we were able to show that our system is responsive to exogenously added chemokines (Fig. 2A). Recently, others have shown that leukocyte adhesion requires oligomerization of chemokines on cell-surface




Fig. 2. The effect of US28R129A chemokine scavenging on chemokine-induced adhesion of MonoMac-6 cells to HUVEC. Addition of increasing amounts of RANTES increased the number of adherent MonoMac-6 cells to TNF $\alpha$ -activated HUVEC monolayers. For the addition of 100 ng/ml RANTES the increase was statistically significant ( $p < 0.05$ ,  $n = 3$ ) (A). A bicistronic message encoding both the US28R129A gene and the green fluorescent protein (GFP) gene was delivered to the HUVECs by retrovirus transduction. Therefore, we assumed that all fluorescent cells also expressed the US28R129A molecule. To investigate whether retrovirus transduction by itself affected ICAM-1 expression levels, the level of cell-surface ICAM-1 was determined by flow cytometry using PE-conjugated monoclonal antibody against ICAM-1. No differences between retrovirus transduced and non-transduced cells were found. Nor did the presence of US28R129A affect the level of ICAM-1 expression ( $n = 1$ ) (B). For the chemokine scavenging assay, transduced HUVEC ( $\sim 40\%$ ) were loaded with 100 ng/ml RANTES 2 h before addition of MonoMac-6 cells. No effect of chemokine scavenging on MonoMac-6 adhesion was found ( $n = 5$ ) (C). All experiments were repeated twice.

expressed proteoglycans which usually occurs at very high chemokine concentrations (Appay et al., 1999). If that is the case, it is unlikely chemokine scavenging by US28R129A can contribute to reduced oligomerization and subsequent adhesion triggering. Moreover, in vitro studies have shown that chemokine-triggered integrin avidity changes occur in subseconds (Grabovsky et al., 2000). Chemokine scavenging may, on its best, delay leukocyte arrest rather than reduce the actual number of adherent leukocytes. We show that at least in vitro, the chemokine scavenging capacity of US28 is insufficient to affect leukocyte arrest. This would indicate that during HCMV infection of endothelial cells, US28 does not play an important role as a chemokine scavenger.

## Acknowledgements

The authors like to thank Henk Moorlag for the isolation and culturing of human umbilical vein endothelial cells and Geert Mesander for the technical assistance with the flow cytometry.

## References

- Alon, R., Feigelson, S., 2002. From rolling to arrest on blood vessels: leukocyte tap dancing on endothelial integrin ligands and chemokines at sub-second contacts. *Semin. Immunol.* 14, 93–104.
- Appay, V., Brown, A., Cribbes, S., Randle, E., Czaplewski, L.G., 1999. Aggregation of RANTES is responsible for its inflammatory properties. Characterization of nonaggregating, noninflammatory RANTES mutants. *J. Biol. Chem.* 274, 27505–27512.
- Billstrom, M.A., Lehman, L.A., Scott Worthen, G., 1999. Depletion of extracellular RANTES during human cytomegalovirus infection of endothelial cells. *Am. J. Respir. Cell Mol. Biol.* 21, 163–167.
- Bodaghi, B., Jones, T.R., Zipeto, D., Vita, C., Sun, L., Laurent, L., Arenzana-Seisdedos, F., Virelizier, J.L., Michelson, S., 1998. Chemokine sequestration by viral chemoreceptors as a novel viral escape strategy: withdrawal of chemokines from the environment of cytomegalovirus-infected cells. *J. Exp. Med.* 188, 855–866.
- Casarosa, P., Bakker, R.A., Verzijl, D., Navis, M., Timmerman, H., Leurs, R., Smit, M.J., 2001. Constitutive signaling of the human cytomegalovirus-encoded chemokine receptor US28. *J. Biol. Chem.* 276, 1133–1137.
- Cross, A.K., Richardson, V., Ali, S.A., Palmer, I., Taub, D.D., Rees, R.C., 1997. Migration responses of human monocytic cell lines to alpha- and beta-chemokines. *Cytokine* 9, 521–528.
- Erl, W., Weber, C., Wardemann, C., Weber, P.C., 1995. Adhesion properties of MonoMac 6, a monocytic cell line with characteristics of mature human monocytes. *Atherosclerosis* 113, 99–107.
- Grabovsky, V., Feigelson, S., Chen, C., Bleijs, D.A., Peled, A., Cinamon, G., Baleux, F., Arenzana-Seisdedos, F., Lapidot, T., van Kooyk, Y., Lobb, R.R., Alon, R., 2000. Subsecond induction of alpha4 integrin clustering by immobilized chemokines stimulates leukocyte tethering and rolling on endothelial vascular cell adhesion molecule 1 under flow conditions. *J. Exp. Med.* 192, 495–506.
- Heemskerk, M.H., Blom, B., Nolan, G., Stegmann, A.P., Bakker, A.Q., Weijer, K., Res, P.C., Spits, H., 1997. Inhibition of T cell and promotion of natural killer cell development by the dominant negative helix loop helix factor Id3. *J. Exp. Med.* 186, 1597–1602.
- Johnston, B., Butcher, E.C., 2002. Chemokines in rapid leukocyte adhesion triggering and migration. *Semin. Immunol.* 14, 83–92.
- Mahalingam, S., Friedland, J.S., Heise, M.T., Rulli, N.E., Meanger, J., Lidbury, B.A., 2003. Chemokines and viruses: friends or foes? *Trends Microbiol.* 11, 383–391.
- Michelson, S., Dal Monte, P., Zipeto, D., Bodaghi, B., Laurent, L., Oberlin, E., Arenzana-Seisdedos, F., Virelizier, J.L., Landini, M.P., 1997. Modulation of RANTES production by human cytomegalovirus infection of fibroblasts. *J. Virol.* 71, 6495–6500.
- Molema, G., Mesander, G., Kroesen, B.J., Helfrich, W., Meijer, D.K., de Leij, L.F., 1998. Analysis of in vitro lymphocyte adhesion and transendothelial migration by fluorescent-beads-based flow cytometric cell counting. *Cytometry* 32, 37–43.
- Randolph-Habecker, J.R., Rahill, B., Torok-Storb, B., Vieira, J., Kolattukudy, P.E., Rovin, B.H., Sedmak, D.D., 2002. The expression of the cytomegalovirus chemokine receptor homolog US28 sequesters biologically active CC chemokines and alters IL-8 production. *Cytokine* 19, 37–46.
- Salazar-Mather, T.P., Hokeness, K.L., 2003. Calling in the troops: regulation of inflammatory cell trafficking through innate cytokine/chemokine networks. *Viral Immunol.* 16, 291–306.
- Streblow, D.N., Soderberg-Naucler, C., Vieira, J., Smith, P., Wakabayashi, E., Ruchti, F., Mattison, K., Altschuler, Y., Nelson, J.A., 1999. The human cytomegalovirus chemokine receptor US28 mediates vascular smooth muscle cell migration. *Cell* 99, 511–520.
- Vaddi, K., Newton, R.C., 1994. Regulation of monocyte integrin expression by beta-family chemokines. *J. Immunol.* 153, 4721–4732.
- Waldhoer, M., Casarosa, P., Rosenkilde, M.M., Smit, M.J., Leurs, R., Whistler, J.L., Schwartz, T.W., 2003. The carboxyl terminus of human cytomegalovirus-encoded 7 transmembrane receptor US28 camouflages agonism by mediating constitutive endocytosis. *J. Biol. Chem.* 278, 19473–19482.
- Waldhoer, M., Kledal, T.N., Farrell, H., Schwartz, T.W., 2002. Murine cytomegalovirus (CMV) M33 and human CMV US28 receptors exhibit similar constitutive signaling activities. *J. Virol.* 76, 8161–8168.