

to the normal squamous epithelial tissues. Typical cytopathic changes identical to those found in the squamous epithelium in vivo were observed throughout the raft following infection with the different viruses. We have now assessed the feasibility of using mononuclear cells (MCs) as viral carriers to infect organotypic epithelial raft cultures. For this purpose, mononuclear cells were isolated from human umbilical cord blood by Ficoll-Hypaque density gradient centrifugation, stimulated with phytohemagglutinin and incubated in the presence of IL-2. MCs were infected with HSV-1, HSV-2 or VV at a multiplicity of infection of approximately 0.01, incubated overnight and then washed to remove the viral inoculum. The MCs were added on top of the raft cultures after the epithelial cells were allowed to differentiate for 6 days. MCs were able to transfer the virus to the epithelium and histological changes characteristic of HSV infection could be observed. When the anti-herpesvirus agents, acyclovir, penciclovir, HPMP (cidofovir) or HPMP-5azaC were added to the culture media at the time the MCs carrying HSV were added on top of the rafts, the epithelium was protected from the virus-induced cytopathic effect in a dose-dependent manner. The antiviral activity was quantified by measuring viral titers by plaque assay. Similarly, HPMP and HPMP-5-azaC were able to protect the epithelial cells from VV spread by the MCs. Our results show the possibility of using MCs as a transfer vehicle for HSV and VV infections of epithelial cells grown in 3D and suggest that MCs could be responsible for the spread of the virus infection.

doi:10.1016/j.antiviral.2007.01.041

### 34

#### Structural Basis for the Expanded Substrate Specificity of Vaccinia Virus Thymidine Kinase: Insight from the Crystal Structure

Mark Prichard<sup>1,\*</sup>, Kathy Keith<sup>1</sup>, Mary Johnson<sup>1</sup>, Norbert Schormann<sup>2</sup>, Alexandra Samal<sup>2</sup>, Debasish Chattopadhyay<sup>2,3</sup>, Earl Kern<sup>1</sup>

<sup>1</sup> Department of Pediatrics, University of Alabama School of Medicine, USA; <sup>2</sup> Center for Biophysical Sciences & Engineering, University of Alabama School of Medicine, USA; <sup>3</sup> Department of Medicine, University of Alabama School of Medicine, USA

Vaccinia virus, like other orthopoxviruses, encodes an enzyme with thymidine kinase (TK) activity. This enzyme is closely related to the human cytosolic enzyme, TK1, and both belong to the type 2 family of TK homologs. Previous studies with *N*-methanocarbathymidine (*N*-MCT), a new antiviral agent with activity against cowpox virus replication, demonstrated that the compound was much less effective in TK negative strains of the virus and suggested that the enzyme could preferentially phosphorylate this compound. A similar TK dependence was also observed with a series of 5-substituted deoxyuridine analogs. These results were unanticipated since the viral TK and TK1 share 70% identity at the amino acid level and were predicted to exhibit very similar substrate specificities. This was investigated

further by purifying both enzymes in bacteria and comparing their relative ability to phosphorylate a number of thymidine analogs. This analysis revealed that the viral enzyme exhibited a marked preference for a few compounds including *N*-MCT and fialuridine. The crystal structure of recombinant TK at 2.9 Å resolution explains the structural basis for these differences. A comparison of this structure with the previously published structure of human TK1 revealed significant differences in the catalytic site. In particular, the active site of the viral enzyme appeared to be more open than the human homolog. These results have important implications in the development of antiviral therapies to orthopoxvirus infections. Thymidine analogs, such as (*N*)-MCT that are selectively phosphorylated by this enzyme in infected cells could be developed as potent and highly selective inhibitors of orthopoxvirus infections.

doi:10.1016/j.antiviral.2007.01.042

### 35

#### Inhibition of an Innate Antiviral Response by Human Cytomegalovirus UL97 Kinase is Antagonized by Maribavir

Mark Prichard\*, Shannon Daily, Amie Perry, Carol Hartline, Earl Kern

Department of Pediatrics, University of Alabama School of Medicine, USA

Nuclear aggresomes are cellular structures that function to degrade highly ordered protein aggregates in the nucleus, such as viral structural proteins and are an important innate antiviral response. The formation of nuclear aggresomes occurs around PML oncogenic domains (PODs, ND10 sites). Many viruses including human cytomegalovirus disrupt PODs, presumably to protect against the sequestration and degradation of viral proteins in the nucleus. One previously described facet of the UL97 negative phenotype was the formation of large nuclear aggregates, which also formed when cells were infected with the wt virus in the presence of the UL97 kinase inhibitor, maribavir (MBV). Interestingly, the kinase was shown to inhibit the nuclear aggregation of pp65 when both proteins were expressed transiently in COS7 cells. Here, we report that the kinase also reduced the nuclear aggregation of the tegument protein pp71, as well as cellular protein that is a marker for nuclear aggresomes. Thus, it appeared that the inhibition of nuclear aggregations was more general than previously suspected. To investigate this further, we examined the effect of the kinase on PODs and observed that it disrupted their formation in a manner similar to IE1, which was used as a positive control. This effect did not occur either with a point mutant of UL97 that has no kinase activity, or if the kinase activity was inhibited by MBV. These results taken together suggest that one important function of UL97 is to inhibit the formation of PODs and the degradation of structural proteins by nuclear aggresomes. In the absence of UL97 kinase activity, at least 90% of the mass of viral structural proteins is sequestered in nuclear aggresomes resulting in a significant impairment of viral morphogenesis. We conclude that the inhibition of the kinase by MBV impairs the ability of the virus to inactivate this innate

antiviral response and is a novel aspect of its mechanism of action.

doi:10.1016/j.antiviral.2007.01.043

### 36

#### **Vaccination as an Antiviral Strategy for Control of Cytomegalovirus (CMV) Disease: A Vectored Vaccine Approach Targeting the UL83 (pp65) Homolog Protects Against Congenital CMV Disease in the Guinea Pig Model**

Mark Schleiss<sup>1,\*</sup>, Juan Lacayo<sup>1</sup>, Yasmine Belkaid<sup>2</sup>, Alistair McGregor<sup>1</sup>, Greg Stroup<sup>3</sup>, Jon Rayner<sup>4</sup>, Kim Alterson<sup>4</sup>, Jeff Chulay<sup>4</sup>, Jonathan Smith<sup>4</sup>

<sup>1</sup> Center for Infectious Diseases and Microbiology Translational Research, University of Minnesota, Department of Pediatrics, Minneapolis, MN, USA; <sup>2</sup> Mucosal Immunology Unit, NIAID, National Institutes of Health, Bethesda, MD, USA; <sup>3</sup> Meridian Biosciences, Cincinnati, OH, USA; <sup>4</sup> AlphaVax, Research Triangle Park, NC, USA

Development of vaccines against cytomegalovirus (CMV) infection is a major priority, both for high-risk immunosuppressed patients, and to protect against disabling congenital infections. Glycoprotein subunit vaccines have demonstrated efficacy against disease in small animal models of congenital CMV infection, but to date there have been no reports of the use of vaccines targeting tegument proteins such as pp65 (UL83), the major target of anti-CMV cytotoxic T-lymphocyte responses, in this setting. We report the use of a propagation-defective, single cycle, RNA replicon vector system, derived from an attenuated strain of the alphavirus Venezuelan equine encephalitis (VEE) virus, to produce virus-like replicon particles (VRP) expressing GP83, the guinea pig CMV (GPCMV) homolog of the human CMV pp65 phosphoprotein encoded by the UL83 gene. Vaccination with VRP GP83 induced antibodies and CD4+ and CD8+ T cell responses in GPCMV-seronegative female guinea pigs. Guinea pigs immunized with VRP-GP83 vaccine, or a VRP vaccine expressing influenza hemagglutinin (VRP-HA), were bred for pregnancy and subsequent GPCMV challenge. After early third trimester challenge with GPCMV, dams previously vaccinated with VRP-GP83 had improved pregnancy outcomes compared to dams previously vaccinated with the VRP-HA control. Among VRP-GP83-vaccinated dams, there were 28 live pups, and 4 dead pups (13% mortality) among 10 evaluable litters, compared to 9 live pups and 12 dead pups (57% mortality) among 8 evaluable litters in the VRP HA vaccinated group ( $p < 0.001$ , Fisher's exact test). The improved pregnancy outcome was accompanied by reduction in maternal whole blood viral load as measured by real-time PCR. These results indicate that cell-mediated immune responses directed against a CMV matrix protein can protect against congenital CMV infection.

doi:10.1016/j.antiviral.2007.01.044

### 37

#### **JuvImmune is a Novel Vaccine Adjuvant that Enhances Protection of Mice from Lethal HSV-2 Infection Following Immunization**

Jane Strasser<sup>1,\*</sup>, Rhonda Cardin<sup>1</sup>, Claudia Chalk<sup>1</sup>, James Ireland<sup>1</sup>, Jeff Fairman<sup>2</sup>, David Bernstein<sup>1</sup>

<sup>1</sup> Division of Infectious Diseases, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA; <sup>2</sup> Juvaris BioTherapeutics, Inc., Pleasanton, CA, USA

Genital herpes simplex type 2 (HSV-2) infection is common and an effective vaccine remains a priority. We therefore examined the efficacy of JuvImmune as a vaccine adjuvant against genital HSV-2 infection in mice. JuvImmune, a complex of lipid carrier and non-coding DNA, enhances protection against several viral and bacterial diseases. In our studies, out-bred Swiss Webster mice (12 animals/group) were immunized IP twice at 3-week intervals with a detergent-inactivated whole HSV-2 vaccine or controls. Groups included placebo, vaccine (VAC) alone, JuvImmune alone, JuvImmune + VAC or MPL + VAC. Three weeks post boost, mice were treated with progesterone and given a lethal intravaginal challenge of HSV-2 ( $5 \times 10^5$  pfu). Vaginal swabs were collected on days 1–4 post challenge and serum was collected at 3 and 5 weeks post immunization. Our preliminary results indicate that animals immunized with JuvImmune + VAC developed higher antibody titers compared to VAC alone or MPL + VAC after the first immunization. Mice were followed for 21 or 30 days in two separate experiments. The JuvImmune + VAC immunization reduced HSV-2 disease symptoms compared to MPL + VAC. In the first experiment, JuvImmune + VAC completely protected mice from death at 21 days compared to a mortality of 40% in the MPL + VAC group ( $P < 0.037$ ). In the second experiment, JuvImmune + VAC protected 83% of the mice from death at day 21, and 75% of mice at 30 days post challenge, compared to only 17% at both 21 days ( $P < 0.003$ ) and 30 days ( $P > 0.036$ ) in the MPL + VAC group. In both experiments, vaginal titers were also reduced at day 2 in the JuvImmune + VAC group compared to all other groups. These results suggest that the JuvImmune adjuvant is superior to the MPL adjuvant in protecting mice from genital HSV-2 infection. Further studies are planned to determine the mechanisms of protection provided by the JuvImmune adjuvant.

**Acknowledgement:** Supported by Contract NO1-AI-15438 from the Virology Branch, NIAID, NIH.

doi:10.1016/j.antiviral.2007.01.045