

# A New Oxygen Sensitivity and Its Potential Application in Photosynthetic H<sub>2</sub> Production

JAMES W. LEE\* AND ELIAS GREENBAUM

*Chemical Sciences Division, Oak Ridge National Laboratory,  
Oak Ridge, TN 37831-6194, E-mail: Leejw@ORNL.gov*

## Abstract

We have discovered a new competitive pathway for O<sub>2</sub> sensitivity in algal H<sub>2</sub> production that is distinct from the O<sub>2</sub> sensitivity of hydrogenase *per se*. This O<sub>2</sub> sensitivity is apparently linked to the photosynthetic H<sub>2</sub> production pathway that is coupled to proton translocation across the thylakoid membrane. Addition of the proton uncoupler carbonyl cyanide-p-trifluoromethoxy-phenylhydrazone eliminates this mode of O<sub>2</sub> inhibition on H<sub>2</sub> photoevolution. This newly discovered inhibition is most likely owing to background O<sub>2</sub> that apparently serves as a terminal electron acceptor in competition with the H<sub>2</sub> production pathway for photosynthetically generated electrons from water splitting. This O<sub>2</sub>-sensitive H<sub>2</sub> production electron transport pathway was inhibited by 3[3,4-dichlorophenyl]1,1-dimethylurea. Our experiments demonstrated that this new pathway is more sensitive to O<sub>2</sub> than the traditionally known O<sub>2</sub> sensitivity of hydrogenase. This discovery provides new insight into the mechanism of O<sub>2</sub> inactivation of hydrogenase and may contribute to the development of a more-efficient and robust system for photosynthetic H<sub>2</sub> production.

**Index Entries:** Oxygen sensitivity; H<sub>2</sub> production; photosynthetic H<sub>2</sub> production; H<sub>2</sub> production pathways; hydrogenase.

## Introduction

Algal photosynthetic hydrogen (H<sub>2</sub>) production by light-activated water splitting is a potentially clean energy resource. However, compared to our knowledge of the pathway of atmospheric CO<sub>2</sub> reduction, and in spite of the potential importance of the hydrogen-producing reaction, relatively little is known concerning the mechanistic pathway of electron flow in hydrogen-producing algae. In green algae, such as *Chlamydomonas reinhardtii*, photoevolution of H<sub>2</sub> and O<sub>2</sub> occurs in the same cell, where the photosynthetically produced O<sub>2</sub> can inhibit the production of H<sub>2</sub> (1). There-

\*Author to whom all correspondence and reprint requests should be addressed.

fore, the application of green algae for  $H_2$  production must address the problem of  $O_2$  sensitivity. Historically, this  $O_2$ -sensitive phenomenon was generally interpreted as the direct  $O_2$  inhibition of hydrogenase activity (2). We report here that the classic interpretation of  $O_2$  sensitivity should be revised. In recent experiments characterizing  $O_2$  tolerance in  $H_2$ -producing wild-type *C. reinhardtii*, we observed a new  $O_2$  sensitivity that is clearly distinct from that of classic  $O_2$  inhibition of hydrogenase. The  $O_2$  sensitivity indicates that there is a competitive electron transport pathway that can redirect electrons from the hydrogenase-catalyzed  $H_2$  production pathway to  $O_2$ . That is to say, suppression of  $H_2$  evolution in the presence of low-level background concentrations of  $O_2$  is owing to the drain of reducing equivalents away from the hydrogenase pathway and toward the reduction of  $O_2$ . Our experiments demonstrated that the competitive pathway mechanism is more sensitive to  $O_2$  than the classic  $O_2$  sensitivity of hydrogenase and can be suppressed by proton uncoupler carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP).

## Materials and Methods

First evidence of the new  $O_2$  sensitivity was obtained from  $H_2$  production assays in *C. reinhardtii* wild-type strain 137c. The assays were conducted using a laboratory-built dual-reactor flow detection system (3). For each assay, 35 mL of algal sample (3  $\mu\text{g Chl/mL}$ ) was placed and sealed in each of the two water-jacketed reactors and held at 20°C with a temperature-controlled water bath. The algal sample was purged with a helium flow (50 mL/min) through the liquid reaction medium. The helium flow served two purposes: (1) to remove  $O_2$  from the algal sample to establish and maintain anaerobic conditions that are necessary for induction of the algal hydrogenase activity and production of  $H_2$ , and (2) to carry the photoproduct  $H_2$  gas product to the hydrogen sensors. After induction of hydrogenase and establishment of steady-state photoevolution of  $H_2$  under the helium atmosphere (which requires about 8 h), a primary standard of 1000 ppm of  $O_2$  in helium replaced the pure helium at the same flow rate (50 mL/min) to characterize the oxygen sensitivity of photoevolution of  $H_2$ . Actinic illumination of 100  $\mu\text{E}/(\text{m}^2\cdot\text{s})$  for the  $H_2$  photoevolution assay was provided by an electronically controlled light emitting diode (LED) light source at a wavelength of 670 nm. As illustrated in Fig. 1, introduction of 0.1000% (1000 ppm) of  $O_2$  dramatically reduced the rate of algal  $H_2$  photoevolution. The steady-state  $H_2$  production rate in the presence of 0.1000%  $O_2$  was 0.33  $\mu\text{mol of } H_2\cdot\text{mg of Chl}^{-1}\cdot\text{h}^{-1}$ , which is only about 2.8% of the full steady-state rate (12  $\mu\text{mol } H_2\cdot\text{mg of Chl}^{-1}\cdot\text{h}^{-1}$ ) before the introduction of the 0.1000%  $O_2$ . In the past, this type of  $H_2$  production decay was commonly interpreted as the inhibition of hydrogenase activity by  $O_2$ . Our results prove that this classic interpretation of oxygen sensitivity on algal  $H_2$  production is not consistent with the data. According to the classic interpretation, the decrease in  $H_2$  production after the introduction of 0.1000%




Fig. 1. Observation of a new  $O_2$  sensitivity to algal  $H_2$  production in *C. reinhardtii*.

$O_2$  is owing to  $O_2$  inhibition of hydrogenase *per se*: that is, loss of hydrogenase activity is the limiting factor for the rate of  $H_2$  photoevolution. If this interpretation were correct, after a brief dark period in the presence of 0.1000%  $O_2$ , one would expect the rate of  $H_2$  photoevolution to be no higher than the inhibited rate ( $0.33 \mu\text{mol of } H_2 \cdot \text{mg of Chl}^{-1} \cdot \text{h}^{-1}$ ) preceding the dark interval. However, the data are quite different from the classic expectation. As shown in Fig. 1, there was a surge of  $H_2$  photoevolution after a 2-h dark period in the continuous presence of 0.1000%  $O_2$ . The peak rate of  $H_2$  photoevolution was about  $15 \mu\text{mol of } H_2 \cdot \text{mg of Chl}^{-1} \cdot \text{h}^{-1}$ , which is about 45 times higher than the classically predicted rate ( $0.33 \mu\text{mol of } H_2 \cdot \text{mg of Chl}^{-1} \cdot \text{h}^{-1}$ ) and well outside the experimental error of the measurement. This assay has now been repeated six times, and all results were consistent with the observation presented in Fig. 1.

This observation clearly indicated that hydrogenase activity was not the limiting factor for  $H_2$  photoevolution at this level of  $O_2$ . There must be an alternative electron transport pathway that takes the photogenerated electrons away from ferredoxin (Fd) to  $O_2$ . The observed reduction of  $H_2$  production after the introduction of 0.1000%  $O_2$  can be explained by such a pathway that competes for electrons with the Fd/hydrogenase-catalyzed  $H_2$  production pathway. This is a significant discovery since it fundamentally redefines the meaning of “oxygen tolerance” in algal  $H_2$  production.




Fig. 2. Stimulation of photosynthetic  $H_2$  production of *C. reinhardtii* 137c following addition of proton uncoupler FCCP in background atmosphere of 1000 ppm of  $O_2$ . Addition of  $5 \mu M$  FCCP produced a dramatic increase in  $H_2$  production followed by a slow decay. The slow decay was owing to a side effect of FCCP known as ADRY, in which FCCP gradually inhibits PSII activity. This experimental result indicates that use of a polypeptide proton channel that does not have the ADRY effect could enhance  $H_2$  production by eliminating the problems of both the proton gradient accumulation and the newly discovered alternative  $O_2$  sensitivity.

Studies with the chemical inhibitor 3[3,4-dichlorophenyl]1,1-dimethylurea (DCMU) and the proton uncoupler FCCP yielded additional evidence for the new  $O_2$  sensitivity. FCCP is a proton uncoupler that can dissipate the proton gradient across the thylakoid membrane in algal cells. As illustrated in Fig. 2, in the presence of 1000 ppm of  $O_2$  after the induction of the hydrogenase enzyme, the steady-state photoevolution of  $H_2$  around the time of 20 h was slightly less than  $1 \mu mol$  of  $H_2 \cdot mg$  of  $Chl^{-1} \cdot h^{-1}$  (cf. Fig. 1 at approx 110 h). After a brief dark period (from 20:20 to 22:20), a burst of  $H_2$  photoevolution appeared, followed by an oscillation in the decay curve. Since both the actinic intensity and the background  $O_2$  concentration (1000 ppm) remained the same, this  $H_2$  production oscillation also indicated that the decay in the rate of  $H_2$  photo-evolution resulted not from  $O_2$  inhibition of the hydrogenase enzyme *per se*, but from a competitive kinetic effect of  $O_2$  on electron transport that is related to the  $H_2$  production process. The addition of  $5 \mu M$  FCCP produced a dramatic removal of  $O_2$  inhibition on  $H_2$  photoevolution. The rate of  $H_2$  production rose to about  $16 \mu mol$  of  $H_2 \cdot mg$  of  $Chl^{-1} \cdot h^{-1}$ . This FCCP-stimulated  $H_2$  production is clearly photo-dependent.

As soon as the actinic light was turned off, H<sub>2</sub> production stopped. The data (Fig. 2) also demonstrated that FCCP-enhanced photoevolution of H<sub>2</sub> can last for more than 4 h, although at a decreasing rate. The decrease in rate of H<sub>2</sub> photoproduction is owing to a secondary effect of FCCP known in the photosynthesis research literature as the acceleration of the deactivation reactions of the water-splitting system Y (ADRY) effect, in which FCCP gradually inhibits photosystem II (PSII) activity by deactivating the photosynthetic water-splitting complex in the S<sub>2</sub> and S<sub>3</sub> states (4). However, FCCP does not have any known effect on hydrogenase *per se*. Therefore, the observed stimulation of H<sub>2</sub> photoevolution by FCCP in the presence of 1000 ppm of O<sub>2</sub> clearly demonstrated that the newly discovered O<sub>2</sub>-sensitive electron transport pathway requires the presence of a proton gradient (or ATP) to operate.

DCMU is a chemical inhibitor that binds at the Q<sub>B</sub> site of PSII and blocks transport of electrons acquired from PSII water splitting to photosystem I. The experimental data (not shown) demonstrated that the addition of DCMU inhibited the burst of H<sub>2</sub> photoevolution after a dark period in the presence of 1000 ppm of background O<sub>2</sub>. This result indicated that > 90% of the electrons that are used in the photoproduction of H<sub>2</sub> are derived from PSII water splitting. Therefore, water is the main source of electrons for the H<sub>2</sub> burst after the dark period in the presence of 1000 ppm of O<sub>2</sub>. Organic reserves such as starch are thus not the main source of electrons in this mode of H<sub>2</sub> production.

The new O<sub>2</sub> sensitivity was further characterized using a series of O<sub>2</sub> concentrations: 10, 100, 300, 1000, 5000, and 10,000 ppm of O<sub>2</sub> in helium. The experimental results showed that the introduction of 100 ppm of O<sub>2</sub> had no significant effect on the steady-state rate of H<sub>2</sub> photoevolution in *C. reinhardtii* wild-type strain 137c. However, the addition of 300 ppm of O<sub>2</sub> began to show an effect. Figure 3 plots the percentage of steady-state H<sub>2</sub> production rate vs background O<sub>2</sub> concentrations in the wild-type 137c. The O<sub>2</sub> concentration that gave 50% inhibition of H<sub>2</sub> photoevolution was about 500 ppm. When the O<sub>2</sub> concentration was raised to 5000 ppm, the inhibition on H<sub>2</sub> production was dramatic and the rate of H<sub>2</sub> photo-evolution decreased to nearly zero (Figs. 3 and 4). However, the hydrogenase in the algal cells was still active even after the continued presence of 5000 ppm of O<sub>2</sub> for more than 10 h. When the actinic was turned on again after hour 198, a small peak of H<sub>2</sub> photoproduction was observed. As illustrated in the expanded scale (Fig. 4B), this H<sub>2</sub> photoproduction peak was clearly above the background noise and/or dark H<sub>2</sub> signal, indicating the presence of active hydrogenase in the algal cells. Therefore, hydrogenase in the wild-type cells can tolerate up to 5000 ppm of O<sub>2</sub>. The newly discovered O<sub>2</sub> sensitivity is about 10 times more sensitive to O<sub>2</sub> than that of the hydrogenase.

## Results and Discussion

The new O<sub>2</sub> sensitivity is apparently linked to the photosynthetic H<sub>2</sub> production pathway that is coupled with proton translocation across the




Fig. 3. Effect of background O<sub>2</sub> concentrations on steady-state H<sub>2</sub> photoevolution.

thylakoid membrane. The addition of the proton uncoupler FCCP can eliminate this mode of O<sub>2</sub> inhibition on H<sub>2</sub> photoevolution. This O<sub>2</sub> inhibition on H<sub>2</sub> production is most likely owing to the competitive uptake of reducing equivalents by background O<sub>2</sub>, with the H<sub>2</sub> production pathway for photosynthetically generated electrons from water splitting. The O<sub>2</sub>-sensitive H<sub>2</sub> production pathway can be inhibited by DCMU. Our experiments demonstrated that the competitive pathway is more sensitive to O<sub>2</sub> than the classic O<sub>2</sub> sensitivity of hydrogenase. These findings redefine the meaning of "oxygen tolerance" in algal H<sub>2</sub> production. As discussed, this O<sub>2</sub> sensitivity apparently represents a new pathway in the photosynthetic H<sub>2</sub> production that is coupled with proton translocation across the thylakoid membrane. As illustrated in Fig. 5A, the site for the reduction of O<sub>2</sub> could be at the RuBisCO enzyme, which can serve as an RuDP (also known as RuBP) carboxylase and/or an RuDP oxygenase in the Calvin cycle. Under conditions for H<sub>2</sub> photoevolution in which CO<sub>2</sub> is not present and ATP is abundant owing to associated photophosphorylation, the Calvin cycle enzymes are fully activated and RuBisCO could act as a strong oxygenase. This hypothesis can explain how FCCP mitigates O<sub>2</sub> inhibition of H<sub>2</sub> photoevolution, since operation of the Calvin cycle requires formation of ATP using the proton gradient across the thylakoid membrane. Another possible site for O<sub>2</sub> interaction could be at Fd, which, according to the classic Mehler reaction, can serve as an electron donor to O<sub>2</sub>. Additional experimental studies with different chemical inhibitors and genetic mutants are under way to elucidate this O<sub>2</sub>-sensitive H<sub>2</sub> production electron transport pathway.






Fig. 4 **(A)** The introduction of 5000 ppm of  $O_2$  dramatically inhibits  $H_2$  production, but the hydrogenase remains active. **(B)** Expanded vertical scale of (A), showing clear peak of  $H_2$  photoevolution after 10 h of continued presence of 5000 ppm of  $O_2$ .





Fig. 5. (A) The newly discovered O<sub>2</sub> sensitivity is likely owing to the background O<sub>2</sub> (at about 1000-ppm levels) acting as a terminal sink, in competition with the Fd/hydrogenase H<sub>2</sub> production pathway, for photoelectronically generated electrons. Illustrated by the blue-dotted arrows is a speculated mechanism of how the background O<sub>2</sub> might interact with the photosynthetic H<sub>2</sub> production pathway at the point(s) of RuBisCO and/or Fd.

(Continued)



**B**



### $H_2$ Production Under Anaerobic Conditions

Fig. 5. (Continued) **(B)** Development of efficient algal  $H_2$  production system by construction and transformation of vector that contains hydrogenase promoter and a piece of synthetic DNA for the polypeptide proton channel. The transformed alga could grow normally using ambient air  $CO_2$  under aerobic conditions without the polypeptide proton channel, which could be expressed only with the induction of the hydrogenase under anaerobic conditions when its function is needed for enhanced  $H_2$  production. This diagram illustrates a predicted mechanism of how photosynthetic  $H_2$  production could be enhanced through the action of the proposed polypeptide proton channel by dissipation of the static proton gradient across the thylakoid membrane so that the photosynthetic electron transport from PSII water splitting to the Fd/hydrogenase  $H_2$  production pathway can become more efficient, and by inactivation of the Calvin cycle so that both the background  $O_2$  and  $CO_2$  could be prevented from acting as a competitive terminal electron sink.

## Potential Application for Enhanced Photosynthetic H<sub>2</sub> Production

The discovery of alternative O<sub>2</sub> sensitivity also provides a new opportunity to develop a more-efficient and robust system for photosynthetic H<sub>2</sub> production. The experimental data with FCCP (Fig. 2) indicate that use of a polypeptide proton channel that does not deactivate the oxidizing equivalents of PSII could enhance H<sub>2</sub> production by eliminating the problems of both the proton-gradient accumulation (5) and the newly discovered alternative O<sub>2</sub> sensitivity. Therefore, we propose to create a “designer” photosynthetic organism for production of H<sub>2</sub> by genetic insertion of programmable proton channels in thylakoid membranes. The genetic insertion of programmable thylakoid-membrane proton channels can be achieved by transformation of a host alga with a genetic vector that contains a hydrogenase promoter-linked CF<sub>1</sub> suppressor or membrane polypeptide proton-channel gene. The envisioned “super” alga that can be created in this way should be able to perform autotrophic photosynthesis using ambient air CO<sub>2</sub> as the carbon source and grow normally under aerobic conditions such as in an open pond. When the algal culture is grown and ready for H<sub>2</sub> production, the CF<sub>1</sub> suppressor or proton-channel gene will then be expressed simultaneously with the induction of the hydrogenase enzyme under anaerobic conditions. The expression of the proton-channel gene should produce polypeptide proton channels in the thylakoid membrane, thus dissipating the proton gradient across the thylakoid membrane without ATP formation (Fig. 5B). The expression of the CF<sub>1</sub> suppressor should create CF<sub>0</sub>, which may act as a free proton channel without the CF<sub>1</sub> cap, thus similarly dissipating the proton gradient across the thylakoid membrane without ATP formation. The free proton-conductive CF<sub>0</sub> or polypeptide proton channels in the thylakoid membrane could provide two advantages for H<sub>2</sub> photoevolution: (1) accumulation of a proton gradient that impedes photosynthetic electron transport from water to Fd/hydrogenase could be prevented, and (2) the newly discovered O<sub>2</sub>-sensitive pathway that competes with the H<sub>2</sub> production pathway for photosynthetically generated electrons could be eliminated.

Therefore, the coexpression of the polypeptide proton channel (or CF<sub>1</sub> suppressor) and hydrogenase genes will make this alga a more-efficient system for production of H<sub>2</sub> by photosynthetic water splitting under anaerobic conditions (Fig. 5B). This organism contains normal mitochondria, which can use reducing power (NADH) from organic reserves (and/or exogenous acetate) to power the cell immediately after returning to aerobic conditions. Therefore, when the algal cell is returned to aerobic conditions after its use under anaerobic conditions for photoevolution of H<sub>2</sub> and O<sub>2</sub>, the cell will stop generating free CF<sub>0</sub> (or polypeptide proton channels) in thylakoid membranes and restore its normal photoautotrophic capability by synthesizing functional thylakoids. Consequently, it should be possible to use this type of genetically transformed organism for repeated cycles of

photoautotrophic culture growth under normal aerobic conditions and efficient production of H<sub>2</sub> and O<sub>2</sub> by photosynthetic water splitting under anaerobic conditions.

## Acknowledgments

We thank C. A. Sanders, B. Forbes, and B. Kusiak for culture media preparation; B. Mathis and A. Jones for secretarial support; M. K. Savage for editorial assistance; and C. D. King and V. W. Purdue for technical illustrations. We also thank Drs. M. Seibert and M. L. Ghirardi for their O<sub>2</sub>-tolerant mutants and informative discussions. This research was supported by the US Department of Energy Hydrogen Program, the DOE Office of Science Young Scientist Award (to J. W. Lee), and the Office of Basic Energy Sciences. Oak Ridge National Laboratory is managed by UT-Battelle, LLC, for the US Department of Energy under contract DE-AC05-00OR22725.

## References

1. Greenbaum, E. and Lee, J. W. (1998) in *BioHydrogen*, Zaborsky, O. R., ed., Plenum Press, New York, NY, pp. 235–241.
2. Ghirardi, M. L., Togasaki, R. K., and Seibert, M. (1997) *Appl. Biochem. Biotechnol.*, **63–65**, 141–151.
3. Lee, J. W., Blankinship, S. L., and Greenbaum, E. (1995), *Appl. Biochem. Biotechnol.*, **51/52**, 379–385.
4. Samuilov, V. D., Barsky, E. L., and Kitashov, A. V. (1995) in *Photosynthesis: from Light to Biosphere*, vol. II, Mathis, P., ed., Kluwer Academic Publishers, The Netherlands, pp. 267–270.
5. Lee, J. W. and Greenbaum, E. (1997), *Fuels and Chemicals from Biomass*, ACS Symposium Series 666, Saha, B. C. and Woodward, J., eds., American Chemical Society, Washington, DC, pp. 209–222.