

Errata

New organotin(IV) derivatives of dipeptides as models for metal–protein interactions: *in vitro* anti-tumour activity

Mala Nath^{1*}, Sandeep Pokharia¹, Xueqing Song², George Eng²,
Marcel Gielen^{3,4**}, Martine Kemmer⁴, Monique Biesemans³,
Rudolph Willem³ and Dick de Vos⁵

¹Department of Chemistry, Indian Institute of Technology–Roorkee, Roorkee 247667, India

²Department of Chemistry and Physics, University of the District of Columbia, Washington, DC 20008, USA

³Vrije Universiteit Brussel (VUB), High Resolution NMR Centre, Room 8G512, Pleinlaan 2, B-1050 Brussels, Belgium

⁴Université Libre de Bruxelles (ULB), Organic Chemistry, Av. F.D. Roosevelt 50, B-1050 Brussels, Belgium

⁵Pharmachemie BV, Medical Department, NL-2003 RN Haarlem, The Netherlands

Appl. Organometal. Chem. 2003; 17(5): 305–314. DOI:10.1002/aoc.451

On p.307 of the above paper, Equations (2) and (3) should have been printed as follows:



In Table 2, p.309, of the same paper, the entries for compound 7 should have been printed as follows:

Table 2. Characteristic IR frequencies (cm^{−1}) of dipeptides and their di- and tri-organotin(IV) complexes^a

Compound	Ligand complex	$\nu(\text{NH})$	$\nu(\text{CO}_{\text{amide}})$	$\nu_{\text{as}}(\text{COO})$	$\nu_{\text{s}}(\text{COO})$	$\Delta\nu$	$\nu_{\text{as}}(\text{Sn}-\text{C})$	$\nu_{\text{s}}(\text{Sn}-\text{C})$	$\nu(\text{Sn}-\text{O})$	$\nu(\text{Sn}-\text{N})/\nu(\text{Sn} \leftarrow \text{N})$
7	Ph ₃ Sn(HL-7)	3379s 3230s 3063s	1679s	1633s	1409s	224	270m	220m	574m	443m

^a s, strong; m, medium; w, weak.