



ORIGINAL ARTICLE

Ionic liquid [tbmim]Cl₂/AlCl₃ under ultrasonic irradiation towards synthesis of 1,4-DHP's



B. Palakshi Reddy, K. Rajesh, V. Vijayakumar *

Organic Chemistry Division, School of Advanced Sciences, VIT University, Vellore 632014, India

Received 10 December 2010; accepted 24 January 2011

Available online 1 February 2011

KEYWORDS

1,4-DHP's;
[tbmim]Cl₂/AlCl₃;
Hantzsch reaction;
Ultrasound irradiation

Abstract One pot, three-component synthesis of 1,4-dihydropyridines using ionic liquid 3,3'-thionyl-bis-1,1'-methylimidazolium chloroaluminate ([tbmim]Cl₂/AlCl₃) was found to be efficient to afford good to excellent yields at room temperature under ultrasonic irradiation. This method has many advantages, which avoids the use of harmful catalysts, occurrence of reactions at room temperature, high yields and methodological simplicity.

© 2011 Production and hosting by Elsevier B.V. on behalf of King Saud University.

1. Introduction

It is well known that Hantzsch 1,4-dihydropyridines (1,4-DHP's) exhibit a wide range of biological properties (Mauzeral and Wertheimer, 1955), acting as potent vasodilators, antihypertensives, bronchodilators, hepatoprotective, antitumor, antimutagenic, geroprotective and antidiabetic agents (Di Stilo et al., 1998). Moreover a number of DHP's acts as calcium antagonists (Janis and Triggle, 1983; Bocker and Guengerich, 1986; Gordeev et al., 1996) and they have been introduced for the treatment of congestive heart failure. In addition to this 1,4-DHP's exhibit several other medicinal applications like neuroprotectant (Klusa, 1995), platelet anti-aggregator activity

(Bretzel et al., 1993), as cerebral antiischemic agents in the treatment of Alzheimer's disease (Bretzel et al., 1992) and as a chemosensitizer in tumor therapy (Boer and Gekeler, 1995).

1,4-DHP's are generally synthesized using Hantzsch method (Hantzsch, 1882), which involves one pot cyclocondensation of aldehydes, dicarbonyls/ β -ketoesters and ammonium acetate/ammonia/primary amine either in refluxing with acetic acid or alcohol leading to low yields with longer reaction time (Loev and Snader, 1965; Sausins and Duburs, 1988). Recently, a number of modified methods have been developed (Evdokimov et al., 2006). Other procedures comprise the use of microwaves (Khadikar et al., 1995), ionic liquids (Ji et al., 2004), high temperatures at reflux (Phillips, 1949), TMSCl–NaI (Sabitha et al., 2003), InCl₃ (Babu and Perumal, 2000), I₂ (Ko et al., 2005), SiO₂/NaHSO₄ (Adharvana and Syamasundar, 2005), SiO₂/HClO₄ (Maheswara et al., 2006), CAN (Ko and Yao, 2006), Na- and Cs-Norit carbons (Perozo-Rondon et al., 2006), tetrabutylammonium hydrogen sulfate (Tewari et al., 2004), fermenting Baker's yeast (Lee, 2005), organocatalysts (Kumar and Maurya, 2007) and metal triflates (Wang et al., 2005). All the above


* Corresponding author. Tel.: +91 4162202332.

E-mail address: kvpsvijayakumar@gmail.com (V. Vijayakumar).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Scheme 1

Table 1 Optimization of reaction conditions for the synthesis of 1,4-DHP's.

Entry	Method	Catalyst [tbmim]Cl ₂ /AlCl ₃	Time (min)	Yield (%)
1	Sonication	1 mmol	40	94
2	Sonication	0 mmol	40	70
3	Reflux	0 mmol	180	80
4	Reflux	0.5 mmol	90	89
5	Stirring	0.5 mmol	150	88
6	Sonication	0.5 mmol	40	94

Reaction conditions: benzaldehyde (10 mmol), ethylacetoacetate (20 mmol), ammonium acetate (10 mmol), ionic liquid.

methods are however associated with several drawbacks, such as unsatisfactory yields, longer duration, difficulty in handling of reagents, toxic solvents, hence the development of an efficient synthesis of Hantzsch 1,4-dihydropyridines seemed to be of prime importance. In green synthesis ionic liquids play a major role. Initially ionic liquids are introduced as alternative green reaction media because of their unique chemical and physical properties of non-volatility, non-flammability, thermal stability, and controlled miscibility. But nowadays ionic liquids have proved to be useful beyond this, showing their importance in controlling reactions as catalysts (Jain et al., 2005).

In organic synthesis ultrasound irradiation has been considered as a clean and useful protocol, compared with the traditional methods. A large number of organic reactions can be carried out in higher yields, shorter reaction time or milder conditions in ultrasonic irradiation (Wilkes, 2002). In continuation of earlier interest on 1,4-DHP's (Rathore et al., 2009; Palakshi Reddy et al., 2009, 2010; Thenmozhi et al., 2009; Fun et al., 2009a,b,c) herein we report an efficient and simple procedure for the synthesis of 1,4-DHP derivatives at room temperature under ultrasonication using ionic liquid [tbmim]Cl₂/AlCl₃ as catalyst (see Scheme 1).

2. Experimental

The melting points were recorded in open capillaries and corrected with benzoic acid. Thin-layer chromatography (TLC) was carried out to monitor the course of the reaction and purity of the product. IR spectra recorded in an Avatar-330 FTIR spectrophotometer (Thermo Nicolet), and only noteworthy absorption levels (reciprocal centimeters) were listed. ¹H NMR spectra were recorded on Bruker AMX 400-MHz or 300-MHz spectrometer using CDCl₃ or DMSO as solvent and TMS as internal standard. Mass spectra were recorded on a LC-MS. Ionic liquid [tbmim]Cl₂/AlCl₃ was prepared according to the literature (Ling et al., 2006).

2.1. General procedure for the synthesis of 1,4-DHP's

A mixture of aromatic aldehyde **1** (1 mmol), ethylacetoacetate/acetylacetone **2** (2 mmol) and ammonium acetate **3** (1 mmol), was taken in a round bottomed flask mixed with [tbmim]Cl₂/AlCl₃ (0.5 mmol), which was irradiated under ultrasonic waves at room temperature for an appropriate time as indicated in Table 2. The progress of the reaction was monitored by TLC. The reaction mixture was then extracted with chloroform (10 mL) leaving behind [tbmim]Cl₂/AlCl₃. The organic

Table 2 Synthesis of various substituted 1,4-dihydropyridines using ionic liquid.

S. No	Ph	R	US (min)	Yield (%)	M.P (°C)
4a	C ₆ H ₅	OC ₂ H ₅	40	94	158–160
4b	4-ClC ₆ H ₄	OC ₂ H ₅	45	95	145–146
4c	4-OCH ₃ C ₆ H ₄	OC ₂ H ₅	50	93	153–155
4d	3-NO ₂ C ₆ H ₄	OC ₂ H ₅	30	96	165–167
4e	4-Pyridyl	OC ₂ H ₅	45	91	185–187
4f	3-Pyridyl	OC ₂ H ₅	50	90	188–190
4g	4-BrC ₆ H ₄	OC ₂ H ₅	45	93	148–150
4h	2-Thienyl	OC ₂ H ₅	50	90	172–173
4i	4-CH ₃ C ₆ H ₄	OC ₂ H ₅	60	88	142–145
4j	4-HOC ₆ H ₄	OC ₂ H ₅	60	90	227–229
4k	3-HOC ₆ H ₄	OC ₂ H ₅	55	87	172–175
4l	3-NO ₂ C ₆ H ₄	CH ₃	35	94	180–182
4m	2-NO ₂ C ₆ H ₄	CH ₃	35	93	195–197
4n	2-Furyl	CH ₃	50	90	169–170
4o	2-Thienyl	CH ₃	55	91	170–172
4p	3-Pyridyl	CH ₃	50	89	262–263
4q	3-ClC ₆ H ₄	CH ₃	40	92	222–224
4r	3-OCH ₃ C ₆ H ₄	CH ₃	50	88	203–205
4s	4-C ₂ H ₅ C ₆ H ₄	CH ₃	55	87	179–181
4t	4-Pyridyl	CH ₃	50	89	245–246
4u	4-BrC ₆ H ₄	CH ₃	40	93	109–111

Reaction conditions: aldehyde (10 mmol), ethylaceto acetate/acetylacetone (20 mmol), ammonium acetate (10 mmol), ionic liquid (0.5 mmol).

layer was then washed with water and dried over anhydrous Na_2SO_4 . Organic solvent was evaporated under reduced pressure and a solid compound was crystallized from absolute ethanol to afford the pure corresponding 1,4-dihydropyridine derivatives **4(a–u)** in excellent yields.

3. Results and discussion

Herein, we report a developed a methodology for the synthesis of 1,4-dihydropyridines (1,4-DHP's) in the presence of $[\text{tbmim}]\text{Cl}_2/\text{AlCl}_3$ under ultrasound-irradiation. Herein we have carried out the reaction of benzaldehyde **1**, ethylacetoacetate/acetylacetone **2** and ammonium acetate **3** catalyzed by $[\text{tbmim}]\text{Cl}_2/\text{AlCl}_3$. Different reaction conditions have been studied for optimization (Table 1).

The use of 1 mmol of ionic liquid does not affect the yield even after increase in reaction time. So 0.5 mmol of ionic liquid is sufficient to yield 1,4-DHP's. It is important to note that in the absence of the ionic liquid and without using ultrasound irradiation, the yield of the reactions decreased with increase in time. Using ionic liquid as a promoter for the synthesis of Hantzsch 1,4-DHP's does not only represent a dramatic improvement at room temperature with regard to yield (87–96%) over conventional thermal heating, but the reaction times are also considerably decreased compared to classical synthesis (3–6 h). Ultrasonic irradiation was very simple and convenient for the synthesis of 1,4-DHP's in room temperature using ionic liquid in ultrasonic cleaner with a frequency of 40 kHz and a nominal power 100 W. In this experiment the ultrasonic technique represented a better procedure with respect to time and yield. After optimizing the conditions, the generality of this method was examined by the reaction of several substituted aldehydes, ethylacetoacetate/acetylacetone and ammonium acetate using $[\text{tbmim}]\text{Cl}_2/\text{AlCl}_3$ as a catalyst under ultrasound irradiation, the results are shown in Table 2.

Herein, we have found that the reactions of aromatic aldehydes having electron-withdrawing groups are fast as compared to the reaction of aldehydes having electron donating groups. Unlike those of ethylacetoacetate products, the reaction of acetylacetone products proceeded at lower rate less efficiently to give 1,4-DHP's. The results obtained in the current method are illustrated in Table 2. All the products obtained were fully characterized by spectroscopic methods, such as IR, ^1H NMR and mass spectroscopy and also by comparison with the reference compounds. Another advantage of the ionic liquid is that it is recyclable. In view of environmentally friendly methodologies, recovery and reuse of the ionic liquid is highly preferable. $[\text{tbmim}]\text{Cl}_2/\text{AlCl}_3$ is easily separated from reaction medium by washing with water and distillation of the solvent under vacuum and it can be reused for subsequent reactions. Recycled ionic liquid shows no loss of efficiency with regard to yield after four successive runs, hence this method can be regarded as a rapid, convenient and environmentally benign method for the synthesis of 1,4-dihydropyridine derivatives.

4. Conclusion

An efficient and environmentally friendly method has been developed for the synthesis of 1,4-dihydropyridines catalyzed by 0.5 mmol of $[\text{tbmim}]\text{Cl}_2/\text{AlCl}_3$ under ultrasonic irradiation. The use of sonication combining ionic liquid in the three-

component condensation at room temperature brings up the merits like milder conditions, low pollution, good yields and simple work-up.

4.1. Spectral data for respective compounds

Compound **4a**: IR (KBr) ν_{max} 3348, 3052, 1692, 1633, 1223. ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 1.26 (t, 6H, $J = 8$ Hz), δ 2.35 (s, 6H), δ 4.10 (m, 4H, $J = 4$ Hz), δ 5.93 (s, 1H), δ 5.76 (s, 1H), δ 7.38–8.14 (Aromatic). MS: m/z 329 (M).

Compound **4b**: IR (KBr) ν_{max} 3345, 3091, 1705. ^1H NMR (400 MHz, CDCl_3): δ 1.22 (t, 6H, $J = 6$ Hz), δ 2.33 (s, 6H), δ 4.09 (m, 4H, $J = 4$ Hz), δ 4.96 (s, 1H), δ 5.70 (s, 1H), δ 7.17–7.25 (Aromatic). MS: m/z 363 (M), 364 (M + 1).

Compound **4c**: IR (KBr) ν_{max} 3342, 3094, 1689. ^1H NMR (400 MHz, CDCl_3): δ 1.23 (t, 6H, $J = 8$ Hz), δ 2.34 (s, 6H), δ 3.72 (s, 3H), δ 4.09 (m, 4H, $J = 4$ Hz), δ 4.94 (s, 1H), δ 5.58 (s, 1H), δ 6.76–7.21 (Aromatic). MS: m/z 360 (M).

Compound **4d**: IR (KBr) ν_{max} 3357, 3092, 1692. ^1H NMR (400 MHz, CDCl_3): δ 1.23 (t, 6H, $J = 8$), δ 2.33 (s, 6H), δ 4.09 (m, 4H, $J = 4$ Hz), δ 5.90 (s, 1H), δ 5.73 (s, 1H), δ 7.38–8.14 (Aromatic). MS: m/z 374 (M).

Compound **4e**: IR (KBr) ν_{max} 3378, 3082, 1698. ^1H NMR (400 MHz, CDCl_3): δ 1.20 (t, 6H, $J = 6$ Hz), δ 2.30 (s, 6H), δ 4.10 (m, 4H, $J = 4$ Hz), δ 5.00 (s, 1H), δ 6.10 (s, 1H), δ 7.20–8.47 (Aromatic). MS: m/z 330 (M).

Compound **4f**: IR (KBr) ν_{max} 3378, 3082, 1698. ^1H NMR (400 MHz, CDCl_3): δ 1.20 (t, 6H, $J = 6$ Hz), δ 2.30 (s, 6H), δ 4.10 (m, 4H, $J = 4$ Hz), δ 5.00 (s, 1H), δ 6.10 (s, 1H), δ 7.20–8.60 (Aromatic). MS: m/z 330 (M).

Compound **4g**: IR (KBr) ν_{max} 3345, 3091, 1689. ^1H NMR (400 MHz, CDCl_3): δ 1.22 (t, 6H, $J = 6$ Hz), δ 2.33 (s, 6H), δ 4.09 (m, 4H, $J = 4$ Hz), δ 4.96 (s, 1H), δ 5.70 (s, 1H), δ 7.17–7.25 (Aromatic). MS: m/z 409 (M + 2).

Compound **4n**: IR (KBr) ν_{max} 3342, 3094, 1689, 1638. ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 2.28 (s, 6H), δ 2.36 (s, 6H), δ 5.29 (s, 1H), δ 6.05 (s, 1H), δ 7.40–8.04 (Aromatic). MS: m/z 330 (M).

Compound **4o**: IR (KBr) ν_{max} 3342, 3094, 1689. ^1H NMR (400 MHz, CDCl_3): δ 2.28 (s, 6H), δ 2.38 (s, 6H), δ 5.29 (s, 1H), δ 6.10 (s, 1H), δ 7.40–8.10 (Aromatic). MS: m/z 314 (M).

Compound **4p**: IR (KBr) ν_{max} 3332, 3096, 1690. ^1H NMR (400 MHz, CDCl_3): δ 2.31 (s, 6H), δ 2.37 (s, 6H), δ 5.15 (s, 1H), δ 5.80 (s, 1H), δ 6.70–7.54 (Aromatic). ^{13}C NMR (CDCl_3): δ 20.0, 29.5, 4.7, 104.5, 141.4, 145.1, 157, 197.8. MS: m/z 259 (M +), 261 (M + 2), 192.

Compound **4q**: IR (KBr) ν_{max} 3332, 3064, 1695. ^1H NMR (400 MHz, CDCl_3): δ 2.32 (s, 6H), δ 2.33 (s, 6H), δ 5.40 (s, 1H), δ 6.10 (s, 1H), δ 6.80–7.80 (Aromatic). ^{13}C NMR (400 MHz, CDCl_3): δ 20.6, 29.7, 35.0, 113.0, 123.7, 128.1, 135.1, 144.1, 150.0, 197.2. MS: m/z 274 (M–1), 275 (M), 276 (M + 1).

Compound **4r**: IR (KBr) ν_{max} 3345, 3090, 1685. ^1H NMR (400 MHz, CDCl_3): δ 2.29 (s, 6H), δ 2.35 (s, 6H), δ 5.12 (s, 1H), δ 5.80 (s, 1H), δ 6.70–7.29 (Aromatic). ^{13}C NMR (CDCl_3 , 400 MHz, ppm): δ 20.4, 30.0, 39.92, 113.7, 119.7, 142.8, 197.8. MS: m/z 299 (M).

Compound **4s**: IR (KBr) ν_{max} 3378, 3082, 1698, 963. ^1H NMR (400 MHz, CDCl_3): δ 2.34 (s, 6H), δ 2.20 (s, 6H), δ 5.11 (s, 1H), δ 5.81 (s, 1H), δ 7.16–7.26 (Aromatic). MS: m/z 303 (M), 304 (M + 1).

Acknowledgment

We thank to Chemistry Division, VIT University, Vellore, India for providing facilities to carry research work.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.arabjc.2011.01.027.

References

- Adharvana, C.M., Syamasundar, K., 2005. Silica gel/NaHSO₄ catalyzed one-pot synthesis of Hantzsch 1,4-dihydropyridines at ambient temperature. *Catal. Commun.* 6, 624.
- Babu, G., Perumal, P.T., 2000. Synthetic applications of indium trichloride catalyzed reactions. *Aldrichim. Acta*, 16.
- Bocker, R.H., Guengerich, E.P.J., 1986. Oxidation of 4-aryl- and 4-alkyl-substituted 2,6-dimethyl-3,5-bis(alkoxycarbonyl)-1,4-dihydropyridines by human liver microsomes and immunochemical evidence for the involvement of a form of cytochrome P-450. *Med. Chem.* 29, 1596.
- Boer, R., Gekeler, V., 1995. Chemosensitizers in tumor therapy: new compounds promise better efficacy. *Drugs Future* 20, 499.
- Bretzel, R.G., Bollen, C.C., Maeser, E., Federlin, K.F., 1992. Trombodipine platelet aggregation inhibitor antithrombotic. *Drugs Future* 17, 465.
- Bretzel, R.G., Bollen, C.C., Maeser, E., Federlin, K.F., 1993. Nephroprotective effects of nitrendipine in hypertensive type I and type II diabetic patients. *Am. J. Kidney Dis.* 21, 53.
- Di Stilo, A., Visentin, A., Clara, C., Gasco, A.M., Ermondi, G., Gasco, A., 1998. New 1,4-Dihydropyridines conjugated to furoxanyl moieties, endowed with both nitric oxide-like and calcium channel antagonist vasodilator activities. *J. Med. Chem.* 41, 5393.
- Evdokimov, N.M., Magedov, I.V., Kireev, A.S., Kornienko, A., 2006. One-step, three-component synthesis of pyridines and 1,4-dihydropyridines with manifold medicinal utility. *Org. Lett.* 8, 899.
- Fun, H.K., Liew, W.C., Palakshi Reddy, B., Sarveswari, S., Vijayakumar, V., 2009a. Diethyl 2,6-dimethyl-4-p-tolyl-1,4-dihydropyridine-3,5-dicarboxylate. *Acta Crystallogr., Sect. E* 65, o2342.
- Fun, H.K., Quah, C.K., Palakshi Reddy, B., Sarveswari, S., Vijayakumar, V., 2009b. Diethyl 2,6-dimethyl-4-p-tolyl-1,4-dihydropyridine-3,5-dicarboxylate. *Acta Crystallogr., Sect. E* 65, o2255.
- Fun, H.K., Goh, J.H., Palakshi Reddy, B., Sarveswari, S., Vijayakumar, V., 2009c. Dimethyl 4-(4-ethoxyphenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylate. *Acta Crystallogr., Sect. E* 65, o2247.
- Gordeev, M.F., Patel, D.V., Gordon, E.M., 1996. Approaches to combinatorial synthesis of heterocycles: a solid-phase synthesis of 1,4-dihydropyridines. *J. Org. Chem.* 61, 924.
- Hantzsch, A., 1882. Ueber die synthese pyridinartiger verbindungen aus acetessigäther und aldehydammoniak. *Liebigs Ann. Chem.* 215, 1.
- Jain, N., Kumar, A., Chauhan, S., Chauhan, S.M.S., 2005. Chemical and biochemical transformations in ionic liquids. *Tetrahedron* 61, 1015.
- Janis, R.A., Triggle, D.J., 1983. New developments in calcium ion channel antagonists. *J. Med. Chem.* 26, 775.
- Ji, S.J., Jiang, Z.Q., Lu, J., Loh, T.P., 2004. Facile Ionic liquids-promoted one-pot synthesis of polyhydroquinoline derivatives under solvent free conditions. *Synlett*, 831.
- Khadikar, B.M., Gaikar, V.G., Chitnavis, A.A., 1995. Aqueous hydrotrope solution as a safer medium for microwave enhanced Hantzsch dihydropyridine ester synthesis. *Tetrahedron Lett.* 36, 8083.
- Klusa, V., 1995. Neuroprotectant, cognition enhancer. *Drugs Future* 20, 135.
- Ko, S., Yao, C.F., 2006. Ceric Ammonium Nitrate (CAN) catalyzes the one-pot synthesis of polyhydroquinoline via the Hantzsch reaction. *Tetrahedron* 62, 7293.
- Ko, S., Sastry, M.N.V., Lin, C., Yao, 2005. Molecular iodine-catalyzed one-pot synthesis of 4-substituted-1,4-dihydropyridine derivatives via Hantzsch reaction. *Tetrahedron Lett.* 46, 5771.
- Kumar, A., Maurya, R.A., 2007. Synthesis of polyhydroquinoline derivatives through unsymmetric Hantzsch reaction using organocatalysts. *Tetrahedron* 63, 1946.
- Lee, J.H., 2005. Synthesis of Hantzsch 1,4-dihydropyridines by fermenting bakers yeast. *Tetrahedron Lett.* 46, 7329.
- Ling, H., Tao, G.H., Liu, W.S., Xiong, W., Tao, W., Kou, Y. Chin., 2006. One-pot synthesis of Lewis acidic ionic liquids for Friedel-Crafts alkylation. *Chem. Lett.* 17, 321.
- Loev, B., Snader, K.M., 1965. The Hantzsch reaction. I. Oxidative dealkylation of certain dihydropyridines. *J. Org. Chem.* 30, 1914.
- Maheswara, M., Siddaiah, V., Rao, Y.K., Tzeng, Y.M., Sridhar, C., 2006. A simple and efficient one-pot synthesis of 1,4-dihydropyridines using heterogeneous catalyst under solvent-free conditions. *J. Mol. Catal.* 260, 179.
- Mauzeral, D., Wertheimer, F.H., 1955. 1-Benzylidihydronicotinamide-a model for reduced DPN. *J. Am. Chem. Soc.* 77, 2261.
- Palakshi Reddy, B., Vijayakumar, V., Suresh, J., Narasimhamurthy, T., Nilantha Lakshman, P.L., 2009. 1,1'-[4-(2-Methoxyphenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-diyl]diethanone. *Acta Crystallogr., Sect. E* 65, o2877.
- Palakshi Reddy, P., Vijayakumar, V., Suresh, J., Narasimhamurthy, T., Nilantha Lakshman, P.L., 2010. Diethyl 4-(2,4-dichlorophenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylate. *Acta Crystallogr., Sect. E* 66, o363.
- Perozo-Rondon, E., Calvino-Casilda, V., Martin-Aranda, R.M., Casal, B., Duran-Valle, C.J., Rojas-Cervantes, M.L., 2006. Catalysis by basic carbons: Preparation of dihydropyridines. *Appl. Surf. Sci.* 252, 6080.
- Phillips, P., 1949. Hantzsch's pyridine synthesis. *J. Am. Chem. Soc.* 71, 4003.
- Rathore, R.S., Palakshi Reddy, B., Vijayakumar, V., Ragavan, R.V., Narasimhamurthy, T., 2009. Hantzsch 1,4-dihydropyridine esters and analogs: candidates for generating reproducible one-dimensional packing motifs. *Acta Crystallogr., Sect. B* 65, o375.
- Sabitha, G., Reddy, G.S.K.K., Reddy, C.S., Yadav, J.S., 2003. A novel TMSI-mediated synthesis of Hantzsch 1,4-dihydropyridines at ambient temperature. *Tetrahedron Lett.* 44, 4129.
- Sausins, A., Duburs, G., 1988. Synthesis of 1,4-dihydropyridines by cyclocondensation reactions. *Heterocycles* 27, 269.
- Tewari, N., Dwivedi, N., Tripathi, R.P., 2004. Tetrabutylammonium hydrogen sulfate catalyzed eco-friendly and efficient synthesis of glycosyl 1,4-dihydropyridines. *Tetrahedron Lett.* 45, 9011.
- Thenmozhi, M., Kavitha, T., Palakshi Reddy, B., Vijayakumar, V., Ponnuswamy, M.N., 2009. 1,1'-[4-(4-Methoxyphenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-diyl]diethanone. *Acta Crystallogr., Sect. E* 65, o2795.
- Wang, L.M., Sheng, J., Zhang, L., Han, J.W., Fan, Z., Tian, H., Qian, C.T., 2005. Facile Yb(OTf)₃ promoted one-pot synthesis of polyhydroquinoline derivatives through Hantzsch reaction. *Tetrahedron* 61, 1539.
- Wilkes, J.S., 2002. A short history of ionic liquids-from molten salts to neoteric solvents. *Green Chem.* 4, 73.