



Mini-Review

Aluminium cycling in the soil-plant-animal-human continuum

Z. Rengel

Soil Science and Plant Nutrition, School of Earth and Geographical Sciences, The University of Western Australia, Crawley WA 6009, Perth, Australia (Fax: +61 8 6488 1050; E-mail: zengel@agric.uwa.edu.au)

Received 26 April 2004; Accepted 10 May 2004; Published online October 2004

Abstract

A critical review of the literature on Al toxicity in plants, animals and humans reveals a similar mode of Al action in all living organisms, namely interference with the secondary messenger system (phosphoinositide and cytosolic Ca^{2+} signalling pathways) and enhanced production of reactive oxygen species resulting in oxidative stress. Aluminium uptake by plants is relatively quick (across the intact plasma membrane in < 30 min and across the tonoplast in < 1 h), despite huge proportion of Al being bound in the cell wall. Aluminium absorption in the animal/human digestive system is low (only about 0.1% of daily Al intake stays in the human body), except when Al is complexed with organic ligands (eg. citrate, tartarate, glutamate). Aluminium accumulates in bones and brain, with Al-citrate and Al-transferrin complexes crossing the blood-brain barrier and accumulating in brain cells. Tea plant and other Al-accumulator plant species contain large amounts of Al in the form of non-toxic organic complexes.

Introduction

Aluminium is the most abundant metal and the third most abundant chemical element in the Earth crust. Most of Al is incorporated into aluminosilicate soil minerals, with only small quantities appearing in soluble forms that can influence living organisms (May & Nordstrom 1991). However, solubilisation of Al-containing minerals is enhanced in acidic environments.

Most soils are being acidified continuously due to (i) imbalances in nitrogen, sulphur and carbon cycles (Bolan & Hedley 2003; Tang & Rengel 2003), (ii) excess uptake of cations over anions (Tang & Rengel 2003), (iii) use of ammonia- and amide-containing fertilizers (Mahler *et al.* 1985), (iv) nitrogen-fixation by legumes (Bolan *et al.* 1991; Coventry & Slattery 1991; Tang & Rengel 2003), and (v) acidic depositions from the atmosphere (Alewell 2003).

Acidic atmospheric depositions (mainly sulphuric and nitric acids) are caused by emissions into atmosphere of sulphur dioxide, nitrogen oxides and ammonia. These acids lower pH, increase Al concentration in surface waters (Bi 2001), decrease biodiversity in

aquatic ecosystems (Driscoll *et al.* 2003), and enhance acidification of sensitive soils (Blaser *et al.* 1999; Herman *et al.* 2001; Jönsson *et al.* 2003), especially in forests because of a filtering effect of tall vegetation (Fowler *et al.* 1999). However, decreased atmospheric emissions in recent decades in Europe (Alewell 2001; Herman *et al.* 2001) and the USA (Gbondo & Driscoll 2002) have resulted in partial, albeit slow, recovery of aquatic and terrestrial ecosystems (Herman *et al.* 2001). Part of the reason for slow recovery is a 'memory effect' of high acidification load from the earlier part of 20th century (1965–1985) (Blaser *et al.* 1999). In addition, soil acidification is still continuing, at least at some locations in Europe (Jönsson *et al.* 2003).

Increasing Al concentration in surface waters due to acidification (Bi 2001) resulted in Al toxicity to fish (Roux *et al.* 1996) and other aquatic organisms (Witters 1998), like benthic invertebrates (Herrmann 2001) when toxic thresholds were exceeded. In soil, Al is toxic to earthworms at acidic pH (Phillips & Bolger 1998) as well as to a range of other soil animals (Markich *et al.* 2002). In terrestrial ecosystems, Al toxicity to plants in acid soils is a major factor limiting

crop production and yield in acid soils worldwide (Foy 1988).

Despite decades of extensive effort to decipher the mechanism(s) of Al toxicity to plants, the primary cause remains largely speculative (Rengel 1992a, 1996; Horst 1995; Matsumoto 2000; Rengel & Zhang 2003). Similarly, there is continuous controversy regarding modes of Al toxicity to animals as well as possible involvement of Al in the Alzheimer's disease and other neurodegenerative diseases in humans (eg. Candy *et al.* 1992; Kruck 1993; Armstrong *et al.* 1996; Savory *et al.* 1996; Makjanic *et al.* 1998; Yokel 2000; Flaten 2001; Campbell 2002). Regardless of the type of living organism, there are similar main reasons for these controversies (i) technical difficulties associated with determining ionic speciation of Al in a given medium, (ii) problems with detecting transport of Al across the plasma membrane of intact cells, and (iii) ascertaining the intracellular localisation of Al.

In this review, I will highlight recent progress in understanding the transport of Al from soil to plants to humans and will critically examine implications of Al location in cells and organelles in Al toxicity. For other aspects of Al toxicity and resistance, readers interested in plants are referred to Kochian (1995), Rengel (1996), Matsumoto (2000), Barcelo & Poschenrieder (2002) and Rengel & Zhang (2003). Aluminium toxicity in animals has been reviewed by Appelberg *et al.* (1992), Battarbee (1994), Nayak & Chatterjee (1999) and Crichton *et al.* (2002), whereas the role of Al as a human neurotoxin has been discussed by Armstrong *et al.* (1996), Srinivasan *et al.* (1999), Yokel (2000), Flaten (2001), Yokel & McNamara (2001), Campbell (2002) and Rondeau (2002).

Symptoms of aluminium toxicity to living organisms

Despite substantial differences in some aspects of Al toxicity in various living organisms, there are certain common points (Strong *et al.* 1996; Yokel 2000). In plants, animals and humans, Al interferes with basic cellular functions, like the phosphoinositide and intracellular Ca^{2+} signalling pathways, which are, as secondary messengers, involved in a myriad of cellular metabolic functions.

In murine strain mice (Haug *et al.* 1994), rat hippocampal slices (Breakwell & Publicover 1994) as well as human neuroblastoma cells (Wood *et al.* 1994), Al exposure disturbs the phosphoinositide signalling

pathway by acting on guanine nucleotide-binding proteins (G-proteins) and a phosphatidylinositol-4,5-diphosphate specific phospholipase C. In addition, Al decreases the accumulation of inositol phosphates [especially inositol-1,4,5-triphosphate (IP_3)] and, by changing the membrane phospholipid composition, can also affect the phospholipase C in murine neuroblastoma cells (Shi & Haug 1992; Shi *et al.* 1993) and wheat root cells (Jones & Kochian 1995).

In murine (Shi & Haug 1992; Haug *et al.* 1994) as well as human neuroblastoma cells (Wood *et al.* 1994) Al disturbs intracellular Ca^{2+} homeostasis and diminishes Ca^{2+} rises, thus affecting the action of cytosolic Ca^{2+} activity ($[\text{Ca}^{2+}]_{\text{cyt}}$) as a secondary messenger. In plant cells, all but one report published so far demonstrated an increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ as a consequence of Al toxicity (Rengel & Zhang 2003), eg. in intact wheat (Zhang *et al.* 1998; Zhang & Rengel 1999) and rye roots (Ma *et al.* 2002), excised barley roots (Nichol & Oliveira 1995), Arabidopsis root hairs (Jones *et al.* 1998a), tobacco suspension cells (Vitarello & Haug 1996) and protoplasts isolated from wheat roots (Lindberg & Strid 1997). On the other hand, Jones *et al.* (1998b) reported a rapid decrease in $[\text{Ca}^{2+}]_{\text{cyt}}$ in BY-2 tobacco cell culture upon exposure to Al. Hence, all plant experimental system tested so far responded to the Al toxicity stress by *altering* the cytosolic Ca^{2+} homeostasis (as predicted by Rengel 1992b).

The other common feature of Al toxicity in plant and animal/human cells is increased production of reactive oxygen species (ROS) resulting in the oxidative stress. In mammalian cells Al can potentiate Fe-induced oxidative stress through increased production of ROS (Crichton *et al.* 2002), which have been implicated in neurological disorders (Strong *et al.* 1996; Campbell 2002). In plants, Al also causes increased production of ROS (Yamamoto *et al.* 2002) as well as lipid peroxidation (Yamamoto *et al.* 2003), with the former being a potential cause of root growth inhibition upon exposure to Al (Yamamoto *et al.* 2003).

Plants

Inhibition of root growth is one of the earliest and most dramatic symptoms exhibited by plants suffering from Al stress; this symptom has been observed within minutes of exposure to micromolar concentrations of Al in solution cultures (Blancaflor *et al.* 1998; Sivaguru & Horst 1998; Ryan *et al.* 1993; Vazquez *et al.* 1999; Zhang *et al.* 1998, 1999; Ahn *et al.* 2001;

Ma *et al.* 2002). The root apex is a critical site of perception and expression of Al toxicity. Inhibition of root growth occurs only after the distal part of the elongation zone within the apex (Sivaguru & Horst 1998; Kollmeier *et al.* 2000) was exposed to Al.

Root growth is a complex and dynamic process, making it likely that a number of biochemical and physiological processes may have already been altered prior to Al-induced inhibition of root growth. Among these processes, the important ones are disturbance of cytoplasmic Ca^{2+} and pH homeostasis (for a recent review see Rengel & Zhang 2003), inhibition of the H^+ -ATPase activity in the plasma membrane of the root tip cells (Ahn *et al.* 2002, 2004; Rengel & Zhang 2003), exacerbated oxidative stress (Yamamoto *et al.* 2002, 2003), and disruption of cytoskeleton dynamics (Sivaguru *et al.* 1999; Schwarzerova *et al.* 2002). These changes may directly or indirectly underlie the observed Al toxicity symptoms on plants. Indeed, neutralisation of the membrane surface charge in the root tips of Al-sensitive wheat genotype (due to a decrease in H^+ -ATPase activity and H^+ transport, Ahn *et al.* 2004) was accompanied by inhibited root growth (Zhang *et al.* 1999), suggesting a link between growth inhibition and alteration of the plasma membrane properties by Al exposure.

Animals and humans

An excellent compilation of metal toxicity for many aquatic and soil animals has recently been published (Markich *et al.* 2002), identifying large differences among animal species in sensitivity to Al toxicity. For example, Al was toxic to benthic invertebrates when inorganic Al concentration reached $0.1\text{--}0.3\text{ mg L}^{-1}$, but large differences in the severity of toxic effects were noted among species (Roux *et al.* 1996).

Different symptoms of Al toxicity were found in different animal species. In freshwater mussels, the filtering action was severely impaired by a short-term (1 h) as well as long-term (15 days) exposure to Al at neutral pH (Kadar *et al.* 2002). In pond snail *Lymnaea stagnalis*, Al altered voltage-activated sodium currents in neurons (Csoti *et al.* 2001), suggesting interference with electrophysiology of ion channels in the plasma membrane. In earthworms, cocoon production was inhibited by Al toxicity (Phillips & Bolger 1998).

Aluminium can be toxic to humans at both environmental and therapeutic levels. The human exposure to Al is primarily dietary, even though occupational (industrial) exposure (eg. Polizzi *et al.* 2002) and

medically administered Al (including medications like antacids, phosphate-complexation treatments like Al trihydroxide, Al-contaminated dialysis fluids, etc.) can represent substantial proportion of Al intake (Yokel & McNamara 2001).

The association between Al and dialysis encephalopathy is well established (eg. see Van Landeghem *et al.* 1997). Administration of Al-containing phosphate-complexation treatment to dialysis patients can cause acute toxicity (Desroches *et al.* 2000) and may be the major factor in renal osteodystrophy (Vanholder *et al.* 2002) and Al accumulation in brain (Candy *et al.* 1992). Also, acute Al toxicity of iatrogenic origin is well documented (Desroches *et al.* 1999).

Aluminium is associated with the inflammatory responses (Platt *et al.* 2001), mediated by interleukins and other inflammatory cytokines (Campbell 2002). In addition, Al increases Fe-induced oxidative injury in brain tissues (Yokel 2000; Campbell 2002; Crichton *et al.* 2002), resulting in lipid peroxidation and cytotoxicity of free-oxygen radicals (Strong *et al.* 1996). Aluminium affects brain neurotransmitters by decreasing the release and subsequent breakdown of 5-hydroxytryptamine (Kumar 2002), by binding to acetylcholine (Matlaba *et al.* 2000) and by increasing acetylcholinesterase activity in brain (Zatta *et al.* 2002).

The neuronal response to Al exposure can be morphologically distinguished into two types: (i) formation of intraneuronal neurofilamentous aggregates, or (ii) induction of significant neurochemical and neurophysiological perturbations without formation of neurofilamentous aggregates (Strong *et al.* 1996).

The neurofibrillary tangles (first described by Alzheimer in 1907) and the senile plaques represent the two principle neuropathologic lesions identified in the brains of patients with Alzheimer's disease. Aluminium contributes to accumulation of insoluble amyloid beta protein (causing senile plaques, Armstrong *et al.* 1996) and hyperphosphorylated tau in brain tissue (leading to formation of neurofibrillary tangles and neuropil threads, Armstrong *et al.* 1996) and to some extent mimics the deficit of cortical cholinergic neurotransmitters (Matlaba *et al.* 2000; Platt *et al.* 2001), all of which are characteristics of Alzheimer's disease (Yokel 2000). However, the Al involvement in Alzheimer's disease and other neurological disorders has remained controversial over the years (eg. Candy *et al.* 1992; Kruck 1993; Armstrong *et al.* 1996; Savory *et al.* 1996; Makjanic

et al. 1998; Yokel 2000; Flaten 2001; Campbell 2002). Epidemiological studies could not provide a conclusive evidence of the causality of the frequently obvious connection between Al and Alzheimer's disease, whereas appropriate physiological studies (eg. employing weakly radioactive ^{26}Al isotope and accelerated mass spectrometry detection, see below) of Al effects under physiological conditions have only recently become possible (eg. Moore *et al.* 2000; Ward *et al.* 2001).

Aluminium speciation

Aluminium makes complexes with numerous ligands, especially oxygen-donor ones. Many of these complexes also tend to hydrolyse and form polymeric species in aqueous solution, resulting in complex coordination chemistry of Al in the environmental samples and biofluids. In addition, complexation constants for Al and common ligands (like citrate, phosphate and hydroxide) are uncertain at best (Harris *et al.* 1996), making reliable Al speciation quite a challenge. Speciation of Al is further hampered by technical difficulties in separating and reliably quantifying various Al species. The time factor also needs to be considered because polynuclear Al species can be depolymerised by organic ligands into uncondensed monomeric Al species over time (Masion *et al.* 2000).

Large differences in toxicity of various Al species (from none to extreme) to living organisms make speciation of Al a necessity. Special attention needs to be paid to bioavailable Al species because they can cause toxicity. All ecotoxicology research should therefore move from expressing total concentrations of Al and other metals to assessing the concentrations of relevant bioavailable metal species (eg. see Witters 1998).

Analytical techniques for determining aluminium speciation in environmental samples and biofluids

Determining Al speciation is a two-step process, comprising separation and quantification. Numerous pitfalls exist along this analytical path, eg. (i) extraneous metal contamination, (ii) unrecognized aspecific binding of metals to proteins, (iii) unwanted interactions with separation equipment such as chromatography columns and ultrafiltration membranes (Van Landeghem *et al.* 1998a), and (iv) interference problems during quantification.

Hybrid techniques (for a review see Sanz 1998b) combine the capacity to separate ionic species with

specific analytical capacity to quantify them. Attempts to avoid the separation step and rely only on the capacity of the ion spray and triple quadrupole-ICP-MS to reliably quantify different Al species present together in a single sample were not successful (Sharp *et al.* 1997).

Separation techniques

Ion chromatography using size exclusion (either by gel filtration or by HPLC, Rollin & Nogueira 1997) and cation exchange column can be useful in separating Al complexes commonly occurring in the environment (eg. fluo, oxalate, citrate) (Kerven *et al.* 1995b; Borrmann & Seubert 1996, Hils *et al.* 1999; Mitrovic & Milacic 2000; Tsunoda *et al.* 2001) and the blood serum (Rollin & Nogueira 1997; Van Landeghem *et al.* 1998b). The size exclusion is especially suitable for separating organic Al complexes (Rollin & Nogueira 1997; Hils *et al.* 1999). Keeping the temperature of a size exclusion/cation exchange column below 10 °C can minimize decomposition of neutral or negatively charged Al-fluoride and Al-citrate complexes (Busch & Seubert 1999).

Anion chromatography using a gradient elution based on chloride start eluent and perchlorate as final eluent can effectively separate neutral or cationic from negatively charged Al complexes (Borrmann & Seubert 1999). A combination of anion and cation chromatography for more complete separation of Al species is recommended (Borrmann & Seubert 1999). An anion-exchange column and fast protein liquid chromatography (FPLC) can be used to separate low-molecular-weight complexes with Al in the serum (Polak *et al.* 2001a) or to separate negatively charged Al-citrate complexes from neutral Al-citrate species and Al^{3+} in solution (Bantan *et al.* 1998). Hydrolysable polyphenol-bound Al complexes can be separated using Amberlite XAD-7 resin, whereas Chelex 100 can be used for separation of cationic Al species in tea infusions (Erdemoglu *et al.* 2000a).

Quantification techniques

After separation of Al species, quantification can be achieved using Inductively Coupled Plasma Atomic Emission Spectroscopy (ICP-AES) (Bantan *et al.* 1998; Berube & Brule 1999), ICP-Mass Spectrometry (ICP-MS) (Sanz 1998a; Berube & Brule 1999; Hils *et al.* 1999; Tsunoda *et al.* 2001), Graphite Furnace Atomic Absorption Spectrometry (GFAAS) (Sanz 1998a; Berube & Brule 1999), Zeeman AAS

(Rollin & Nogueira 1997) or Electrothermal AAS (ET-AAS) (Van Landeghem *et al.* 1997; Mitrovic *et al.* 1998; Van Landeghem *et al.* 1998b; Mitrovic & Milacic 2000; Polak *et al.* 2001a, b). Electrospray tandem Mass Spectrometry (ES-MS-MS) can also be used to detect low-molecular-weight Al complexes (eg. Al-phosphate, Al-citrate, and ternary Al-citrate-phosphate complexes, Polak *et al.* 2001a), or to reliably quantify Al-malate and Al-citrate in the same sample (Polak *et al.* 2001b). Flame atomic absorption spectrometry was also used occasionally (Erdemoglu *et al.* 2000a, b), even though this technique is less sensitive to Al than other atomic spectroscopy techniques (Rengel 1996).

The techniques like ICP-MS, quadrupole-ICP-MS and double focusing ICP-MS (DF-ICP-MS) offer a significant increase in sensitivity to Al over ICP-AES. In addition, DF-ICP-MS is four times more sensitive than quadrupole-ICP-MS in detecting Al in biofluids (Marchante *et al.* 1999). Using DF-ICP-MS, detectable levels of Al were found in the blood serum from uraemic patients, but not in healthy subjects (Bayon *et al.* 1999).

Other techniques used for quantification of Al species in a range of environmental samples and biofluids are nuclear magnetic resonance (^{27}Al NMR) spectroscopy (Kerven *et al.* 1995a; Masion *et al.* 2000; Wu & Wang 2000; Perry & Shafran 2001; Song *et al.* 2003) and various electrochemical methods. More prominent electrochemical methods are oscillographic for labile monomeric Al species (Bi *et al.* 1999), amperometry for 'free Al' [$\text{Al}^{3+} + \text{Al}(\text{OH})^{2+}$] (Downard *et al.* 1997), automated potentiometry for hydrolysis of Al ions in solutions (Perry & Shafran 2001), derivative adsorption chronopotentiometry on a hanging mercury drop electrode for various Al fractions, especially monomeric Al (Wang *et al.* 2001a), and adsorption stripping voltammetry for monomeric Al in water (Wang *et al.* 2001b) and for complexes between Al and acetylcholine *in situ* (Matlaba *et al.* 2000). For further information on electrochemical as well as atomic spectroscopy techniques used to determine Al species after the separation step, readers are referred to recent reviews (Das & Chakraborty 1997; Sanz 1998b; Van Landeghem *et al.* 1998a; Bi *et al.* 2001b).

Post separation-column reaction with Tiron (Borrmann & Seubert 1996; Busch & Seubert 1999) or pyrocatechol violet (Flaten & Lund 1997) followed by UV photometry is frequently employed. Pyrocatechol violet can also determine Al_{13} complex (Hils *et al.*

1999) that is extremely toxic to plants (Parker *et al.* 1989; Kinraide 1997).

Reactive Al in water is often determined using 8-hydroxyquinoline method followed by 15-s extraction of chelated forms into an organic solvent (Dixon & Gardner 1998). Keeping reaction times with 8-hydroxyquinoline short (3 s) allows separation of reactive Al from the non-toxic AlF^{2+} complexes (Fairman & Sanz 1996), even though the success may not be guaranteed (Simpson *et al.* 1997). Flow-injection analysis can be critical in ensuring precise control of the reaction time, which is important for determining Al species in equilibrium systems (Simpson *et al.* 1997; Pyrzynska *et al.* 2000). Using 8-hydroxyquinoline-5-sulphonic acid in the post column mix with the separation buffer after separation of cations via capillary electrophoresis can result in increased sensitivity, especially for studying Al-F complexes (Zhu & Kok 1998). However, the 8-hydroxyquinoline method (and its variants), as well as some other chromogenic and fluorometric analytical techniques, eg. ferron (Jallah & Smyth 1998), pyrocatechol violet (Flaten & Lund 1997; Hils *et al.* 1999), chrome-azurol S (Simpson *et al.* 1997) or morin (Lian *et al.* 2003), can only be useful in kinetic reaction studies to operationally define Al fractions, but actual speciation within these fractions is frequently uncertain. Even when such separation of Al species was achieved (Tsunoda *et al.* 2001), the sensitivity of ICP-MS as a quantification technique was better than that of the 8-hydroxyquinoline-5-sulphonic acid method.

Computation of aluminium speciation

In addition to a range of analytical techniques for separation and quantification of Al species, researchers resorted to using computer programs, like MINEQL (e.g. Bi 2001; Bi *et al.* 2001a) and GEOCHEM (e.g. Kerven *et al.* 1995a; Stevens *et al.* 1997; Jallah & Smyth 1998; Naumann & Horst 2003), to calculate the activities of individual Al species in aqueous systems and biofluids based on an assumed equilibrium condition and on the set of thermodynamic constants for the reactions under study (Parker *et al.* 1995). However, uncertainties associated with the thermodynamic constants (Harris *et al.* 1996) make computational speciation of complex, Al-containing aqueous solutions imprecise, yielding only rough estimates of activities of Al species.

Aluminium species in environmental samples and biofluids

Toxicity of aluminium species

Bioavailability and thus toxicity of various Al species to living organisms is different. Aluminate ion $\text{Al}(\text{OH})_4^-$ is non-toxic to fish (Poleo & Hytterod 2003) and plants (Kinraide 1990), even though relatively high aluminate concentration at $\text{pH} > 9.0$ can be toxic to wheat (Ma *et al.* 2003). Aluminium trihydroxide is non-toxic to plants (e.g. Yoshimura *et al.* 2003) and humans (it is used frequently as a phosphate-complexing agent, e.g. Van Landeghem *et al.* 1998b). The monomeric Al species $\text{Al}(\text{OH})^{2+}$ and $\text{Al}(\text{OH})_2^+$ are non-toxic to plants (Kinraide 1997), but no definite information could be found for animal/human cells (even though these Al species were included in the sum of Al species presumed toxic to aquatic organisms, Bi 2001). Aluminium-phosphate is non-toxic to plant (Takita *et al.* 1999) and animal/human cells (because it is present in serum of healthy individuals, Polak *et al.* 2001a). Aluminium complexes with sulphate are harmless to plants (Kinraide 1997) and presumably also to animals and humans.

Aluminium complexes with organic acids are non-toxic to plants (see Matsumoto 2000). However, in animals and humans, Al-citrate complex is present in substantial amounts in serum of healthy individuals (Bantan *et al.* 1999b; Polak *et al.* 2001a), but is also (i) the main Al complex in the cerebrospinal fluid of acutely Al-intoxicated dialysis patients (Van Landeghem *et al.* 1997), and (ii) one of the two major forms of Al taken up across the plasma membrane of brain cells (Yokel *et al.* 2002). Hence, Al-citrate in animal/human cells may not be toxic itself, but may represent a vehicle for delivering Al to sensitive sites, whereby upon changes in Al speciation, toxicity may occur. Among other Al-organic complexes, lipophilic ones (like Al-acetylacetonate and Al-maltolate) are cytotoxic to murine neuroblastoma cells (Zatta & Zambenedetti 1996), Al-maltolate is toxic to neuronal and astroglial cells (Levesque *et al.* 2000), and Al-lactate causes toxicity in mice brain by affecting acetylcholine metabolism (Zatta *et al.* 2002). In contrast, binding of Al to organic ligands eliminates Al toxicity to aquatic organisms (Bi *et al.* 2001a).

Regarding Al-F complexes, early reports indicated their non-toxicity to plants (Cameron *et al.* 1986; MacLean *et al.* 1992), but recent work suggested increasing toxicity of Al-F complexes with their in-

creasing positive charge (Kinraide 1997; Stevens *et al.* 1997, 1998). The Al-F complexes are non-toxic to fish and other aquatic organisms (Bi *et al.* 2001a). In human cells, the Al-F complex resulted in the lowest Al accumulation and had the lowest toxic effect on viability of neuronal and astroglial cells compared to the other three Al complexes studied (maltolate, lactate and chloride) (Levesque *et al.* 2000).

It remains to be established whether non-toxicity of certain Al species and complexes is related to (i) a lack of uptake across the plasma membrane, or (ii) decreased (or an absence of) the positive charge (eg. plant toxicity of ions generally increases with positive charge, Kinraide 1997). Indeed, polynuclear hydroxy-Al complexes with a high positive charge (eg. Al_{13} , $\text{AlO}_4\text{Al}_{12}(\text{OH})_{24}(\text{H}_2\text{O})_{12}^{7+}$, Parker & Bertsch 1992) are more toxic to plants than monomeric, less charged species (Comin *et al.* 1999). The Al_{13} species is also toxic to bacteria (Amonette *et al.* 2003), but no information could be found on its potential toxicity to animal/human cells.

Aluminium speciation in water

Aluminium in water is mainly bound to the colloidal matter (55%) compared to the soluble fraction (23%), with the rest being in a particulate form (Gundersen & Steinnes 2003). However, depending on the source of raw water (water destined for drinking, but before treatment), up to 80% of Al can be in the particulate form (Schintu *et al.* 2000). After water treatment (coagulation, flocculation and filtration, which results in 'finished water' that is used for drinking) most Al (between 40 and 62%) was in the monomeric form (Schintu *et al.* 2000).

Polyaluminium chloride or aluminium sulphate is regularly used in treatment of water for drinking. New coagulant has been produced recently by combining polyaluminium chloride and polysilicic acid (polyaluminium silicate chloride). It has an increased coagulation capacity and results in a lower proportion of Al_{13} species compared with polyaluminium chloride (Song *et al.* 2003), and may therefore be better to use.

The regular drinking water may contain substantial amounts of Al (average about $70 \mu\text{g L}^{-1}$). However, bioavailability of Al from drinking water is estimated to be low, with only 0.3% of the total Al amount ingested in water being absorbed into the body (Yokel & McNamara 2001). This appears to be a small percentage given potentially high proportion of bioavailable monomeric Al forms in drinking water (eg. up to

62%, Schintu *et al.* 2000). However, even with a high bioavailability of Al in drinking water, Al absorption from water may still be a small proportion of the total amount of Al ingested. For example, regular antacid users ingest gram amounts of Al daily, which may be three to four orders of magnitude greater than ingestion of Al from drinking water (see Flaten 2001; Yokel & McNamara 2001).

Out of a range of Al species in drinking water in the Quebec study [total Al, total dissolved Al, monomeric organic Al, monomeric inorganic Al, polymeric Al, Al^{3+} , $\text{Al}(\text{OH})^{2+}$, AlF^{2+} , $\text{AlH}_3\text{SiO}_4^{2+}$, $\text{Al}(\text{SO}_4)^+$], a significant relationship was found between the long-term exposure of subjects to monomeric organic Al and Alzheimer's disease (Gauthier *et al.* 2000). Many other studies (nine out of thirteen studies considered) also found a positive relationship between Al in drinking water and Alzheimer's disease (for a review see Flaten 2001).

Aluminium speciation in plants

In the plant sap of *Sempervivum tectorum* and *Sansevieria trifasciata*, Al-citrate and Al-aconitate were the only low-molecular-weight Al complexes detected (Bantan *et al.* 1999a). In contrast, type of Al species supplied in the root medium influenced the Al form transported in the xylem sap of Al-tolerant Chinese cabbage (*Brassica rapa pekinensis*). When Al^{3+} was the predominant Al form in the nutrient solution, Al complexation to malate occurred in the root tissue, and Al-malate was the dominant form in the xylem sap (70%), with Al^{3+} being the other fraction (30%), whereas supplying Al to the solution as Al-citrate or Al-malate resulted in these same forms being exclusively present in the xylem sap (Polak *et al.* 2001b).

Aluminium speciation in food

Aluminium speciation in food has been reported relatively rarely (for a review see Marchante *et al.* 1999), even though a need for more research of that type has been emphasised. Clearly, the variety of foods and a range of Al-complexing ligands represent a challenge in trying to speciate Al in realistic food mixtures. Speciation of Al in simulated food solutions (pH 4–8) suggested that most Al is present in the organic complexes, most of which have relatively high bioavailability and thus potential toxicity to humans (Bi 1996). Most work on Al species in food solutions

was done on dietary organic acids (Venturini & Berthoin 1998, 2001; Desroches *et al.* 1999; Desroches *et al.* 2000; Dayde *et al.* 2003); see the section on Al uptake by animal and humans.

Cooking and preparing food in Al-containing containers result in migration of Al into the food, especially in acidic and salty environments (Gramiccioni *et al.* 1996). Al-complexing agents (e.g. organic acids, fluoride, OH^- , etc.) can also increase release of Al from Al-containing cooking utensils (Bi 1996).

Aluminium speciation in human and animal biofluids

The main low-molecular-weight Al complexes in blood serum of healthy human subjects were Al-citrate, Al-phosphate and ternary Al-citrate-phosphate complexes (Bantan *et al.* 1999b; Polak *et al.* 2001a). The relative ratio of these complexes varied between subjects tested, but no specific difference was found between patients on continuous ambulatory peritoneal dialysis and healthy subjects (Polak *et al.* 2001a).

In serum, Al is mainly transported in the complex with high-molecular-weight Fe-transporting protein transferrin (Harris *et al.* 1996), whereas albumin (Harris *et al.* 1996; Rollin & Nogueira 1997) and ferritin (Ward *et al.* 2001) do not appear to have a role in Al binding in serum. Increased Fe load in humans may decrease Al binding to transferrin; this effect is not due to interference by Fe in Al absorption in the gastrointestinal tract, but rather due to direct Fe competition with Al for binding to transferrin in blood (Van Landeghem *et al.* 1998b). Hence, a negative correlation (albeit relatively weak) was found between transferrin saturation with Fe and Al levels in serum (Van Landeghem *et al.* 1998b).

Occupational exposure to Al can increase the proportion of Al bound to high-molecular-weight complexes (largely transferrin) compared to Al bound in low-molecular weight complexes in serum (Rollin & Nogueira 1997). Microultrafiltration through a membrane filter (e.g. with a cut-off point of 30 kDa) can be used to effectively separate the transferrin-type high-molecular-weight Al complexes from low-molecular-weight ones in serum (Polak *et al.* 2001a). However, even a small proportion of Al bound to low-molecular-weight complexes at high total Al concentration caused by occupational exposure means that relatively high amounts of low-molecular-weight Al complexes are present in serum of such patients compared with healthy individuals. High content of low-molecular-weight Al complexes in serum sug-

gests high toxicity because of ease by which these complexes can be transferred into cells (Rollin & Nogueira 1997).

The main form of Al in the cerebrospinal fluid of acutely Al-intoxicated dialysis patients is Al-citrate/silicate (Van Landeghem *et al.* 1997), without any Al-transferrin complexes detected. This result is in contrast to Al-transferrin complex being the important form of Al uptake into brain cells across the blood-brain barrier (Van Landeghem *et al.* 1998a; Yokel 2000, Yokel & McNamara 2001).

Analytical techniques for measuring aluminium in cells and tissues

Analytical techniques for quantifying Al in cells and tissues can be divided into two groups (i) for measuring Al *in situ* in tissue sections and individual cells, and (ii) for determining total Al content in cells, tissues and organs.

Important *in situ* techniques used for measuring Al accumulation in plant and animal/human tissues, cells and cell compartments are: energy-dispersive X-ray analysis (EDXMA, see e.g. Peri & Good 1990; Memon *et al.* 1981; Marienfeld & Stelzer 1993; Ownby 1993; Marienfeld *et al.* 1995; Echlin 1996), particle induced X-ray emission (PIXE, Watt 1996; Schofield *et al.* 1998; Irigaray *et al.* 2000; Clark *et al.* 2001), secondary ion mass spectrometry (SIMS, Candy *et al.* 1992; Chaussidon *et al.* 1993; Lazof *et al.* 1994; Jones *et al.* 1997; Mangabeira *et al.* 1999; Florent *et al.* 2001; Kishikawa *et al.* 2003) and laser microprobe mass analysis (LAMMA, Peri & Good 1990; Kuhn *et al.* 1995; Iancu *et al.* 1996; Reusche *et al.* 1999; Marienfeld *et al.* 2000). Sometimes a combination of techniques is used for a maximum effect, eg. to determine Al in neuritic plaques in Alzheimer's patients, PIXE (for trace element concentrations) was combined with Scanning Transmission Ion Microscopy (for imaging features in the tissue), and Rutherford Backscattering Spectrometry (to characterize the tissue matrix) (Watt 1996).

Techniques for measuring total Al in plant and animal specimens are: Graphite Furnace Atomic Absorption Spectroscopy (GFAAS, Zhang & Taylor 1989, 1990; McDonald-Stephens & Taylor 1995; Archambault *et al.* 1996; Vitorello & Haug 1996; Ranau *et al.* 1999; Taylor *et al.* 2000; Baffi *et al.* 2002), Inductively Coupled Plasma Atomic Emission Spectroscopy (ICP-AES) (Asano *et al.* 2002; Baffi *et al.*

2002; Rahil *et al.* 2002; Chan & Lo 2003; Juranovic *et al.* 2003; Lesniewicz & Zyrnicki 2003; Piczak *et al.* 2003), ICP-Mass Spectrometry (ICP-MS) (Baffi *et al.* 2002; Nystrom *et al.* 2002; Adamo *et al.* 2003; Chan & Lo 2003; Noel *et al.* 2003); colourimetric and fluorometric techniques (Reid *et al.* 1996; Vitorello & Haug 1996, 1997; Yoshimura *et al.* 2003), ion chromatography (Rincón & Gonzales 1992), and ^{27}Al NMR (Nagata *et al.* 1992; Ma *et al.* 1997; Watanabe *et al.* 1998; Xia & Rayson 1998; Shen *et al.* 2002; Yoshimura *et al.* 2003).

Techniques for measurement of aluminium *in situ*

The theoretical principles of the *in situ* analytical techniques are beyond the scope of the present review; readers can find more information in Rengel (1996). Briefly, all these techniques rely on bombardment of the specimen with a beam of either ions (SIMS and PIXE), electrons (EDXMA) or photons (LAMMA), causing either ejection of surface atoms/ions that are analysed by mass spectrometry (SIMS and LAMMA) or knocking out of some shell electrons that give rise to element-specific primary X-rays (EDXMA and PIXE). The EDXMA and SIMS are superior to the other two and will be considered further.

Potential advantages of SIMS are high elemental sensitivity, 3-D analysis and capacity to provide isotopic information, whereas a disadvantage is non-uniform ion ejection (sputtering) that may cause different cell constituents to sputter at different rates (Linton *et al.* 1980), resulting in lateral resolution of SIMS being insufficient for identification of sub-cellular structures. However, Lazof *et al.* (1994) estimated that lateral resolution for SIMS was in the order of 2 μm (as for comparison, the cell wall of epidermal root cells might be several micrometers thick, Marschner 1995). Other methods used to study concentration of Al *in situ* can potentially have better spatial resolution (down to 0.1 μm for EDXMA, Linton *et al.* 1980) but, due to limitation of sample preparations and the tear-drop shape of the primary electron beam penetrating biological samples, the minimal limit of lateral resolution for EDXMA may be considerably greater, possibly exceeding even the diameter of the individual cell (see discussion in Lazof *et al.* 1994, 1997). In contrast, Marienfeld & Stelzer (1993) suggested that the spatial resolution of EDXMA is sufficient for distinguishing between cytoplasm and the nucleus, but that small vesicles in meristematic and

differentiating cells may interfere with measurements of the cytoplasmic mineral content.

Assuming no spectral interferences, a detection limit for Al of around $1 \mu\text{g g}^{-1}$ (37 nmol g^{-1}) can be achieved by SIMS (Lazof *et al.* 1994, 1997), which is at least one (Linton *et al.* 1980) or even two orders of magnitude lower than for EDXMA (Marienfeld & Stelzer 1993).

There are several recent advancements in the traditional SIMS technique. Resonance Ionisation Mass Spectrometry (RIMS) is superior to SIMS in Al quantification because RIMS alleviates the isobaric interference problems with compounds like CNH (Jones *et al.* 1997). Time-of-Flight SIMS is also more sensitive than traditional SIMS and is expected to be important in quantifying Al concentrations in biological specimens (Kishikawa *et al.* 2003).

The new nanoSIMS technology, currently used in the author's laboratory to determine Al transport across the wheat root-cell plasma membrane, is considerably better (lateral resolution of 50 nm and sensitivity about $0.1 \mu\text{g Al g}^{-1}$ tissue) than SIMS. This lateral resolution allows distinguishing Al accumulated in the cell wall from Al that has crossed the plasma membrane and is situated in the cytosol or in subcellular compartments.

Localisation of aluminium in animal and human cells and tissues in situ

The dynamic SIMS technique was used to detect Al accumulation in cortical pyramidal neurons of dialysis patients (without Alzheimer's diagnosis) who were given large doses of Al-containing phosphate-binding agents (Candy *et al.* 1992). Electrothermal atomic absorption spectroscopy revealed increased Al content in brains of Alzheimer's patients compared to controls (Kruck 1993). Aluminium (as indicated by the fluorescent morin staining) was concentrated in white matter of the medial striatum, corpus callosum, and cingulate bundle (Platt *et al.* 2001) (see also Senitz & Bluethner 1990). The nuclear microscopy technique, however, failed to detect Al in pyramidal neurons of Alzheimer's disease patients (Makjanic *et al.* 1998), leading authors to suggest that the previous findings of Al in neurofibrillary tangles were caused by contamination or elemental redistribution during preparation of tissue for analysis.




Figure 1. Aluminium accumulation in the wheat (*Triticum aestivum*) root tips. After Al exposure at different temperatures, roots were desorbed in 0.5 mM citrate (pH 4.5) for 30 min. Compiled from Zhang & Taylor (1989, 1990).

Uptake of aluminium

Characterisation of Al transport across biological membranes has been hindered by the complex aqueous coordination chemistry of Al and the lack of sensitive analytical techniques for detecting the low levels of Al associated with subcellular compartments. Hence, the long exposure times (>24 h) and concentrations of Al high enough to cause precipitation or polymerisation of Al had to be used. For plant cells, there are additional challenges because the cell wall (which accumulates large amounts of Al) cannot generally be separated from the cell symplasm (which has only a relatively low concentration of Al). Desorption of Al from the cell wall cannot be achieved anywhere near to completion in roots (Figure 1) and algal cells (Table 1) regardless of the desorption method used. Hence, the rates of Al uptake were grossly overestimated (Figure 1, Table 2).

Recent studies on Al uptake across the plasma membrane of plant cells (Reid *et al.* 1996; Taylor *et al.* 1996, 2000) have focused on the giant-cell green alga *Chara corallina*, in which surgical separation of the cell wall from the cytoplasm can be achieved, thus eliminating problems inherent in measuring exceedingly small amounts of intracellular Al in the very high background of apoplasmic Al. In brief, a single giant cylindrical algal cell (approx. 0.8 mm diameter and up to 80 mm length) is mounted into a 3-compartment chamber and the middle compartment exposed to the Al-containing solution. Upon termination of the uptake period, cell ends which were not in contact with




Figure 2. Short-term accumulation of ^{26}Al within subcellular fractions (cell wall, cytoplasm and vacuole) of single cells of *Chara corallina* as measured by Accelerated Mass Spectrometry. Cells were exposed to $50\ \mu\text{M}$ AlCl_3 in $0.4\ \text{mM}$ CaCl_2 spiked with $75\ \text{pCi}$ of ^{26}Al (pH 4.3), followed by desorption for 30 min in $1.0\ \text{mM}$ citric acid (4°C , pH 4.3) (to remove exchangeable Al from the cell wall) and surgical separation of cell wall, cytoplasm and vacuole. Note different units for ^{26}Al accumulation in the cell wall (top graph) and the cytoplasm or vacuole (bottom two graphs). Values represent the mean $\pm$ SE of three replicates. Modified from Taylor *et al.* (2000).

Table 1. The ratio of Al content in the cell wall and the symplasm of giant-cell alga *Chara corallina*. Cells were exposed to artificial pond water containing 10 to $100\ \mu\text{M}$ Al for up to 7 days in a number of experiments. After termination of Al treatment, desorption of cell wall Al was performed by a method indicated, followed by surgical separation of the cell wall and the intracellular content. Modified from Reid *et al.* (1996).

Al content in cell wall / / Al content in the symplasm	Desorption method
60	1.5 h 5 mM citrate/ $\text{Ca}(\text{OH})_2$, pH 5.0
22	4.5 h 10 mM citrate/ $\text{Ca}(\text{OH})_2$, pH 4.4
28	2 h 10 mM CaCl_2 + 1 mM citrate/ NaOH , pH 4.5
>20	2 h 1 mM LaCl_3 + 2 mM citrate/ $\text{Ca}(\text{OH})_2$, pH 4.4
30	2 h 1 mM $\text{La}(\text{NO}_3)_3$, pH 4.4

Table 2. Rates of Al uptake across the plasma membrane of plant cells

Species	Method	Al uptake rate $\mu\text{mol g}^{-1} \text{FW h}^{-1}$	References
<i>Triticum aestivum</i> (wheat)	GFAAS	0.3–0.4	Zhang & Taylor (1989, 1990); Archambault <i>et al.</i> (1996)
<i>Glycine max</i> (soybean)	SIMS	0.3	Lazof <i>et al.</i> (1994)
<i>Chara chorallina</i>	fluorimetry	$0.01\text{--}0.3 \times 10^{-3}$	Reid <i>et al.</i> (1996)
<i>Chara chorallina</i>	^{26}Al and AMS	up to 0.005×10^{-3}	Taylor <i>et al.</i> (1996, 2000)

GFAAS = Graphite Furnace Atomic Absorption Spectroscopy.

AMS = Accelerator Mass Spectrometry.

Al solution at all are cut, syringe needle inserted, and cell content flushed out by rapidly injecting deionized water through the cell lumen (Reid *et al.* 1996). Such advantage of physical separation of cell wall and the cytosol (Reid *et al.* 1996) (Table 1), especially when coupled with increased sensitivity due to usage of radioactive isotope ^{26}Al (Taylor *et al.* 2000), may represent the best model system for quantifying the magnitude and timing of Al flux across the plasma membrane of plant cells (Table 2).

Labelling with the radioactive isotope of aluminium

An accurate quantification of kinetics of transmembrane uptake of Al in intact organisms, cells, organelles and/or membrane vesicles can be achieved using a suitable tracer. The ^{26}Al is the only Al isotope suitable for tracer work (e.g. Taylor *et al.* 2000). It is a weak β and γ emitter, with the long half-life (approximately 7.3×10^5 y). The conventional mass spectrometry is not sensitive enough for measuring ^{26}Al . In contrast, Accelerator Mass Spectrometry (AMS) allows acceleration of atomic and molecular ions to MeV energies as opposed to KeV, providing sufficient sensitivity to resolve ^{26}Al from ^{27}Al at atom ratios as low as 10^{-14} and a detection limit in the 10^{-17} g range (see Taylor *et al.* 2000). Hence, combining ^{26}Al with AMS detection provides sufficient sensitivity to detect Al in subcellular fractions isolated from cells exposed to realistic environmental Al concentrations for a short period of time.

The ^{26}Al isotope has been used to monitor Al uptake by plant species *Gleditsia triacanthos* and *Pinus taeda* (Schaedle *et al.* 1989) and by internodal cells of the giant algae *Chara corallina* (Taylor *et al.* 1996, 2000). The ^{26}Al -AMS is also beginning to play a role in animal and human physiology, including studies of Al absorption in the human gastrointestinal tract (Priest *et al.* 1998; Jugdaohsingh *et al.* 2000; Moore

et al. 2000) and uptake of Al by brain and other animal and human tissues (Yokel & McNamara 2001).

Aluminium species transported across the plasma membrane and existing in the symplasm

It is unclear at present which Al species cross the plasma membrane due to uncertainty in Al speciation. Negative charges in the cell wall of plant cells as well as on the plasma membrane of all cells would alter the Al speciation in vicinity of these cellular structures compared with Al speciation in the bulk solution. Therefore, determining the identity of Al complexes in contact with the plasma membrane and the time-course of their transfer into the cytosol are beyond the limits of the current experimental techniques.

Upon transfer of Al species from outside of the plasma membrane into the symplasm, speciation of Al is altered because of higher pH (near-neutral) in the cytosol compared to the medium outside the cell. Hence, cytosolic Al is expected to be complexed with hydroxyl ions, high-molecular-weight substances, phosphate-containing compounds, and/or Ca-dependent regulatory cytosolic protein calmodulin (see Martin 1988; Taylor 1991; Rengel 1996). Nevertheless, a small proportion of cytosolic Al may remain in the free ionic form in dynamic equilibrium with a larger pool of complexed Al. Increasing the pool of complexed Al over time would inevitably result in an increase in the pool of the free ionic Al that may potentially affect cellular structures and processes after a prolonged exposure to Al. Concentrations of free Al in the nanomolar to sub-nanomolar range *in vitro* have affected polymerization of tubulin (Macdonald *et al.* 1987; Martin 1988) and activity of protein kinase C (see Rengel, 1996); these concentrations of free Al may be expected to occur in the cytosol of Al-intoxicated cells.

Aluminium uptake across the plasma membrane of plant cells

Transport of ^{26}Al across the plasma membrane of *Chara corallina* showed a clear maximum at pH 4.3 ($176 \text{ ng m}^{-2} \text{ min}^{-1}$) (Taylor *et al.* 2000), with rates of transport approximately an order of magnitude lower at pH 3.7 and 5.2, suggesting that Al^{3+} may be the main species transported. A decrease in uptake at pH 5.2 was caused by a lowered proportion of Al^{3+} , whereas lower uptake at pH 3.7 was likely due to competition with H^{+} (cf. Kinraide 1993).

When total Al was kept constant, but $\{\text{Al}^{3+}\}$ was reduced in the presence of sulphate and citrate, rates of Al transport across the plasma membrane of *Chara corallina* increased by 2.1-fold in the presence of SO_4^{2-} and by 2.0-fold in the presence of citric acid (Taylor *et al.* 2000), suggesting that complexes such as AlSO_4^{+} , AlCit_{H-1}^{-} , or AlCit^0 may be readily transported across the plasma membrane. Indeed, in animal and human cells, Al-citrate is one of the two major forms of Al taken up across the plasma membrane of brain cells (Yokel *et al.* 2002).

In contrast to animal cells, plant cells have the vacuole as a relatively large organelle where toxic ions are stored to decrease their metabolic impact (Rengel 1997). In *Chara corallina*, transport of Al into the vacuole was limited during the first 30 min of exposure, then increased to a rate that represented a large portion of Al across the plasma membrane (Figure 2) (Taylor *et al.* 2000). This suggests that (i) Al may need to accumulate in the cytoplasm before transport to vacuole can occur, and/or (ii) induction of a transport system capable of moving Al across the tonoplast may be required. Given predictions about the limited solubility of Al in the cytoplasm and its propensity to bind to biological ligands (Martin 1988; Taylor 1991; Rengel 1996; Taylor *et al.* 2000), such a transport system is likely to facilitate transport of organic Al complexes into the vacuole.

Aluminium absorption in the animal/human digestive tract and uptake by cells and tissues

The average daily Al intake by US adults is 8–9 mg for males and around 7 mg for females (Pennington & Schoen 1995). In Italian population, daily Al intake ranges from 2.5 to 6.3 mg (Gramiccioni *et al.* 1996), and in UK is 3.4 mg on average (Ysart *et al.* 2000). The 'normal' daily intake of Al is defined as 5 mg Al by the World Health Organisation (WHO 1996). About 0.1% of daily amounts of Al taken stays in the

body (less than $10 \mu\text{g}$), whereas the rest is excreted (Priest *et al.* 1998).

Bioavailability of Al (which determines Al uptake, retention in the body and thus Al toxicity) is vastly different in different sources, ranging from 0.012% in antiperspirant applications and 0.3% in drinking water to 1.5% from inhaled airborne soluble Al and virtually 100% from intramuscular injections (e.g. some vaccines contain Al) (Priest *et al.* 1998; Stauber *et al.* 1999; Yokel 2000; Yokel & McNamara 2001). Bioavailability of Al in food, however, is influenced by numerous factors (see section on Al speciation in food) and would have to be determined on a case-by-case basis.

Dietary acids, by their intrinsic acidity and co-ordinating capacity, can extend solubility of Al^{3+} and thus increase Al bioavailability during longer sections of the gastrointestinal tract, and also help Al^{3+} diffusion across the intestinal epithelium through the formation of neutral complex species (Desroches *et al.* 2000; Dayde *et al.* 2003) (in small intestine alkaline pH would normally eliminate any Al absorption). Hence, these dietary acids, e.g. glutamic acid, an essential amino acid and a common food additive (Dayde *et al.* 2003); malic acid, added to many beverages and processed foods as well as present in vegetables and fruits (Venturini & Berthon 2001) and tartaric acid, found commonly in fruits and industrial drinks and food (Desroches *et al.* 2000) can serve as organic ligands, enhancing Al absorption in the gastrointestinal tract and thus aggravating Al toxicity. It is therefore advisable to avoid simultaneous ingestion of these organic ligands when using any therapeutic aluminium salt (Venturini & Berthon 2001).

Absorption of Al from drinking water in the gastrointestinal tract of rats was dependent not only on the concentration of Al in water, but even more so on the Al-binding capacity of food components in the stomach determining the Al bioavailability (Glynn *et al.* 1999a). Indeed, when the stomach content of rats was fractionated for Al, large portion of Al ingested in acidic water (pH 2.5–3.0) was found to be insoluble, even when citrate, fluoride and silicate as Al-complexing agents were included in the water (Glynn *et al.* 2001). In other studies with rats and mice, however, addition of fluoride caused higher Al absorption from the gastrointestinal tract (Allain *et al.* 1996). Hence, bioavailability of Al cannot be measured in water or food before ingestion because large changes may occur in the gastrointestinal tract.

The study employing radioactive ^{26}Al isotope and quantification by accelerated mass spectrometry showed that only 0.06 to 0.1% of ingested Al was absorbed in the gastrointestinal tract of healthy aged patients after 1 h of drinking Al-laced fruit juice. However, the proportion of Al absorbed was 1.64 times greater in aged-matched Alzheimer's sufferers, suggesting an increased capacity of Alzheimer's disease patients to take up Al from food (Moore *et al.* 2000).

Aluminium concentration in blood serum is increased upon administering Al to the animals (Glynn *et al.* 1999b). Over time, Al accumulates in liver (Lote *et al.* 1995; Glynn *et al.* 1999b; Swegert *et al.* 1999; Esparza *et al.* 2000), brain (Candy *et al.* 1992; Swegert *et al.* 1999), heart (Swegert *et al.* 1999) and especially in bones (Glynn *et al.* 2001; Vanholder *et al.* 2002), affecting their mechanical properties (Hui & Ellers 1999).

An important component of Al uptake into the brain cells across the blood-brain barrier is likely to be transferrin-receptor mediated endocytosis (Van Landeghem *et al.* 1998a; Yokel 2000; Yokel & McNamara 2001), even though Al-transferrin complexes are not always found in the cerebrospinal fluid (Van Landeghem *et al.* 1997). An additional pathway of Al entry is uptake of Al-citrate complex into the mouse brain endothelial cells via a carrier, probably a monocarboxylate transporter or organic anion transporter (Yokel *et al.* 2002). Interestingly, uptake of Al-citrate is 70% greater than uptake of citrate, even though citrate is one of primary substrates for transporters suggested to be involved in Al-citrate transport. Neither the monocarboxylate transporter 1 (MCT 1) nor anion exchanger was involved in Al-citrate uptake by brain cells (Yokel *et al.* 2002).

Aluminium is effluxed from the brain tissues as Al-citrate (Yokel 2000), probably via a carrier (Yokel & McNamara 2001). However, Al accumulates in the brain tissues, potentially causing compounded damage (Yokel 2000).

Excretion of Al from the body can be via faeces in healthy humans (Priest *et al.* 1998), but is also renal [and may be enhanced by citrate (Lote *et al.* 1995) or fluoride (Chiba *et al.* 2002)], with <2% being excreted in bile (Yokel & McNamara 2001). Intravenous infusion of citrate can facilitate renal excretion of Al (Lote *et al.* 1995).

Plant species that accumulate aluminium in cells and tissues

Many native plant species grow vigorously on acid soils that contain high concentrations of bioavailable Al. For these plant species, Al can be considered a beneficial element (Kinraide 1993; Osaki *et al.* 1997) because growth is stimulated when a high concentration of Al is present in the growing medium (Watanabe *et al.* 1997; Watanabe & Osaki 2002). These plant species can be divided into Al-excluders and Al-accumulators (for a review see Watanabe & Osaki 2002), the latter being of more interest to discussion here. Al-accumulators can have leaf Al concentrations of up to 37 g Al kg^{-1} dry weight (DW) (Masunaga *et al.* 1998), which are concentrations similar to nitrogen and potassium as the two most abundant essential nutrients present in plant tissue.

The Al-accumulator species belong to various families (eg. Euphorbiaceae, Melastomataceae, Rubiaceae, Theaceae) (Owour & Cheruiyot 1989; Masunaga *et al.* 1997, 1998); many are shrubs and trees, but some are herbaceous species. Examples of Al-accumulator plants are *Camellia sinensis* (tea, Owour & Cheruiyot 1989), *Hydrangea macrophylla* (hydrangea, Takeda *et al.* 1985; Osaki *et al.* 1997; Naumann & Horst 2003), *Fagopyron esculentum* (buckwheat, Osaki *et al.* 1997; Shen *et al.* 2002), *Aporosa frutescens*, *Aporosa maingayi*, *Euphorbia acuminata*, *Melastoma laurinum*, *M. corymbosus* (Masunaga *et al.* 1998), *M. malabatricum* (Osaki *et al.* 1997; Watanabe *et al.* 1997; Osaki *et al.* 1998), *Tibouchina pulchra* (Domingos *et al.* 2003), *Dioclea guianensis* (Izaguirre & Flores 1995) and *Stylosanthes elegans* (Izaguirre *et al.* 2002).

The Al-accumulators may accumulate Al in roots (Osaki *et al.* 1997; Izaguirre & Flores 1995; Izaguirre *et al.* 2002), but exclude it completely from root nodules of N_2 -fixing plant species (Izaguirre *et al.* 2002). Most Al-accumulators accumulate Al in leaves, especially old ones (Owour & Cheruiyot 1989; Memon *et al.* 1981; Nagata *et al.* 1992; Masunaga *et al.* 1998; Shen *et al.* 2002). However, accumulation of Al in young, metabolically most productive leaves may occur, for example in *Melastoma malabatricum* (Osaki *et al.* 1997, 1998; Watanabe *et al.* 1997, 1998) with up to 10 g Al kg^{-1} DW in old and up to 7 g Al kg^{-1} DW in new leaves.

In contrast to other Al-accumulators, in hydrangea Al accumulates in sepals (Takeda *et al.* 1985; Naumann & Horst 2003). Sufficient transport of Al-citrate

into sepals of hydrangea is essential to achieve the commercially desirable colour change from original pink to blue by complexation of the anthocyanin delphinidine (Naumann & Horst 2003).

In Al-accumulators, Al is usually complexed with organic acids or other organic compounds to make it non-toxic. The predominant Al form is Al-catechins in leaves of tea plant (Nagata *et al.* 1992), Al-citrate in hydrangea leaves (Ma *et al.* 1997) and Al-oxalate in buckwheat (Shen *et al.* 2002) and *Melastoma malabatricum* leaves (Watanabe *et al.* 1998), rendering the high total tissue concentrations non-phytotoxic to the cell cytosol. In addition, the cytosol is protected by Al accumulating predominantly in the cell wall (Memon *et al.* 1981; Shen *et al.* 2002) or vacuoles (Echlin 1996; Shen *et al.* 2002) of tea and buckwheat leaf cells. Binding of Al in the cell wall is mainly to pectic substances, as shown in *Melastoma malabatricum* (Watanabe *et al.* 1998).

In hydrangea, Al-citrate was the main transporting Al form in the xylem sap (Naumann & Horst 2003) as well as the main Al form accumulating in leaves (Ma *et al.* 1997). In contrast, in *Melastoma malabatricum*, Al-citrate was the main Al transporting form in the xylem sap, whereas the main form in leaves was Al-oxalate (Watanabe & Osaki 2001, 2002).

Aluminium in the continuum of soil / tea plant / tea drink / humans

Tea drink is the second (after water) most widely drunk beverage in the world. Given that tea plant *Camellia sinensis* is grown on acidic soils and is the Al-accumulator, there is an obvious need to clarify the potential transfer of Al from soil to tea leaves to tea infusions and into a human body.

Increasing soil acidity results in higher bioavailability of Al and higher uptake of Al by tea roots. Indeed, Al accumulation in tea leaves was correlated with CaCl₂-extractable (bioavailable) Al in subsoil, and was especially prominent when soil pH dropped below 5.0 (Dong *et al.* 1999). High Al content was present in tea leaves coming from Chinese provinces with highly acidic soils (Wong *et al.* 1998).

In tea leaves, the difference between Al concentration in young and old leaves can be substantial (Table 3) (Nagata *et al.* 1992). Hence, Al content in various tea preparations (e.g. green, oolong, black, brick tea) differs depending on the maturity of leaves collected (only a bud and the two youngest leaves are used for green tea, whereas more mature leaves are

Table 3. Aluminium concentration in various parts of *Camellia sinensis* (tea) cultivars grown under diverse environmental conditions.

Plant part	Al content (g kg^{-1} dry weight)	
	Adapted from Nagata <i>et al.</i> (1992)	Adapted from Wong <i>et al.</i> (2003)
Buds	0.07	1–2
Young leaves	0.3	2.4–11
Mature leaves	5.5	8–15
Old leaves	8.3	12–18

collected for other types of tea, Wong *et al.* 1998). The presence of branches (eg. in brick tea) can contribute to Al content in brick tea being more than 4 times higher than in green tea (Wong *et al.* 2003).

Brick tea infusions can have total Al concentration of 15 mg Al L⁻¹ after 1 h of infusion, whereas other types of tea contain between 0.6 and 1.9 mg Al L⁻¹ (Wong *et al.* 2003). Hence, drinking just 0.3 L of brick tea per day would supply the amount of Al estimated as 'normal' (5 mg L⁻¹, WHO 1996). The extraction of Al from tea leaves (1% w/v mixture) is about 35% in the first infusion and about 14% in the second for most common tea types (green, black, oolong) (Wong *et al.* 1998).

Aluminium species in tea infusions

Although high total Al concentrations can be found in tea infusions (like 0.1 mM), most of that Al is complexed so strongly that even Chelex 100 cannot exchange Al from these complexes (Alberti *et al.* 2003), resulting in low chemical and biological reactivity of Al in tea infusions. About 10% of total Al in tea infusions may be bound to fluoride (Erdemoglu *et al.* 2000b). However, the proportion of Al-F complexes increases significantly when simulated gastric juice (pH 2.5) was added to tea infusion (Liang *et al.* 1999). The Al-F complexes decrease absorption of Al by neural cells (Levesque *et al.* 2000).

In black tea infusions, 10–19% of total Al was present as cationic species, whereas 28–33% was present as hydrolysable polyphenol complexes (Erdemoglu *et al.* 2000a). In green tea infusions, the most prevalent complex was Al-oxalate (Liang *et al.* 1999). In contrast, organic compounds of large molecular size were found to complex Al in infusions from some types of tea (Flaten & Lund 1997).

Conclusions

Bioavailability of Al increases in acidic environments. Different chemical species of Al have different toxicity to living organisms, making Al speciation a necessity. Complexes between some organic compounds and Al have high bioavailability and can cross the plasma membrane. The main source of Al in human diet is food, whereas water provides only a minor fraction of Al absorbed. Aluminium accumulates in bones, but also in brain where it causes neurotoxicity. The relationship between common neurological disorders and dietary or environmental exposure to Al is strenuous and requires clarification. Options for minimising medicinal exposure to Al are urgently needed. Further work is required on clarifying various aspects of the transport of Al in the soil-water-plant-human continuum.

Acknowledgements

The research in the author's lab has been supported by the Australian Research Council, Grains Research and Development Corporation, and Australian Academy of Science. The support from the Alexander von Humboldt Fellowship during the write-up stage is appreciated.

References

- Adamo P, Giordano S, Vingiani S, Cobianchi RC, Violante P. 2003 Trace element accumulation by moss and lichen exposed in bags in the city of Naples (Italy). *Env Poll* **122**, 91–103.
- Ahn SJ, Rengel Z, Matsumoto H. 2004 Aluminum-induced plasma membrane surface potential and H^+ -ATPase activity in near-isogenic wheat lines differing in tolerance to aluminum. *New Phytol* **162**, 71–79.
- Ahn SJ, Sivaguru M, Chung GC, Rengel Z, Matsumoto H. 2002 Aluminium-induced growth inhibition is associated with impaired efflux and influx of H^+ across the plasma membrane in root apices of squash (*Cucurbita pepo*). *J Exp Bot* **53**, 1959–1966.
- Ahn SJ, Sivaguru M, Osawa H, Chung GC, Matsumoto H. 2001 Aluminum inhibits the H^+ -ATPase activity by permanently altering the plasma membrane surface potentials in squash roots. *Plant Physiol* **126**, 1381–1390.
- Alberti G, Biesuz R, Profumo A, Pesavento M. 2003 Determination of the total concentration and speciation of Al(III) in tea infusions. *J Inorg Chem* **97**, 79–88.
- Alewel C. 2001 Predicting reversibility of acidification: The European sulfur story. *Water Air Soil Poll* **131**, 1271–1276.
- Alewel C. 2003 Acid input into the soil from acid rain. In: Rengel Z. ed. *Handbook of Soil Acidity*. New York: Marcel Dekker, Inc: 83–115.
- Allain P, Gauchard F, Krari N. 1996 Enhancement of aluminum digestive absorption by fluoride in rats. *Res Comm Mol Pathol Pharmacol* **91**, 225–231.
- Amonette JE, Russell CK, Carosino KA, Robinson NL, Ho JT. 2003 Toxicity of Al to *Desulfovibrio desulfuricans*. *Appl Env Microbiol* **69**, 4057–4066.
- Appelberg M, Degerman E, Norrgren L. 1992 Effects of acidification and liming on fish in Sweden: A review. *Finn Fish Res* **13**, 77–91.
- Archambault DJ, Zhang G, Taylor GJ. 1996 A comparison of the kinetics of aluminum (Al) uptake and distribution in roots of wheat (*Triticum aestivum*) using different aluminum sources. A revision of the operational definition of symplastic Al. *Physiol Plant* **98**, 578–586.
- Armstrong RA, Winsper SJ, Blair JA. 1996 Aluminium and Alzheimer's disease: review of possible pathogenic mechanisms. *Dementia* **7**, 1–9.
- Asano R, Suzuki K, Otsuka T, Otsuka M, Sakurai H. 2002 Concentrations of toxic metals and essential minerals in the mane hair of healthy racing horses and their relation to age. *J Vet Med Sci* **64**, 607–610.
- Baffi C, Bettinelli M, Beone GM, Spezia S. 2002 Comparison of different analytical procedures in the determination of trace elements in lichens. *Chemosphere* **48**, 299–306.
- Bantan T, Milacic R, Mitrovic B, Pihlar B. 1999a Combination of various analytical techniques for speciation of low molecular weight aluminium complexes in plant sap. *Fresenius' J Anal Chem* **365**, 545–552.
- Bantan T, Milacic R, Mitrovic B, Pihlar B. 1999b Investigation of low molecular weight Al complexes in human serum by fast protein liquid chromatography (FPLC)-ETAAS and electrospray (ES)-MS-MS techniques. *J Anal Atom Spectr* **14**, 1743–1748.
- Bantan T, Milacic R, Pihlar B. 1998 Possibilities for speciation of Al-citrate and other negatively charged Al complexes by anion-exchange FPLC-ICP-AES. *Talanta* **46**, 227–235.
- Barcelo J, Poschenrieder C. 2002 Fast root growth responses, root exudates, and internal detoxification as clues to the mechanisms of aluminium toxicity and resistance: a review. *Env Exp Bot* **48**, 75–92.
- Battarbee RW. 1994 Diatoms, lake acidification and the surface water acidification programme (SWAP): A review. *Hydrobiologia* **274**, 1–7.
- Bayon MM, Soldado CAB, Blanco GE, Garcia AJI, Sanz MA. 1999 Capabilities of fast protein liquid chromatography coupled to a double focusing inductively coupled plasma mass spectrometer for trace metal speciation in human serum. *J Anal Atom Spectr* **14**, 947–951.
- Berube D, Brule DG. 1999 A field aluminium speciation method to study the aluminium hazard in water. *Fresenius' J Anal Chem* **363**, 566–570.
- Bi S. 1996 A model describing the complexing effect in the leaching of aluminum from cooking utensils. *Env Poll* **92**, 85–89.
- Bi S. 2001 Speciation of aluminum in the stream waters from the Susquehanna River watershed, Chesapeake Bay. *Environ Geol* **40**, 300–304.
- Bi S, Li X, Gan N, Ye L, Liang H, Dai L, Cao M, Chen Y. 1999 Direct determination of labile monomeric aluminum in natural waters by a.c. oscillopolarography in the presence of rubeanic acid. *Anal Lett* **32**, 1435–1446.
- Bi SP, An SQ, Yang M, Chen T. 2001 Dynamics of aluminum speciation in forest-well drainage waters from the Rhode River watershed, Maryland. *Env Intern* **26**, 377–380.
- Bi SP, Yang XD, Zhang FP, Wang XL, Zou GW. 2001 Analytical methodologies for aluminium speciation in environmental and

- biological samples: A review. *Fresenius' J Anal Chem* **370**, 984–996.
- Blancaflor EB, Jones DL, Gilroy S. 1998 Alterations in the cytoskeleton accompany aluminum-induced growth inhibition and morphological changes in primary roots of maize. *Plant Physiol* **118**, 159–172.
- Blaser P, Zysset M, Zimmermann S, Luster J. 1999 Soil acidification in southern Switzerland between 1987 and 1997: A case study based on the critical load concept. *Env Sci Technol* **33**, 2383–2389.
- Bolan NS, Hedley MJ. 2003 Role of carbon, nitrogen and sulfur cycles in soil acidification. In: Rengel Z. ed. *Handbook of Soil Acidity*. New York: Marcel Dekker: 29–56.
- Bolan NS, Hedley MJ, White RE. 1991 Processes of soil acidification during nitrogen cycling with emphasis on legume based pasture. *Plant Soil* **134**, 53–63.
- Borrmann G, Seubert A. 1996 Aluminum speciation by liquid chromatography concerning hydro- and geo-chemical aspects. *Anal Chim Acta* **332**, 233–239.
- Borrmann G, Seubert A. 1999 Aluminum speciation by means of anion chromatography and coupled anion/cation chromatography. *Anal Chim Acta* **386**, 77–88.
- Breakwell NA, Publicover SJ. 1994 Prolonged enhancement of synaptic transmission in area CA1 of rat hippocampal slices induced by NaF/AlCl₃ does not require NMDA receptor activation but is suppressed by inhibitors of phosphoinositide-mediated signalling pathways. *Brain Res* **633**, 72–76.
- Busch M, Seubert A. 1999 Influence of column temperature on the ion chromatographic separation of aluminum species. *Anal Chim Acta* **399**, 223–235.
- Cameron RS, Ritchie GSP, Robson AD. 1986 Relative toxicities of inorganic aluminium complexes to barley. *Soil Sci Soc Am J* **50**, 1231–1236.
- Campbell A. 2002 The potential role of aluminium in Alzheimer's disease. *Nephrol Dial Transpl* **17**, 17–20.
- Candy JM, McArthur FK, Oakley AE, Taylor GA, Chen CPL-H, Mountfort SA, Thompson JE, Chalker PR, Bishop HE et al. 1992 Aluminium accumulation in relation to senile plaque and neurofibrillary tangle formation in the brains of patients with renal failure. *J Neurol Sci* **107**, 210–218.
- Chan YY, Lo SCL. 2003 Analysis of Ling Zhi (*Ganoderma lucidum*) using dynamic reaction cell ICP-MS and ICP-AES. *J Anal Atom Spectr* **18**, 146–150.
- Chaussidon M, Netter P, Kessler M, Membre H, Fener P, Delons S, Albaredo F. 1993 Dialysis-associated arthropathy: Secondary ion mass spectrometry evidence of aluminum silicate in beta-2-microglobulin amyloid synovial tissue and articular cartilage. *Nephron* **65**, 559–563.
- Chiba J, Kusumoto M, Shirai S, Ikawa K, Sakamoto S. 2002 The influence of fluoride ingestion on urinary aluminum excretion in humans. *Tohoku J Exp Med* **196**, 139–149.
- Clark BM, St CLL, Mangelson NF, Rees LB, Grant PG, Bench GS. 2001 Characterization of mycobiont adaptations in the foliose lichen *Xanthoparmelia chlorochroa* (Parmeliaceae). *Am J Bot* **88**, 1742–1749.
- Comin JJ, Barloy J, Bourrie G, Trolard F. 1999 Differential effects of monomeric and polymeric aluminium on the root growth and on the biomass production of root and shoot of corn in solution culture. *Eur J Agron* **11**, 115–122.
- Coventry DR, Slattery WJ. 1991 Acidification of soil associated with lupins grown in a crop rotation in north-eastern Victoria. *Aust J Agric Res* **42**, 391–397.
- Crichton RR, Wilmet S, Legssyer R, Ward RJ. 2002 Molecular and cellular mechanisms of iron homeostasis and toxicity in mammalian cells. *J Inorg Chem* **91**, 9–18.
- Csoti T, Gyori J, Salanki J, Erdelyi L. 2001 pH-dependent actions of aluminum on voltage-activated sodium currents in snail neurons. *Neurotoxicology* **22**, 109–116.
- Das AK, Chakraborty R. 1997 Electrothermal atomic absorption spectrometry in the study of metal ion speciation. *Fresenius' J Anal Chem* **357**, 1–17.
- Dayde S, Brumas V, Champmartin D, Rubini P, Berthon G. 2003 Aluminum speciation studies in biological fluids. Part 9. A quantitative investigation of aluminum(III)-glutamate complex equilibria and their potential implications for aluminum metabolism and toxicity. *J Inorg Chem* **97**, 104–117.
- Desroches S, Biron F, Berthon G. 1999 Aluminum speciation studies in biological fluids. Part 5. A quantitative investigation of Al(III) complex equilibria with desferrioxamine, 2,3-dihydroxybenzoic acid, Tiron, CP20 (L1), and CP94 under physiological conditions, and computer-aided assessment of the aluminum-mobilizing capacities of these ligands in vivo. *J Inorg Chem* **75**, 27–35.
- Desroches S, Dayde S, Berthon G. 2000 Aluminum speciation studies in biological fluids. Part 6. Quantitative investigation of aluminum(III)-tartrate complex equilibria and their potential implications for aluminum metabolism and toxicity. *J Inorg Chem* **81**, 301–312.
- Dixon E, Gardner M. 1998 Reactive aluminum in UK surface waters. *Chem Spec Bioavail* **10**, 11–17.
- Domingos M, Klumpp A, Rinaldi MCS, Modesto IF, Klumpp G, Delitti WBC. 2003 Combined effects of air and soil pollution by fluoride emissions on *Tibouchina pulchra* Cogn., at Cubatao, SE Brazil, and their relations with aluminium. *Plant Soil* **249**, 297–308.
- Dong D, Xie Z, Du Y, Liu C, Wang S. 1999 Influence of soil pH on aluminum availability in the soil and aluminum in tea leaves. *Commun Soil Sci Plant Anal* **30**, 873–883.
- Downard AJ, Powell KJ, Money SD. 1997 Speciation determination of aluminium (Al³⁺) by flow injection amperometry via the 1,2-dihydroxyanthraquinone-3-sulfonic acid complex. *Anal Chim Acta* **349**, 111–120.
- Driscoll CT, Driscoll KM, Mitchell MJ, Raynal DJ. 2003 Effects of acidic deposition on forest and aquatic ecosystems in New York State. *Env Poll* **123**, 327–336.
- Echlin P. 1996 Distribution of aluminium in the cells of developing leaves of the tea plant as measured by low-temperature X-ray microanalysis. *Scanning* **18**, 197–198.
- Erdemoglu SB, Pyrzyniska K, Gucer S. 2000 Speciation of aluminum in tea infusion by ion-exchange resins and flame AAS detection. *Anal Chim Acta* **411**, 81–89.
- Erdemoglu SB, Turkdemir H, Gucer S. 2000 Determination of total and fluoride bound aluminium in tea infusions by ion selective electrode and flame atomic absorption spectrometry. *Anal Lett* **33**, 1513–1529.
- Esparza JL, Gomez M, Domingo JL, Del Castillo D, Hernandez M. 2000 Age-related differences on aluminium mobilization by chelating agents in aluminium-loaded uraemic rats. *J Ind Microbiol Biotechnol* **87**, 33–38.
- Fairman B, Sanz MA. 1996 Flow injection-mini-column technique with ICP-AES detection for the isolation and preconcentration of the fast reactive aluminium fraction in waters. *Fresenius' J Anal Chem* **355**, 757–762.
- Flaten AK, Lund W. 1997 Speciation of aluminium in tea infusions studied by size exclusion chromatography with detection by post-column reaction. *Sci Total Env* **207**, 21–28.

- Flaten TP. 2001 Aluminium as a risk factor in Alzheimer's disease, with emphasis on drinking water. *Brain Res Bull* **55**, 187–196.
- Florent C, Tekaya L, Escaig F, Labejof L, Mouthon G, Galle P. 2001 Analytical microscopy observations of rat enterocytes after oral administration of soluble salts of lanthanides, actinides and elements of group III-A of the periodic chart. *Cell Mol Biol* **47**, 419–425.
- Fowler D, Cape JN, Coyle M, Flechard C, Kuylensstierna J, Hicks K, Derwent D, Johnson C, Stevenson D. 1999 The global exposure of forests to air pollutants. *Water Air Soil Poll* **116**, 5–32.
- Foy CD. 1988 Plant adaptation to acid, aluminum-toxic soils. *Commun Soil Sci Plant Anal* **19**, 959–987.
- Gauthier E, Fortier I, Courchesne F, Pepin P, Mortimer J, Gauvreau D. 2000 Aluminum forms in drinking water and risk of Alzheimer's disease. *Env Res* **84**, 234–246.
- Gbondo TSS, Driscoll CT. 2002 Retrospective analysis of the response of soil and stream chemistry of a northern forest ecosystem to atmospheric emission controls from the 1970 and 1990 Amendments of the Clean Air Act. *Env Sci Technol* **36**, 4714–4720.
- Glynn AW, Sparen A, Danielsson LG, Sundstrom B, Jorhem L. 1999 Concentration-dependent absorption of aluminum in rats exposed to labile aluminum in drinking water. *J Toxicol Env Health* **56**, 501–512.
- Glynn AW, Sparen A, Danielsson LG, Sundstrom B, Jorhem L. 2001 The influence of complexing agents on the solubility and absorption of aluminium in rats exposed to aluminium in water. *Food Addit & Contam* **18**, 515–523.
- Glynn AW, Thuvander A, Sundstrom B, Sparen A, Danielsson LG, Jorhem L. 1999 Does aluminium stimulate the immune system in male rats after oral exposure? *Food Addit & Contam* **16**, 129–135.
- Gramiccioni L, Ingrao G, Milana MR, Santaroni P, Tomassi G. 1996 Aluminium levels in Italian diets and in selected foods from aluminium utensils. *Food Addit & Contam* **13**, 767–774.
- Gundersen P, Steinnes E. 2003 Influence of pH and TOC concentration on Cu, Zn, Cd, and Al speciation in rivers. *Water Res* **37**, 307–318.
- Harris WR, Berthon G, Day JP, Exley C, Flaten TP, Forbes WF, Kiss T, Orvig C, Zatta PF. 1996 Speciation of aluminum in biological systems. *J Toxicol Env Health* **48**, 543–568.
- Haug A, Shi B, Vitorello V. 1994 Aluminum interaction with phosphoinositide-associated signal transduction. *Archives Toxicol* **68**, 1–7.
- Herman F, Smidt S, Huber S, Englisch M, Knoflacher M. 2001 Evaluation of pollution-related stress factors for forest ecosystems in central Europe. *Env Sci Poll Res Intern* **8**, 231–242.
- Herrmann J. 2001 Aluminum is harmful to benthic invertebrates in acidified waters, but at what threshold(s)? *Water Air Soil Poll* **130**, 837–842.
- Hils A, Grote M, Janssen E, Eichhorn J. 1999 Speciation of trace amounts of aluminium in percolating water of forest soil by online coupling HPLC-ICP-MS. *Fresenius' J Anal Chem* **364**, 457–461.
- Horst WJ. 1995 The role of the apoplast in aluminium toxicity and resistance of higher plants: a review. *Z Pflanzenern Boden* **158**, 419–428.
- Hui CA, Ellers O. 1999 Effect of high aluminum consumption on mechanics and composition of furculae of free-ranging coots. *Env Toxicol Chem* **18**, 970–975.
- Iancu TC, Perl DP, Sternlieb I, Lerner A, Leshinsky E, Kolodny EH, Hsu A, Good PF. 1996 The application of laser microprobe mass analysis to the study of biological material. *BioMetals* **9**, 57–65.
- Irigaray JL, Barbotteau Y, Chassot E, Oudadesse H. 2000 Trace elements cartography by the PIXE method at the interfaces of an implant coated with bioglass. *J Trace Microprobe Tech* **18**, 499–503.
- Izaguirre MML, Flores S. 1995 Symbiotic nitrogen fixation in *Dioclea guianensis* Benth., a shade-tolerant and aluminium-accumulator legume species native of tropical cloud forests. *Symbiosis* **19**, 65–83.
- Izaguirre MML, Flores S, Oropeza T. 2002 Aluminum tolerance in nodulated N₂-fixing legumes species native to two contrasting savanna sites. *Plant Soil* **245**, 163–168.
- Jallah JK, Smyth TJ. 1998 Assessment of rhizotoxic aluminum in soil solutions by computer and chromogenic speciation. *Commun Soil Sci Plant Anal* **29**, 37–50.
- Jones DL, Gilroy S, Larsen PB, Howell SH, Kochian LV. 1998a Effect of aluminum on cytoplasmic Ca²⁺ homeostasis in root hairs of *Arabidopsis thaliana* (L.). *Planta* **206**, 378–387.
- Jones DL, Kochian LV. 1995 Aluminum inhibition of the inositol 1,4,5-triphosphate signal transduction pathway in wheat roots: A role in aluminum toxicity? *Plant Cell* **7**, 1913–1922.
- Jones DL, Kochian LV, Gilroy S. 1998b Aluminum induces a decrease in cytosolic calcium concentration in BY-2 tobacco cell cultures. *Plant Physiol* **116**, 81–89.
- Jones OR, Perks RM, Abraham CJ, Telle HH, Oakley AE. 1997 A comparison of the techniques of secondary ion mass spectrometry and resonance ionization mass spectrometry for the analysis of potentially toxic element accumulation in neural tissue. *Rapid Communications in Mass Spectrometry* **11**, 179–183.
- Jönsson U, Rosengren U, Thelin G, Nihlgård B. 2003 Acidification-induced chemical changes in coniferous forest soils in southern Sweden 1988–1999. *Env Poll* **123**, 75–83.
- Jugdaohsingh R, Reffitt DM, Oldham C, Day JP, Fifield LK, Thompson RP, Powell JJ. 2000 Oligomeric but not monomeric silica prevents aluminum absorption in humans. *Am J Clin Nutr* **71**, 944–949.
- Juranovic I, Breinhoelder P, Steffan I. 2003 Determination of trace elements in pumpkin seed oils and pumpkin seeds by ICP-AES. *J Anal Atom Spectr* **18**, 54–58.
- Kadar E, Salanki J, Powell J, White KN, McCrohan CR. 2002 Effect of sub-lethal concentrations of aluminium on the filtration activity of the freshwater mussel *Anodonta cygnea* L. at neutral pH. *Acta Bot Hung* **53**, 485–493.
- Kerven GL, Larsen PL, Bell LC, Edwards DG. 1995 Quantitative ²⁷Al NMR spectroscopic studies of Al(III) complexes with organic acid ligands and their comparison with GEOCHEM predicted values. *Plant Soil* **171**, 35–39.
- Kerven GL, Ostatek-Boczynski Z, Edwards DG, Asher CJ, Owczkin J. 1995 Chromatographic techniques for the separation of Al and associated organic ligands present in soil solution. In: Date RA, Grundon NJ, Rayment GE, Probert ME. eds. *Plant-Soil Interactions at Low pH*. Dordrecht: Kluwer Academic Publishers: 47–52.
- Kinraide TB. 1990 Assessing the rhizotoxicity of the aluminate ion, Al(OH)₄⁻. *Physiol Plant* **94**, 1620–1625.
- Kinraide TB. 1993 Aluminum enhancement of plant growth in acid rooting media. A case of reciprocal alleviation of toxicity by two toxic cations. *Physiol Plant* **88**, 619–625.
- Kinraide TB. 1997 Reconsidering the rhizotoxicity of hydroxyl, sulphate, and fluoride complexes of aluminium. *J Exp Bot* **48**, 1115–1124.
- Kishikawa Y, Gong H, Kitaoka T, Amemiya T, Takaya K, Tozu M, Hoshi T, Ohashi Y. 2003 Elements and organic substances in epiretinal proliferative tissue excised during vitreous surgery:

- Analysis by time-of-flight secondary-ion mass spectrometry. *J Electron Microsc* **52**, 349–354.
- Kochian LV. 1995 Cellular mechanisms of aluminum toxicity and resistance in plants. *Annu Rev Plant Physiol Plant Mol Biol* **46**, 237–260.
- Kollmeier M, Felle HH, Horst WJ. 2000 Genotypical differences in aluminum resistance of maize are expressed in the distal part of the transition zone. Is reduced basipetal auxin flow involved in inhibition of root elongation by aluminum? *Plant Physiol* **122**, 945–956.
- Kruck TPA. 1993 Aluminium – Alzheimer's link? *Nature* **363**, 119.
- Kuhn AJ, Bauch J, Schroeder WH. 1995 Monitoring uptake and contents of Mg, Ca and K in Norway spruce as influenced by pH and Al, using microprobe analysis and stable isotope labelling. *Plant Soil* **168–169**, 135–150.
- Kumar S. 2002 Aluminium-induced changes in the rat brain serotonin system. *Food and Chemical Toxicology* **40**, 1875–1880.
- Lazof DB, Goldsmith JG, Linton RW. 1997 The *in situ* analysis of intracellular aluminium in plants. *Prog Bot* **58**, 112–149.
- Lazof DB, Goldsmith JG, Rufty TW, Linton RW. 1994 Rapid uptake of aluminum into cells of intact soybean root tips. A microanalytical study using secondary ion mass spectrometry. *Plant Physiol* **106**, 1107–1114.
- Lesniewicz A, Zyrmicki W. 2003 Extraction of trace elements from spruce needles. *Commun Soil Sci Plant Anal* **34**, 629–643.
- Levesque L, Mizzen CA, McLachlan DR, Fraser PE. 2000 Ligand specific effects on aluminum incorporation and toxicity in neurons and astrocytes. *Brain Res* **877**, 191–202.
- Lian HZ, Kang Yf, Bi SP, Yasin A, Shao DL, Chen YJ, Dai Lm, Tian LC. 2003 Morin applied in speciation of aluminium in natural waters and biological samples by reversed-phase high-performance liquid chromatography with fluorescence detection. *Anal Bioanal Chem* **376**, 542–548.
- Liang JY, Chung RS, Lin HC. 1999 Speciation of aluminum and fluorine in a blend of green tea infusion and human gastric juice by ion chromatography and ^{19}F -NMR. *Food Sci Agric Chem* **1**, 215–222.
- Lindberg S, Strid H. 1997 Aluminium induces rapid changes in cytosolic pH and free calcium and potassium concentrations in root protoplasts of wheat (*Triticum aestivum*). *Physiol Plant* **99**, 405–414.
- Linton RW, Walker SR, DeVries CR, Ingram P, Shelburne JD. 1980 Ion microanalysis of cells. *Scanning Elec Microsc* **II**, 583–596.
- Lote CJ, Willmott K, Wood JA, Thewles A, Freeman M. 1995 Renal excretion of aluminium in the rat: Effect of citrate infusion. *Human Exp Toxicol* **14**, 945–948.
- Ma G, Rengasamy P, Rathjen AJ. 2003 Phytotoxicity of aluminium to wheat plants in high-pH solutions. *Aust J Exp Agric* **43**, 497–501.
- Ma JF, Hiradate S, Nomoto K, Iwashita T, Matsumoto H. 1997 Internal detoxification mechanism of Al in *Hydrangea*. Identification of Al form in the leaves. *Plant Physiol* **113**, 1033–1039.
- Ma QF, Rengel Z, Kuo J. 2002 Aluminium toxicity in rye (*Secale cereale*): Root growth and dynamics of cytoplasmic Ca^{2+} in intact root tips. *Ann Bot* **89**, 241–244.
- Macdonald TL, Humphreys WG, Martin RB. 1987 Promotion of tubulin assembly by aluminum ion *in vitro*. *Science* **236**, 183–186.
- MacLean DC, Hansen KS, Schneider RE. 1992 Amelioration of aluminium toxicity in wheat by fluoride. *New Phytol* **121**, 81–88.
- Mahler RL, Halvorson AR, Koehler FE. 1985 Long-term acidification of farmland in northern Idaho and eastern Washington. *Commun Soil Sci Plant Anal* **16**, 83–95.
- Makjanic J, McDonald B, Chen CPL-H, Watt F. 1998 Absence of aluminium in neurofibrillary tangles in Alzheimer's disease. *Neurosci Lett* **240**, 123–126.
- Mangabeira P, Mushrifah I, Escaig F, Laffray D, Franca MG, Galle P. 1999 Use of SIMS microscopy and electron probe X-ray microanalysis to study the subcellular localization of aluminium in *Vicia faba* roots cells. *Cell Mol Biol* **45**, 413–422.
- Marchante GJM, Sariago MC, Garcia AJI, Sanz MA. 1999 Multielemental trace analysis of biological materials using double focusing inductively coupled plasma mass spectrometry detection. *Anal Chim Acta* **400**, 307–320.
- Marienfeld S, Lehmann H, Stelzer R. 1995 Ultrastructural investigations and EDX-analyses of Al-treated oat (*Avena sativa*) roots. *Plant Soil* **171**, 167–173.
- Marienfeld S, Schmohl N, Klein M, Schroeder WH, Kuhn AJ, Horst WJ. 2000 Localisation of aluminium in root tips of *Zea mays* and *Vicia faba*. *J Plant Physiol* **156**, 666–671.
- Marienfeld S, Stelzer R. 1993 X-ray microanalyses in roots of Al-treated *Avena sativa* plants. *J Plant Physiol* **141**, 569–573.
- Markich SJ, Warne MSJ, Westbury AM, Roberts CJ. 2002 A compilation of data on the toxicity of chemicals to species in Australasia. Part 3: Metals. *Australas J Ecotoxicol* **8**, 1–137.
- Marschner H. 1995 Mineral Nutrition of Higher Plants, 2nd ed. London: Academic Press.
- Martin RB. 1988 Bioinorganic chemistry of aluminum. In: Sigel H. ed. *Metal Ions in Biological Systems. Vol. 24. Aluminum and its Role in Biology*. New York, USA: Marcel Dekker: 1–57.
- Masion A, Vilge RA, Rose J, Stone WEE, Teppen BJ, Rybacki D, Bottero JY. 2000 Coagulation-flocculation of natural organic matter with Al salts: Speciation and structure of the aggregates. *Env Sci Technol* **34**, 3242–3246.
- Masunaga T, Kubota D, Hotta M, Wakatsuki T. 1997 Nutritional characteristics of mineral elements in tree species of tropical rain forest, West Sumatra, Indonesia. *Soil Sci Plant Nutr* **43**, 405–418.
- Masunaga T, Kubota D, Hotta M, Wakatsuki T. 1998 Nutritional characteristics of mineral elements on leaves of tree species in tropical rain forest, West Sumatra, Indonesia. *Soil Sci Plant Nutr* **44**, 315–329.
- Matlaba P, Daya S, Nyokong T. 2000 Interaction of the neurotransmitter acetylcholine with aluminium, calcium and sodium. *Pharm Pharmacol Comm* **6**, 201–205.
- Matsumoto H. 2000 Cell biology of aluminum toxicity and tolerance in higher plants. *Int Rev Cytol* **200**, 1–46.
- May HM, Nordstrom DK. 1991 Assessing the solubilities and reaction kinetics of aluminous minerals in soils. In: Ulrich B, Sumner ME. eds. *Soil Acidity*. Berlin: Springer Verlag: 125–148.
- McDonald-Stephens JL, Taylor GJ. 1995 Kinetics of aluminum uptake by cell suspensions of *Phaseolus vulgaris* L. *J Plant Physiol* **145**, 327–334.
- Memon AR, Chino M, Yatazawa M. 1981 Microdistribution of aluminum and manganese in the tea leaf tissue as revealed by X-ray microanalysis. *Commun Soil Sci Plant Anal* **12**, 441–452.
- Mitrovic B, Milacic R. 2000 Speciation of aluminium in forest soil extracts by size exclusion chromatography with UV and ICP-AES detection and cation exchange fast protein liquid chromatography with ETAAS detection. *Sci Total Env* **258**, 183–194.
- Mitrovic B, Milacic R, Pihlar B, Simoncic P. 1998 Speciation of trace amounts of aluminium in environmental samples by cation-exchange FPLC – ETAAS. *Analusis* **26**, 381–388.
- Moore PB, Day JP, Taylor GA, Ferrier IN, Fifield LK, Edwardson JA. 2000 Absorption of aluminium-26 in Alzheimer's disease, measured using accelerator mass spectrometry. *Dem Geriatric Cogn Disorders* **11**, 66–69.

- Nagata T, Hayatsu M, Kosuge N. 1992 Identification of aluminium forms in tea leaves by ^{27}Al NMR. *Phytochemistry* **31**, 1215–1218.
- Naumann A, Horst WJ. 2003 Effect of aluminium supply on aluminium uptake, translocation and blueing of *Hydrangea macrophylla* (Thunb.) Ser. cultivars in a peatclay substrate. *J Hort Sci Biotechnol* **78**, 463–469.
- Nayak P, Chatterjee AK. 1999 Biochemical view of aluminum-induced neurohazards. *J Env Biol* **20**, 77–84.
- Nichol BE, Oliveira LA. 1995 Effects of aluminum on the growth and distribution of calcium in roots of an aluminum-sensitive cultivar of barley (*Hordeum vulgare*). *Can J Bot* **73**, 1849–1858.
- Noel L, Leblanc JC, Guerin T. 2003 Determination of several elements in duplicate meals from catering establishments using closed vessel microwave digestion with inductively coupled plasma mass spectrometry detection: Estimation of daily dietary intake. *Food Addit & Contam* **20**, 44–56.
- Nystrom RC, Lindh U, Thelin S, Lindquist O, Friman G, Ilback NG. 2002 Trace element changes in sclerotic heart valves from patients undergoing aortic valve surgery. *Biol Trace Elem Res* **88**, 9–24.
- Osaki M, Watanabe T, Ishizawa T, Nilnond C, Nuyim T, Sittibush C, Tadano T. 1998 Nutritional characteristics in leaves of native plants grown in acid sulfate, peat, sandy podzolic, and saline soils distributed in Peninsular Thailand. *Plant Soil* **201**, 175–182.
- Osaki M, Watanabe T, Tadano T. 1997 Beneficial effect of aluminum on growth of plants adapted to low pH soils. *Soil Sci Plant Nutr* **43**, 551–563.
- Ownby JD. 1993 Mechanisms of reaction with aluminium-treated roots. *Physiol Plant* **87**, 371–380.
- Owour PO, Cheruiyot DKA. 1989 Effects of nitrogen fertilizers on the aluminium contents of mature tea leaf and extractable aluminium in the soil. *Plant Soil* **119**, 342–345.
- Parker DR, Bertsch PM. 1992 Formation of the 'Al₁₃' tridecameric polycation under diverse synthesis conditions. *Env Sci Technol* **26**, 914–921.
- Parker DR, Chaney RL, Norvell WA. 1995 Chemical equilibrium models: applications to plant nutrition research. In: Loeppert RH, Schwab AP, Goldberg S. eds. *Chemical Equilibrium and Reaction Models*. Madison, WI, USA: Soil Science Society of America Inc.: 163–200.
- Parker DR, Kinraide TB, Zelazny LW. 1989 On the phytotoxicity of polynuclear hydroxy-aluminum complexes. *Soil Sci Soc Am J* **53**, 789–796.
- Pennington JAT, Schoen SA. 1995 Estimates of dietary exposure to aluminium. *Food Addit & Contam* **12**, 119–128.
- Peri DP, Good PF. 1990 Microprobe studies of aluminum accumulation in association with human central nervous system disease. *Env Geochem Health* **12**, 97–102.
- Perry CC, Shafran KL. 2001 The systematic study of aluminium speciation in medium concentrated aqueous solutions. *J Inorg Chem* **87**, 115–124.
- Phillips DR, Bolger T. 1998 Sublethal toxic effects of aluminium on the earthworm *Eisenia fetida*. *Pedobiologia* **42**, 125–130.
- Piczak K, Lesniewicz A, Zyrnicki W. 2003 Metal concentrations in deciduous tree leaves from urban areas in Poland. *Env Monitor Assessm* **86**, 273–287.
- Platt B, Fiddler G, Riedel G, Henderson Z. 2001 Aluminium toxicity in the rat brain: Histochemical and immunocytochemical evidence. *Brain Res Bull* **55**, 257–267.
- Polak TB, Milacic R, Mitrovic B, Benedik M. 2001a Speciation of low molecular weight Al complexes in serum of CAPD patients. *J Pharm Biomed Anal* **26**, 189–201.
- Polak TB, Milacic R, Pihlar B, Mitrovic B. 2001b The uptake and speciation of various Al species in the *Brassica rapa pekinensis*. *Phytochemistry* **57**, 189–198.
- Poleo ABS, Hytterod S. 2003 The effect of aluminium in Atlantic salmon (*Salmo salar*) with special emphasis on alkaline water. *J Inorg Chem* **97**, 89–96.
- Polizzi S, Pira E, Ferrara M, Bugiani M, Papaleo A, Albera R, Palmi S. 2002 Neurotoxic effects of aluminium among foundry workers and Alzheimer's disease. *Neurotoxicology* **23**, 761–774.
- Priest ND, Talbot RJ, Newton D, Day JP, King SJ, Fifield LK. 1998 Uptake by man of aluminium in a public water supply. *Human Exp Toxicol* **17**, 296–301.
- Pyrzynska K, Gucer S, Bulska E. 2000 Flow-injection speciation of aluminium. *Water Res* **34**, 359–365.
- Rahil KR, Bolann BJ, Myking A, Ulvik RJ. 2002 Multi-element analysis of trace element levels in human autopsy tissues by using inductively coupled atomic emission spectrometry technique (ICP-AES). *J Trace Elem Med Biol* **16**, 15–25.
- Ranau R, Oehlenschlaeger J, Steinhart H. 1999 Determination of aluminium in the edible part of fish by GFAAS after sample pre-treatment with microwave activated oxygen plasma. *Fresenius' J Anal Chem* **364**, 599–604.
- Reid RJ, Rengel Z, Smith FA. 1996 Membrane fluxes and comparative toxicities of aluminium, scandium and gallium. *J Exp Bot* **47**, 1881–1888.
- Rengel Z. 1992a Role of calcium in aluminium toxicity. *New Phytol* **121**, 499–513.
- Rengel Z. 1992b Disturbance of Ca²⁺ homeostasis as a primary trigger in the Al toxicity syndrome. *Plant Cell Environ* **15**, 931–938.
- Rengel Z. 1996 Tansley Review No. 89: Uptake of aluminium by plant cells. *New Phytol* **134**, 389–406.
- Rengel Z. 1997 Mechanisms of plant resistance to toxicity of aluminium and heavy metals. In: Basra AS, Basra RK. eds. *Mechanisms of Environmental Stress Resistance in Plants*. Amsterdam: Harwood Academic Publishers: 241–276.
- Rengel Z, Zhang WH. 2003 Role of dynamics of intracellular calcium in aluminium toxicity syndrome. *New Phytol* **159**, 295–314.
- Reusche E, Gerke P, Krueger S, Rohwer J, Lindner B, Rob PM. 1999 Long-term organic brain syndrome and stroke-like brain stem symptoms in undiagnosed dialysis-associated encephalopathy. *Deutsche Mediz Wochenschr* **124**, 176–181.
- Rincón M, Gonzales RA. 1992 Aluminum partitioning in intact roots of aluminum-tolerant and aluminum-sensitive wheat (*Triticum aestivum* L.) cultivars. *Plant Physiol* **99**, 1021–1028.
- Rollin HB, Nogueira CMCA. 1997 Identification of aluminum fractions in serum using the techniques of high performance liquid chromatography, ultrafiltration and Zeeman atomic absorption spectrometry. *Eur J Clin Chem Clin Biochem* **35**, 215–222.
- Rondeau V. 2002 A review of epidemiologic studies on aluminum and silica in relation to Alzheimer's disease and associated disorders. *Rev Environ Health* **17**, 107–121.
- Roux DJ, Jooste SHJ, Mackay HM. 1996 Substance-specific water quality criteria for the protection of South African freshwater ecosystems: Methods for derivation and initial results for some inorganic toxic substances. *South Af J Sci* **92**, 198–206.
- Ryan PR, DiTomaso JM, Kochian LV. 1993 Aluminium toxicity in roots: an investigation of spatial sensitivity and the role of the root cap. *J Exp Bot* **44**, 437–446.
- Sanz MA. 1998 The chemical speciation of aluminium and silicon in human serum. *Analysis* **26**, M76–M80.

- Sanz MA. 1998 Trace element analytical speciation in biological systems: Importance, challenges and trends. *Spectrochim Acta Part B Atom Spectrosc* **53**, 197–211.
- Savory J, Exley C, Forbes WF, Huang Y, Joshi JG, Kruck T, McLachlan DRC, Wakayama I. 1996 Can the controversy of the role of aluminum in Alzheimer's disease be resolved? What are the suggested approaches to this controversy and methodological issues to be considered? *J Toxicol Env Health* **48**, 615–635.
- Schaedle M, Thornton FC, Raynal DJ, Tepper HB. 1989 Response of tree seedlings to aluminum. *Tree Physiol* **5**, 337–356.
- Schintu M, Meloni P, Contu A. 2000 Aluminum fractions in drinking water from reservoirs. *Ecotoxicol Env Safety* **46**, 29–33.
- Schofield RMS, Pallon J, Fiskesjo G, Karlsson G, Malmqvist KG. 1998 Aluminum and calcium distribution patterns in aluminum-intoxicated roots of *Allium cepa* do not support the calcium-displacement hypothesis and indicate signal-mediated inhibition of root growth. *Planta* **205**, 175–180.
- Schwarzerova K, Zelenkova S, Nick P, Opatrny Z. 2002 Aluminum-induced rapid changes in the microtubular cytoskeleton of tobacco cell lines. *Plant Cell Physiol* **43**, 207–216.
- Senitz D, Bluethner K. 1990 Presence of aluminum in cerebral vessels in cases of Alzheimer disease. *Zb Allg Pathol Pathol Anat* **136**, 329–335.
- Sharp BL, Sulaiman AB, Taylor KA, Green BN. 1997 Observations on the use of ion spray mass spectroscopy for elemental speciation in aqueous solutions without recourse to chromatography. *J Anal Atom Spectr* **12**, 603–609.
- Shen R, Ma JF, Kyo M, Iwashita T. 2002 Compartmentation of aluminum in leaves of an Al-accumulator *Fagopyrum esculentum* Moench. *Planta* **215**, 394–398.
- Shi B, Chou K, Haug A. 1993 Aluminum impacts elements of the phosphoinositide signalling pathway in neuroblastoma cells. *Mol Cell Biochem* **121**, 109–118.
- Shi B, Haug A. 1992 Aluminum interferes with signal transduction in neuroblastoma cells. *Pharm Toxicol* **71**, 308–313.
- Simpson SL, Powell KJ, Nilsson NHS. 1997 Flow injection determination of Al^{3+} and $Al_{13}O_4(OH)_{24}(H_2O)_{12}^{7+}$ species using a 1,3-s reaction with 8-quinolinol-derivatised fractogel. *Anal Chim Acta* **343**, 19–32.
- Sivaguru M, Baluska F, Volkmann D, Felle HH, Horst WJ. 1999 Impacts of aluminum on the cytoskeleton of the maize root apex. *Plant Physiol* **119**, 1073–1082.
- Sivaguru M, Horst WJ. 1998 The distal part of the transition zone is the most aluminum-sensitive apical root zone of maize. *Plant Physiol* **116**, 155–163.
- Song YH, Luan ZK, Tang HX. 2003 Preparation and characterisation of polyaluminium silicate chloride coagulant. *Env Technol* **24**, 319–327.
- Srinivasan PT, Viraraghavan T, Subramanian KS. 1999 Aluminum in drinking water: An overview. *Water S A (Pretoria)* **25**, 47–55.
- Stauber JL, Florence TM, Davies CM, Adams MS, Buchanan SJ. 1999 Bioavailability of Al in alum-treated drinking water. *Am Water Works Ass J* **91**, 84–93.
- Stevens DP, McLaughlin MJ, Alston AM. 1997 Phytotoxicity of aluminium-fluoride complexes and their uptake from solution culture by *Avena sativa* and *Lycopersicon esculentum*. *Plant Soil* **192**, 81–93.
- Stevens DP, McLaughlin MJ, Alston AM. 1998 Phytotoxicity of the fluoride ion and its uptake from solution culture by *Avena sativa* and *Lycopersicon esculentum*. *Plant Soil* **200**, 119–129.
- Strong MJ, Garruto RM, Joshi JG, Mundy WR, Shafer TJ. 1996 Can the mechanisms of aluminum neurotoxicity be integrated into a unified scheme? *J Toxicol Env Health* **48**, 599–613.
- Swegert CV, Dave KR, Katyare SS. 1999 Effect of aluminium-induced Alzheimer like condition on oxidative energy metabolism in rat liver, brain and heart mitochondria. *Mechanism Age Dev* **112**, 27–42.
- Takeda K, Kubota R, Yagioka C. 1985 Copigments in the blueing of sepal colour of the *Hydrangea macrophylla*. *Phytochemistry* **24**, 1207–1209.
- Takita E, Koyama H, Hara T. 1999 Organic acid metabolism in aluminum-phosphate utilizing cells of carrot (*Daucus carota* L.). *Plant Cell Physiol* **40**, 489–495.
- Tang C, Rengel Z. 2003 Role of plant cation/anion uptake ratio in soil acidification. In: Rengel Z, ed. *Handbook of Soil Acidity*. New York: Marcel Dekker: 57–81.
- Taylor GJ. 1991 Current views of the aluminum stress response; the physiological basis of tolerance. *Curr Top Plant Biochem Physiol* **10**, 57–93.
- Taylor GJ, Hunter DB, Stephens J, Bertsch PM, Elmore D, Rengel Z, Reid RJ. 1996 Direct measurement of Al transport across the plasma membrane of *Chara corallina*. *Proc 4th Intern Sym Plant-Soil Inter Low pH*, O15.
- Taylor GJ, McDonald-Stephens JL, Hunter DB, Bertsch PM, Elmore D, Rengel Z, Reid RJ. 2000 Direct measurement of aluminum uptake and distribution in single cells of *Chara corallina*. *Plant Physiol* **123**, 987–996.
- Tsunoda KI, Umemura T, Ohshima K, Aizawa SI, Yoshimura E, Satake KI. 2001 Determination and speciation of aluminum in environmental samples by cation exchange high-performance liquid chromatography with high resolution ICP-MS detection. *Water Air Soil Poll* **131**, 1589–1594.
- Van Landeghem GF, De Broe ME, Haese PCD. 1998 Al and Si: Their speciation, distribution, and toxicity. *Clinical Biochem* **31**, 385–397.
- Van Landeghem GF, Haese PCD, Lamberts LV, Barata JD, De Broe ME. 1997 Aluminium speciation in cerebrospinal fluid of acutely aluminium-intoxicated dialysis patients before and after desferrioxamine treatment; a step in the understanding of the element's neurotoxicity. *Nephrol Dial Transpl* **12**, 1692–1698.
- Van Landeghem GF, Haese PCD, Lamberts LV, Djukanovic L, Pejanovic S, Goodman WG, De Broe ME. 1998 Low serum aluminum values in dialysis patients with increased bone aluminum levels. *Clin Nephrol* **50**, 69–76.
- Vanholder R, Cornelis R, Dhondt A, Lameire N. 2002 The role of trace elements in uraemic toxicity. *Nephrol Dial Transpl* **17**, 2–8.
- Vazquez MD, Poschenrieder C, Corrales I, Barcelo J. 1999 Change in apoplastic aluminum during the initial growth response to aluminum by roots of a tolerant maize variety. *Plant Physiol* **119**, 435–444.
- Venturini SM, Berthon G. 1998 Aluminum speciation studies in biological fluids. Part 4. A new investigation of aluminum-succinate complex formation under physiological conditions, and possible implications for aluminum metabolism and toxicity. *J Inorg Chem* **71**, 135–145.
- Venturini SM, Berthon G. 2001 Aluminum speciation studies in biological fluids. Part 7. A quantitative investigation of aluminum(III)-malate complex equilibria and their potential implications for aluminum metabolism and toxicity. *J Inorg Chem* **85**, 143–154.
- Vitarello VA, Haug A. 1997 An aluminum-morin fluorescence assay for the visualization and determination of aluminum in cultured cells of *Nicotiana tabacum* L. cv. BY-2. *Plant Sci* **122**, 35–42.
- Vitarello VA, Haug A. 1996 Short-term aluminium uptake by tobacco cells: Growth dependence and evidence for internalization in a discrete peripheral region. *Physiol Plant* **97**, 536–544.

- Wang X, Bi S, Gan N, Wei Z. 2001 Aluminum speciation with adsorptive pyrocatechol violet-Al(III) complex by derivative adsorption chronopotentiometry. *Electroanalysis* **13**, 1279–1286.
- Wang X, Lei J, Bi S, Gan N, Wei Z. 2001 Determination of the speciation of aluminum(III) in natural waters by adsorption stripping voltammetry and complexation with AlIII-solochrome violet RS. *Anal Chim Acta* **449**, 35–44.
- Ward RJ, Zhang Y, Crichton RR. 2001 Aluminium toxicity and iron homeostasis. *J Inorg Chem* **87**, 9–14.
- Watanabe T, Osaki M. 2001 Influence of aluminum and phosphorus on growth and xylem sap composition in *Melastoma malabathricum* L. *Plant Soil* **237**, 63–70.
- Watanabe T, Osaki M. 2002 Mechanisms of adaptation to high aluminum condition in native plant species growing in acid soils: A review. *Commun Soil Sci Plant Anal* **33**, 1247–1260.
- Watanabe T, Osaki M, Tadano T. 1997 Aluminum-induced growth stimulation in relation to calcium, magnesium, and silicate nutrition in *Melastoma malabathricum* L. *Soil Sci Plant Nutr* **43**, 827–837.
- Watanabe T, Osaki M, Yoshihara T, Tadano T. 1998 Distribution and chemical speciation of aluminum in the Al accumulator plant, *Melastoma malabathricum* L. *Plant Soil* **201**, 165–173.
- Watt F. 1996 Nuclear microscope analysis in Alzheimer's and Parkinson's disease: A review. *Cell Mol Biol* **42**, 17–26.
- WHO 1996 Guidelines for drinking-water quality. Health Criteria and Other Supporting Information, vol. 2, 2nd ed. Geneva: World Health Organization.
- Witters HE. 1998 Chemical speciation dynamics and toxicity assessment in aquatic systems. *Ecotoxicol Env Safety* **41**, 90–95.
- Wong MH, Fung KF, Carr HP. 2003 Aluminium and fluoride contents of tea, with emphasis on brick tea and their health implications. *Toxicol Lett* **137**, 111–120.
- Wong MH, Zhang ZQ, Wong JWC, Lan CY. 1998 Trace metal contents (Al, Cu and Zn) of tea: Tea and soil from two tea plantations, and tea products from different provinces of China. *Env Geochem Health* **20**, 87–94.
- Wood PC, Wojcikiewicz RJH, Burgess J, Castleden CM, Nahorski SR. 1994 Aluminium inhibits muscarinic agonist-induced inositol 1,4,5-trisphosphate production and calcium mobilization in permeabilized SH-SY5Y human neuroblastoma cells. *J Neurochem* **62**, 2219–2223.
- Wu CW, Wang MK. 2000 Effect of acetate, oxalate, and citrate anions on the formation of Al_3 and aluminum hydroxide in partially neutralized aqueous aluminum salt system. *Food Sci Agric Chem* **2**, 15–19.
- Xia H, Rayson GD. 1998 Investigation of aluminum binding to a *Datura innoxia* material using ^{27}Al NMR. *Env Sci Technol* **32**, 2688–2692.
- Yamamoto Y, Kobayashi Y, Devi SR, Rikiishi S, Matsumoto H. 2002 Aluminum toxicity is associated with mito-chondrial dysfunction and the production of reactive oxygen species in plant cells. *Plant Physiol* **128**, 63–72.
- Yamamoto Y, Kobayashi Y, Devi SR, Rikiishi S, Matsumoto H, Abe J. 2003 Oxidative stress triggered by aluminum in plant roots. *Plant Soil* **255**, 239–243.
- Yokel RA. 2000 The toxicology of aluminum in the brain: a review. *Neurotoxicology* **21**, 813–828.
- Yokel RA, McNamara PJ. 2001 Aluminium toxicokinetics: An updated mini review. *Pharm Toxicol* **88**, 159–167.
- Yokel RA, Wilson M, Harris WR, Halestrap AP. 2002 Aluminum citrate uptake by immortalized brain endothelial cells: Implications for its blood-brain barrier transport. *Brain Res* **930**, 101–110.
- Yoshimura E, Nakagawa T, Mori S. 2003 Distribution and chemical forms of aluminum in roots of aluminum-tolerant rice cultivar grown with or without phosphate pretreatment. *Commun Soil Sci Plant Anal* **34**, 1549–1555.
- Ysart G, Miller P, Croasdale M, Crews H, Robb P, Baxter M, De L'Argy C, Harrison N. 2000 1997 UK Total Diet Study: Dietary exposures to aluminium, arsenic, cadmium, chromium, copper, lead, mercury, nickel, selenium, tin and zinc. *Food Addit & Contam* **17**, 775–786.
- Zatta P, Ibn Lkhayat IM, Zambenedetti P, Kilyen M, Kiss T. 2002 In vivo and in vitro effects of aluminum on the activity of mouse brain acetylcholinesterase. *Brain Res Bull* **59**, 41–45.
- Zatta P, Zambenedetti P. 1996 Aluminum speciation and morphological differentiation in murine neuroblastoma cells. *Biol Trace Elem Res* **51**, 77–85.
- Zhang G, Taylor GJ. 1989 Kinetics of aluminum uptake by excised roots of aluminum-tolerant and aluminum-sensitive cultivars of *Triticum aestivum* L. *Plant Physiol* **91**, 1094–1099.
- Zhang G, Taylor GJ. 1990 Kinetics of aluminum uptake in *Triticum aestivum* L. Identity of the linear phase of aluminum uptake by excised roots of aluminum-tolerant and aluminum-sensitive cultivars. *Plant Physiol* **94**, 577–584.
- Zhang W-H, Rengel Z. 1999 Aluminium induces an increase in cytoplasmic calcium in intact wheat root apical cells. *Aust J Plant Physiol* **26**, 401–409.
- Zhang W-H, Rengel Z, Kuo J. 1998 Determination of intracellular Ca^{2+} in cells of intact wheat roots: Loading of acetoxymethyl ester of Fluo-3 under low temperature. *Plant J* **15**, 147–151.
- Zhu R, Kok WT. 1998 Determination of trace metal ions by capillary electrophoresis with fluorescence detection based on post-column complexation with 8-hydroxyquinoline-5-sulphonic acid. *Anal Chim Acta* **371**, 269–277.