

# Iron- and 2-oxoglutarate-dependent Dioxygenases: an emerging group of molecular targets for nickel toxicity and carcinogenicity

Haobin Chen · Max Costa

Received: 17 October 2008 / Accepted: 7 December 2008 / Published online: 19 December 2008  
© Springer Science+Business Media, LLC. 2008

**Abstract** Nickel compounds are important occupational and environmental pollutants. Chronic exposure to these pollutants has been connected with increased risks of respiratory cancers and cardiovascular diseases. However, it is still not clear what are the specific molecular targets for nickel toxicity and carcinogenicity. Here, we propose that the iron- and 2-oxoglutarate-dependent dioxygenase family enzymes are important intracellular targets that mediate the toxicity and carcinogenicity of nickel. In support of this hypothesis, our data show that three different classes of enzymes in this iron- and 2-oxoglutarate-dependent dioxygenase family, including HIF-prolyl hydroxylase PHD2, histone demethylase JHDM2A/JMJD1A, and DNA repair enzyme ABH3, are all highly sensitive to nickel inhibition. Inactivation of these enzymes accounts for a number of deleterious effects caused by nickel in cells, namely hypoxia-mimic stress and aberrant epigenetic changes. Future studies on nickel's effects on these iron- and 2-oxoglutarate-dependent dioxygenases would deepen our understanding on nickel toxicity and carcinogenicity.

**Keywords** Nickel · Dioxygenase · Iron · JHDM2A/JMJD1A · ABH3 · HIF · Epigenetic · Histone methylation

Nickel (Ni) compounds are important occupational and environmental pollutants. Epidemiological studies have provided evidence showing a strong correlation between worksite exposure to Ni compounds and an increased incidence of nasal and lung cancers in nickel refinery workers (Polednak 1981; Roberts et al. 1984; Roberts et al. 1989). Recent studies have also indicated that nickel, a common component present in fine particulate matter of ambient air, plays an important role in development of cardiovascular diseases in susceptible human populations (Lippmann et al. 2006). Using in vitro cell models and animal models, nickel compounds have been found to generate various types of deleterious effects, including chromosomal aberrations, DNA strand breaks, excessive reactive oxygen species production, impaired DNA repair, hypoxia-mimic stress, aberrant epigenetic changes, and signaling cascade activation (reviewed by Lu et al. 2005). However, it is still not clear what are the specific molecular targets for nickel toxicity and carcinogenicity. Here, we propose that the iron- and 2-oxoglutarate-dependent dioxygenase family enzymes are important intracellular targets that mediate the toxicity and carcinogenicity of nickel. Preliminary data in support of this hypothesis are provided in this article.

---

H. Chen · M. Costa (✉)  
Department of Environmental Medicine, New York  
University School of Medicine, 57 Old Forge Road,  
Tuxedo, New York, NY 10987, USA  
e-mail: max.costa@nyumc.org

The first piece of supportive evidence came from our studies on nickel-induced hypoxia-mimic stress. One of the most prominent changes in cells following nickel exposure is an accumulation of hypoxia inducible factor 1 alpha (HIF1 $\alpha$ ), a transcriptional factor important for cell adaption to low oxygen tension environment (Costa et al. 2005). The intracellular protein levels of HIF1 $\alpha$  are mainly regulated through protein stability. Under normoxia conditions, HIF1 $\alpha$  is constantly synthesized but is rapidly degraded through proteasome pathway, while it becomes stabilized and trans-activated under hypoxic conditions. A number of iron-containing dioxygenases, such as prolyl hydroxylase domain proteins 1–3 (PHD1–3), have been found to actively hydroxylate several proline residues located in the oxygen-dependent degradation domain (ODDD) of HIF1 $\alpha$  in the presence of oxygen, iron, 2-oxoglutarate, and ascorbic acid (Stolze et al. 2006). Hydroxylation of these proline residues leads to the association of HIF-1 $\alpha$  with a Von-Hippel-Lindau (VHL) E3 ubiquitin ligase complex, and consequently causes an ubiquitin-dependent degradation of this protein (Stolze et al. 2006). Our previous studies have indicated that nickel replaces the iron in these HIF-prolyl hydroxylases, and causes inhibition of their enzymatic activity and stabilization of HIF1 $\alpha$  (Davidson et al. 2005; Davidson et al. 2006). An inhibiting concentration 50 (IC<sub>50</sub>) of nickel was found to be 22  $\mu$ M for PHD2 at the presence of 100  $\mu$ M Fe<sup>2+</sup>, indicating this enzyme is highly sensitive to nickel inhibition (Davidson et al. 2006). These findings prompted us to ask the question whether other enzymes that belong to the same dioxygenase family as HIF-prolyl hydroxylases could have similar sensitivity to nickel inhibition.

Our studies on epigenetic effects of nickel led to the discovery of a new class of enzymes that belong to the same dioxygenase family as HIF-prolyl hydroxylases, histone H3 lysine 9 (H3K9) demethylases (Chen et al. 2006a). In our previous work, we studied the epigenetic effects of nickel on gene expression in detail by using G12 cells, which is a Chinese hamster V79-derived cell clone possessing a single copy of the bacterial *gpt* (xanthine guanine phosphoribosyltransferase) transgene near the telomere of chromosome 1. We found that exposure of G12 cells to nickel compounds silenced the *gpt* transgene via epigenetic mechanisms (reviewed by

Costa et al. 2005). In the nickel-induced *gpt*-inactivated cell clones, the promoter of this transgene was associated with decreases in histone acetylation and H3K4 methylation as well as increases in dimethylated histone H3K9 (H3K9me2), DNA methylation, and chromatin condensation (Chen et al. 2006a; Lee et al. 1995; Yan et al. 2003). Since H3K9me2 is found to be a critical mark for establishment of DNA methylation and long-term gene silencing (Jackson et al. 2004), this modification is likely to play an important role in nickel-induced DNA methylation and gene silencing (Chen et al. 2006a). During our studies on the mechanism by which nickel increases the global level of H3K9me2 in cells, we found that hypoxia and several other hypoxic-mimic agents, such as iron-chelator (deferrioxamine) and dimethylallylglycine (DMOG; an analog of 2-oxoglutarate), can also effectively increase this modification in cells by sharing a similar pattern as HIF-1 $\alpha$  induction (Chen et al. 2006a). Using an in vitro histone demethylation assay, we demonstrated that histone H3K9me2 demethylase activity exists in cells and is dependent on iron and 2-oxoglutarate, and that nickel inhibits this demethylase activity (Chen et al. 2006a). It is now known that JMJD2A-D and JHDM2A/JMJD1A which all belong to the iron- and 2-oxoglutarate-dependent dioxygenase family, demethylate H3K9me2 by catalyzing the generation of highly reactive oxygen species in the presence of oxygen, iron, 2-oxoglutarate, and ascorbic acid. These resultant species attack the methyl groups on histone lysines and produce unstable oxidized intermediates that spontaneously release formaldehyde, resulting in the removal of methyl groups from histone lysines (Klose and Zhang 2007; Shi and Whetstone 2007).

To date, besides HIF-prolyl hydroxylases and histone demethylases, this family of iron- and 2-oxoglutarate-dependent dioxygenases also includes, but is not limited to, HIF-1 $\alpha$  asparagine hydroxylases (FIH-1) and the human homologues of bacterial DNA repair enzyme Alkb (ABH1-9) (Ozer and Bruick 2007). FIH-1 hydroxylates the Asn(803) residue of HIF-1 $\alpha$  and prevents the binding of histone acetyltransferase p300 to this site, which consequently inhibits the transcriptional activity of HIF-1 $\alpha$  (Lando et al. 2002). Similar as Alkb, human homologues ABH2 and ABH3 catalyze oxidative demethylation reaction to repair alkylated lesions in DNA, such as


1-methyladenine or 3-methylcytosine, in an iron- and 2-oxoglutarate-dependent fashion (Aas et al. 2003; Duncan et al. 2002). After knowing what other members of this iron- and 2-oxoglutarate-dependent dioxygenase family are, we chose to test the sensitivity of JHDM2A and ABH3 to nickel inhibition.

To this end, we expressed Flag-tagged JHDM2A recombinant protein in insect cells and purified it using anti-Flag antibody affinity chromatography (Fig. 1a). Consistent with a previous report (Yamane et al. 2006), JHDM2A specifically demethylates both H3K9me2 and me1, but not H3K9me3 (Fig. 1b). Similarly, 6×His-tagged ABH3 was expressed in bacterial cells and purified by Ni-NTA (nickel-nitrilotriacetic acid) chromatography (Fig. 1c). In vitro assay showed that the purified 6×His-tagged ABH3 actively demethylated 3-methylcytosines located in the synthesized oligonucleotides (Fig. 1d). We next studied the nickel inhibitory kinetics of purified Flag-JHDM2A and 6×His-ABH3 recombinant proteins. By incubating Flag-JHDM2A with different concentrations of nickel ions at the presence of 100 μM Fe(II), we found that nickel inhibited Flag-JHDM2A demethylase activity in a dose-dependent fashion with an  $IC_{50} = 25 \mu M$  (Fig. 2a–b). Similarly, nickel also inhibited the demethylase activity of 6×His-ABH3 in a dose-dependent manner with an  $IC_{50} = 75 \mu M$  (Fig. 2c–d). In contrast, purified aconitase, a Krebs cycle enzyme that binds iron in a form of iron-sulfur cluster, was not inhibited by up to 5 mM  $NiCl_2$  (Fig. 2e). Taken together, these results provided direct evidence showing that the members of this iron- and 2-oxoglutarate-dependent dioxygenase family have similar sensitivity to nickel inhibition.

Why do these iron- and 2-oxoglutarate-dependent dioxygenases have similar sensitivity to nickel inhibition, even though they are involved in many different biological processes in cells? This is likely due to the fact that these enzymes utilize a similar mechanism to bind the cofactor iron. An analysis of the crystal structures of JMJD2A, ABH2-3, PHD2, and FIH-1 reveals that these enzymes all use a 2-histidines-1-carboxylate triad motif to bind iron (Chen et al. 2006b; Dann et al. 2002; McDonough et al. 2006; Sundheim et al. 2006; Yang et al. 2008). It is estimated that the affinity constant of Ni(II) binding to this iron-binding motif is at least three orders of magnitude greater than that of Fe(II)

(Davidson et al. 2006). Based on this estimation, it can predict that nickel ions will effectively compete with iron ions for binding to this motif, and that iron ions will not be able to replace nickel ions once nickel ions bind to this motif first. This research is currently ongoing in our group. Different from our opinion, another group proposes that nickel exposure depletes intracellular ascorbate and consequently leads to inhibition of HIF-prolyl hydroxylases (Salnikow et al. 2004). This theory was based on the findings that nickel exposure decreased ascorbate uptake in an in vitro cell culture system, and that addition of ascorbate in the culture medium blocked nickel-induced HIF transactivation in cells. However, a different interpretation of these findings should be considered. First of all, it is commonly believed that an increase in ascorbate levels in culture medium promotes the availability of Fe(II) to the active site of HIF-prolyl hydroxylases (Knowles et al. 2003), which can effectively compete with Ni(II) for binding. Secondly, different from an in vitro tissue culture system that has only 1–5 μM ascorbate in medium and undetectable amount of this compound ( $<0.5 \mu M$ ) in cells (Quievryn et al. 2002), it is estimated that the intracellular ascorbate level in humans is about 1 mM (Meister 1994). It is questionable how nickel exposure in humans can lower the intracellular ascorbate concentration to such a level that causes inhibition of these Fe- and 2-oxoglutarate-dependent dioxygenases. It is also difficult to reconcile this theory with our in vitro assay results that nickel inhibits JHDM2A and ABH3 demethylase activity at the presence of 2 mM ascorbic acid in each reaction mixture. In contrast with the estimated 1 mM intracellular ascorbate level in humans, the bio-available Fe level (also called the labile Fe pool) in cells is estimated to be just about 0.8 μM (Konijn et al. 1999) and is tightly controlled by iron responsive proteins (IRPs) (Muckenthaler et al. 2008). Because there is no known defensive mechanism against nickel ions in cells, nickel ions that enter cells would have unrestricted access to the iron-binding sites of these Fe- and 2-oxoglutarate-dependent dioxygenases.

In summary, the iron- and 2-oxoglutarate-dependent dioxygenase family enzymes are emerging molecular targets for nickel toxicity and carcinogenicity. Because the members of this dioxygenase family are involved in many different biological




**Fig. 1** Purification of Flag-tagged JHDM2A and 6xHis-tagged ABH3 recombinant proteins and characterization of their demethylase activity. **a** Purification of Flag-tagged JHDM2A recombinant protein from insect cells using Anti-Flag antibody affinity chromatography. The eluate was separated on a 4–20% SDS-polyacrylamide gradient gel. The Coomassie blue staining of the gel is shown here. **b** Detection of Flag-JHDM2A demethylase activity in vitro. Different amounts of purified Flag-JHDM2A recombinant protein were used in an in vitro histone demethylase assay, which contained 100 µM Fe(II), 1 mM 2-oxoglutarate, and 2 mM ascorbic acid in each reaction (Chen et al. 2006a). After the reactions, histones were separated by electrophoresis and transferred to PVDF membranes. Immunoblotting was then performed using specific antibodies to histone H3 and different methyl levels of histone H3K9. **c** Purification of 6xHis-tagged ABH3 recombinant protein from bacterial cells using Ni-NTA affinity chromatography. The eluate was separated in a 4–20% SDS-polyacrylamide gradient gel. The Coomassie blue staining of the gel is shown here. **d** Detection of 6xHis-tagged ABH3

demethylase activity in vitro. The assay was performed essentially as previously described by Lee et al. (Lee et al. 2005). In brief, purified 6xHis-tagged ABH3 was incubated with a <sup>32</sup>P end-labeled single-stranded oligonucleotide (24 nucleotides in length) containing one 3-methylcytosine in its HpaII cutting sequence. After incubation in a reaction buffer containing 100 µM Fe(II), 1 mM 2-oxoglutarate, and 2 mM ascorbic acid, the single-stranded oligonucleotide was purified and annealed with its complementary sequence, which was then subject to digestion with methylation-sensitive restriction enzyme HpaII. The reaction products were separated on 20% denaturing polyacrylamide gels. The radioactivity on the gels was detected by autoradiography. The appearance of 7-oligonucleotide fragments indicates the removal of the methyl groups at the 3-N position of cytosines and subsequent cutting of 24-oligonucleotides by HpaII. Single-stranded oligonucleotides of the same sequence but without a 3-methylcytosine were used as a positive control, while the assay without addition of 6xHis-tagged ABH3 was used as a negative control and performed in parallel

processes in cells, their inhibition by nickel could have a broad and complex impact on cells. Future research should aim to elucidate the mechanism by

which nickel inhibits these enzymes as well as to understand the biological consequences of these enzymes inhibition by nickel.



**Fig. 2** A kinetic study on Ni inhibition of JHDM2A and ABH3 demethylase activity. **a** Purified Flag-JHDM2A was assayed for its demethylase activity in the presence of different concentrations of Ni(II) ions as indicated. The assay with addition of EDTA, a chelator of divalent metals, was performed in parallel as a negative control. **b** Data quantification of Fig. 2a. **c** Purified 6×His-tagged ABH3 was assayed for its demethylase activity in the presence of different concentrations of Ni(II) ions as indicated. **d** Data quantification

of Fig. 2c. **e** Purified aconitase, an Fe–S cluster-containing enzyme, was incubated with different concentrations of Ni(II) ions for 4 h. After incubation, the aconitase activity was measured immediately as previously described (Chen et al. 2005). Aconitase activity is presented as that relative to levels in the control samples. Each bar represents the mean (±SD) from three samples per treatment. \*Statistically significant change ( $P < 0.05$ ) compared to control samples

**Acknowledgments** We thank Dr. Yi Zhang for providing the Flag-tagged JHDM2A insect cell expression vector and Dr. Timothy R. O'Connor for the 6×His-tagged ABH3 bacterial expression vector. This work was supported by grant numbers ES005512 and ES014454 from the National Institute of Environmental Health Sciences.

## References

Aas PA, Otterlei M, Falnes PO et al (2003) Human and bacterial oxidative demethylases repair alkylation damage in both RNA and DNA. *Nature* 421:859–863. doi:10.1038/nature01363

- Chen H, Davidson T, Singleton S et al (2005) Nickel decreases cellular iron level and converts cytosolic aconitase to iron-regulatory protein 1 in A549 cells. *Toxicol Appl Pharmacol* 206:275–287. doi:[10.1016/j.taap.2004.11.011](https://doi.org/10.1016/j.taap.2004.11.011)
- Chen H, Ke Q, Kluz T et al (2006a) Nickel ions increase histone H3 lysine 9 dimethylation and induce transgene silencing. *Mol Cell Biol* 26:3728–3737. doi:[10.1128/MCB.26.10.3728-3737.2006](https://doi.org/10.1128/MCB.26.10.3728-3737.2006)
- Chen Z, Zang J, Whetstone J et al (2006b) Structural insights into histone demethylation by JMJD2 family members. *Cell* 125:691–702. doi:[10.1016/j.cell.2006.04.024](https://doi.org/10.1016/j.cell.2006.04.024)
- Costa M, Davidson TL, Chen H et al (2005) Nickel carcinogenesis: epigenetics and hypoxia signaling. *Mutat Res* 592:79–88. doi:[10.1016/j.mrfmmm.2005.06.008](https://doi.org/10.1016/j.mrfmmm.2005.06.008)
- Dann CEIII, Bruck RK, Deisenhofer J (2002) Structure of factor-inhibiting hypoxia-inducible factor 1: an asparaginyl hydroxylase involved in the hypoxic response pathway. *Proc Natl Acad Sci USA* 99:15351–15356. doi:[10.1073/pnas.202614999](https://doi.org/10.1073/pnas.202614999)
- Davidson T, Chen H, Garrick MD et al (2005) Soluble nickel interferes with cellular iron homeostasis. *Mol Cell Biochem* 279:157–162. doi:[10.1007/s11010-005-8288-y](https://doi.org/10.1007/s11010-005-8288-y)
- Davidson TL, Chen H, Di Toro DM et al (2006) Soluble nickel inhibits HIF-prolyl-hydroxylases creating persistent hypoxic signaling in A549 cells. *Mol Carcinog* 45:479–489. doi:[10.1002/mc.20176](https://doi.org/10.1002/mc.20176)
- Duncan T, Treweek SC, Koivisto P et al (2002) Reversal of DNA alkylation damage by two human dioxygenases. *Proc Natl Acad Sci USA* 99:16660–16665. doi:[10.1073/pnas.262589799](https://doi.org/10.1073/pnas.262589799)
- Jackson JP, Johnson L, Jasencakova Z et al (2004) Dimethylation of histone H3 lysine 9 is a critical mark for DNA methylation and gene silencing in *Arabidopsis thaliana*. *Chromosoma* 112:308–315. doi:[10.1007/s00412-004-0275-7](https://doi.org/10.1007/s00412-004-0275-7)
- Klose RJ, Zhang Y (2007) Regulation of histone methylation by demethylation and demethylation. *Nat Rev Mol Cell Biol* 8:307–318. doi:[10.1038/nrm2143](https://doi.org/10.1038/nrm2143)
- Knowles HJ, Raval RR, Harris AL et al (2003) Effect of ascorbate on the activity of hypoxia-inducible factor in cancer cells. *Cancer Res* 63:1764–1768
- Konijn AM, Glickstein H, Vaisman B et al (1999) The cellular labile iron pool and intracellular ferritin in K562 cells. *Blood* 94:2128–2134
- Lando D, Peet DJ, Whelan DA et al (2002) Asparagine hydroxylation of the HIF transactivation domain a hypoxic switch. *Science* 295:858–861. doi:[10.1126/science.1068592](https://doi.org/10.1126/science.1068592)
- Lee YW, Klein CB, Kargacin B et al (1995) Carcinogenic nickel silences gene expression by chromatin condensation and DNA methylation: a new model for epigenetic carcinogens. *Mol Cell Biol* 15:2547–2557
- Lee DH, Jin SG, Cai S et al (2005) Repair of methylation damage in DNA and RNA by mammalian AlkB homologues. *J Biol Chem* 280:39448–39459. doi:[10.1074/jbc.M509881200](https://doi.org/10.1074/jbc.M509881200)
- Lippmann M, Ito K, Hwang JS et al (2006) Cardiovascular effects of nickel in ambient air. *Environ Health Perspect* 114:1662–1669
- Lu H, Shi X, Costa M et al (2005) Carcinogenic effect of nickel compounds. *Mol Cell Biochem* 279:45–67. doi:[10.1007/s11010-005-8215-2](https://doi.org/10.1007/s11010-005-8215-2)
- McDonough MA, Li V, Flashman E et al (2006) Cellular oxygen sensing: crystal structure of hypoxia-inducible factor prolyl hydroxylase (PHD2). *Proc Natl Acad Sci USA* 103:9814–9819. doi:[10.1073/pnas.0601283103](https://doi.org/10.1073/pnas.0601283103)
- Meister A (1994) Glutathione-ascorbic acid antioxidant system in animals. *J Biol Chem* 269:9397–9400
- Muckenthaler MU, Galy B, Hentze MW (2008) Systemic iron homeostasis and the iron-responsive element/iron-regulatory protein (IRE/IRP) regulatory network. *Annu Rev Nutr* 28:197–213. doi:[10.1146/annurev.nutr.28.061807.155521](https://doi.org/10.1146/annurev.nutr.28.061807.155521)
- Ozer A, Bruck RK (2007) Non-heme dioxygenases: cellular sensors and regulators jelly rolled into one? *Nat Chem Biol* 3:144–153. doi:[10.1038/nchembio863](https://doi.org/10.1038/nchembio863)
- Polednak AP (1981) Mortality among welders, including a group exposed to nickel oxides. *Arch Environ Health* 36:235–242
- Quiévryn G, Messer J, Zhitkovich A (2002) Carcinogenic chromium(VI) induces cross-linking of vitamin C to DNA in vitro and in human lung A549 cells. *Biochemistry* 41:3156–3167. doi:[10.1021/bi011942z](https://doi.org/10.1021/bi011942z)
- Roberts RS, Julian JA, Muir DC et al. (1984) Cancer mortality associated with the high-temperature oxidation of nickel subsulfide. *IARC Sci Publ* 53:23–35
- Roberts RS, Julian JA, Muir DC et al (1989) A study of mortality in workers engaged in the mining, smelting, and refining of nickel. II: mortality from cancer of the respiratory tract and kidney. *Toxicol Ind Health* 5:975–993
- Salnikow K, Donald SP, Bruck RK et al (2004) Depletion of intracellular ascorbate by the carcinogenic metals nickel and cobalt results in the induction of hypoxic stress. *J Biol Chem* 279:40337–40344. doi:[10.1074/jbc.M403057200](https://doi.org/10.1074/jbc.M403057200)
- Shi Y, Whetstone JR (2007) Dynamic regulation of histone lysine methylation by demethylases. *Mol Cell* 25:1–14. doi:[10.1016/j.molcel.2006.12.010](https://doi.org/10.1016/j.molcel.2006.12.010)
- Stolze IP, Mole DR, Ratcliffe PJ (2006) Regulation of HIF: prolyl hydroxylases. *Novartis Found Symp* 272:15–25. doi:[10.1002/9780470035009.ch3](https://doi.org/10.1002/9780470035009.ch3) (discussion 25–36)
- Sundheim O, Vagbo CB, Bjoras M et al (2006) Human ABH3 structure and key residues for oxidative demethylation to reverse DNA/RNA damage. *EMBO J* 25:3389–3397. doi:[10.1038/sj.emboj.7601219](https://doi.org/10.1038/sj.emboj.7601219)
- Yamane K, Toumazou C, Tsukada Y et al (2006) JHDM2A, a JmJC-containing H3K9 demethylase, facilitates transcription activation by androgen receptor. *Cell* 125:483–495. doi:[10.1016/j.cell.2006.03.027](https://doi.org/10.1016/j.cell.2006.03.027)
- Yan Y, Kluz T, Zhang P et al (2003) Analysis of specific lysine histone H3 and H4 acetylation and methylation status in clones of cells with a gene silenced by nickel exposure. *Toxicol Appl Pharmacol* 190:272–277. doi:[10.1016/S0014-008X\(03\)00169-8](https://doi.org/10.1016/S0014-008X(03)00169-8)
- Yang CG, Yi C, Duguid EM et al (2008) Crystal structures of DNA/RNA repair enzymes AlkB and ABH2 bound to dsDNA. *Nature* 452:961–965. doi:[10.1038/nature06889](https://doi.org/10.1038/nature06889)