

ERRATUM

HANS SELYE, ÁRPÁD SOMOGYI and PÁL VÉGH, Inflammation, topical stress and the concept of pluricausal diseases. *Biochem. Pharmac. suppl.* 17, 107 (1968).

The table on page 117 should read as follows:

TABLE 2. EFFECT OF 5-HT ON PODITIS ELICITED BY VARIOUS TOPICAL IRRITANTS

Group	Treatment	×	Inflammation		Necrosis	
			Control	5-HT	Control	5-HT
1	Alcohol	40%	0.2 ± 0.1	0.8 ± 0.2*	1.0 ± 0.2	1.4 ± 0.2
2	Formaldehyde	1%	0.6 ± 0.2	2.5 ± 0.5†	0	0
3	Allylthiocyanate	200 µg	1.0 ± 0	2.1 ± 0.2*	0	0
4	Mustard oil	1%	0.2 ± 0.1	0.7 ± 0.1*	0	0.1 ± 0.1
5	Dioxane	60%	0.2 ± 0.1	1.6 ± 0.2‡	1.2 ± 0.1	2.4 ± 0.2‡
6	Iodoacetic acid	200 µg	1.8 ± 0.1	3.0 ± 0‡	0	0.2 ± 0.1
7	NaCl	20 mg	0.1 ± 0.1	0.9 ± 0.2‡	0.7 ± 0.3	1.3 ± 0.4
8	Naf	1 mg	0	1.2 ± 0.2‡	0	0.4 ± 0.2*
9	Acetic acid	0.5%	0.5 ± 0.2	0.5 ± 0.2	0	0.6 ± 0.2*
10	Xylene	20%	1.0 ± 0.1	1.1 ± 0.1	0.4 ± 0.2	2.0 ± 0.2‡
11	Ox bile (desiccated)	4%	0.8 ± 0.1	0.7 ± 0.1	0.1 ± 0.1	1.2 ± 0.2‡
12	CdCl ₂	300 µg	2.4 ± 0.2	2.7 ± 0.1	0.3 ± 0.1	1.4 ± 0.3†
13	InCl ₃	800 µg	2.4 ± 0.3	1.5 ± 0.2	0.6 ± 0.2	1.8 ± 0.2†
14	ZnCl ₂	600 µg	1.4 ± 0.2	1.9 ± 0.3	0.6 ± 0.2	2.0 ± 0.3‡
15	Carrageenin (crude)	2 mg	0.9 ± 0.1	0.7 ± 0.1	0	0
16	Carrageenin (Lambda)	500 µg	2.2 ± 0.2	2.2 ± 0.2	0	0
17	Lycopodium	2%	0	0	0	0
18	Allylamine	1.5 mg	1.0 ± 0.3	1.9 ± 0.3	2.5 ± 0.2	2.9 ± 0.1
19	Croton oil (in Mazola oil)	1%	2.7 ± 0.1	2.6 ± 0.1	0.8 ± 0.2	0.7 ± 0.3
20	AgNO ₃	300 µg	1.1 ± 0.2	1.6 ± 0.2	0.3 ± 0.1	0
21	HCl	0.04 N	0.9 ± 0.2	0.9 ± 0.2	0	0
22	CaCl ₂	3.5 mg	1.3 ± 0.1	1.0 ± 0.1	1.2 ± 0.3	0.4 ± 0.2

* P > 0.05; † P > 0.01; ‡ P < 0.001.