

STIMULATION OF SECRETION BY THE T₈₄ COLONIC EPITHELIAL CELL LINE WITH DIETARY FLAVONOLS

TOAN D. NGUYEN,* ANDREW T. CANADA,† GREGORY G. HEINTZ, THOMAS W. GETTYS and JONATHAN A. COHN

Departments of Medicine and † Anesthesiology, Duke University School of Medicine and Durham V.A. Medical Center, Durham, NC, U.S.A.

(Received 26 July 1990; accepted 3 January 1991)

Abstract—Flavonols are dietary compounds widely distributed in plants and characterized by a 2-phenyl-benzo(α)pyrane nucleus possessing hydroxyl and ketone groups at positions 3 and 4, respectively. Kaempferol, quercetin, and myricetin are flavonols that are further mono-, di-, or trihydroxylated on the phenyl ring, respectively. To test whether these ingested flavonols might exert a direct secretory effect on intestinal epithelial cells, monolayers of the T₈₄ colonocyte cell line were mounted in Ussing chambers and examined for ion transport response. Twenty minutes after addition of 100 μ M quercetin to either the serosal or mucosal side, the short-circuit current change was maximal at 16.6 μ A/cm². Kaempferol was less potent than quercetin, while myricetin and glycosylated quercetin (rutin) did not induce secretion. The secretion induced by quercetin did not seem to be mediated by the reactive oxygen species generated by quercetin through auto-oxidation and/or redox cycling (superoxide, hydrogen peroxide, and the hydroxyl radical) because it was neither enhanced by iron, nor inhibited by desferrioxamine B or catalase (alone or in combination with superoxide dismutase). Like vasoactive intestinal peptide, quercetin induced a secretory response that was inhibited by barium chloride and bumetanide, and which exhibited synergism with carbachol. Quercetin also stimulated a modest increase in intracellular cAMP levels and the phosphorylation of endogenous protein substrates for cAMP-dependent protein kinase. Thus, quercetin is a potent stimulus of colonocyte secretion that resembles secretagogues which act via a cAMP-mediated signaling pathway.

Flavonoids constitute a class of compounds which contain the basic structural feature of a 2-phenyl-benzo(α)pyrane nucleus (Fig. 1). Either as free aglycones or more commonly glycosylated at carbons 3, 4, or 7, these compounds are universally distributed among vascular plants where they may serve as natural transport regulators for the plant growth substance auxin [1]. Flavonols are a subgroup of the flavonoids, characterized by a hydroxyl group at position 3 and a ketone at position 4. Flavonols can be further hydroxylated at positions 3', 4' or 5' on the β phenyl ring to yield the 4'-monohydroxy-flavonol kaempferol, the 3',4'-dihydroxy-flavonol quercetin, and the 3',4',5'-trihydroxy-flavonol myricetin (Fig. 1). Both quercetin and myricetin produce reactive oxygen species (superoxide, hydrogen peroxide, and hydroxyl radical) through auto-oxidation and redox cycling [2–4]. Since reactive oxygen species may induce intestinal secretion [5, 6], we examined the possibility that flavonols might act on the intestinal epithelial cell to stimulate ion transport.

In this study, the colonic epithelial cell line T₈₄ was used as a model to study the effects of flavonols on the enterocyte. These cells grow as well-differentiated monolayers which exhibit vectorial chloride secretion when mounted in Ussing chambers and exposed to a variety of neurohormonal stimuli [7]. Chloride secretion, monitored by a change in

the short-circuit current (I_{sc}) necessary to nullify the potential difference across the cell monolayer, is stimulated by agents which increase cAMP, such as vasoactive intestinal peptide (VIP) or prostaglandin E₁ [8, 9], and also by agents which act through Ca²⁺-mediated pathways, such as carbachol, histamine and calcium ionophores [10, 11]. Chloride secretion occurs through Cl[−] channels located on the apical membrane of confluent monolayers [12], and is the result of the Cl[−] electrochemical gradient generated by the concerted action of three transporters: the basolateral Na⁺,K⁺,Cl[−] co-transporter, the Na⁺,K⁺-ATPase pump, and K⁺ channels [9]. The Na⁺ and K⁺ imported into the cell by the co-transporter are recycled to the extracellular compartment, respectively, by the Na⁺,K⁺-ATPase pump and by at least two distinct K⁺ efflux channels (activated separately by VIP and by carbachol). Because the secretory response in T₈₄ cells is well characterized and reflects the direct action of secretagogues on the enterocyte, we chose this model to characterize the effects of dietary flavonols on colonic secretion.

MATERIALS AND METHODS

Chemicals. Quercetin, kaempferol, myricetin, rutin, barium chloride, superoxide dismutase (SOD) (from bovine erythrocytes), catalase (from bovine liver), desferrioxamine B, carbachol, and Fe(III)-EDTA were obtained from Sigma, St. Louis, MO. Bumetanide was provided by Biomol Research Laboratories, Plymouth Landing, PA. VIP was from Peninsula, Belmont, CA. ³²P as inorganic phosphate

* Correspondence: Toan D. Nguyen, M.D., Building 2, 2nd Floor, V.A. Medical Center, 508 Fulton St., Durham, NC 27705.




Fig. 1. Chemical structures of flavonoids (A), flavonols (B), kaempferol (C), quercetin (D), and myricetin (E).

was obtained from ICN, Irvine, CA. Culture medium was obtained from the Tissue Culture Facility of the University of North Carolina, Chapel Hill, NC, or from Gibco, Grand Island, NY.

Growth and maintenance of T_{84} cells. T_{84} cells were provided by Dr. K. Dharmasathaphorn (University of California, San Diego). These cells were cultured at 5% CO_2 and 37° in a 1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F-12 medium supplemented with 5% (v/v) newborn calf serum. Cells were seeded onto collagen-coated Nuclepore filters previously glued onto plastic rings (surface area: 2.9 cm^2 , approximately 10^6 cells/filter). These filters were then set on glass beads to allow medium (supplemented with 5000 units/L

penicillin, and 5000 $\mu\text{g/L}$ streptomycin sulfate) to bathe both sides of the cell monolayer. The confluent monolayers used for secretory studies were maintained for at least 7 days after the filters were seeded.

Secretory studies. The cell monolayers on the filter/ring units were mounted in a modified Ussing chamber as previously described [7], and both sides of the monolayer were bathed with a Ringer's solution containing 115 mM NaCl, 1.2 mM CaCl_2 , 1.2 mM MgCl_2 , 0.4 mM KH_2PO_4 , 2.5 mM K_2HPO_4 , 25 mM NaHCO_3 , and 10 mM glucose. Quercetin, myricetin, and kaempferol were dissolved in ethanol and added to the Ringer's solution at a 1:100 dilution (final concentrations of ethanol 1%). The medium was warmed to 37° with a circulating water jacket and gently mixed and oxygenated with a constant inflow of 95% O_2 /5% CO_2 . During secretory studies, spontaneous tissue potential differences were short-circuited by an automatic voltage clamp (model DVC-1000, World Precision Instruments, New Haven, CT) with Ag-AgCl electrodes, and the current necessary to maintain this short circuit (I_{sc}) recorded at 1-min intervals. Instrument calibration was performed prior to each experiment using a filter/ring unit without cells. All comparative studies used matched pairs of monolayers seeded at the same time and studied concurrently. Spot checks of the T_{84} monolayers after completion of the experiments showed that cells exposed to 100 μM quercetin for $> 1 \text{ hr}$ could still exclude trypan blue.

Cellular protein phosphorylation. T_{84} cell protein phosphorylation responses were studied using methods previously described [13]. Briefly, cells were labeled with $^{32}\text{P}_i$ in a phosphate-free buffer, exposed to 100 μM quercetin for 5 min, scraped from the filters, and homogenized with a glass-Teflon homogenizer. Phosphoproteins contained in the supernatant fraction after centrifugation at 436,000 g for 15 min were precipitated with 10% trichloroacetic acid and resolved by two-dimensional polyacrylamide gel electrophoresis. Phosphoproteins were detected by autoradiography using X-OMAT AR5 film exposed at room temperature.

cAMP Assay. T_{84} cells grown to confluence on filters were washed twice with Ringer's solution and immersed in 15 mL of Ringer's solution supplemented with 0.2 mM 3-isobutyl-1-methylxanthine and 10 mM glucose, warmed to 37° , and equilibrated with 95% O_2 , 5% CO_2 . Quercetin (final concentration 100 μM) or VIP (final concentration 1 nM) was added to the solution, and after different time periods, the filters containing the cells were transferred into 12 mm \times 75 mm plastic tubes and placed in liquid nitrogen. The cAMP was then extracted by boiling the cells for 7 min in 1 mL of 5 mM KH_2PO_4 , 5 mM K_2HPO_4 , 1 mM EDTA, and 0.1 mM 3-isobutyl-1-methylxanthine. The supernatant resulting from centrifugation at 15,000 g for 7.5 min was assayed for cAMP according to the method described by Gettys *et al.* [14].

RESULTS

Stimulation of secretion with flavonols. As shown in Fig. 2, addition of 100 μM quercetin to the mucosal




Fig. 2. Secretory effect of quercetin. T₈₄ monolayers were grown to confluence on semipermeable membranes, mounted on modified Ussing chambers, and stimulated with quercetin as described in Materials and Methods. The resulting Isc's were recorded every minute and the resulting means and SEM (alternated for clarity in panels C and D) shown. Panel A (top left): Incubation with 100 μM quercetin added to the mucosal compartment (1–36 min: $N = 47$; 37–66 min: $N = 20$ –46). Panel B (top right): Incubation with either 10 or 100 μM quercetin added to the mucosal compartment (three matched pairs). Panel C (bottom left): Incubation with either 100 or 300 μM quercetin added to the mucosal compartment (three matched pairs). Panel D (bottom right): Incubation with 100 μM quercetin added to either the serosal or the mucosal side of the T₈₄ cell monolayer. Within each of the seven matched pairs, the Isc increases were normalized using the maximal Isc increase; the resulting means and SEM are shown [mean maximal Isc response: $53 \pm 8.7 \mu\text{A}$ ($18 \mu\text{A}/\text{cm}^2$)].

side of the cell monolayer produced an increase in Isc which peaked after 15–20 min to a maximum value above baseline of $16.6 \mu\text{A}/\text{cm}^2$ (SEM = $0.9 \mu\text{A}/\text{cm}^2$, $N = 48$, $48.1 \mu\text{A}/\text{filter}$). The Isc was unaffected in control filters exposed to ethanol alone at a final concentration of 1%.

A threshold response to quercetin was obtained at 10 μM (Fig. 2B). The maximal Isc increase produced by 300 μM quercetin was the same as that produced by 100 μM , but the response was more rapid in onset and shorter in duration (Fig. 2C). As shown in Fig. 2D, quercetin stimulated a similar secretory response whether added to the serosal or the mucosal side of the monolayer. However, the maximal response elicited from the serosal side was only 68% of the maximal response obtained from the mucosal side ($P = 0.01$ for a smaller serosal response, paired two-tailed t -test with 6 df, mean maximal mucosal Isc response: $11.9 \pm 1.9 \mu\text{A}/\text{cm}^2$).

The effects of flavonols structurally related to quercetin were also investigated. The maximal Isc increase observed with 100 μM kaempferol was $26 \pm 2\%$ of the maximal increase obtained with 100 μM quercetin (three paired experiments). For myricetin, secretion was detected only at 300 μM and not at 100 μM . A minimal response was obtained with a 300 μM concentration of the glycosylated quercetin-rutinoside (rutin) (data not shown).

Effects of modulators of the metabolism of reactive oxygen species on quercetin-stimulated chloride secretion. It is possible that the reactive oxygen species (superoxide, hydrogen peroxide, or hydroxyl radical) produced by quercetin upon auto-oxidation and/or redox cycling may mediate its secretory effect. In this case, the Isc response should be altered by compounds or enzymes which modulate the production or degradation of these species. Quercetin-induced secretion was therefore studied after cells were preincubated with superoxide dismutase (SOD) (to enhance the conversion of superoxide to hydrogen peroxide), catalase (to enhance the conversion of hydrogen peroxide to water and oxygen), iron (to facilitate the Haber–Weiss reaction in which hydrogen peroxide is converted to hydroxyl radical), or desferroxamine B (to chelate iron and prevent the Haber–Weiss reaction). If quercetin-induced secretion is mediated by hydrogen peroxide, then catalase should inhibit the secretory response to quercetin. Similarly, if secretion is mediated by hydroxyl radical, then the Isc response should be inhibited by desferroxamine and enhanced by Fe^{3+} . Finally, if extracellular superoxide is the key mediator, then secretion should be blocked by the combination of SOD and catalase. As shown in Table 1, little effect on quercetin-induced secretion was observed after preincubation with 50 μM

Table 1. Effects of modulators of the metabolism of reactive oxygen species on quercetin-induced secretion in T₈₄ cells

Agent	Maximal Isc (% control)	N	P (\neq 100)
SOD (470 units/mL) + catalase (450 units/mL)	104 \pm 12.6	6	0.78
Catalase (450 units/mL)	117 \pm 7.8	6	0.08
Desferroxamine B (50 μ M)	97 \pm 5.6	6	0.68
Fe ³⁺ (50 μ M)	75 \pm 9.2	6	0.04

T₈₄ cells mounted in modified Ussing chambers were incubated with the different modulators of reactive oxygen species metabolism for 10–15 min, and quercetin was added to a final concentration of 100 μ M. In each matched pair, the maximal short-circuit current (Isc) change induced by quercetin in the presence of the modulator is expressed as the percentage of the maximal Isc change induced by quercetin in the absence of the modulator. Values are means \pm SEM. The mean maximal Isc changes in the control monolayers were, respectively, 21.9 \pm 3.3, 15 \pm 2.5, 15.8 \pm 1.66, and 18.3 \pm 2.7 μ A/cm² for the experiments studying the effects of SOD plus catalase, catalase, desferroxamine B, and Fe³⁺. P values were calculated using two-tailed *t*-tests with 5 df.

desferroxamine B or a combination of 470 units/mL SOD and 450 units/mL catalase. A slight enhancement in secretion, which did not reach statistical significance ($0.1 > P > 0.05$), was noted with 450 units/mL catalase, while 50 μ M Fe(III)-EDTA produced a significant ($P < 0.05$) inhibition of quercetin-induced secretion. However, as discussed previously, these last two effects were opposite of what was expected should either hydrogen peroxide or the hydroxyl radical mediate secretion. In the aggregate, these findings do not support a role for reactive oxygen species in the secretory response of quercetin.

Intracellular mechanism of secretion. T₈₄ cell apical chloride secretion is dependent on the chloride gradient across the mucosal membrane of the cell. This gradient is generated by the basolateral Na⁺,K⁺,Cl⁻ co-transporter with the imported Na⁺ and K⁺ recycled outside the cell by the Na⁺,K⁺-ATPase pump and K⁺ efflux channels [15]. The role of these transport systems in quercetin-induced chloride secretion was studied using bumetanide, which inhibits the Na⁺,K⁺,Cl⁻ co-transporter, and barium chloride, which inhibits a VIP-responsive K⁺ channel [10, 16]. Table 2 demonstrates the inhibitory effects of 0.3 mM bumetanide and 6 mM BaCl₂ on the secretory response elicited by 100 μ M quercetin. Compared with matched controls, the Isc response obtained 20 min after addition of quercetin was only 19 \pm 1% and 31 \pm 5% of the expected response for bumetanide and barium chloride, respectively. Thus, quercetin-induced secretion appears to require active Na⁺,K⁺,Cl⁻ co-transport and K⁺ efflux mechanisms.

One approach to determining which intracellular signalling pathway(s) mediates the quercetin secretory response is to evaluate whether quercetin exhibits synergism when administered with other secretagogues. The interactions between carbachol, VIP, and quercetin were therefore studied. In Fig. 3A, cells were exposed either to carbachol (final concentration 10 μ M) or to quercetin (final concentration 50 μ M). After 15 min, quercetin was added to the cells previously exposed to carbachol and vice versa. In these matched pairs, the effect of

Table 2. Effects of ion transport inhibitors on quercetin-induced secretion in T₈₄ cells

Agent	Isc (% control)	N	P (\neq 100)
Bumetanide (0.3 mM)	19 \pm 1	4	0.0001
Barium chloride (6 mM)	31 \pm 5	4	0.0007

T₈₄ cells were incubated for 15 min with either bumetanide or barium chloride, and quercetin was added to a final concentration of 100 μ M. In each matched pair, the Isc change induced by quercetin after 20 min in the presence of the inhibitor is expressed as the percentage of the Isc change induced by quercetin alone. Bumetanide was dissolved in 0.1 M NaOH and added to the cells at a 1:100 dilution; the corresponding control also contained 1:100 dilution of 0.1 M NaOH. Values are means \pm SEM. The maximal Isc responses in the control monolayers were 12.1 \pm 0.59 and 13.4 \pm 1.2 μ A/cm², respectively, for the experiments studying the effects of bumetanide and barium chloride. P values were calculated using two-tailed *t*-tests with 3 df.

quercetin followed by carbachol (or carbachol followed by quercetin) in one monolayer can be compared to the initial effect of quercetin (or carbachol) alone in the other. Analyzed in this fashion, quercetin alone produced an Isc increase of 9.4 \pm 0.4 μ A/cm² (27 \pm 1.2 μ A, N = 3) after 15 min, while carbachol produced an Isc increase of 2.3 \pm 1 μ A/cm² (7 \pm 3.5 μ A, N = 3) after 3 min. Carbachol added to cells responding maximally to quercetin produced an additional Isc increase of 34.9 \pm 5 μ A/cm² (101 \pm 14.4 μ A) resulting in a combined total Isc increase of 44.4 \pm 5 μ A/cm² (129 \pm 14.4 μ A). Quercetin added to cells previously exposed to carbachol produced an Isc increase of 19.1 \pm 1.7 μ A/cm² (55 \pm 4.8 μ A). However, in the latter case, quercetin was added after completion of the carbachol response and the interaction between quercetin and carbachol may be suboptimal. In additional experiments, the secretory response produced by a simultaneous dose of carbachol and




Fig. 3. Interactions between quercetin, carbachol and VIP. Pairs of confluent T₈₄ monolayers were mounted in modified Ussing chambers and their secretory responses (mean Isc and SEM) compared. Panel A (left panel): In each pair, one monolayer was exposed to 10 μM carbachol [C] (serosal surface) or to 50 μM quercetin [Q] (mucosal surface); after 15 min, 50 μM quercetin was added to the monolayer previously exposed to carbachol and vice versa. Additional monolayers were also exposed to a combined concentration of 10 μM carbachol and 50 μM quercetin [Q&C]. The following symbols are used: (●) quercetin followed by carbachol ($N = 3$), (○) carbachol followed by quercetin ($N = 3$), (△) quercetin and carbachol added simultaneously ($N = 5$). Panel B (right panel): In each pair, one monolayer was exposed to 1 nM VIP (serosal surface) and the other to 50 μM quercetin [Q] (mucosal surface). After 15 min, 50 μM quercetin was added to the monolayer previously exposed to VIP and vice versa. The following symbols are used: (●) quercetin followed by VIP ($N = 3$), (○) VIP followed by quercetin ($N = 3$).

quercetin was also evaluated. A peak Isc with an intermediate value of $23.6 \pm 2.2 \mu\text{A}/\text{cm}^2$ ($69 \pm 6.5 \mu\text{A}$, $N = 5$) was obtained 14 min after addition of the combined secretagogues. In all the different sequences, the Isc increases produced by cells exposed to the combination of quercetin and carbachol were greater than the sum of the individual Isc changes produced by quercetin and carbachol ($9.4 \pm 2.3 = 11.7 \mu\text{A}/\text{cm}^2$). The difference in the degrees of synergism probably reflects the different timing of the maximal effect of quercetin and carbachol. In contrast, as shown in Fig. 3B, when the interaction between quercetin and VIP was analyzed in the same fashion, no such synergism, but a possible inhibitory effect was demonstrated. The synergism between carbachol and quercetin, but not between VIP and quercetin, suggests that quercetin may induce secretion through pathways related to the ones activated by VIP, but not by carbachol. This possibility was explored further in the following phosphorylation studies.

Phosphorylation and intracellular cAMP studies. Previous studies have shown that T₈₄ cells exhibit distinct phosphorylation responses to stimuli acting via cAMP or via Ca^{2+} [13, 17]. Phosphoproteins p83, p29, and p23 are examples of proteins exhibiting increased phosphorylation in cells stimulated by agents which act via Ca^{2+} , such as carbachol, histamine, and ionomycin. By contrast, phosphoproteins p37, p18, and p23 exhibit increased phosphorylation in cells exposed to agents which act via cAMP, such as VIP and forskolin. Each of these five phosphoproteins showed increased labeling in monolayers stimulated with forskolin plus carbachol (Fig. 4, comparing panels A and B). When monolayers were stimulated with 100 μM quercetin, only three of these five phosphorylation responses were observed: quercetin stimulated the phosphorylation of p37, p18, and p23, but not p29 or p83 (Fig. 4, comparing panels C and D). These results

suggest that quercetin activates intracellular signaling mechanisms mediated by cAMP, but not by Ca^{2+} .

In an attempt to study whether quercetin induces the generation of cAMP, this second messenger was measured in cells exposed to 100 μM quercetin for different time periods ranging from 1 to 20 min. Surprisingly, as shown in Table 3, only a modest increase in cAMP levels was detected. These increases were minimal when compared with the mean 85-fold increase in cAMP demonstrated with control monolayers exposed to 1 nM VIP for the same time periods (data not shown).

DISCUSSION

We have demonstrated that quercetin is a potent stimulator of ion transport in T₈₄ cells. When added to either the mucosal or serosal side of the cell monolayer, quercetin produced a concentration-dependent increase in Isc, which peaked 15–20 min after the addition of 100 μM quercetin. The observed Isc response is consistent with the possibility that quercetin can act directly on the enterocyte to stimulate electrogenic Cl^- secretion. Of the related hydroxylated flavonols, kaempferol was less potent than quercetin, while myricetin and the glycosylated quercetin-rutinoside (rutin) produced minimal responses. Considering the effective mucosal–serosal barrier, the observation that quercetin can act from either side of the monolayer suggests that its effect is not initially mediated by cellular surface components selectively present on either side of the cell (e.g. receptors). It is also possible that quercetin, being a small hydrophobic molecule, penetrated the cell and produced its effect intracellularly. The similar time courses of the secretory responses produced by the addition of quercetin to either side of the cell do not support the possibility that the effect of quercetin is localized to one side of the cell




Fig. 4. Protein phosphorylation induced by quercetin. As described in Materials and Methods, monolayers were labeled with ^{32}P and exposed to different stimuli for 5 min. Soluble phosphoproteins were then resolved using two-dimensional gel electrophoresis by isoelectric point from pH 4 to pH 7.5 (from left to right) and by size (from top to bottom), and detected by radioautography. Panels A and B: Compared to a control incubation (panel A), at least five proteins showed increased labeling after stimulation with 10 μM forskolin and 100 μM carbachol (panel B): p18, p23, p29, p37, and p83. Panels C and D: Compared to a control incubation (panel C), exposure to 100 μM quercetin for 5 min (panel D) resulted in the increased labeling of p18, p23 and p37. The labeling of p29 and p83 was not increased. Similar results were obtained in three additional experiments.

Table 3. Effect of 100 μ M quercetin on cAMP production in T₈₄ cells

Incubation time (min)	cAMP production (% control)	N (df)	P (\neq 100)
1	94 \pm 6	6 (5)	0.19
5	152 \pm 31	6 (5)	0.07
10	143 \pm 13	8 (7)	0.006
15	116 \pm 12	8 (7)	0.11
20	126 \pm 13	6 (5)	0.05

Confluent T₈₄ cell monolayers were exposed to 100 μ M quercetin for the indicated time periods, and the levels of cAMP in the corresponding cell homogenates were determined as outlined in Materials and Methods. In each paired experiment, cAMP production in cells exposed to quercetin is expressed as the percentage of the cAMP production in control cells not exposed to quercetin. Values are means \pm SEM. The mean cAMP production values in control cells for the time periods 1, 5, 10, 15, and 20 min were respectively, 10.5 \pm 0.8, 7.4 \pm 0.9, 15.9 \pm 1.3, 12.1 \pm 0.8, and 11.4 \pm 1.2 pmol/filter. P values were calculated using one-tailed *t*-tests with the indicated df.

and that quercetin needs to be transported transcellularly when added to the other side.

We initially postulated that the reactive oxygen species produced by flavonol auto-oxidation and/or redox cycling may be the ultimate mediators of quercetin-induced secretion. However, because the secretory response to quercetin was not enhanced by Fe³⁺ and was not inhibited by desferrioxamine B or catalase (alone or in combination with SOD), we were unable to substantiate a role for either superoxide, hydrogen peroxide, or the hydroxyl radical in the action of quercetin. The possibility that quercetin-induced secretion is independent of reactive oxygen species is further supported by the observation that kaempferol, which does not generate reactive oxygen species, stimulates secretion, while the converse is true of myricetin, a potent generator of reactive oxygen species [4]. When studies were performed to identify the intracellular mechanism responsible for quercetin-induced secretion, quercetin resembled other secretagogues known to stimulate T₈₄ cells via cAMP: (a) quercetin-induced secretion was inhibited by barium chloride and bumetanide, (b) quercetin was synergistic with carbachol, but not with VIP, and (c) exposure of intact T₈₄ cells to quercetin resulted in the phosphorylation of endogenous protein substrates for cAMP-dependent protein kinase. However, quercetin, at concentrations capable of stimulating cellular secretion, did not increase cAMP levels to the extent demonstrated by VIP. It is possible that, similar to the case of the adenosine analogue 5'-(*N*-ethyl)-carboxamido-adenosine [18], there may be a shift in the concentration-response curve when stimulation of cAMP is studied instead of secretion. However, we have not been able to explore this possibility because of the poor solubility and potential toxicity of high concentrations of quercetin [19]. It still remains possible that the modest increase in cAMP produced by quercetin was sufficient to stimulate the phosphorylation of the substrates for cAMP-dependent protein kinase responsible for controlling chloride secretion. That the interaction between quercetin and the cAMP pathway may be

complex is further suggested by a possible inhibition of VIP-induced secretion by quercetin (e.g. less potent activation by quercetin of a pathway shared with VIP). Should this be the case, quercetin may prove to be useful as a probe for further studies of the intracellular mechanisms regulating intestinal secretion.

Additional studies will be required to judge whether flavonols are physiologically important stimuli of intestinal secretion. Flavonols are found in the edible portions of many fruits and vegetables and the average intake is estimated to be 100 mg/day [20, 21], with vegetarians consuming significantly larger amounts. Quercetin is the most abundant dietary flavonol, and it therefore seems plausible that luminal concentrations of quercetin may reach 50–100 μ M (15–30 mg/L) depending on the disposition of this flavonol after ingestion. Should these concentrations of quercetin be obtained in the intestinal lumen, the findings presented in this study indicate that this flavonol could stimulate a significant secretory response. In this regard, it seems noteworthy that dietary supplementation with fruits and vegetables has proved to be useful in the clinical management of chronic constipation [22]. Even though it has been assumed that the benefit of fruits and vegetables results from secretory as well as osmotic mechanisms, and from the contained fiber, scant information exists concerning the identity of the substances that may function as secretory stimuli. The present study raises the possibility that the secretory action of quercetin and other flavonols may account for the beneficial effects of fruits and vegetables for patients with constipation.

Acknowledgements—This research was funded in part by the Department of Veterans Affairs, by NIH Grants DK 40506 (T.D.N.), DK 40701 (J.A.C.) and DK 42486 (T.W.G.), by NIEHS Grant IS 04752 (A.T.C.), and by a grant from the Cystic Fibrosis Foundation (J.A.C.). The authors are indebted to Margaret Wolfe, Jolanta Kole, and Wei Wang for their technical assistance, and to Dr. Helen Berschneider for her help in setting up the Ussing chambers.

REFERENCES

- Jacobs M and Rubery PH, Naturally occurring auxin transport regulators. *Science* **241**: 346–349, 1988.
- Hodnick WF, Kung FS, Roettger WJ, Bohmont CW and Pardini RS, Inhibition of mitochondrial respiration and production of toxic oxygen radicals by flavonoids. A structure–activity study. *Biochem Pharmacol* **35**: 2345–2357, 1986.
- Hodnick WF, Kalyanaraman B, Pritsos CA and Pardini RS, The production of hydroxyl and semiquinone free radicals during the autoxidation of redox active flavonoids. In: *Oxygen Radicals in Biology and Medicine* (Eds. Simic MG, Taylor KA, Ward JF and VonSonntag C), pp. 149–153. Plenum Press, New York, 1988.
- Canada A, Gianella E, Nguyen TD and Mason RP, The production of reactive oxygen species by dietary flavonols. *Free Radic Biol Med* **9**: 441–449, 1990.
- Bern MJ, Sturbaum CW, Karayalcin SS, Berschneider HM, Wachsman JT and Powell DW, Immune system control of rat and rabbit colonic electrolyte transport: Role of prostaglandins and enteric nervous system. *J Clin Invest* **83**: 1810–1820, 1989.
- Karayalcin SS, Sturbaum CW, Dixon MU and Powell DW, Hydrogen peroxide is the reactive oxygen species (ROS) which stimulates colonic electrolyte secretion. *Gastroenterology* **94**: A216(Abtract), 1988.
- Dharmasathaphorn K, McRoberts JA, Mandel KG, Tisdale LD and Masui H, A human colonic tumor cell line that maintains vectorial electrolyte transport. *Am J Physiol* **246**: G204–G208, 1984.
- Weymer A, Huott P, Liu W, McRoberts JA and Dharmasathaphorn K, Chloride secretory mechanism induced by prostaglandin E_1 in a colonic epithelial cell line. *J Clin Invest* **76**: 1828–1836, 1985.
- Dharmasathaphorn K, Mandel KG, Masui H and McRoberts JA, Vasoactive intestinal polypeptide-induced chloride secretion by a colonic epithelial cell line: Direct participation of a basolaterally localized Na^+ , K^+ , Cl^- cotransport system. *J Clin Invest* **75**: 462–471, 1985.
- Dharmasathaphorn K and Pandolf SJ, Mechanism of chloride secretion induced by carbachol in a colonic epithelial cell line. *J Clin Invest* **77**: 348–354, 1986.
- Dharmasathaphorn K, Cohn J and Beuerlein G, Multiple calcium-mediated effector mechanisms regulate chloride secretory responses in T_{84} cells. *Am J Physiol* **256**: C1224–C1230, 1989.
- Mandel KG, Dharmasathaphorn K and McRoberts JA, Characterization of a cyclic AMP-activated Cl^- transport pathway in the apical membrane of a human colonic epithelial cell line. *J Biol Chem* **261**: 704–712, 1986.
- Cohn JA, Vasoactive intestinal peptide stimulates protein phosphorylation in a colonic epithelial cell line. *Am J Physiol* **253**: G420–G424, 1987.
- Gettys TW, Okonogi K, Tarry WC, Johnston J, Horton C and Taylor IL, Examination of relative rates of cAMP synthesis and degradation in crude membranes of adipocytes treated with hormones. *Second Messengers Phosphoproteins* **13**: 37–49, 1990.
- Fondacaro JD, Intestinal ion transport and diarrheal disease. *Am J Physiol* **250**: G1–G8, 1986.
- Mandel KG, McRoberts JA, Beuerlein G, Foster ES and Dharmasathaphorn K, Ba^{2+} inhibition of VIP and A23187-stimulated Cl^- secretion by T_{84} cell monolayers. *Am J Physiol* **250**: C486–C494, 1986.
- Cohn JA, Protein kinase C mediates cholinergically regulated protein phosphorylation in a Cl^- secreting epithelium. *Am J Physiol* **258**: C227–C233, 1990.
- Barrett KE, Cohn JA, Huott PA, Wasserman SI and Dharmasathaphorn K, Immune-related intestinal chloride secretion. II. Effect of adenosine on T_{84} cell line. *Am J Physiol* **258**: C902–C912, 1990.
- Canada AT, Watkins WD and Nguyen T, The toxicity of flavonoids to guinea pig enterocytes. *Toxicol Appl Pharmacol* **99**: 357–361, 1989.
- Brown JP, A review of the genetic effects of naturally occurring flavonoids, anthraquinones and related compounds. *Mutat Res* **75**: 243–277, 1980.
- Kuhnau J, The flavonoids. A class of semi-essential food components: Their role in human nutrition. *World Rev Nutr Diet* **24**: 117–191, 1976.
- Spiro HM, *Clinical Gastroenterology*. Macmillan, New York, 1983.