

Subject index Volume 70

- A_{2B} adenosine receptors**
Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601
- A₃ adenosine receptor agonist**
A novel adenosine analog, thio-Cl-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918
- 17-AAG**
17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580
- ABC transporter**
MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056
- ABCA1**
A novel LXR-α activator identified from the natural product *Gynostemma pentaphyllum*, 1298
- ABCB1**
Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695
- ABCG2**
Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695
- ABCG2/BCRP**
Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949
- Abnormal mitosis**
Tangutorine induces p21 expression and abnormal mitosis in human colon cancer HT-29 cells, 287
- Acanmyotoxin**
Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807
- Acanthophis***
Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807
- Acetaminophen**
Transport deficient (TR⁻) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832
- Acetylcholine**
Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
- Acetylcholinesterase**
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Actin cytoskeleton**
Calpain activation contributes to oxidative stress-induced pancreatic acinar cell injury, 1241
- Activator protein-1**
Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200
6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211
- Active site**
The role of phenylalanine 483 in cytochrome P450 2D6 is strongly substrate dependent, 1253
- Active Vitamin D**
Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479
- Acyclic nucleoside phosphonates**
Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894
- Adducts**
Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593
- Adenine nucleotides**
Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y₁₃ receptor, 266
- Adenosine**
Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355
Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883
- Adenosine A_{2B} receptor**
Evidence against adenosine analogues being agonists at the growth hormone secretagogue receptor, 598
- Adenosine A₃ receptor**
Human adenosine A₃ receptor leads to intracellular Ca²⁺ mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487
- Adenylate cyclase**
Gα_q-coupled receptor signaling enhances adenylate cyclase type 6 activation, 113
- Adhesion**
Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30
- Adipocyte**
Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90
Inhibition of adipogenesis by RGD-dependent disintegrin, 1469
- Adipogenesis**
Inhibition of adipogenesis by RGD-dependent disintegrin, 1469

- Adiponectin
Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPAR γ , 249
- Adiponectin receptor
Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPAR γ , 249
- Adipose tissue
Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPAR γ , 249
- Adrenal medullary cells
Effect of prolonged exposure to milnacipran on norepinephrine transporter in cultured bovine adrenal medullary cells, 1389
- Adrenergic agonists
Role of cytosolic phospholipase A₂ in the enhancement of α_2 -adrenoceptor-mediated vasoconstriction by the thromboxane-mimetic U46619 in the porcine isolated ear artery: Comparison with vasopressin-enhanced responses, 1200
- α_2 b-adrenergic receptor
Cloning and functional characterization of the rat α_2 B-adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606
- Adriamycin
Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066
- Agmatine
IRAS, a candidate for I₁-imidazoline receptor, mediates inhibitory effect of agmatine on cellular morphine dependence, 1079
- Agonist concentration
Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170
- Agouti related protein
Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308
- AhR
Activation of the aryl hydrocarbon receptor by berberine in HepG2 and H4IIE cells: Biphasic effect on CYP1A1, 925
- Airway smooth muscle
Thrombin induces DNA synthesis and phosphoinositide hydrolysis in airway smooth muscle by activation of distinct receptors, 959
- AKT
17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580
- Albumin
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Alcohol
Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593
- Aldehydes
Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593
- Aldose reductase inhibitors
Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653
- Alkaline comet assay
The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125
- Alkyl gallates
Effects of alkyl gallates on P-glycoprotein function, 1262
- Allosteric modulation
Allosteric modulation of ligand-gated ion channels, 1267
- Allosteric modulator
Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382
- Alpha #6
Low-molecular-weight fucoidan enhances the proangiogenic phenotype of endothelial progenitor cells, 1167
- Alternative pathway of bilirubin degradation
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
- 8-Aminoquinoline
Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158
- Amphotericin B
An effect of antibiotic amphotericin B on ion transport across model lipid membranes and tonoplast membranes, 668
- AMPK
Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- Amyloid
The binding of xanthone derivatives to transthyretin, 1861
- ANF
Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686
- Angiogenesis
Low-molecular-weight fucoidan enhances the proangiogenic phenotype of endothelial progenitor cells, 1167
- ANIT hepatotoxicity
Limited protective role of V-PYRRO/NO against cholestasis produced by alpha-naphthylisothiocyanate in mice, 144
- Anorexia
Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308
- Antagonist
A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382
Inhibitory effects of dimethylacetyl- β -cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506
- Anthocyanidins
Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure-activity relationship and molecular mechanisms involved, 417
- Anthracycline derivatives
Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431
- Anthracyclines
Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161
- Anti-cancer
Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37
- Anti-cancer drug
Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells, 676
- Anticancer drugs
Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431
- Anti-cancer therapy
Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987

Antidepressant

The effects of desmethylimipramine on cyclic AMP-stimulated gene transcription in a model cell system, 762

Anti-diabetic agent

S-15261, a new anti-hyperglycemic agent, reduces hepatic glucose production through direct and insulin-sensitizing effects, 527

Antifungal

Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82

Anti-inflammation

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330

Anti-inflammatory

The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552

Antinociception

Chronic Δ^9 -tetrahydrocannabinol treatment produces antinociceptive tolerance in mice without altering Protein Kinase A activity in mouse brain and spinal cord, 152

Antioxidant

Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242

Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618

Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642

Mitochondria-targeted peptide prevents mitochondrial depolarization and apoptosis induced by *tert*-butyl hydroperoxide in neuronal cell lines, 1796

Antioxidants

Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192

Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371

Antioxidative

The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552

Antivenom

Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794

Aorta

Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334

AP-1

Amphiphilic pyridinium salts block TNF α /NF κ B signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381

Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330

Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B, 1352

Apigenin

Oral flavonoids delay recovery from experimental autoimmune encephalomyelitis in SJL mice, 220

Apn1p

Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59

ApoE^{-/-} mice

Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334

Apolipoprotein E

A novel LXR- α activator identified from the natural product *Gynostemma pentaphyllum*, 1298

Apoptosis

Thymidine phosphorylase suppresses apoptosis induced by microtubule-interfering agents, 13

Cell apoptosis induced by a synthetic carbazole compound LCY-2-CHO is mediated through activation of caspase and mitochondrial pathways, 102

Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229

Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242

Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258

Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453

Mechanism of action of a novel "combi-triazene" engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511

TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570

17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580

Bcl-2 mediated inhibition of erucylphosphocholine-induced apoptosis depends on its subcellular localisation, 837

A novel adenosine analog, thio-C1-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918

Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066

Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH₂-terminal kinase-dependent activation of mitochondrial pathway, 1398

Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685

The effect of inhibition of Ca²⁺-independent phospholipase A₂ on chemotherapeutic-induced death and phospholipid profiles in renal cells, 1697

Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883

Apoptotic cell death

Induction of apoptotic cell death by 2'-hydroxycinnamaldehyde is involved with ERK-dependent inactivation of NF- κ B in TNF- α -treated SW620 colon cancer cells, 1147

APS

Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

Arabinosylcytosine

Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426

Arachidonic acid

Role of Ca²⁺-independent phospholipase A₂ and cytochrome P-450 in store-operated calcium entry in 3T6 fibroblasts, 733

- Aryl hydrocarbon receptor
Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse Ah^{b-1} receptor, 1623
- Ascorbate
Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343
- Aspirin
NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Astrocyte
Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590
- ATDC5
Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70
- Atherosclerosis
Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334
Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185
Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192
Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735
- ATP hydrolysis
Interactions of mefloquine with ABC proteins, MRP1 (ABCC1) and MRP4 (ABCC4) that are present in human red cell membranes, 500
- ATP synthesis
Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82
- ATP-sensitive potassium channel
Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (–)-epigallocatechin-3-gallate, 1560
- Autonomic nervous system
Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371
- AZT
Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90
The effect of AZT and chloroquine on the activities of ricin and a saporin–transferrin chimeric toxin, 560
- B⁰AT1
Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756
- Baculovirus-mediated expression system
Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019
- Base excision repair
Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59
Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458
- Bcl-2
Bcl-2 mediated inhibition of erucylphosphocholine-induced apoptosis depends on its subcellular localisation, 837
- Bcl-2 family
Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229
- BCRP
Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695
- Benzo[a]pyrene
Inhibition of osteoclast differentiation by polycyclic aryl hydrocarbons is dependent on cell density and RANKL concentration, 300
- Benzo[c]phenanthridine
Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37
- Berberine
Activation of the aryl hydrocarbon receptor by berberine in HepG2 and H4IIE cells: Biphasic effect on CYP1A1, 925
Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176
- Betaine
Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883
- Bile acid
Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685
- Bile acids
Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
- Biliary
Transport deficient (TR[–]) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832
- Bilirubin
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
- Binding affinity
The binding of xanthone derivatives to transthyretin, 1861
- Bioenergetics
The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361
- β-Lactams
Inactivators in competition. How to deal with them ... and not!, 1417
- β-Linkage
Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382
- β1
HDAC inhibition prevents NF-κB activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes IκBα, 394
- β2
HDAC inhibition prevents NF-κB activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes IκBα, 394
- β5
HDAC inhibition prevents NF-κB activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes IκBα, 394
- Blood pressure
Protein kinase C isoforms as specific targets for modulation of vascular smooth muscle function in hypertension, 1537
- Breast cancer
Inhibition of Sp1 activity by a decoy PNA–DNA chimera prevents urokinase receptor expression and migration of breast cancer cells, 1277
- Breast cancer resistance protein
Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627

- Bufalin**
Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479
- Butyrate**
HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394
- Butyrylcholinesterase knockout mice**
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Ca²⁺ entry**
Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca²⁺ signals through thiol modification in neutrophils, 1320
- Ca²⁺ homeostasis**
Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740
- Ca²⁺ influx**
Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740
- Ca²⁺ luminescence**
Motilin and erythromycin-A share a common binding site in the third transmembrane segment of the motilin receptor, 879
- Ca²⁺ release**
Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca²⁺ signals through thiol modification in neutrophils, 1320
- Ca²⁺-permeable channel**
Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740
- Ca-ATPase**
Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634
- Caco-2**
HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394
- Caco-2 cells**
Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695
Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851
- Cadmium**
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
- Calcineurin**
The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369
- Calcium**
 $G\alpha_q$ -coupled receptor signaling enhances adenylate cyclase type 6 activation, 113
Evaluation of the neuronal apoptotic pathways involved in cytoskeletal disruption-induced apoptosis, 470
Protein kinase C isoforms as specific targets for modulation of vascular smooth muscle function in hypertension, 1537
Inhibition of store-operated calcium entry-mediated superoxide generation by histamine trifluoromethyltoluide independent of histamine receptors, 1613
- Calcium channels**
Role of Ca²⁺-independent phospholipase A₂ and cytochrome P-450 in store-operated calcium entry in 3T6 fibroblasts, 733
- Calcium entry**
Non-stimulated Ca²⁺ leak pathway in cerebellar granule neurones, 786
- Calcium leak**
Non-stimulated Ca²⁺ leak pathway in cerebellar granule neurones, 786
- Calcium signal**
Non-stimulated Ca²⁺ leak pathway in cerebellar granule neurones, 786
- Calcium signaling**
AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752
- Calcium-activated potassium channel**
Tamoxifen alters gating of the BK α subunit and mediates enhanced interactions with the avian β subunit, 47
- Calcium-mediated cell damage**
Calpain activation contributes to oxidative stress-induced pancreatic acinar cell injury, 1241
- Calpain**
Calpain activation contributes to oxidative stress-induced pancreatic acinar cell injury, 1241
- Calpains**
Evaluation of the neuronal apoptotic pathways involved in cytoskeletal disruption-induced apoptosis, 470
- cAMP**
A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
- cAMP overshoot**
IRAS, a candidate for I₁-imidazoline receptor, mediates inhibitory effect of agmatine on cellular morphine dependence, 1079
- Camptothecins**
The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125
- Cancer**
Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969
- Cancer chemotherapy**
Drugs targeting mitochondrial functions to control tumor cell growth, 1
Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458
- Cannabinoid**
Chronic Δ^9 -tetrahydrocannabinol treatment produces antinociceptive tolerance in mice without altering Protein Kinase A activity in mouse brain and spinal cord, 152
- Capacitative calcium entry**
Role of Ca²⁺-independent phospholipase A₂ and cytochrome P-450 in store-operated calcium entry in 3T6 fibroblasts, 733
- Carbazole**
Cell apoptosis induced by a synthetic carbazole compound LCY-2-CHO is mediated through activation of caspase and mitochondrial pathways, 102
- Carboxylesterase**
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Cardiac glycosides**
Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851
- Cardioactive steroid**
Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851
- Cardiotonic steroid**
Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479
- Carnitine**
Regulation of testis-specific carnitine transporter (octn3) gene by proximal *cis*-acting elements Sp1 in mice, 858

Caspase

Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229

Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH₂-terminal kinase-dependent activation of mitochondrial pathway, 1398

Caspases

Cell apoptosis induced by a synthetic carbazole compound LCY-2-CHO is mediated through activation of caspase and mitochondrial pathways, 102

Evaluation of the neuronal apoptotic pathways involved in cytoskeletal disruption-induced apoptosis, 470

Cationic therapeutic drugs

Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851

Celecoxib

Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658

Cell adhesion

Resveratrol inhibits type II phosphatidylinositol 4-kinase: A key component in pathways of phosphoinositide turn over, 1048

Cell adhesion molecule

Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192

Cell cycle

TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570

Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725

Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176

The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288

Cell cycle arrest

A novel adenosine analog, thio-CI-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918

Cell cycle inhibition

Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824

Cell damage

Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740

Cell death

Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590

Cell injury

Synergistic effects of hydrogen peroxide and ethanol on cell viability loss in PC12 cells by increase in mitochondrial permeability transition, 317

Cell proliferation

Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453

The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288

Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735

Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744

Cell signaling

Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443

Cell surface molecules

Prostaglandin E₂ induces the expression of functional inhibitory CD94/NKG2A receptors in human CD8⁺ T lymphocytes by a cAMP-dependent protein kinase A type I pathway, 714

Cell-penetrating cationic peptides

Mitochondria-targeted peptide prevents mitochondrial depolarization and apoptosis induced by *tert*-butyl hydroperoxide in neuronal cell lines, 1796

Cellular resistance

Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426

Cephalosporin antibiotics

Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104

Cerebellar granule neurones

Non-stimulated Ca²⁺ leak pathway in cerebellar granule neurones, 786

cGMP

Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686

Channel site

Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382

Chemokine binding

AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752

Chemoprevention

Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969

EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568

Chemotaxis

AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752

Chick biventer

Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794

Chick biventer cervicis nerve-muscle

Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807

Chicken

Tamoxifen alters gating of the BK α subunit and mediates enhanced interactions with the avian β subunit, 47

Chlamydia pneumoniae

In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222

Chloroquine

The effect of AZT and chloroquine on the activities of ricin and a saporin-transferrin chimeric toxin, 560

Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158

CHO cell

The effects of desmethylimipramine on cyclic AMP-stimulated gene transcription in a model cell system, 762

CHO cells

Evidence against adenosine analogues being agonists at the growth hormone secretagogue receptor, 598

Cholestasis

Limited protective role of V-PYRRO/NO against cholestasis produced by alpha-naphthylisothiocyanate in mice, 144

Chondrocyte

Lysophosphatidic acid stimulates cell proliferation in rat chondrocytes, 1764

- Chondrocytes
 Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70
 Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200
- Chromanols
 The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361
- Chromatin
 A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911
- Chromium
 Hypoxia impedes the formation of chromium DNA-adducts in a cell-free system, 1814
- CINOD
 A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343
- cis*-Diamminedichloroplatinum II (cisplatin)
 Serum thymic factor, FTS, attenuates cisplatin nephrotoxicity by suppressing cisplatin-induced ERK activation, 1408
- Cisplatin
 The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137
 EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
 Specific adducts recognised by a monoclonal antibody against cisplatin-modified DNA, 1717
 Hypoxia impedes the formation of chromium DNA-adducts in a cell-free system, 1814
 Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823
- c-Jun N-terminal kinase
 6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211
- C-Jun N-terminal kinase
 Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200
- Cocaine
 The interaction of methylphenidate and bupropion with the dopamine transporter is different than other substrates and ligands, 461
- Cocaine esterase
 Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Colon adenocarcinoma cells
 NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
- Colon cancer
 17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580
 Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658
 Salicylic acid modulates oxidative stress and glutathione peroxidase activity in the rat colon, 888
 NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
 Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
 Induction of apoptotic cell death by 2'-hydroxycinnamaldehyde is involved with ERK-dependent inactivation of NF- κ B in TNF- α -treated SW620 colon cancer cells, 1147
- Combination of siRNAs
 Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424
- Competition
 Inactivators in competition. How to deal with them ... and not!, 1417
- Competition-binding studies
 The binding of xanthone derivatives to transthyretin, 1861
- Conjugated eicosapentaenoic acid (cEPA)
 Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453
- Contractile function
 Role of cytosolic phospholipase A₂ in the enhancement of α_2 -adrenoceptor-mediated vasoconstriction by the thromboxane-mimetic U46619 in the porcine isolated ear artery: Comparison with vasopressin-enhanced responses, 1200
- Coptisine
 Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176
- Coumarin 7-hydroxylation
 A novel *CYP2A6*20* allele found in African-American population produces a truncated protein lacking enzymatic activity, 801
- Covalent protein binding
 Role of benoxaprofen and flunoxaprofen acyl glucuronides in covalent binding to rat plasma and liver proteins in vivo, 937
- COX inhibition
 Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334
- COX-1
 Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969
- COX-2
 Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658
 Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969
 New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330
- COX-2 independent targets
 Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969
- Coxib therapy
 A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343
- CREB
 The effects of desmethylimipramine on cyclic AMP-stimulated gene transcription in a model cell system, 762
- Crocetin
 Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192
- Curcumin
 The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552
 Curcumin (diferuloylmethane) inhibits constitutive NF- κ B activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma, 700
- Curve analysis
 Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170

- CXCR4 antagonist
AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752
- Cyclic AMP
The effects of desmethylimipramine on cyclic AMP-stimulated gene transcription in a model cell system, 762
- Cyclic AMP assays
Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601
- Cyclin kinase inhibitor p21
Tangutorine induces p21 expression and abnormal mitosis in human colon cancer HT-29 cells, 287
- Cyclodextrin
Inhibitory effects of dimethylacetyl- β -cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506
- Cycloheximide
Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse Ah^{b-1} receptor, 1623
- Cyclooxygenase
Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30
Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334
- Cyclooxygenase (COX) inhibitor
A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343
- Cyclooxygenase-2
Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure-activity relationship and molecular mechanisms involved, 417
Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl) hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages, 1772
- Cyclopiazonic acid
Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca²⁺ signals through thiol modification in neutrophils, 1320
- Cyclosporin A
Effective prevention of lethal acute graft-versus-host disease by combined immunosuppressive therapy with prodigiosin and cyclosporine A, 1518
- CYL-26z
Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901
- CYP
Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870
- CYP1B1
Inhibition of osteoclast differentiation by polycyclic aryl hydrocarbons is dependent on cell density and RANKL concentration, 300
- CYP2C
CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096
- Cystic fibrosis
Amphiphilic pyridinium salts block TNF α /NF κ B signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381
- Cytochrome *c*
Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229
- Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242
- Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258
- Cytochrome P450
A novel CYP2A6*20 allele found in African-American population produces a truncated protein lacking enzymatic activity, 801
Activation of the aryl hydrocarbon receptor by berberine in HepG2 and H4IIE cells: Biphasic effect on CYP1A1, 925
- Cytochrome P450 2A5
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
- Cytochrome P450 2D6
The role of phenylalanine 483 in cytochrome P450 2D6 is strongly substrate dependent, 1253
- Cytokines
Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200
Controlling cytokine signaling by constitutive inhibitors, 649
- Cytoreprotection
Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590
- Cytoskeleton
Evaluation of the neuronal apoptotic pathways involved in cytoskeletal disruption-induced apoptosis, 470
- Cytotoxicity
Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453
Prostaglandin E₂ induces the expression of functional inhibitory CD94/NKG2A receptors in human CD8⁺ T lymphocytes by a cAMP-dependent protein kinase A type I pathway, 714
Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823
- Death adder
Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthopis* sp. Seram, 1807
- Degranulation
Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901
- Dehydroepiandrosterone
Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPAR γ , 249
- Delphinidin
Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure-activity relationship and molecular mechanisms involved, 417
- Deoxycytidine kinase
Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426
- Diabetes
Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355
- Diabetes complication
Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653
- Dictyostelium*
Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells, 676

Dietary factor

Ursolic acid promotes the release of macrophage migration inhibitory factor via ERK2 activation in resting mouse macrophages, 1497

DIF-1

Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells, 676

DIF-3

Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells, 676

Difference spectra

Orientation of oxazolidinones in the active site of monoamine oxidase, 407

Differentiation therapy

Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431

Digestive tract

Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756

Digitalis

Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851

Digitoxigenin derivatives

Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851

Dihydrocoumarin

Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019

Dimethylacetyl- β -cyclodextrin

Inhibitory effects of dimethylacetyl- β -cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506

Dipeptidyl peptidase IV

Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22

Dipeptidyl-peptidase IV

Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-[[(3-hydroxyadamantan-1-yl)amino]acetyl]-pyrrolidine-2-carbonitrile), 134

Disintegrin

Inhibition of adipogenesis by RGD-dependent disintegrin, 1469

Dissociated glucocorticoids

Dissociated glucocorticoids equipotently inhibit cytokine- and cAMP-induced matrix degrading proteases in rat mesangial cells, 433

Divalent cations

Allosteric modulation of ligand-gated ion channels, 1267

DNA adduct

Specific adducts recognised by a monoclonal antibody against cis-platin-modified DNA, 1717

DNA adducts

Hypoxia impedes the formation of chromium DNA-adducts in a cell-free system, 1814

DNA and RNA incorporation

Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121

DNA damage

Mechanism of action of a novel "combi-triazene" engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511

DNA methyltransferase

Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121

DNA polymerase

Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453

DNA polymerase β

Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458

DNA repair

Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458

DNA replication

Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453

The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288

DNA strand breaks

The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125

Docking simulation

The binding of xanthone derivatives to transthyretin, 1861

Dominant negative

Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685

Dopamine

The interaction of methylphenidate and bupropion with the dopamine transporter is different than other substrates and ligands, 461
Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371

Doxorubicin

Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568

Drug binding motif

Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37

Drug bioactivation

Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870

Drug discovery

Systems and integrative biology as alternative guises for pharmacology: Prime time for an *iPharm* concept?, 1707

Drug interaction

CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096

Drug metabolism

CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096

Drug resistance

MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β -D-glucuronate), 1056

Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158

Drug transport

Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851

Drug-protein adducts

Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870

Drug-resistance

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

- E₂17β-G**
MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056
- EAATs**
Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70
Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744
- EAE**
Oral flavonoids delay recovery from experimental autoimmune encephalomyelitis in SJL mice, 220
- EC₅₀**
Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170
- EGCG**
Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (–)-epigallocatechin-3-gallate, 1560
- EGF**
EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- EGFR signaling**
Mechanism of action of a novel “combi-triazene” engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511
- Eicosanoids**
Role of Ca²⁺-independent phospholipase A₂ and cytochrome P-450 in store-operated calcium entry in 3T6 fibroblasts, 733
- Electron ionization-mass spectrometry**
The effect of inhibition of Ca²⁺-independent phospholipase A₂ on chemotherapeutic-induced death and phospholipid profiles in renal cells, 1697
- Emodin**
Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229
- Endothelial cells**
Effect of statins on the proteasomal activity in mammalian endothelial and vascular smooth muscle cells, 520
Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593
- Endothelial progenitor cells**
Low-molecular-weight fucoidan enhances the proangiogenic phenotype of endothelial progenitor cells, 1167
- Endothelium**
Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30
- Endotoxin**
Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535
- eNOS**
Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343
- Enzyme inactivation**
Inactivators in competition. How to deal with them ... and not!, 1417
- Enzyme inhibitor**
Inhibitory effect of conjugated eicosapentaenoic acid on mammalian DNA polymerase and topoisomerase activities and human cancer cell proliferation, 453
- Epidermal growth factor receptor**
Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- Epirubicin**
Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725
- Epithelium**
Amphiphilic pyridinium salts block TNFα/NFκB signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381
- Epitope mapping**
A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
- Epoxyeicosatrienoic acids**
Role of Ca²⁺-independent phospholipase A₂ and cytochrome P-450 in store-operated calcium entry in 3T6 fibroblasts, 733
- ER**
Bcl-2 mediated inhibition of erucylphosphocholine-induced apoptosis depends on its subcellular localisation, 837
- Erbitux™**
EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- ERK**
Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066
Induction of apoptotic cell death by 2'-hydroxycinnamaldehyde is involved with ERK-dependent inactivation of NF-κB in TNF-α-treated SW620 colon cancer cells, 1147
- ERK activation**
Serum thymic factor, FTS, attenuates cisplatin nephrotoxicity by suppressing cisplatin-induced ERK activation, 1408
- ERK1/2**
Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF-κB, 1352
Role of mechanical and redox stress in activation of mitogen-activated protein kinases in primary cultured rat hepatocytes, 1785
- ERM proteins**
Interferon-γ-induced STAT1-mediated membrane retention of NHE1 and associated proteins ezrin, radixin and moesin in HT-29 cells, 1312
- Erucylphosphocholine**
Bcl-2 mediated inhibition of erucylphosphocholine-induced apoptosis depends on its subcellular localisation, 837
- Erythrocyte membranes**
Interactions of mefloquine with ABC proteins, MRP1 (ABCC1) and MRP4 (ABCC4) that are present in human red cell membranes, 500
- Erythroid differentiation**
Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431
- Erythromycin-A**
Motilin and erythromycin-A share a common binding site in the third transmembrane segment of the motilin receptor, 879
- ESR**
Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343
- Estrogen**
Tamoxifen alters gating of the BK α subunit and mediates enhanced interactions with the avian β subunit, 47
- Estrone sulfate**
Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104
- ET-743**
Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59

Ethanol

Synergistic effects of hydrogen peroxide and ethanol on cell viability loss in PC12 cells by increase in mitochondrial permeability transition, 317

Etodolac

Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658

Experimental therapeutics

Thymidine phosphorylase suppresses apoptosis induced by microtubule-interfering agents, 13

Extracellular matrix

Dissociated glucocorticoids equipotently inhibit cytokine- and cAMP-induced matrix degrading proteases in rat mesangial cells, 433

Extracellular matrix (ECM)

Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593

Extracellular signal-regulated kinase

Lysophosphatidic acid stimulates cell proliferation in rat chondrocytes, 1764

Extracellular signal-regulated kinase 2

Ursolic acid promotes the release of macrophage migration inhibitory factor via ERK2 activation in resting mouse macrophages, 1497

Farnesol

Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735

Farnesyl transferase inhibitor

Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735

Farnesylation

Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735

Fatty acid

Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177

Fatty acid oxidation

S-15261, a new anti-hyperglycemic agent, reduces hepatic glucose production through direct and insulin-sensitizing effects, 527

FGF-2

Low-molecular-weight fucoidan enhances the proangiogenic phenotype of endothelial progenitor cells, 1167

Fibrates

Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653

Fibronectin

Rat sinusoidal liver endothelial cells (SECs) produce pro-fibrotic factors in response to adducts formed from the metabolites of ethanol, 1593

Fibrosis

N-Acetyl-L-cysteine suppresses TGF- β signaling at distinct molecular steps: The biochemical and biological efficacy of a multifunctional, antifibrotic drug, 1026

Flavonoids

Oral flavonoids delay recovery from experimental autoimmune encephalomyelitis in SJL mice, 220
Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627

Flexible docking

Orientation of oxazolidinones in the active site of monoamine oxidase, 407

fMLP

Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca^{2+} signals through thiol modification in neutrophils, 1320

Fucoidan

Low-molecular-weight fucoidan enhances the proangiogenic phenotype of endothelial progenitor cells, 1167

G protein

$\text{G}\alpha_q$ -coupled receptor signaling enhances adenylate cyclase type 6 activation, 113

G protein-coupled receptor

Human adenosine A_3 receptor leads to intracellular Ca^{2+} mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487

GABA_A receptor

Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [^{35}S] TBPS binding to the mammalian GABA_A receptor, 1382

Gallate-ester moiety

Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (–)-epigallocatechin-3-gallate, 1560

GATA1

Cloning and functional characterization of the rat $\alpha 2\text{B}$ -adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606

GC-A

Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686

Gcn5p

A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911

Geldanamycin

Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse $\text{Ah}^{\text{b-1}}$ receptor, 1623

Gene expression

Limited protective role of V-PYRRO/NO against cholestasis produced by alpha-naphthylisothiocyanate in mice, 144
Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824
Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019
Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

Gene regulation

TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570

Genetic polymorphism

A novel *CYP2A6*20* allele found in African-American population produces a truncated protein lacking enzymatic activity, 801

Genipin

Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH_2 -terminal kinase-dependent activation of mitochondrial pathway, 1398

Ghrelin

Evidence against adenosine analogues being agonists at the growth hormone secretagogue receptor, 598

GHSR1a

Evidence against adenosine analogues being agonists at the growth hormone secretagogue receptor, 598

 $\text{G}_{i/o}$ protein

Human adenosine A_3 receptor leads to intracellular Ca^{2+} mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487

Gimatecan

The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125

- GIP**
Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-{[(3-hydroxyadamantan-1-yl)amino]acetyl}-pyrrolidine-2-carbonitrile), 134
- Glabridin**
The licorice root derived isoflavan glabridin increases the function of osteoblastic MC3T3-E1 cells, 363
- GLP-1**
Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22
Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-{[(3-hydroxyadamantan-1-yl)amino]acetyl}-pyrrolidine-2-carbonitrile), 134
- Gluconeogenesis**
S-15261, a new anti-hyperglycemic agent, reduces hepatic glucose production through direct and insulin-sensitizing effects, 527
- Gluconeogenic**
S-15261, a new anti-hyperglycemic agent, reduces hepatic glucose production through direct and insulin-sensitizing effects, 527
- Glucose**
Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- GluRs**
Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744
- Glutamate**
Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70
Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744
- Glutamate receptors**
Allosteric modulation of ligand-gated ion channels, 1267
- Glutathione**
The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552
Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870
- Glutathione peroxidase**
Salicylic acid modulates oxidative stress and glutathione peroxidase activity in the rat colon, 888
- Glycation**
Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185
- Glycolytic and lipogenic gene expression**
S-15261, a new anti-hyperglycemic agent, reduces hepatic glucose production through direct and insulin-sensitizing effects, 527
- Glycooxidation**
Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185
- G-protein-coupled receptor kinases**
Peptide inhibitors of G protein-coupled receptor kinases, 1001
- Graft-versus-host disease**
Effective prevention of lethal acute graft-versus-host disease by combined immunosuppressive therapy with prodigiosin and cyclosporine A, 1518
- Growth arrest**
Tangutorine induces p21 expression and abnormal mitosis in human colon cancer HT-29 cells, 287
Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431
- GSH**
MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056
- Guanylyl cyclase**
Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686
- Gunn rat**
Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR⁻ rats, 1088
- Gynosaponin**
A novel LXR-α activator identified from the natural product *Gynostemma pentaphyllum*, 1298
- Gynostemma pentaphyllum*
A novel LXR-α activator identified from the natural product *Gynostemma pentaphyllum*, 1298
- Gypenoside**
A novel LXR-α activator identified from the natural product *Gynostemma pentaphyllum*, 1298
- hA_{2B}CHO cells**
Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601
- Haem oxygenase 1**
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
- HAT inhibitor**
A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911
- HDAC**
HDAC inhibition prevents NF-κB activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes IκBα, 394
- HEK 293 cells**
Evidence against adenosine analogues being agonists at the growth hormone secretagogue receptor, 598
- HEK-293 cells**
Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601
- Hepatocytes**
Role of mechanical and redox stress in activation of mitogen-activated protein kinases in primary cultured rat hepatocytes, 1785
Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883
- Hepatotoxicity**
Transport deficient (TR⁻) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832
- HepG2**
Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242
- ³H]glutamate accumulation**
Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744
- [³H]Glutamate accumulation**
Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70
- Hill coefficient**
Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170
- Histone acetyltransferase**
A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911
- HIV entry**
AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752
- HL-60 cells**
A novel adenosine analog, thio-Cl-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918
- HMC-1 cells**
Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601

- HNF-4 α**
Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548
- Honokiol**
Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443
- Hormone**
Tamoxifen alters gating of the BK α subunit and mediates enhanced interactions with the avian β subunit, 47
- HPB**
Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618
- [2(1H)-Pyrimidinone riboside]**
Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121
- HSP70**
Serum thymic factor, FTS, attenuates cisplatin nephrotoxicity by suppressing cisplatin-induced ERK activation, 1408
- HT-29 cells**
Interferon- γ -induced STAT1-mediated membrane retention of NHE1 and associated proteins ezrin, radixin and moesin in HT-29 cells, 1312
- HT29 cells**
Tangutorine induces p21 expression and abnormal mitosis in human colon cancer HT-29 cells, 287
- HT-29 human**
NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
- HTMT**
Inhibition of store-operated calcium entry-mediated superoxide generation by histamine trifluoromethyltoluide independent of histamine receptors, 1613
- 5-HT₄ receptor**
A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
- Human**
Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695
Prostaglandin E₂ induces the expression of functional inhibitory CD94/NKG2A receptors in human CD8⁺ T lymphocytes by a cAMP-dependent protein kinase A type I pathway, 714
- Human lymphoid cells**
Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426
- Human paraoxonase-3**
Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019
- Human placenta**
Effects of nicotine-specific antibodies, Nic311 and Nic-IgG, on the transfer of nicotine across the human placenta, 1664
- Human plasma**
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Human red blood cells**
Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634
- Human telomerase**
Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894
- Humanized mice**
Human adenosine A₃ receptor leads to intracellular Ca²⁺ mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487
- Hydrogen peroxide**
Synergistic effects of hydrogen peroxide and ethanol on cell viability loss in PC12 cells by increase in mitochondrial permeability transition, 317
- 2'-Hydroxycinnamaldehyde**
Induction of apoptotic cell death by 2'-hydroxycinnamaldehyde is involved with ERK-dependent inactivation of NF- κ B in TNF- α -treated SW620 colon cancer cells, 1147
- Hyperglycemia**
Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653
- Hypoxia**
The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137
Hypoxia impedes the formation of chromium DNA-adducts in a cell-free system, 1814
- ICP-MS**
Specific adducts recognised by a monoclonal antibody against cisplatin-modified DNA, 1717
- IFN-beta**
The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231
- IGF**
EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- I κ B**
Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443
- I κ B α**
Curcumin (diferuloylmethane) inhibits constitutive NF- κ B activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma, 700
- I κ B α**
HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394
- IKK**
Curcumin (diferuloylmethane) inhibits constitutive NF- κ B activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma, 700
Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443
- IL-1 β**
HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394
- IL-2**
The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369
- IL-6**
New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330
- Imidazoline receptor**
IRAS, a candidate for I₁-imidazoline receptor, mediates inhibitory effect of agmatine on cellular morphine dependence, 1079
- Immunoassay**
Specific adducts recognised by a monoclonal antibody against cisplatin-modified DNA, 1717
- Immunoblot**
Role of benoxaprofen and flunoxaprofen acyl glucuronides in covalent binding to rat plasma and liver proteins in vivo, 937

In vivo efficacy

Mechanism of action of a novel "combi-triazene" engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511

Inducible nitric oxide synthase

6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211

Inflammation

Limited protective role of V-PYRRO/NO against cholestasis produced by alpha-naphthylisothiocyanate in mice, 144

Amphiphilic pyridinium salts block TNF α /NF κ B signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381

Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535

Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590

Inhibition of store-operated calcium entry-mediated superoxide generation by histamine trifluoromethyltoluide independent of histamine receptors, 1613

Inhibitors of signaling

Controlling cytokine signaling by constitutive inhibitors, 649

iNOS

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330

Inositol trisphosphate

Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y₁₃ receptor, 266

Inside-out vesicles

Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634

Insulin

Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177

Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685

Insulin secretion

Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22

Insulin-like growth factor receptor

EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568

Integrative biology

Systems and integrative biology as alternative guises for pharmacology: Prime time for an *iPharm* concept?, 1707

Integrin

Inhibition of adipogenesis by RGD-dependent disintegrin, 1469

Interferon- γ

Interferon- γ -induced STAT1-mediated membrane retention of NHE1 and associated proteins ezrin, radixin and moesin in HT-29 cells, 1312

Inhibition of interferon- γ signaling by a mercurio-substituted dihydrodipsoralen in murine keratinocytes, 1726

Interindividual variability

A novel CYP2A6*20 allele found in African-American population produces a truncated protein lacking enzymatic activity, 801

Interleukin 8

Amphiphilic pyridinium salts block TNF α /NF κ B signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381

Intestinal mapping

Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695

Intestine

Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851

Inverse agonists

Pharmacological characterization of novel adenosine ligands in recombinant and native human A_{2B} receptors, 1601

Involucrin

Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824

Ion-exchange chromatography

Specific adducts recognised by a monoclonal antibody against cisplatin-modified DNA, 1717

iPharm

Systems and integrative biology as alternative guises for pharmacology: Prime time for an *iPharm* concept?, 1707

IRAS

IRAS, a candidate for I₁-imidazoline receptor, mediates inhibitory effect of agmatine on cellular morphine dependence, 1079

Irinotecan

Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673

Iron chelator

Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642

Isoquinoline alkaloid

Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176

JAK/STAT signalling

Controlling cytokine signaling by constitutive inhibitors, 649

Janus kinase 2

6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211

JNK

Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066

K562

Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and antiproliferative activity in K562 human leukemic cells, 676

K562 cells

The effect of AZT and chloroquine on the activities of ricin and a saporin-transferrin chimeric toxin, 560

Effects of anthracycline derivatives on human leukemia K562 cell growth and differentiation, 1431

KB-C2 cells

Effects of alkyl gallates on P-glycoprotein function, 1262

Keratinocytes

Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824

Kidney

Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR⁻ rats, 1088

Kinase

Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685

Kinetics

Orientation of oxazolidinones in the active site of monoamine oxidase, 407
Inactivators in competition. How to deal with them ... and not!, 1417

Kir6.2

Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (–)-epigallocatechin-3-gallate, 1560

L₀

MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056

LAF237

Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-{[(3-hydroxyadamantan-1-yl)amino]acetyl}-pyrrolidine-2-carbonitrile), 134

LCY-2-CHO

Cell apoptosis induced by a synthetic carbazole compound LCY-2-CHO is mediated through activation of caspase and mitochondrial pathways, 102

LDL oxidation

Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019

Leptin

Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPARγ, 249

Leukocyte

Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30

Ligand-gated ion channels

Allosteric modulation of ligand-gated ion channels, 1267

Ligands

The interaction of methylphenidate and benztropine with the dopamine transporter is different than other substrates and ligands, 461

Linoleic acid

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPARγ-, dependent pathways in stimulated macrophages, 1330

Lipid peroxidation

The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361

Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634

Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642

Lipodystrophy

Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90

Lipopolysaccharide

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure–activity relationship and molecular mechanisms involved, 417

Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535

6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211

The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231

Inhibitory effects of dimethylacetyl-β-cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506

Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl) hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages, 1772

Lipoproteins

Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185

Liposomes

An effect of antibiotic amphotericin B on ion transport across model lipid membranes and tonoplast membranes, 668

5-Lipoxygenase

Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30

Lipoxygenase

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPARγ-, dependent pathways in stimulated macrophages, 1330

Liver

N-Acetyl-L-cysteine suppresses TGF-β signaling at distinct molecular steps: The biochemical and biological efficacy of a multifunctional, antifibrotic drug, 1026

Transport deficient (TR[−]) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832

Liver protein targets

Role of benoxaprofen and flunoxaprofen acyl glucuronides in covalent binding to rat plasma and liver proteins in vivo, 937

Liver X receptor

A novel LXR-α activator identified from the natural product *Gynostemma pentaphyllum*, 1298

Low concentrations of siRNA

Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424

5-LOX/COX inhibition

Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30

LS180 cells

Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695

LTC₄

MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056

Lung

Amphiphilic pyridinium salts block TNFα/NFκB signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381

Lung infection

In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222

Luteolin

Oral flavonoids delay recovery from experimental autoimmune encephalomyelitis in SJL mice, 220

In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222

Lysophosphatidic acid

Lysophosphatidic acid stimulates cell proliferation in rat chondrocytes, 1764

Macrophage

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330

Ursolic acid promotes the release of macrophage migration inhibitory factor via ERK2 activation in resting mouse macrophages, 1497

Inhibitory effects of dimethylacetyl- β -cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506

Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl) hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages, 1772

Macrophage migration inhibitory factor

Ursolic acid promotes the release of macrophage migration inhibitory factor via ERK2 activation in resting mouse macrophages, 1497

Magnesium sulfate

Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634

Malaria

Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158

MAP kinase

AMD3465, a monomacrocyclic CXCR4 antagonist and potent HIV entry inhibitor, 752

Ellagic acid blocks activation of pancreatic stellate cells, 869

Mastoparan-1

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

Matrix metalloproteinase

Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535

Matrix metalloproteinase-9

Dissociated glucocorticoids equipotently inhibit cytokine- and cAMP-induced matrix degrading proteases in rat mesangial cells, 433

Matrix metalloproteinases

Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200

MC1626

A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911

MCL

Curcumin (diferuloylmethane) inhibits constitutive NF- κ B activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma, 700

MDR

Effects of alkyl gallates on P-glycoprotein function, 1262

MDR1

Distribution of breast cancer resistance protein (BCRP/ABCG2) mRNA expression along the human GI tract, 695

MDR1 silencing

Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424

MDR1/ABCB1

Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949

Mechanical stress

Role of mechanical and redox stress in activation of mitogen-activated protein kinases in primary cultured rat hepatocytes, 1785

Mechanical stretch

Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740

Mechanism

Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82

Medica

Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548

Mefloquine

Interactions of mefloquine with ABC proteins, MRP1 (ABCC1) and MRP4 (ABCC4) that are present in human red cell membranes, 500

Melanocortin receptor

Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308

Membrane transport

Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161

Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627

Transport of dietary phenethyl isothiocyanate is mediated by multidrug resistance protein 2 but not P-glycoprotein, 640

Metabolism

Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177

Activation of the aryl hydrocarbon receptor by berberine in HepG2 and H4IIE cells: Biphasic effect on CYP1A1, 925

Metatarsals

Functional expression of particular isoforms of excitatory amino acid transporters by rodent cartilage, 70

6-Methoxydihydrosanguinarine (6ME)

Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242

Methylation

Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883

6-(Methylsulfinyl)hexyl isothiocyanate

6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages, 1211

Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl) hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages, 1772

Microtubule

Thymidine phosphorylase suppresses apoptosis induced by microtubule-interfering agents, 13

Milnacipran

Effect of prolonged exposure to milnacipran on norepinephrine transporter in cultured bovine adrenal medullary cells, 1389

Mitochondria

Drugs targeting mitochondrial functions to control tumor cell growth, 1
Cell apoptosis induced by a synthetic carbazole compound LCY-2-CHO is mediated through activation of caspase and mitochondrial pathways, 102

- Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82
- Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- Bcl-2 mediated inhibition of erucylphosphocholine-induced apoptosis depends on its subcellular localisation, 837
- Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH₂-terminal kinase-dependent activation of mitochondrial pathway, 1398
- The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361
- Mitochondrial membrane permeability**
- Synergistic effects of hydrogen peroxide and ethanol on cell viability loss in PC12 cells by increase in mitochondrial permeability transition, 317
- Mitochondrial permeability transition**
- Mitochondria-targeted peptide prevents mitochondrial depolarization and apoptosis induced by *tert*-butyl hydroperoxide in neuronal cell lines, 1796
- Mitogen-activated protein kinase**
- Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure-activity relationship and molecular mechanisms involved, 417
- Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl) hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages, 1772
- Mitoneet**
- Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- Mitoxantrone**
- Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161
- Mitoxantrone transport**
- Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627
- Mivacurium**
- Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- MK-571**
- Interactions of mefloquine with ABC proteins, MRP1 (ABCC1) and MRP4 (ABCC4) that are present in human red cell membranes, 500
- Modified lipoproteins**
- Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185
- Molecular dynamics**
- Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37
- Monoamine oxidase**
- Orientation of oxazolidinones in the active site of monoamine oxidase, 407
- Monoclonal antibody**
- A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
- Motilin fragments**
- Motilin and erythromycin-A share a common binding site in the third transmembrane segment of the motilin receptor, 879
- Motilin receptor**
- Motilin and erythromycin-A share a common binding site in the third transmembrane segment of the motilin receptor, 879
- Mouse**
- In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222
- MRP1**
- MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β-D-glucuronate), 1056
- MRP1/ABCC1**
- Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949
- MRP2**
- Transport of dietary phenethyl isothiocyanate is mediated by multidrug resistance protein 2 but not P-glycoprotein, 640
- Mrp2**
- Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR⁻ rats, 1088
- Transport deficient (TR⁻) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832
- MRP2/ABCC2**
- Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949
- Mrp4**
- Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR⁻ rats, 1088
- MtDNA**
- Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90
- MTP**
- Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548
- Multi-drug resistance**
- Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725
- Multidrug resistance**
- Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424
- Multidrug resistance**
- Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161
- Multidrug resistance-associated proteins 1 and 4**
- Interactions of mefloquine with ABC proteins, MRP1 (ABCC1) and MRP4 (ABCC4) that are present in human red cell membranes, 500
- Muscarinic receptor**
- Gα_q-coupled receptor signaling enhances adenylate cyclase type 6 activation, 113
- Muscarinic receptors**
- Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
- Muscular dystrophy**
- Protective effects of Ca²⁺ handling drugs against abnormal Ca²⁺ homeostasis and cell damage in myopathic skeletal muscle cells, 740
- Mutagenesis**
- Motilin and erythromycin-A share a common binding site in the third transmembrane segment of the motilin receptor, 879
- Myotoxin**
- Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807
- Na⁺, K⁺-ATPase**
- Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851
- nAChRs**
- Allosteric modulation of ligand-gated ion channels, 1267

NADPH oxidase

Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH₂-terminal kinase-dependent activation of mitochondrial pathway, 1398

Naproxen

A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343

Necrosis

Calpain activation contributes to oxidative stress-induced pancreatic acinar cell injury, 1241

Nephrotoxicity

Serum thymic factor, FTS, attenuates cisplatin nephrotoxicity by suppressing cisplatin-induced ERK activation, 1408

Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823

N-Ethylmaleimide

Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca²⁺ signals through thiol modification in neutrophils, 1320

Neuroblastoma

The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125

Neurodegenerative diseases

Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642

Neuropathic pain

Use-dependent blockade of Ca_v2.2 voltage-gated calcium channels for neuropathic pain, 489

Neuroprotection

Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371

Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642

Neurotoxin

Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794

Neutrophil

Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535

Neutrophils

Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901

Distinct effects of *N*-ethylmaleimide on formyl peptide- and cyclopiazonic acid-induced Ca²⁺ signals through thiol modification in neutrophils, 1320

NFAT

The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369

NF-E2

Cloning and functional characterization of the rat α 2B-adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606

NF- κ B

17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580

Curcumin (diferuloylmethane) inhibits constitutive NF- κ B activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma, 700

Induction of apoptotic cell death by 2'-hydroxycinnamaldehyde is involved with ERK-dependent inactivation of NF- κ B in TNF- α -treated SW620 colon cancer cells, 1147

NF κ B

Amphiphilic pyridinium salts block TNF α /NF κ B signaling and constitutive hypersecretion of interleukin-8 (IL-8) from cystic fibrosis lung epithelial cells, 381

NF- κ B

HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394

NF- κ B

Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B, 1352

Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443

NF κ B

New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330

NF- κ B

The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369

NHE1

Interferon- γ -induced STAT1-mediated membrane retention of NHE1 and associated proteins ezrin, radixin and moesin in HT-29 cells, 1312

Nicotine

Effects of nicotine-specific antibodies, Nic311 and Nic-IgG, on the transfer of nicotine across the human placenta, 1664

Nicotine C-oxidation

A novel *CYP2A6**20 allele found in African-American population produces a truncated protein lacking enzymatic activity, 801

Nicotine-specific antibodies

Effects of nicotine-specific antibodies, Nic311 and Nic-IgG, on the transfer of nicotine across the human placenta, 1664

Nicotinic acetylcholine receptors

Allosteric modulation of ligand-gated ion channels, 1267

Nitric oxide

Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177

Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343

Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590

Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618

Redox regulation: A new challenge for pharmacology, 811

Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B, 1352

Inhibition of interferon- γ signaling by a mercurio-substituted dihydrodropsoralen in murine keratinocytes, 1726

Nitric oxide donor

A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343

Nitric oxide (NO)

A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods, 1343

Nitric oxide synthase

Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618

- NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231
- Nitric oxide-releasing aspirin
NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
- NK109
Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37
- N*-Monomethyl-S1P
Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258
- Non-competitive inhibition
Peptide inhibitors of G protein-coupled receptor kinases, 1001
- Norepinephrine transporter
Effect of prolonged exposure to milnacipran on norepinephrine transporter in cultured bovine adrenal medullary cells, 1389
- Norepinephrine
Role of protein turnover in the activation of p38 mitogen-activated protein kinase in rat pinealocytes, 1840
- NOS2
NO-donating aspirin inhibits both the expression and catalytic activity of inducible nitric oxide synthase in HT-29 human colon cancer cells, 993
- NPR-B
Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686
- NSAIDs
Non-COX-2 targets and cancer: Expanding the molecular target repertoire of chemoprevention, 969
- Nuclear factor- κ B
Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618
- Nuclear receptor
Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479
- Nucleoside analog
Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987
- Nucleoside kinase
Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987
- Nucleoside phosphorylation
Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987
- Nucleoside transport
Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426
- Nucleoside transporter
Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355
- Nucleotide excision repair
Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59
- Null hypothesis
Systems and integrative biology as alternative guises for pharmacology: Prime time for an *iPharm* concept?, 1707
- O⁶-alkylguanine transferase
Mechanism of action of a novel "combi-triazene" engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511
- ob/ob mice
Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22
- OCT3
Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851
- OCTN2
Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756
- octn3
Regulation of testis-specific carnitine transporter (octn3) gene by proximal *cis*-acting elements Sp1 in mice, 858
- Octyl gallate
In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222
- Oligonucleotide
Specific adducts recognised by a monoclonal antibody against cis-platin-modified DNA, 1717
- Opioid dependence
IRAS, a candidate for I₁-imidazoline receptor, mediates inhibitory effect of agmatine on cellular morphine dependence, 1079
- Oral glucose tolerance test
Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22
- Organic anion transporter
Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104
- Organic cation transporter 2
Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823
- Organic ion transporter
Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756
- Osteoarthritis
Retinoic acid blocks pro-inflammatory cytokine-induced matrix metalloproteinase production by down-regulating JNK-AP-1 signaling in human chondrocytes, 200
- Osteoblastic MC3T3-E1 cells
The licorice root derived isoflavan glabridin increases the function of osteoblastic MC3T3-E1 cells, 363
- Osteoclast
Inhibition of osteoclast differentiation by polycyclic aryl hydrocarbons is dependent on cell density and RANKL concentration, 300
- Ouabain
Effect of the digitoxigenin derivative, INCICH-D7, on Na⁺, K⁺-ATPase, 851
- Oxaliplatin
Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658
- Oxazolidinone
Orientation of oxazolidinones in the active site of monoamine oxidase, 407
- Oxidative stress
Redox regulation: A new challenge for pharmacology, 811
Evidence for induced microsomal bilirubin degradation by cytochrome P450 2A5, 1527
Novel potential neuroprotective agents with both iron chelating and amino acid-based derivatives targeting central nervous system neurons, 1642
Role of mechanical and redox stress in activation of mitogen-activated protein kinases in primary cultured rat hepatocytes, 1785
Mitochondria-targeted peptide prevents mitochondrial depolarization and apoptosis induced by *tert*-butyl hydroperoxide in neuronal cell lines, 1796

- Oxidoreductase
Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653
- Oxygen
Hypoxia impedes the formation of chromium DNA-adducts in a cell-free system, 1814
- P2X receptors
Allosteric modulation of ligand-gated ion channels, 1267
- P2Y₁₃ receptor
Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y₁₃ receptor, 266
- p38 MAPK
Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258
- p38 Signaling
Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824
- p38MAPK
Role of protein turnover in the activation of p38 mitogen-activated protein kinase in rat pinealocytes, 1840
- p42/44MAPK
Role of protein turnover in the activation of p38 mitogen-activated protein kinase in rat pinealocytes, 1840
- p53/bax
Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242
- p*-Aminohippurate
Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104
- Pancreatic β -cell
Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (–)-epigallocatechin-3-gallate, 1560
- Pancreatic fibrosis
Ellagic acid blocks activation of pancreatic stellate cells, 869
- Pancreatic stellate cells
Ellagic acid blocks activation of pancreatic stellate cells, 869
- Pancreatitis
Ellagic acid blocks activation of pancreatic stellate cells, 869
- PAR
Thrombin induces DNA synthesis and phosphoinositide hydrolysis in airway smooth muscle by activation of distinct receptors, 959
- Paraoxonase
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Parkinson's disease
Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371
- Patch-clamp
An effect of antibiotic amphotericin B on ion transport across model lipid membranes and tonoplast membranes, 668
- PBMC
Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949
- PC12 cells
Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258
Synergistic effects of hydrogen peroxide and ethanol on cell viability loss in PC12 cells by increase in mitochondrial permeability transition, 317
- PEPT1
Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756
- Peptide nucleic acid
Inhibition of Sp1 activity by a decoy PNA–DNA chimera prevents urokinase receptor expression and migration of breast cancer cells, 1277
- Permeability transition pore
Drugs targeting mitochondrial functions to control tumor cell growth, 1
- Peroxynitrite
Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343
Redox regulation: A new challenge for pharmacology, 811
- Pertussis toxin-sensitive G-protein
Lysophosphatidic acid stimulates cell proliferation in rat chondrocytes, 1764
- P-glycoprotein
Transport of anthracyclines and mitoxantrone across membranes by a flip-flop mechanism, 161
Transport of dietary phenethyl isothiocyanate is mediated by multidrug resistance protein 2 but not P-glycoprotein, 640
Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725
Effects of alkyl gallates on P-glycoprotein function, 1262
- P-gp expression
Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424
- Phage display
Identification of a drug target motif: An anti-tumor drug NK109 interacts with a PNxxxxP, 37
- Pharmacogenetics
CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096
- Pharmacokinetics
Inhibition of dipeptidyl peptidase IV by novel inhibitors with pyrazolidine scaffold, 22
Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823
- Pharmacology
Systems and integrative biology as alternative guises for pharmacology: Prime time for an *iPharm* concept?, 1707
- Phenethyl isothiocyanate
Transport of dietary phenethyl isothiocyanate is mediated by multidrug resistance protein 2 but not P-glycoprotein, 640
- Phenylalanine 483
The role of phenylalanine 483 in cytochrome P450 2D6 is strongly substrate dependent, 1253
- Phenytoin
CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096
- Phosphoinositide 3-kinase γ
Human adenosine A₃ receptor leads to intracellular Ca²⁺ mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487
- Phospholipase A₂
Calpain activation contributes to oxidative stress-induced pancreatic acinar cell injury, 1241
The effect of inhibition of Ca²⁺-independent phospholipase A₂ on chemotherapeutic-induced death and phospholipid profiles in renal cells, 1697
Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807

Phospholipase D

Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901

Phospholipids

The effect of inhibition of Ca^{2+} -independent phospholipase A_2 on chemotherapeutic-induced death and phospholipid profiles in renal cells, 1697

Phosphorylation

Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121

Peptide inhibitors of G protein-coupled receptor kinases, 1001

PIAS

Controlling cytokine signaling by constitutive inhibitors, 649

Pineal

Role of protein turnover in the activation of p38 mitogen-activated protein kinase in rat pinealocytes, 1840

PKC

$\text{G}\alpha_q$ -coupled receptor signaling enhances adenylate cyclase type 6 activation, 113

PKG

Differential regulation of NPR-B/GC-B by protein kinase c and calcium, 686

Platelet-derived growth factor

Ellagic acid blocks activation of pancreatic stellate cells, 869

Platinum(IV)

The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137

PNA-DNA chimera

Inhibition of Sp1 activity by a decoy PNA-DNA chimera prevents urokinase receptor expression and migration of breast cancer cells, 1277

Polyene antibiotics

An effect of antibiotic amphotericin B on ion transport across model lipid membranes and tonoplast membranes, 668

Polygodial

Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82

Polymorphism

Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308

Polyol pathway

Fibrates inhibit aldose reductase activity in the forward and reverse reactions, 1653

Polyphenol

Ellagic acid blocks activation of pancreatic stellate cells, 869

Polyphenols

Effects of alkyl gallates on P-glycoprotein function, 1262

Porcine

Role of cytosolic phospholipase A_2 in the enhancement of α_2 -adrenoceptor-mediated vasoconstriction by the thromboxane-mimetic U46619 in the porcine isolated ear artery: Comparison with vasopressin-enhanced responses, 1200

PPADS (pyridoxal-5'-phosphate-6-azo-phenyl-2,4-disulfonate)

Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y_{13} receptor, 266

PPAR α

Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548

PPAR γ

Effect of DHEA on endocrine functions of adipose tissue, the involvement of PPAR γ , 249

Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548

Preeclampsia

Effect of magnesium sulfate on the calcium-stimulated adenosine triphosphatase activity and lipid peroxidation of red blood cell membranes from preeclamptic women, 1634

Pregnancy

Effects of nicotine-specific antibodies, Nic311 and Nic-IgG, on the transfer of nicotine across the human placenta, 1664

Primaquine

Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158

Primary cultured hepatocytes

Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870

Procaine

Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673

Prodigiosin

Effective prevention of lethal acute graft-versus-host disease by combined immunosuppressive therapy with prodigiosin and cyclosporine A, 1518

Pro-inflammatory cytokines

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

Proliferation

Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035
Lysophosphatidic acid stimulates cell proliferation in rat chondrocytes, 1764

Promoter

Cloning and functional characterization of the rat $\alpha 2\text{B}$ -adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606
Regulation of testis-specific carnitine transporter (octn3) gene by proximal *cis*-acting elements Sp1 in mice, 858

Prooxidative

The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552

Prostaglandins

Inhibition of cyclooxygenase with indomethacin phenethylamide reduces atherosclerosis in apoE-null mice, 334
Salicylic acid modulates oxidative stress and glutathione peroxidase activity in the rat colon, 888

Protease inhibitor

Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-{[(3-hydroxyadamantan-1-yl)amino]acetyl}-pyrrolidine-2-carbonitrile), 134

Proteases

Dissociated glucocorticoids equipotently inhibit cytokine- and cAMP-induced matrix degrading proteases in rat mesangial cells, 433

Proteasome

Effect of statins on the proteasomal activity in mammalian endothelial and vascular smooth muscle cells, 520
Role of protein turnover in the activation of p38 mitogen-activated protein kinase in rat pinealocytes, 1840

Proteasome

HDAC inhibition prevents NF- κB activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I $\kappa\text{B}\alpha$, 394

Protein absorption

Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756

Protein degradation

Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse Ah $^{\text{b-1}}$ receptor, 1623

Protein Kinase A

Chronic Δ^9 -tetrahydrocannabinol treatment produces antinociceptive tolerance in mice without altering Protein Kinase A activity in mouse brain and spinal cord, 152

Proteinase-activated receptors

Thrombin induces DNA synthesis and phosphoinositide hydrolysis in airway smooth muscle by activation of distinct receptors, 959

Proton transport

An effect of antibiotic amphotericin B on ion transport across model lipid membranes and tonoplast membranes, 668

Psoriasis

Inhibition of interferon- γ signaling by a mercurio-substituted dihydrodipsoralen in murine keratinocytes, 1726

Purine

Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987

Purines

Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y₁₃ receptor, 266

PUVA

Inhibition of interferon- γ signaling by a mercurio-substituted dihydrodipsoralen in murine keratinocytes, 1726

PXR

Expression of the drug transporters *MDR1/ABCB1*, *MRP1/ABCC1*, *MRP2/ABCC2*, *BCRP/ABCG2*, and *PXR* in peripheral blood mononuclear cells and their relationship with the expression in intestine and liver, 949

Pyridoxal phosphate derivatives

Synthesis of pyridoxal phosphate derivatives with antagonist activity at the P2Y₁₃ receptor, 266

Quantitative structure activity relationships

Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627

Quercetin

In vivo treatment of acute *Chlamydia pneumoniae* infection with the flavonoids quercetin and luteolin and an alkyl gallate, octyl gallate, in a mouse model, 1222

Quiescent cells

The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137

Quinazolines

Mechanism of action of a novel "combi-triazene" engineered to possess a polar functional group on the alkylating moiety: Evidence for enhancement of potency, 511

Racial difference

A novel *CYP2A6**20 allele found in African-American population produces a truncated protein lacking enzymatic activity, 801

Rad1p

Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59

RALBP1

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

Raltitrexed

Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458

RANKL

Inhibition of osteoclast differentiation by polycyclic aryl hydrocarbons is dependent on cell density and RANKL concentration, 300

Ras

Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B, 1352

RAW 264.7 macrophages

Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase by phenolic (3*E*)-4-(2-hydroxyphenyl)but-3-en-2-one in RAW 264.7 macrophages, 618

Reactive acyl glucuronide

Role of benoxaprofen and flunoxaprofen acyl glucuronides in covalent binding to rat plasma and liver proteins in vivo, 937

Reactive oxygen

N-Acetyl-L-cysteine suppresses TGF- β signaling at distinct molecular steps: The biochemical and biological efficacy of a multifunctional, antifibrotic drug, 1026

Reactive oxygen species

The relationship between the anti-inflammatory effects of curcumin and cellular glutathione content in myelomonocytic cells, 552

Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192

Neuroprotective effects of TEMPOL in central and peripheral nervous system models of Parkinson's disease, 1371

Genipin-induced apoptosis in hepatoma cells is mediated by reactive oxygen species/c-Jun NH₂-terminal kinase-dependent activation of mitochondrial pathway, 1398

Mitochondria-targeted peptide prevents mitochondrial depolarization and apoptosis induced by *tert*-butyl hydroperoxide in neuronal cell lines, 1796

Reactive oxygen species (ROS)

Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90

Reactive oxygen species-mediated induction of apoptosis by a plant alkaloid 6-methoxydihydrosanguinarine in HepG2 cells, 242

Receptor

Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170

Inhibition of insulin/IGF-1 receptor signaling enhances bile acid toxicity in primary hepatocytes, 1685

Receptor desensitization

Peptide inhibitors of G protein-coupled receptor kinases, 1001

Receptor signaling

N-Acetyl-L-cysteine suppresses TGF- β signaling at distinct molecular steps: The biochemical and biological efficacy of a multifunctional, antifibrotic drug, 1026

Recombinant expression

Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308

Redox regulation

Redox regulation: A new challenge for pharmacology, 811

Renal secretion

Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104

Renal tubular epithelial cells

Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat, 1823

Repeat addition processivity

Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894

Resistance reversing agent

Primaquine synergises the activity of chloroquine against chloroquine-resistant *P. falciparum*, 1158

Resistin

Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

Respiratory burst

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

Respiratory chain

Drugs targeting mitochondrial functions to control tumor cell growth, 1

Respiratory complexes

Site specific alterations of adipose tissue mitochondria in 3'-azido-3'-deoxythymidine (AZT)-treated rats: An early stage in lipodystrophy?, 90

Reverse transcriptase

Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894

Reversible metabolism

Role of benoxaprofen and flunoxaprofen acyl glucuronides in covalent binding to rat plasma and liver proteins in vivo, 937

Rheumatoid arthritis

Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744

RhoA

Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901

RhoGDI

Inhibition of phospholipase D activation by CYL-26z in formyl peptide-stimulated neutrophils involves the blockade of RhoA activation, 901

Ribonucleotide reductase

The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288

Ricin

The effect of AZT and chloroquine on the activities of ricin and a saporin-transferrin chimeric toxin, 560

RLIP76

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

RNG

Cloning and functional characterization of the rat α 2B-adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606

Rolling

Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30

ROS

Emodin induces apoptosis in human lung adenocarcinoma cells through a reactive oxygen species-dependent mitochondrial signaling pathway, 229

Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066

Roscovitine

Effects of the cyclin-dependent kinase inhibitor CYC202 (*R*-roscovitine) on the physiology of cultured human keratinocytes, 824

Rosiglitazone

Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

Rosmarinic acid

Inhibitory effects of rosmarinic acid on adriamycin-induced apoptosis in H9c2 cardiac muscle cells by inhibiting reactive oxygen species and the activations of c-Jun N-terminal kinase and extracellular signal-regulated kinase, 1066

Saccharomyces cerevisiae

Low cytotoxicity of ecteinascidin 743 in yeast lacking the major endonucleolytic enzymes of base and nucleotide excision repair pathways, 59

S-adenosylhomocysteine

Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: Protection by betaine, 1883

SAR

Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [35 S] TBPS binding to the mammalian GABA_A receptor, 1382

Second messengers

Role of cytosolic phospholipase A₂ in the enhancement of α ₂-adrenoceptor-mediated vasoconstriction by the thromboxane-mimetic U46619 in the porcine isolated ear artery: Comparison with vasopressin-enhanced responses, 1200

Selectin

Licofelone, a novel 5-LOX/COX-inhibitor, attenuates leukocyte rolling and adhesion on endothelium under flow, 30

Selectivity

Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176

Sepsis

A synthesized cationic tetradecapeptide from hornet venom kills bacteria and neutralizes lipopolysaccharide in vivo and in vitro, 209

Targeting neutrophil collagenase/matrix metalloproteinase-8 and gelatinase B/matrix metalloproteinase-9 with a peptidomimetic inhibitor protects against endotoxin shock, 535

Inhibitory effects of dimethylacetyl- β -cyclodextrin on lipopolysaccharide-induced macrophage activation and endotoxin shock in mice, 1506

Serine-proteases

Evaluation of the neuronal apoptotic pathways involved in cytoskeletal disruption-induced apoptosis, 470

Serotonin norepinephrine reuptake inhibitor

Effect of prolonged exposure to milnacipran on norepinephrine transporter in cultured bovine adrenal medullary cells, 1389

Serum thymic factor (FTS)

Serum thymic factor, FTS, attenuates cisplatin nephrotoxicity by suppressing cisplatin-induced ERK activation, 1408

Sf9 cell

Cloning, high level expression of human paraoxonase-3 in Sf9 cells and pharmacological characterization of its product, 1019

SHP

Controlling cytokine signaling by constitutive inhibitors, 649

Signal transduction

TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570

Sequential induction of heme oxygenase-1 and manganese superoxide dismutase protects cultured astrocytes against nitric oxide, 590

Resveratrol inhibits type II phosphatidylinositol 4-kinase: A key component in pathways of phosphoinositide turn over, 1048

Bile acid-induced proliferation of a human colon cancer cell line is mediated by transactivation of epidermal growth factor receptors, 1035

Protein kinase C isoforms as specific targets for modulation of vascular smooth muscle function in hypertension, 1537

Signaling pathways

Role of mechanical and redox stress in activation of mitogen-activated protein kinases in primary cultured rat hepatocytes, 1785

Simvastatin

Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185

siRNA

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

Site-directed mutagenesis

The role of phenylalanine 483 in cytochrome P450 2D6 is strongly substrate dependent, 1253

Skeletal muscle

Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794

Contrasting effects of insulin and cellular differentiation on expression of the novel insulin receptor substrate APS in skeletal muscle, 1309

- Skin**
Inhibition of interferon- γ signaling by a mercurio-substituted dihydrodropsoralen in murine keratinocytes, 1726
- Smad**
N-Acetyl-L-cysteine suppresses TGF- β signaling at distinct molecular steps: The biochemical and biological efficacy of a multifunctional, antifibrotic drug, 1026
- Small interference RNA**
The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288
- Small interfering RNA**
Modulation of *MDR1* gene expression in multidrug resistant MCF7 cells by low concentrations of small interfering RNAs, 1424
- Smooth muscle cell**
Isothiazole dioxide derivative 6n inhibits vascular smooth muscle cell proliferation and protein farnesylation, 1735
- Smooth muscle cells**
Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355
Effect of statins on the proteasomal activity in mammalian endothelial and vascular smooth muscle cells, 520
- SN-38**
MRP1 mutated in the L₀ region transports SN-38 but not leukotriene C₄ or estradiol-17 (β -D-glucuronate), 1056
- Snake**
Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794
- SOCE**
Inhibition of store-operated calcium entry-mediated superoxide generation by histamine trifluoromethyltoluide independent of histamine receptors, 1613
- SOCs**
Controlling cytokine signaling by constitutive inhibitors, 649
- Solid tumour models**
The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137
- Sp1**
Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426
Regulation of testis-specific carnitine transporter (octn3) gene by proximal *cis*-acting elements Sp1 in mice, 858
- Sp1 transcription factor**
Inhibition of Sp1 activity by a decoy PNA-DNA chimera prevents urokinase receptor expression and migration of breast cancer cells, 1277
- Species difference**
Human adenosine A₃ receptor leads to intracellular Ca²⁺ mobilization but is insufficient to activate the signaling pathway via phosphoinositide 3-kinase γ in mice, 1487
- Sphingosine-1-phosphate (S1P)**
Involvement of p38 MAP kinase-mediated cytochrome *c* release on sphingosine-1-phosphate (S1P)- and *N*-monomethyl-S1P-induced cell death of PC12 cells, 258
- Src-family tyrosine kinases**
The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231
- Stably transfected cell lines**
Drug specificity and intestinal membrane localization of human organic cation transporters (OCT), 1851
- Stalk-cell differentiation**
Structural requirements of *Dictyostelium* differentiation-inducing factors for their stalk-cell-inducing activity in *Dictyostelium* cells and anti-proliferative activity in K562 human leukemic cells, 676
- Staphylococcus aureus BlaR**
Inactivators in competition. How to deal with them ... and not!, 1417
- Staphylococcus aureus PBP2a**
Inactivators in competition. How to deal with them ... and not!, 1417
- Statins**
Effect of statins on the proteasomal activity in mammalian endothelial and vascular smooth muscle cells, 520
Effect of simvastatin on the oxidation of native and modified lipoproteins, 1185
EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- Stimulation**
Z-analysis: A new approach to analyze stimulation curves with intrinsic basal stimulation, 170
- Stress response**
Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- Structure activity relationships**
Structure activity relationships and quantitative structure activity relationships for the flavonoid-mediated inhibition of breast cancer resistance protein, 627
- Subcellular localization**
The ribonucleotide reductase subunit M2B subcellular localization and functional importance for DNA replication in physiological growth of KB cells, 1288
- Substance P**
Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-{[(3-hydroxyadamantan-1-yl)amino]acetyl}-pyrrolidine-2-carbonitrile), 134
- Substrate selectivity**
The role of phenylalanine 483 in cytochrome P450 2D6 is strongly substrate dependent, 1253
- Substrates**
The interaction of methylphenidate and benztrapine with the dopamine transporter is different than other substrates and ligands, 461
- Succinylcholine**
Butyrylcholinesterase, paraoxonase, and albumin esterase, but not carboxylesterase, are present in human plasma, 1673
- Sugar**
Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382
- Suicide gene therapy**
Phosphorylation of 9- β -D-arabinofuranosylguanine monophosphate by *Drosophila melanogaster* guanylate kinase, 987
- Superoxide**
Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343
Redox regulation: A new challenge for pharmacology, 811
Inhibition of store-operated calcium entry-mediated superoxide generation by histamine trifluoromethyltoluide independent of histamine receptors, 1613
- Surface plasmon resonance**
A high-affinity monoclonal antibody with functional activity against the 5-hydroxytryptaminergic (5-HT₄) receptor, 1009
- Survivin**
Combination of cyclooxygenase-2 inhibitors and oxaliplatin increases the growth inhibition and death in human colon cancer cells, 658
- Synovial fibroblasts**
Excitatory amino acid transporters expressed by synovial fibroblasts in rats with collagen-induced arthritis, 1744
- T cells**
Oral flavonoids delay recovery from experimental autoimmune encephalomyelitis in SJL mice, 220

- The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369
- TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570
- T lymphocytes
- Prostaglandin E₂ induces the expression of functional inhibitory CD94/NKG2A receptors in human CD8⁺ T lymphocytes by a cAMP-dependent protein kinase A type I pathway, 714
 - Resveratrol inhibits type II phosphatidylinositol 4-kinase: A key component in pathways of phosphoinositide turn over, 1048
- T24 bladder carcinoma
- Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121
- Tamoxifen
- Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725
- Tangutorine
- Tangutorine induces p21 expression and abnormal mitosis in human colon cancer HT-29 cells, 287
- TarcevaTM
- EGFR blockade by cetuximab alone or as combination therapy for growth control of hepatocellular cancer, 1568
- Taurine chloramine
- Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B, 1352
- TBPS
- Radioligand binding studies of caloposide and novel congeners with contrasting effects upon [³⁵S] TBPS binding to the mammalian GABA_A receptor, 1382
- TCDD
- Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse Ah^{b-1} receptor, 1623
- Telomerase inhibition
- Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894
- Testis
- Regulation of testis-specific carnitine transporter (octn3) gene by proximal *cis*-acting elements Sp1 in mice, 858
- Tetrahydrocannabinol
- CYP2C-catalyzed delta(9)-tetrahydrocannabinol metabolism: Kinetics, pharmacogenetics and interaction with phenytoin, 1096
- THC
- Chronic Δ^9 -tetrahydrocannabinol treatment produces antinociceptive tolerance in mice without altering Protein Kinase A activity in mouse brain and spinal cord, 152
- Thiazolidinedione
- Receptor-independent actions of PPAR thiazolidinedione agonists: Is mitochondrial function the key?, 177
- Thiazolidinediones
- Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355
 - Transcriptional suppression of human microsomal triglyceride transfer protein by hypolipidemic insulin sensitizers, 1548
- Thio-Cl-IB-MECA
- A novel adenosine analog, thio-Cl-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918
- Thiol oxidation
- The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361
- Thiols
- Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthesis, 343
- Thrombin
- Thrombin induces DNA synthesis and phosphoinositide hydrolysis in airway smooth muscle by activation of distinct receptors, 959
- Thymidine phosphorylase
- Thymidine phosphorylase suppresses apoptosis induced by microtubule-interfering agents, 13
- Thymineless stress
- Determination of apoptosis, uracil incorporation, DNA strand breaks, and sister chromatid exchanges under conditions of thymidylate deprivation in a model of BER deficiency, 1458
- Tienilic acid
- Kinetics of tienilic acid bioactivation and functional generation of drug-protein adducts in intact rat hepatocytes, 1870
- Tissue plasminogen activator
- Dissociated glucocorticoids equipotently inhibit cytokine- and cAMP-induced matrix degrading proteases in rat mesangial cells, 433
- TNF- α
- HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394
- TNF- α
- Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation, 1443
- Tocopheryl quinone
- The protection of bioenergetic functions in mitochondria by new synthetic chromanols, 1361
- Tolerance
- Chronic Δ^9 -tetrahydrocannabinol treatment produces antinociceptive tolerance in mice without altering Protein Kinase A activity in mouse brain and spinal cord, 152
- Toll-like receptor
- The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231
 - New class of linoleic acid metabolites biosynthesized by corn and rice lipoxygenases: Suppression of proinflammatory mediator expression via attenuation of MAPK- and Akt-, but not PPAR γ -, dependent pathways in stimulated macrophages, 1330
- Topoisomerase I inhibitors
- The novel lipophilic camptothecin analogue gimatecan is very active in vitro in human neuroblastoma: A comparative study with SN38 and topotecan, 1125
- Topoisomerase II α
- Modulation of epirubicin cytotoxicity by tamoxifen in human breast cancer cell lines, 725
- Toxins
- The effect of AZT and chloroquine on the activities of ricin and a saporin-transferrin chimeric toxin, 560
- TR⁻ rat
- Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR⁻ rats, 1088
 - Transport deficient (TR⁻) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity, 1832
- TRAIL
- 17-Allylamino-17-demethoxygeldanamycin overcomes TRAIL resistance in colon cancer cell lines, 580
- Transcription
- Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479
- Transcription factor
- Cytosine arabinoside affects multiple cellular factors and induces drug resistance in human lymphoid cells, 426
 - Suppression of vascular cell adhesion molecule-1 expression by crocetin contributes to attenuation of atherosclerosis in hypercholesterolemic rabbits, 1192

Transcription factors

Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: Structure–activity relationship and molecular mechanisms involved, 417

TCR pathway involves ICBP90 gene down-regulation via E2F binding sites, 570

Cloning and functional characterization of the rat α 2B-adrenergic receptor gene promoter region: Evidence for binding sites for erythropoiesis-related transcription factors GATA1 and NF-E2, 606

Transfection

Depletion of RLIP76 sensitizes lung cancer cells to doxorubicin, 481

Transferrin

The effect of AZT and chloroquine on the activities of ricin and a saporin–transferrin chimeric toxin, 560

Transplacental transfer

Effects of nicotine-specific antibodies, Nic311 and Nic-IgG, on the transfer of nicotine across the human placenta, 1664

Transport

Human organic anion transporter hOAT3 is a potent transporter of cephalosporin antibiotics, in comparison with hOAT1, 1104

Transporter

The interaction of methylphenidate and benztropine with the dopamine transporter is different than other substrates and ligands, 461

Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract, 1756

Transthyretin

The binding of xanthone derivatives to transthyretin, 1861

TRAP

Inhibition of osteoclast differentiation by polycyclic aryl hydrocarbons is dependent on cell density and RANKL concentration, 300

TRAP assay

Inhibition of human telomerase by diphosphates of acyclic nucleoside phosphonates, 894

TRIF

The regulation of the expression of inducible nitric oxide synthase by Src-family tyrosine kinases mediated through MyD88-independent signaling pathways of Toll-like receptor 4, 1231

Troglitazone

Effect of thiazolidinediones on equilibrative nucleoside transporter-1 in human aortic smooth muscle cells, 355

Tropisetron

The 5-HT₃ receptor antagonist tropisetron inhibits T cell activation by targeting the calcineurin pathway, 369

TSA

HDAC inhibition prevents NF- κ B activation by suppressing proteasome activity: Down-regulation of proteasome subunit expression stabilizes I κ B α , 394

Tumour microenvironment

The influence of tumour microenvironmental factors on the efficacy of cisplatin and novel platinum(IV) complexes, 1137

Uncoupler

Inhibition of the mitochondrial ATP synthesis by polygodial, a naturally occurring dialdehyde unsaturated sesquiterpene, 82

Up-regulation

Up-regulation of Mrp4 expression in kidney of Mrp2-deficient TR[−] rats, 1088

Effect of prolonged exposure to milnacipran on norepinephrine transporter in cultured bovine adrenal medullary cells, 1389

Uptake

The interaction of methylphenidate and benztropine with the dopamine transporter is different than other substrates and ligands, 461

Urate

Interactions of peroxynitrite with uric acid in the presence of ascorbate and thiols: Implications for uncoupling endothelial nitric oxide synthase, 343

Urokinase receptor

Inhibition of Sp1 activity by a decoy PNA–DNA chimera prevents urokinase receptor expression and migration of breast cancer cells, 1277

Ursolic acid

Ursolic acid promotes the release of macrophage migration inhibitory factor via ERK2 activation in resting mouse macrophages, 1497

Use-dependent block

Use-dependent blockade of Ca_v2.2 voltage-gated calcium channels for neuropathic pain, 489

Vascular inflammation

Redox regulation: A new challenge for pharmacology, 811

Vascular smooth muscle

Protein kinase C isoforms as specific targets for modulation of vascular smooth muscle function in hypertension, 1537

Vascular smooth muscle cell

Double blockade of cell cycle progression by coptisine in vascular smooth muscle cells, 1176

Venom

Isolation and pharmacological characterisation of papuatoxin-1, a postsynaptic neurotoxin from the venom of the Papuan black snake (*Pseudechis papuanus*), 794

Isolation and characterisation of acanmyotoxin-2 and acanmyotoxin-3, myotoxins from the venom of the death adder *Acanthophis* sp. Seram, 1807

Vitamin D receptor

Enhancement of ligand-dependent Vitamin D receptor transactivation by the cardiotonic steroid bufalin, 1479

Vitamin E

Salicylic acid modulates oxidative stress and glutathione peroxidase activity in the rat colon, 888

Voltage-gated calcium channels

Use-dependent blockade of Ca_v2.2 voltage-gated calcium channels for neuropathic pain, 489

V-PYRRO/NO

Limited protective role of V-PYRRO/NO against cholestasis produced by alpha-naphthylisothiocyanate in mice, 144

Weight regulation

Functional analysis of the Ala67Thr polymorphism in agouti related protein associated with anorexia nervosa and leanness, 308

Wnt signaling

A novel adenosine analog, thio-Cl-IB-MECA, induces G₀/G₁ cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells, 918

Xanthone

The binding of xanthone derivatives to transthyretin, 1861

Xenoeostrogen

Tamoxifen alters gating of the BK α subunit and mediates enhanced interactions with the avian β subunit, 47

Xenopus oocyte

Inhibitory modulation of ATP-sensitive potassium channels by gallate-ester moiety of (−)-epigallocatechin-3-gallate, 1560

Yeast

A novel Gcn5p inhibitor represses cell growth, gene transcription and histone acetylation in budding yeast, 911

Zebularine

Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells, 121