

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/biochempharm

Corrigendum

Corrigendum to “A comparison of the cyclooxygenase inhibitor-NO donors (CINOD), NMI-1182 and AZD3582, using in vitro biochemical and pharmacological methods” [Biochem. Pharmacol. 70/9 (2005) 1343–1351]

Delano V. Young^{a,*}, Edward D. Cochran^a, Vijay Dhawan^a, Richard A. Earl^b, James L. Ellis^a, David S. Garvey^b, David R. Janero^a, Subhash P. Khanapure^b, L. Gordon Letts^{a,b}, Terry L. Melim^a, Madhavi G. Murty^a, Matthew J. Shumway^a, Shiow-Jyi Wey^b, Irina S. Zemtseva^a, William M. Selig^a

^a Department of Biology, NitroMed Inc., 125 Spring St., Lexington, MA 02421, USA

^b Department of Chemistry, NitroMed Inc., 125 Spring St., Lexington, MA 02421, USA

The authors regret that in the above article Fig. 11 shown on p. 1349 was printed incorrectly. It is now reproduced correctly here. The authors apologise for any confusion this may have caused.




Fig. 11 – NMI-1182 produced more absolute total NO_x, much of it as nitrite, than AZD3582; most of the AZD3582 NO_x was in the form of nitrate, not nitrite in human liver S9 homogenates. Liver S9 homogenate incubations of test compounds (added at 70 μM) for NO_x and nitrite were performed as described in detail in Section 2. The results were the averages ± S.E.M. of three experiments (each performed in triplicate); **p* < 0.01 and ****p* < 0.001 comparing total NO_x from NMI-1182 vs. AZD3582 (closed symbols) or nitrite released by NMI-1182 vs. AZD3582 (open symbols) at the indicated times, by two-way ANOVA, Bonferroni post test.

DOI of original article: 10.1016/j.bcp.2005.08.004.

* Corresponding author. Tel.: +1 781 266 4184; fax: +1 781 274 8083.

E-mail address: dyoung@nitromed.com (D.V. Young).

0006-2952/\$ – see front matter © 2005 Elsevier Inc. All rights reserved.

doi:10.1016/j.bcp.2005.12.011