

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/biochempharm

Subject index Volume 71

14-3-3ζ

14-3-3ζ interacts with human thromboxane receptors and is involved in the agonist-induced activation of the extra-cellular-signal-regulated kinase, 624

A3AR functionally humanized mice

Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294

Abacavir 5'-monophosphate

N⁶-Methyl-AMP aminohydrolase activates N⁶-substituted purine acyclic nucleoside phosphonates, 1370

ABCA1

Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605

ABCB1

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

ABCC

Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470

ABCC1

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

ABCC2

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

ABCC3

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

Acetylcholinesterase

Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497

Actinomycetes

The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981

Action potentials

Diethyl pyrocarbonate, a histidine-modifying agent, directly stimulates activity of ATP-sensitive potassium channels in pituitary GH₃ cells, 615

Activating protein-1

Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by N-lauroyl-D-erythro-sphingosine and N-lauroyl-D-erythro-C₂₀-sphingosine, 1229

Adenosine A₁ receptor

Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand-receptor interaction, 865

Adenosine A3 receptor

Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294

Adenosine kinase

Structure-activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671

Adenylyl cyclase

Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ-thio]triphosphates and nucleoside 5'-[β,γ-imido]triphosphates, 89

Adipocyte

Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184

Adipocytokine

Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184

Adrenalectomy

Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155

Affinity index

Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858

Affinity label

Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629

AG1478

Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422

AGC1

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Agonist

Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663

AICAR

AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637

Airways

A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223

Akt

AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637

- Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735
- Aldose reductase inhibitor**
Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338
- Aldosterone**
Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175
- Aldoxime**
Oxygen-insensitive enzymatic reduction of oximes to imines, 354
- Alkylphosphocholines**
Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114
- Allergy**
Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by N-lauroyl-D-erythro-sphingosine and N-lauroyl-D-erythro-C₂₀-sphingosine, 1229
- Allopurinol**
Superoxide radical production by allopurinol and xanthine oxidase, 1747
- Alpha class**
Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629
- Amidoxime**
Oxygen-insensitive enzymatic reduction of oximes to imines, 354
- 4-Aminobutyrate transaminase**
Inhibition of rabbit brain 4-aminobutyrate transaminase by some taurine analogues: A kinetic analysis, 1510
- 4-Aminobutyrate transaminase inhibitors**
Inhibition of rabbit brain 4-aminobutyrate transaminase by some taurine analogues: A kinetic analysis, 1510
- δ-Aminolevulinic acid**
Mechanisms involved in δ-aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: Relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin, 42
- 4-Aminoquinolines**
Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61
- 21-Aminosteroid U74389G**
The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366
- Amitriptyline**
Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA_B receptor expression and function in an animal model of neuropathic pain, 196
- AMPA receptor potentiators**
A role for AMPA receptors in mood disorders, 1273
- AMP-activated kinase**
AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637
- Amphotericin**
Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779
- AMPK**
AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637
- Angiotensin**
Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319
- Angiotensin II**
Cross-talk between angiotensin II and glucagon receptor signaling mediates phosphorylation of mitogen-activated protein kinases ERK 1/2 in rat glomerular mesangial cells, 1711
- Animal Models**
The role of pharmacodynamic research in the assessment and development of new antibacterial drugs, 1057
- Antagonist**
Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663
- Antagonist/inverse agonist**
The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103
- Anthocyanin**
Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184
- Anthracyclin analog**
Inhibitory effect of DA-125, a new anthracyclin analog anti-tumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21
- Anthracycline**
Contribution of an unidentified sodium-dependent nucleoside transport system to the uptake and cytotoxicity of anthracycline in mouse M5076 ovarian sarcoma cells, 565
- Antraquinone**
Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136
- Antibacterial**
Antibacterial drug discovery—Then, now and the genomics future, 901
Natural products — The future scaffolds for novel antibiotics?, 919
Structure-based drug design meets the ribosome, 1016
The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042
Crystallizing new approaches for antimicrobial drug discovery, 1048
The role of pharmacodynamic research in the assessment and development of new antibacterial drugs, 1057
Use of genomics to select antibacterial targets, 1066
When will the genomics investment pay off for antibacterial discovery?, 1096
- Antibacterial agents**
Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996
- Antibacterial discovery**
Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996
- Antibiotic**
Understanding the longevity of the β-lactam antibiotics and of antibiotic/β-lactamase inhibitor combinations, 930
Natural products — The future scaffolds for novel antibiotics?, 919
Glycopeptides: Update on an old successful antibiotic class, 968
Use of genomics to select antibacterial targets, 1066
- Antibiotic efflux**
Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949

- The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073
- Antibiotic potentiating agent
Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949
- Antibiotic resistance
Commensals upon us, 893
Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949
The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042
Surveillance uncovers the smoking gun for resistance emergence, 1036
- Antibiotic synergy
Antibacterial drug discovery—Then, now and the genomics future, 901
- Antibiotics
Antibacterial drug discovery—Then, now and the genomics future, 901
The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981
Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996
Empirical antibacterial drug discovery—Foundation in natural products, 1006
Structure-based drug design meets the ribosome, 1016
Crystallizing new approaches for antimicrobial drug discovery, 1048
- Anticancer activity
Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24), 426
- Antichlamydial activity
Inhibitory effect of dietary phenolic compounds on *Chlamydia pneumoniae* in cell cultures, 735
- Antimalarial agents
Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61
- Antimicrobial
The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042
- Antimicrobial agents
Unmet medical needs in antibacterial therapy, 991
- Antimicrobial drug discovery
Crystallizing new approaches for antimicrobial drug discovery, 1048
- Antimicrobial peptide
Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289
- Antimicrobial research
Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026
- Antimonials
Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257
- Antioxidant
Antioxidative function and biodistribution of $[\text{Gd}@\text{C}_{82}(\text{OH})_{22}]_n$ nanoparticles in tumor-bearing mice, 872
- Anti-oxidant enzymes
Suppression of survival in human SKBR3 breast carcinoma in response to metal-chelator complexes is preferential for copper-dithiocarbamate, 722
- Antipsychotics
Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521
- Antitumor drug resistance
Expression of γ -glutamyltransferase in cancer cells and its significance in drug resistance, 231
- AOAA
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- AP-1
MZF1 possesses a repressively regulatory function in ERCC1 expression, 761
Molecular targets of dietary agents for prevention and therapy of cancer, 1397
Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, 1602
- Apoptosis
Cell cycle arrest and proapoptotic effects of the anticancer cyclodepsipeptide serratamolide (AT514) are independent of p53 status in breast cancer cells, 32
Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257
Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24), 426
Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530
Schisandrin B enhances doxorubicin-induced apoptosis of cancer cells but not normal cells, 584
Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742
Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791
Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114
Hydroxymethylglutaryl-coenzyme A reductase inhibitors induce apoptosis in human cardiac myocytes in vitro, 1324
Molecular targets of dietary agents for prevention and therapy of cancer, 1397
Blockade of IGF-1 receptor tyrosine kinase has antineoplastic effects in hepatocellular carcinoma cells, 1435
The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540
Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562
- Arachidonic acid
Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂- α , 144
Ceramide-1-phosphate activates cytosolic phospholipase A₂ α directly and by PKC pathway, 850
- aralar1
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Aripiprazole
Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521

- Artocarpol A**
Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735
- Arylpiperazines**
Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61
- Aspartate aminotransferase**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Aspartate–glutamate carrier**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Assay**
Natural products — The future scaffolds for novel antibiotics?, 919
- Astrocytes**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- AT₁ receptor**
Conformational induction is the key process for activation of the AT₁ receptor, 464
- ATM**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- Atorvastatin**
Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605
- ATP**
CysLT₁ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115
Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y₁ receptor in a Ca²⁺ independent manner in differentiated human skeletal muscle cells, 1497
- ATP-binding cassette transporter A1**
Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453
- Axon outgrowth**
Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240
- AZT**
3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342
- B cell malignancy**
Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742
- Bacterial cell wall biosynthesis**
Targeting the forgotten transglycosylases, 957
- Baculovirus**
Agonist potency differentiates G protein activation and Ca²⁺ signalling by the orexin receptor type 1, 827
- 6-Benzylthioinosine analogues**
6-Benzylthioinosine analogues: Promising anti-toxoplasmaic agents as inhibitors of the mammalian nucleoside transporter ENT1 (es), 69
- Berberine**
Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806
- Bilirubin**
Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693
- Binding**
Characterization of the melatonergic MT₃ binding site on the NRH:quinone oxidoreductase 2 enzyme, 74
- Binding mode**
Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338
- Binding stoichiometry**
Characterization of the inhibitory activity of tenatoprazole on the gastric H⁺,K⁺-ATPase in vitro and in vivo, 837
- Binding studies**
Synthesis and antinociceptive activity of cyclic endomorphin-2 and morphiceptin analogs, 188
- Biodistribution**
Antioxidative function and biodistribution of [Gd@C₈₂(OH)₂₂]_n nanoparticles in tumor-bearing mice, 872
- Biofilms**
Unmet medical needs in antibacterial therapy, 991
- Bioinformatics**
Use of genomics to select antibacterial targets, 1066
When will the genomics investment pay off for antibacterial discovery?, 1096
- Biological therapies**
ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479
- Blood**
Metabolism of oxidants by blood from different mouse strains, 1753
- Blood pressure**
Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175
- BMA**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Bone loss**
Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818
- Brain-derived neurotrophic factor (BDNF)**
A role for AMPA receptors in mood disorders, 1273
- Breast cancer**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- Buprenorphine (BUP)**
The effect of methadone and buprenorphine on human placental aromatase, 1255
- Ca²⁺**
Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449
- Ca²⁺ influx**
N-(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- Caco-2**
Modulation of intestinal barrier properties by miltefosine, 486

- The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581
- Caco-2 cells**
Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570
Effects of capsaicin on P-gp function and expression in Caco-2 cells, 1727
- Calcitonin gene related peptide**
A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223
- Calcitonin receptor**
Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818
- Calcium**
Agonist potency differentiates G protein activation and Ca^{2+} signalling by the orexin receptor type 1, 827
- Calcium imaging**
Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173
- Calcium signalling**
Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y_1 receptor in a Ca^{2+} independent manner in differentiated human skeletal muscle cells, 1497
- Calmodulin**
Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449
- cAMP**
Serotonin induces pulmonary artery smooth muscle cell migration, 386
Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219
The Cdc42 inhibitor secramine B prevents cAMP-induced K^+ conductance in intestinal epithelial cells, 1720
- Camptothecins**
Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791
- Cancer**
Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136
Molecular targets of dietary agents for prevention and therapy of cancer, 1397
- Cannabidiol**
The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146
- Cannabinoids**
The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146
- Capsaicin**
Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173
Effects of capsaicin on P-gp function and expression in Caco-2 cells, 1727
- Carbapenemases**
The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073
- Carbon monoxide**
CO from enhanced HO activity or from CORM-2 inhibits both O_2^- and NO production and downregulates HO-1 expression in LPS-stimulated macrophages, 307
- Carbonylation**
Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610
- Carbovir 5'-monophosphate**
 N^6 -Methyl-AMP aminohydrolase activates N^6 -substituted purine acyclic nucleoside phosphonates, 1370
- Carboxylesterases**
Hydrolytic metabolism of pyrethroids by human and other mammalian carboxylesterases, 657
- Cardiac myocytes**
Hydroxymethylglutaryl-coenzyme A reductase inhibitors induce apoptosis in human cardiac myocytes in vitro, 1324
- Cardiac remodeling**
Hydroxymethylglutaryl-coenzyme A reductase inhibitors induce apoptosis in human cardiac myocytes in vitro, 1324
- Cardiomyocytes**
Schisandrin B enhances doxorubicin-induced apoptosis of cancer cells but not normal cells, 584
- Caspase**
Schisandrin B enhances doxorubicin-induced apoptosis of cancer cells but not normal cells, 584
Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742
- Caspase-10**
Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
- Caspase-8**
Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
- Caspases**
Cell cycle arrest and proapoptotic effects of the anticancer cyclodepsipeptide serratomolide (AT514) are independent of p53 status in breast cancer cells, 32
- Catecholamine**
 N -(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- Cathepsin G**
Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287
- CBI**
Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683
- CCR5**
Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- CD36**
Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605
- Cdc42**
The Cdc42 inhibitor secramine B prevents cAMP-induced K^+ conductance in intestinal epithelial cells, 1720
- CDK**
Anti-Tat and anti-HIV activities of trimers of n -alkylglycines, 596
- Cdx2**
The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581
- Cell cycle**
Molecular targets of dietary agents for prevention and therapy of cancer, 1397

- Blockade of IGF-1 receptor tyrosine kinase has antineoplastic effects in hepatocellular carcinoma cells, 1435
- Cell cycle arrest
- Cell cycle arrest and proapoptotic effects of the anticancer cyclodepsipeptide serratamolide (AT514) are independent of p53 status in breast cancer cells, 32
- Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24), 426
- Cell differentiation
- Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219
- Cell motility
- Relative effects of phenolic constituents from *Yucca schidigera* Roehl. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479
- Cell sensitivity
- Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791
- Cell wall antibiotics
- Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996
- Cell-penetrating peptide
- Overcoming methotrexate resistance in breast cancer tumour cells by the use of a new cell-penetrating peptide, 416
- Cellular iron metabolism
- Mobilization of iron from cells by hydroxyquinoline-based chelators, 214
- Ceramide
- Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by *N*-lauroyl-D-erythro-sphingosine and *N*-lauroyl-D-erythro-C₂₀-sphingosine, 1229
- Ceramide-1-phosphate
- Ceramide-1-phosphate activates cytosolic phospholipase A2 α directly and by PKC pathway, 850
- C-FLIP
- Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
- c-Fos
- Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818
- Chalcones
- Inhibition of TNF α -induced activation of nuclear factor κ B by kava (*Piper methysticum*) derivatives, 1206
- Chemical modification
- Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629
- Chemoattractant receptors
- The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488
- Chemokine
- Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- Chemosensitivity
- Methionine dependence of tumours: A biochemical strategy for optimizing paclitaxel chemosensitivity in vitro, 772
- Chemotherapeutic agents
- Epidermal growth factor receptor inhibitor (PD168393) potentiates cytotoxic effects of paclitaxel against androgen-independent prostate cancer cells, 751
- Chemotherapy
- 6-Benzylthioinosine analogues: Promising anti-toxoplasmic agents as inhibitors of the mammalian nucleoside transporter ENT1 (es), 69
- Role of GADD34 in modulation of cisplatin cytotoxicity, 239
- Contribution of an unidentified sodium-dependent nucleoside transport system to the uptake and cytotoxicity of anthracycline in mouse M5076 ovarian sarcoma cells, 565
- Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136
- Chimeric receptor
- Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294
- Chlamydia pneumoniae*
- Inhibitory effect of dietary phenolic compounds on *Chlamydia pneumoniae* in cell cultures, 735
- Chloride secretion
- The Cdc42 inhibitor secramine B prevents cAMP-induced K⁺ conductance in intestinal epithelial cells, 1720
- Chloroquine
- NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1
- Cholesterol
- Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453
- Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605
- Cholesterol metabolism
- Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114
- Cholesteryl esters
- Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114
- Cholinergic system
- Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497
- Chromaffin cells
- N*-(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- Chronic lymphocytic leukemia
- The pattern of deoxycytidine- and deoxyguanosine kinase activity in relation to messenger RNA expression in blood cells from untreated patients with B-cell chronic lymphocytic leukemia, 882
- Chronopharmacotherapy
- Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155
- Cigarette smoke
- Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, 1602
- Circadian rhythm
- Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155

Cisplatin

Differential response of normal, dedifferentiated and transformed thyroid cell lines to cisplatin treatment, 50
Role of GADD34 in modulation of cisplatin cytotoxicity, 239

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

cis-Platin

Suppression of survival in human SKBR3 breast carcinoma in response to metal-chelator complexes is preferential for copper-dithiocarbamate, 722

Cisplatin resistance

MZF1 possesses a repressively regulatory function in ERCC1 expression, 761

cJun N-terminal kinase

Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268

Cl⁻ channel

Serotonin induces pulmonary artery smooth muscle cell migration, 386

Class 1 integrins

The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073

Clearance prediction

In vitro-in vivo correlation for drugs and other compounds eliminated by glucuronidation in humans: Pitfalls and promises, 1531

Clinical trial

Natural products — The future scaffolds for novel antibiotics?, 919

c-Myc

Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248

Cognition

The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103

Co(III) complex

Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683

Colchicine

Fibroblast heterogeneity in collagenolytic response to colchicine, 574

Collagenase

Fibroblast heterogeneity in collagenolytic response to colchicine, 574

Colon cancer

The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540

Combinatorial libraries

Anti-Tat and anti-HIV activities of trimers of *n*-alkylglycines, 596

Commensal

Commensals upon us, 893

Computational chemistry

Structure-based drug design meets the ribosome, 1016

Conformational induction

Conformational induction is the key process for activation of the AT₁ receptor, 464

Conformational selection

Conformational induction is the key process for activation of the AT₁ receptor, 464

Constitutive activity

Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663

Constitutively active

Conformational induction is the key process for activation of the AT₁ receptor, 464

Constraint-based modeling

Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026

CORM-2

CO from enhanced HO activity or from CORM-2 inhibits both O₂⁻ and NO production and downregulates HO-1 expression in LPS-stimulated macrophages, 307

Corticosterone

Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155

CPT-11

Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155

Cyanidin

Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184

Cyclic AMP

Niacin induces PPAR γ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646
Agonist potency differentiates G protein activation and Ca²⁺ signalling by the orexin receptor type 1, 827

Cyclooxygenase

Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂- α , 144

Cyclooxygenase-2

A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223

CYP19/aromatase

The effect of methadone and buprenorphine on human placental aromatase, 1255

CYP1A1

The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366

CYP1A2

The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366

Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377

CYP2C8

Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377

CYP2D6

Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377

Catalytic roles of CYP2D6.10 and CYP2D6.36 enzymes in mexiletine metabolism: In vitro functional analysis of recombinant proteins expressed in *Saccharomyces cerevisiae*, 1386

CYP2D6*10

Catalytic roles of CYP2D6.10 and CYP2D6.36 enzymes in mexiletine metabolism: In vitro functional analysis of recombinant proteins expressed in *Saccharomyces cerevisiae*, 1386

CYP2D6*36

Catalytic roles of CYP2D6.10 and CYP2D6.36 enzymes in mexiletine metabolism: In vitro functional analysis of recombinant proteins expressed in *Saccharomyces cerevisiae*, 1386

- CYP3A4**
Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377
- CYP3A6**
The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366
- Cysteine**
Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629
- Cysteinyl-glycine**
Expression of γ -glutamyltransferase in cancer cells and its significance in drug resistance, 231
- Cytochrome P450**
Oxygen-insensitive enzymatic reduction of oximes to imines, 354
The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366
- Cytomegalovirus**
Pyridine *N*-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122
- Cytoskeleton**
Serotonin induces pulmonary artery smooth muscle cell migration, 386
- Cytosolic phospholipase A2 α**
Ceramide-1-phosphate activates cytosolic phospholipase A2 α directly and by PKC pathway, 850
- Cytostatics**
Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268
Blockade of IGF-1 receptor tyrosine kinase has anti-neoplastic effects in hepatocellular carcinoma cells, 1435
- Cytotoxicity**
Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136
Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289
Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610
- DARPP-32**
A role for AMPA receptors in mood disorders, 1273
- de novo resistance**
Contribution of an unidentified sodium-dependent nucleoside transport system to the uptake and cytotoxicity of anthracycline in mouse M5076 ovarian sarcoma cells, 565
- Death receptors**
Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
- Delbavancin**
Glycopeptides: Update on an old successful antibiotic class, 968
- 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂**
Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A₂ receptor gene in human erythroleukemia cells, 1308
- Deoxycytidine kinase**
The pattern of deoxycytidine- and deoxyguanosine kinase activity in relation to messenger RNA expression in blood cells from untreated patients with B-cell chronic lymphocytic leukemia, 882
- Deoxyguanosine kinase**
The pattern of deoxycytidine- and deoxyguanosine kinase activity in relation to messenger RNA expression in blood cells from untreated patients with B-cell chronic lymphocytic leukemia, 882
- Depression**
A role for AMPA receptors in mood disorders, 1273
- Depsipeptide**
Cell cycle arrest and proapoptotic effects of the anticancer cyclodepsipeptide serratamolide (AT514) are independent of p53 status in breast cancer cells, 32
- Development**
The role of pharmacodynamic research in the assessment and development of new antibacterial drugs, 1057
- Dexamethasone**
Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693
- Diabetes mellitus**
Cross-talk between angiotensin II and glucagon receptor signaling mediates phosphorylation of mitogen-activated protein kinases ERK 1/2 in rat glomerular mesangial cells, 1711
- Diabetic plasma**
Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338
- Dibenzodiazepine**
Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98
- Dichloroacetate**
Evaluation of dichloroacetate treatment in a murine model of hereditary tyrosinemia type 1, 1648
- Dietary agents**
Molecular targets of dietary agents for prevention and therapy of cancer, 1397
- Diethyl pyrocarbonate (DEPC)**
Diethyl pyrocarbonate, a histidine-modifying agent, directly stimulates activity of ATP-sensitive potassium channels in pituitary GH₃ cells, 615
- Digoxin transport**
Effects of capsaicin on P-gp function and expression in Caco-2 cells, 1727
- Dithiothreitol**
Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562
- DMPP**
N-(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- DNA microarray**
Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184
- DNA minor groove alkylator**
Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683
- DNA repair**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- DNA-PK**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- Docetaxel**
Epidermal growth factor receptor inhibitor (PD168393) potentiates cytotoxic effects of paclitaxel against androgen-independent prostate cancer cells, 751

Double-stranded RNA

Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum, 702

Doxorubicin

Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248

Schisandrin B enhances doxorubicin-induced apoptosis of cancer cells but not normal cells, 584

Mitochondrial localization and activity of P-glycoprotein in doxorubicin-resistant K562 cells, 1162

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

Drimanial

Pharmacological characterisation of the plant sesquiterpenes polygodial and drimanial as vanilloid receptor agonists, 1248

Drug delivery

Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248

Overcoming methotrexate resistance in breast cancer tumour cells by the use of a new cell-penetrating peptide, 416

Drug discovery

Antibacterial drug discovery—Then, now and the genomics future, 901

The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103

Drug metabolism

Metabolism of nilutamide in rat lung, 377

Drug resistance

Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61

Increased lysosomal uptake of methotrexate-polyglutamates in two methotrexate-resistant cell lines with distinct mechanisms of resistance, 203

Overcoming methotrexate resistance in breast cancer tumour cells by the use of a new cell-penetrating peptide, 416

Drug response

Fibroblast heterogeneity in collagenolytic response to colchicine, 574

Drug transport

Increased lysosomal uptake of methotrexate-polyglutamates in two methotrexate-resistant cell lines with distinct mechanisms of resistance, 203

Drug-binding site

Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338

Early growth response gene (Egr-1)

Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806

Efflux

Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910

Efflux pump inhibitor

Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949

Efflux pumps

Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949

Unmet medical needs in antibacterial therapy, 991

EGFR

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

Blockade of IGF-1 receptor tyrosine kinase has anti-neoplastic effects in hepatocellular carcinoma cells, 1435

EGFR receptor

Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422

Eicosanoid

Involvement of tumor necrosis factor (TNF)- α in phorbol ester 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin edema in mice, 1331

Elastase

Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287

Elastin-like polypeptide

Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248

Empiric screening

Empirical antibacterial drug discovery—Foundation in natural products, 1006

Endothelin-1

Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175

Energy metabolism

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530

EPIs

Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910

Epithelial integrity

Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289

Epoxyquinoid

Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634

Epoxyquinone A monomer

Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634

Equol

Estrogen receptor-independent inhibition of tumor necrosis factor- α gene expression by phytoestrogen equol is mediated by blocking nuclear factor- κ B activation in mouse macrophages, 136

ERCC1

MZF1 possesses a repressively regulatory function in ERCC1 expression, 761

ERK

Differential response of normal, dedifferentiated and transformed thyroid cell lines to cisplatin treatment, 50

14-3-3 ζ interacts with human thromboxane receptors and is involved in the agonist-induced activation of the extracellular-signal-regulated kinase, 624

The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540

- ERK1/2
Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y₁ receptor in a Ca²⁺ independent manner in differentiated human skeletal muscle cells, 1497
- Erythrocyte
Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779
Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799
- ESBLs
The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073
- Essential gene
Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026
- Essentiality
Use of genomics to select antibacterial targets, 1066
- Estrogen
Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAR cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349
- Estrogen formation
The effect of methadone and buprenorphine on human placental aromatase, 1255
- Estrogen receptors
Isoflavone genistein and daidzein up-regulate LPS-induced inducible nitric oxide synthase activity through estrogen receptor pathway in RAW264.7 cells, 108
Quaternary ammonium-linked glucuronidation of *trans*-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358
Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta, 1459
- Ethnopharmacology
Inhibition of TNF α -induced activation of nuclear factor κ B by kava (*Piper methysticum*) derivatives, 1206
- Etoposide
Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248
- Extended ternary complex model
Conformational induction is the key process for activation of the AT₁ receptor, 464
- Extracellular matrix
Inhibitory effect of DA-125, a new anthracyclin analog antitumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21
- Extracellular matrix proteins
ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479
- Extracellular signal-regulated kinase (ERK)
Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806
- Fermentation
Empirical antibacterial drug discovery—Foundation in natural products, 1006
- Ferrochelatase
Mechanisms involved in δ -aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: Relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin, 42
- Fibroblast heterogeneity
Fibroblast heterogeneity in collagenolytic response to colchicine, 574
- Filipin
Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779
- Flavonoids
Inhibitory effect of dietary phenolic compounds on *Chlamydia pneumoniae* in cell cultures, 735
Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570
- Fluoroquinolones
Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910
- Flux balance analysis
Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026
- Formyl peptide receptors
The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488
- Forsythia koreana*
Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198
- Foxy
Oxidative metabolism of 5-methoxy-N,N-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377
- Fumarylacetoacetate hydrolase
Evaluation of dichloroacetate treatment in a murine model of hereditary tyrosinemia type 1, 1648
- Functional activity
Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521
- Fungal metabolite
Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634
- Fungi
The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981
- Fura-2
Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173
- G protein
Agonist potency differentiates G protein activation and Ca²⁺ signalling by the orexin receptor type 1, 827
Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449
- G protein-coupled receptor
Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294
Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y₂ and P2Y₄ receptors, 540

GABA

Inhibition of rabbit brain 4-aminobutyrate transaminase by some taurine analogues: A kinetic analysis, 1510

γ -Aminobutyric acid (GABA)-C receptor stimulation increases prolactin (PRL) secretion in cultured rat anterior pituitary cells, 1705

GABA_B receptor

Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA_B receptor expression and function in an animal model of neuropathic pain, 196

GABA_B receptor subunits

Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA_B receptor expression and function in an animal model of neuropathic pain, 196

GABA-C receptor

γ -Aminobutyric acid (GABA)-C receptor stimulation increases prolactin (PRL) secretion in cultured rat anterior pituitary cells, 1705

GADD34

Role of GADD34 in modulation of cisplatin cytotoxicity, 239

Gamma-glutamyltransferase expression

Expression of γ -glutamyltransferase in cancer cells and its significance in drug resistance, 231

Gastrointestinal tract infections

Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289

[Gd@C₈₂(OH)₂₂]_n nanoparticles

Antioxidative function and biodistribution of [Gd@C₈₂(OH)₂₂]_n nanoparticles in tumor-bearing mice, 872

Gene expression

Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497

Gene expression profile

Microarray profiling of gene expression in human adipocytes in response to anthocyanins, 1184

Genetic polymorphism

Catalytic roles of CYP2D6.10 and CYP2D6.36 enzymes in mexiletine metabolism: In vitro functional analysis of recombinant proteins expressed in *Saccharomyces cerevisiae*, 1386

Genomic and nongenomic glucocorticoid effects

Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530

Genomics

Antibacterial drug discovery—Then, now and the genomics future, 901

Use of genomics to select antibacterial targets, 1066

When will the genomics investment pay off for antibacterial discovery?, 1096

GH₃ cells

Diethyl pyrocarbonate, a histidine-modifying agent, directly stimulates activity of ATP-sensitive potassium channels in pituitary GH₃ cells, 615

Gimatecan

Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791

GLU_{A1}

A role for AMPA receptors in mood disorders, 1273

Glucagon

Cross-talk between angiotensin II and glucagon receptor signaling mediates phosphorylation of mitogen-activated protein kinases ERK 1/2 in rat glomerular mesangial cells, 1711

Glucocorticoid receptor

Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693

Glucuronidation

In vitro–in vivo correlation for drugs and other compounds eliminated by glucuronidation in humans: Pitfalls and promises, 1531

Glutathione

Expression of γ -glutamyltransferase in cancer cells and its significance in drug resistance, 231

Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257

Redox modifications of protein–thiols: Emerging roles in cell signaling, 551

Suppression of survival in human SKBR3 breast carcinoma in response to metal–chelator complexes is preferential for copper–dithiocarbamate, 722

Several glutathione S-transferase isozymes that protect against oxidative injury are expressed in human liver mitochondria, 1619

Glutathione conjugate

Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268

Glutathione disulfide

Metabolism of oxidants by blood from different mouse strains, 1753

Glutathione S-transferase

Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268

Several glutathione S-transferase isozymes that protect against oxidative injury are expressed in human liver mitochondria, 1619

Glutathione transferase

Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629

Glutathionylation

Metabolism of oxidants by blood from different mouse strains, 1753

Glycerol 3-phosphate shuttle

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Glycopeptide

Glycopeptides: Update on an old successful antibiotic class, 968

Glyoxal

Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610

Gonadotropins

The effects of prostaglandin F_{2 α} treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

GPCR

Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319

The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103

- G-protein**
Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- GPRK**
Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219
- Growth hormone**
Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470
- GSK3**
AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637
- H₂DCFDA**
Suppression of survival in human SKBR3 breast carcinoma in response to metal-chelator complexes is preferential for copper-dithiocarbamate, 722
- H₃ receptor**
The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103
- Haemoglobin sulfhydryl groups**
Metabolism of oxidants by blood from different mouse strains, 1753
- Haemolysis**
Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779
- Heart failure**
Superoxide radical production by allopurinol and xanthine oxidase, 1747
- HEK-293 cells**
Block of hERG channel by ziprasidone: Biophysical properties and molecular determinants, 278
Agonist potency differentiates G protein activation and Ca²⁺ signalling by the orexin receptor type 1, 827
- Heme oxygenase**
CO from enhanced HO activity or from CORM-2 inhibits both O₂⁻ and NO production and downregulates HO-1 expression in LPS-stimulated macrophages, 307
- Hemin**
Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799
- Hemolysis**
Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799
- Heparin**
Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287
- Hepatocellular carcinoma**
ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479
- Hepatocyte nuclear factor**
Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693
- Hepatocytes**
The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366
- Hepatotoxicity**
Anti-tumour activity in non-small cell lung cancer models and toxicity profiles for novel ruthenium(II) based organo-metallic compounds, 408
- Hereditary tyrosinemia type 1**
Evaluation of dichloroacetate treatment in a murine model of hereditary tyrosinemia type 1, 1648
- hERG**
Block of hERG channel by ziprasidone: Biophysical properties and molecular determinants, 278
- Hexadecylphosphocholine**
Modulation of intestinal barrier properties by miltefosine, 486
- High density lipoprotein**
Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453
- Histamine**
The challenge of drug discovery of a GPCR target: Analysis of preclinical pharmacology of histamine H₃ antagonists/inverse agonists, 1103
- Histamine receptors**
Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219
- Histamine-trifluoromethyltoluide**
N-(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- Histone H2AX**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- HIV-1**
Anti-Tat and anti-HIV activities of trimers of *n*-alkylglycines, 596
- HM74a**
Niacin induces PPAR γ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646
- Homocysteine**
Methionine dependence of tumours: A biochemical strategy for optimizing paclitaxel chemosensitivity in vitro, 772
- Homology modeling**
Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y₂ and P2Y₄ receptors, 540
- Hormonal regulation**
Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470
- Hormone refractory**
Epidermal growth factor receptor inhibitor (PD168393) potentiates cytotoxic effects of paclitaxel against androgen-independent prostate cancer cells, 751
- Hospital-acquired gram-negative rods**
Surveillance uncovers the smoking gun for resistance emergence, 1036
- Hot-plate test**
Synthesis and antinociceptive activity of cyclic endomorphin-2 and morphiceptin analogs, 188
- HT1080 cell**
Inhibitory effect of DA-125, a new anthracyclin analog anti-tumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21
- Human immunodeficiency virus**
Pyridine *N*-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122

Human liver

Several glutathione S-transferase isozymes that protect against oxidative injury are expressed in human liver mitochondria, 1619

Human liver microsomes

Quaternary ammonium-linked glucuronidation of *trans*-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358

Human metabolism

Hydrolytic metabolism of pyrethroids by human and other mammalian carboxylesterases, 657

Human neoplasia

Expression of γ -glutamyltransferase in cancer cells and its significance in drug resistance, 231

Human peripheral blood mononuclear cells

Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530

Human placenta

The effect of methadone and buprenorphine on human placental aromatase, 1255

Human serum albumin

Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338

Human skeletal muscle cells

Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y₁ receptor in a Ca²⁺ independent manner in differentiated human skeletal muscle cells, 1497

Hydrogen peroxide

Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610

Hydrolysable tannin

Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570

Hydrophilic

Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand–receptor interaction, 865

Hydrophobic

Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand–receptor interaction, 865

Hydroxyl radical

Hydroxyl radical scavenging reactivity of proton pump inhibitors, 1337

Hydroxyl radical scavenger

Hydroxyl radical scavenging reactivity of proton pump inhibitors, 1337

4-Hydroxytamoxifen

Quaternary ammonium-linked glucuronidation of *trans*-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358

5-HT

Serotonin induces pulmonary artery smooth muscle cell migration, 386

5-HT1B/1D receptor

Serotonin induces pulmonary artery smooth muscle cell migration, 386

5-HT2 receptors

Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521

5-HT2C isoforms

Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521

5-HT4 receptor

Serotonin induces pulmonary artery smooth muscle cell migration, 386

5-HTT

Serotonin induces pulmonary artery smooth muscle cell migration, 386

17 β -Hydroxysteroid dehydrogenase type I (17 β -HSD I)

Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAR cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349

Hypertrophy

Hydroxymethylglutaryl-coenzyme A reductase inhibitors induce apoptosis in human cardiac myocytes in vitro, 1324

IFN-gamma

Inhibition of sphingolipid synthesis impairs cellular activation, cytokine production and proliferation in human lymphocytes, 126

IGF-1 receptor

Blockade of IGF-1 receptor tyrosine kinase has antineoplastic effects in hepatocellular carcinoma cells, 1435

IkappaB

Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634

IkappaB kinase

Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634

IL-6

Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530

Immunity

Commensals upon us, 893

In silico modeling

Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026

In vitro–in vivo correlation

In vitro–in vivo correlation for drugs and other compounds eliminated by glucuronidation in humans: Pitfalls and promises, 1531

In vivo

Antioxidative function and biodistribution of [Gd@C₈₂(OH)₂₂]_n nanoparticles in tumor-bearing mice, 872

Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663

Inducible nitric oxide

Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198

Infection

Commensals upon us, 893

Inflammation

The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366

Relative effects of phenolic constituents from *Yucca schidigera* Roez. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479

Inhibition

Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287

Inhibition of acid secretion

Characterization of the inhibitory activity of tenatoprazole on the gastric H^+ , K^+ -ATPase in vitro and in vivo, 837

Inhibitors

Characterization of the melatoninergic MT_3 binding site on the NRH:quinone oxidoreductase 2 enzyme, 74

Binding kinetics and duration of in vivo action of novel prolyl oligopeptidase inhibitors, 683

Understanding the longevity of the β -lactam antibiotics and of antibiotic/ β -lactamase inhibitor combinations, 930

iNOS

Isoflavone genistein and daidzein up-regulate LPS-induced inducible nitric oxide synthase activity through estrogen receptor pathway in RAW264.7 cells, 108

Interaction

Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287

Interleukin-4

Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by N-lauroyl-D-erythro-sphingosine and N-lauroyl-D-erythro- C_{20} -sphingosine, 1229

Intestinal absorption

Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570

Intestinal epithelial cells

Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289

Intestinal metaplasia

The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581

Intracellular iron chelators

Mobilization of iron from cells by hydroxyquinoline-based chelators, 214

Intracellular P-gp

Mitochondrial localization and activity of P-glycoprotein in doxorubicin-resistant K562 cells, 1162

Invasion

Inhibitory effect of DA-125, a new anthracyclin analog antitumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21

ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479

Inverse agonist

Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663

Ionising radiation

Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683

Iron chelator toxicity

Mobilization of iron from cells by hydroxyquinoline-based chelators, 214

Ischemia-reperfusion

Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia-reperfusion, 441

Isoflavone

Isoflavone genistein and daidzein up-regulate LPS-induced inducible nitric oxide synthase activity through estrogen receptor pathway in RAW264.7 cells, 108

Isoforms

Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A_2 receptor gene in human erythroleukemia cells, 1308

Isoform-specific residues

The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299

IVPM

The role of pharmacodynamic research in the assessment and development of new antibacterial drugs, 1057

JAK/STAT pathway

JAK/STAT signal transduction: Regulators and implication in hematological malignancies, 713

Jurkat cells

Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562

K562

Inhibition of $TNF\alpha$ -induced activation of nuclear factor κB by kava (*Piper methysticum*) derivatives, 1206

Kaposi's sarcoma

Relative effects of phenolic constituents from *Yucca schidigera* Roetzl. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479

 K_{ATP} -channel

Diethyl pyrocarbonate, a histidine-modifying agent, directly stimulates activity of ATP-sensitive potassium channels in pituitary GH_3 cells, 615

Kava

Inhibition of $TNF\alpha$ -induced activation of nuclear factor κB by kava (*Piper methysticum*) derivatives, 1206

KCNQ1/KCNE3

The Cdc42 inhibitor secramine B prevents cAMP-induced K^+ conductance in intestinal epithelial cells, 1720

Kinase

Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449

L929 cells

Ceramide-1-phosphate activates cytosolic phospholipase $A2\alpha$ directly and by PKC pathway, 850

 β -Lactam

Understanding the longevity of the β -lactam antibiotics and of antibiotic/ β -lactamase inhibitor combinations, 930

 β -Lactamase

Understanding the longevity of the β -lactam antibiotics and of antibiotic/ β -lactamase inhibitor combinations, 930

Lactones

Inhibition of $TNF\alpha$ -induced activation of nuclear factor κB by kava (*Piper methysticum*) derivatives, 1206

Laminin-5

ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479

Lanthanum

Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24), 426

Lead discovery

The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981

Leukaemia

Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257

Ligand

Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98
Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta, 1459

Ligand-macromolecules interaction

Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858

Lipopolysaccharide

A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223

Lipoxygenase

Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂-α, 144

Liver mitochondria

3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342

Local anesthetic drug

The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299

Locomotor activity

Synthesis and antinociceptive activity of cyclic endomorphin-2 and morphiceptin analogs, 188

LTR

Anti-Tat and anti-HIV activities of trimers of *n*-alkylglycines, 596

Lung cancer

Anti-tumour activity in non-small cell lung cancer models and toxicity profiles for novel ruthenium(II) based organo-metallic compounds, 408

Lung carcinoma

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

Lysosomes

Increased lysosomal uptake of methotrexate-polyglutamates in two methotrexate-resistant cell lines with distinct mechanisms of resistance, 203
Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136

M5076 cell

Contribution of an unidentified sodium-dependent nucleoside transport system to the uptake and cytotoxicity of anthracycline in mouse M5076 ovarian sarcoma cells, 565

Macrophage

Isoflavone genistein and daidzein up-regulate LPS-induced inducible nitric oxide synthase activity through estrogen receptor pathway in RAW264.7 cells, 108
Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257
Different roles of liver X receptor α and β in lipid metabolism: Effects of an α-selective and a dual agonist in mice deficient in each subtype, 453
Niacin induces PPARγ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646

Malaria

Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61

Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779

Malate dehydrogenase

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Malate-α-ketoglutarate carrier

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Malate-aspartate shuttle

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Maleylacetoacetate isomerase

Evaluation of dichloroacetate treatment in a murine model of hereditary tyrosinemia type 1, 1648

MAP kinase (ERK1/2)

Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240

MAP kinases

Molecular targets of dietary agents for prevention and therapy of cancer, 1397

MAPK

Serotonin induces pulmonary artery smooth muscle cell migration, 386

Maraviroc

Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163

Mas

Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319

Matrix metalloproteinase

Inhibitory effect of DA-125, a new anthracycline analog anti-tumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21

Matrix metalloproteinases

Fibroblast heterogeneity in collagenolytic response to colchicine, 574

MDA-MB-231 cells

Overcoming methotrexate resistance in breast cancer tumour cells by the use of a new cell-penetrating peptide, 416

MDR1

Effects of capsaicin on P-gp function and expression in Caco-2 cells, 1727

Mechanism

Understanding the longevity of the β-lactam antibiotics and of antibiotic/β-lactamase inhibitor combinations, 930
Glycopeptides: Update on an old successful antibiotic class, 968

Mechanism-of-action

Empirical antibacterial drug discovery—Foundation in natural products, 1006

MEK1/2 (mitogen-activated protein kinase kinase 1/2)

Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806

Melatonin

NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1
Characterization of the melatonergic MT₃ binding site on the NRH:quinone oxidoreductase 2 enzyme, 74

- Membrane fluidity**
Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799
- Membrane microdomains**
Inhibition of sphingolipid synthesis impairs cellular activation, cytokine production and proliferation in human lymphocytes, 126
- Membrane stabilizer**
Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799
- 5-MeO-DIPT**
Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377
- 5-MeO-IPT**
Oxidative metabolism of 5-methoxy-*N,N*-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377
- Mesangial cells**
Cross-talk between angiotensin II and glucagon receptor signaling mediates phosphorylation of mitogen-activated protein kinases ERK 1/2 in rat glomerular mesangial cells, 1711
- Metabolism**
Use of constraint-based modeling for the prediction and validation of antimicrobial targets, 1026
- Metalloenzyme**
The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042
- Methadone**
The effect of methadone and buprenorphine on human placental aromatase, 1255
- Methicillin-resistant *Staphylococcus aureus***
Surveillance uncovers the smoking gun for resistance emergence, 1036
- Methionine-dependence**
Methionine dependence of tumours: A biochemical strategy for optimizing paclitaxel chemosensitivity in vitro, 772
- Methotrexate**
Increased lysosomal uptake of methotrexate-polyglutamates in two methotrexate-resistant cell lines with distinct mechanisms of resistance, 203
Overcoming methotrexate resistance in breast cancer tumour cells by the use of a new cell-penetrating peptide, 416
- Methyl-adenosine**
Structure–activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671
- Mexiletine**
Catalytic roles of CYP2D6.10 and CYP2D6.36 enzymes in mexiletine metabolism: In vitro functional analysis of recombinant proteins expressed in *Saccharomyces cerevisiae*, 1386
- Microarray**
Role of GADD34 in modulation of cisplatin cytotoxicity, 239
Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497
- Microphysiometer**
Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422
- Microsomes**
Oxygen-insensitive enzymatic reduction of oximes to imines, 354
- Migration**
Serotonin induces pulmonary artery smooth muscle cell migration, 386
- Miltefosine**
Modulation of intestinal barrier properties by miltefosine, 486
- MIP-1 β**
Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- Mitochondria**
Oxygen-insensitive enzymatic reduction of oximes to imines, 354
Mitochondrial localization and activity of P-glycoprotein in doxorubicin-resistant K562 cells, 1162
Distribution of tocopheryl quinone in mitochondrial membranes and interference with ubiquinone-mediated electron transfer, 1589
Several glutathione S-transferase isozymes that protect against oxidative injury are expressed in human liver mitochondria, 1619
- Mitochondrial toxicity**
3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342
- Modulation**
Role of GADD34 in modulation of cisplatin cytotoxicity, 239
- Molecular docking**
Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- Molecular pharmacology**
NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1
Characterization of the melatonergic MT₃ binding site on the NRH:quinone oxidoreductase 2 enzyme, 74
- Monocytes**
Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605
Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818
- Mono-Mac 6 cells**
Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562
- Montelukast**
CysLT₁ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115
- Mouse**
Involvement of tumor necrosis factor (TNF)- α in phorbol ester 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin edema in mice, 1331
- Mouse N2a neuroblastoma**
Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240
- Mouse strains**
Metabolism of oxidants by blood from different mouse strains, 1753
- Mrp**
Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470
- MRS2179**
Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y₁ receptor in a Ca²⁺ independent manner in differentiated human skeletal muscle cells, 1497

MRSA

Commensals upon us, 893

MT₃

Characterization of the melatonergic MT₃ binding site on the NRH:quinone oxidoreductase 2 enzyme, 74

Multidrug resistance

Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268

Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910

The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146

Mitochondrial localization and activity of P-glycoprotein in doxorubicin-resistant K562 cells, 1162

Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470

Multidrug resistant *Klebsiella pneumoniae*

The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073

Mur pathway

Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996

Mutagenesis

Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand–receptor interaction, 865

Use of genomics to select antibacterial targets, 1066

Mycobacterium tuberculosis

Surveillance uncovers the smoking gun for resistance emergence, 1036

Structure–activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671

Myeloma

Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742

Myocardial energy metabolism

Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia–reperfusion, 441

Myocardium

Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175

MZF1

MZF1 possesses a repressively regulatory function in ERCC1 expression, 761

N⁶-Cyclopropyl-2,6-diamino-9-[2-(phosphonomethoxy)ethyl]purine

N⁶-Methyl-AMP aminohydrolase activates N⁶-substituted purine acyclic nucleoside phosphonates, 1370

N⁶-Methyl-AMP

N⁶-Methyl-AMP aminohydrolase activates N⁶-substituted purine acyclic nucleoside phosphonates, 1370

N⁶-Methyl-AMP aminohydrolase

N⁶-Methyl-AMP aminohydrolase activates N⁶-substituted purine acyclic nucleoside phosphonates, 1370

NADPH-oxidase

The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488

Natural products

Natural products — The future scaffolds for novel antibiotics?, 919

The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981

Empirical antibacterial drug discovery—Foundation in natural products, 1006

NER

MZF1 possesses a repressively regulatory function in ERCC1 expression, 761

Neurofilament heavy chain

Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240

Neuronal and astrocytic shuttle mechanisms

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Neurons

Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399

Neuropathic pain

Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA_B receptor expression and function in an animal model of neuropathic pain, 196

Neuropathy target esterase

Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240

Neutrophils

The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488

Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735

New targets

Empirical antibacterial drug discovery—Foundation in natural products, 1006

NF- κ B

Inhibition of transcription factor NF- κ B signaling proteins IKK β and p65 through specific cysteine residues by epoxyquinone A monomer: Correlation with its anti-cancer cell growth activity, 634

NF- κ B

Inhibitory effect of DA-125, a new anthracycline analog antitumor agent, on the invasion of human fibrosarcoma cells by down-regulating the matrix metalloproteinases, 21

Estrogen receptor-independent inhibition of tumor necrosis factor- α gene expression by phytoestrogen equol is mediated by blocking nuclear factor- κ B activation in mouse macrophages, 136

The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366

Redox modifications of protein–thiols: Emerging roles in cell signaling, 551

Pyridine N-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122

- Inhibition of TNF α -induced activation of nuclear factor κ B by kava (*Piper methysticum*) derivatives, 1206
- Molecular targets of dietary agents for prevention and therapy of cancer, 1397
- Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, 1602
- NGFI-B**
- Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98
- N-Glucuronidation**
- Quaternary ammonium-linked glucuronidation of *trans*-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358
- Niacin**
- Niacin induces PPAR γ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646
- Nicotinic acetylcholine receptor**
- N-(4-Trifluoromethylphenyl)amide group of the synthetic histamine receptor agonist inhibits nicotinic acetylcholine receptor-mediated catecholamine secretion, 670
- Nilutamide**
- Metabolism of nilutamide in rat lung, 377
- Nitric oxide**
- CO from enhanced HO activity or from CORM-2 inhibits both O $_2^-$ and NO production and downregulates HO-1 expression in LPS-stimulated macrophages, 307
- Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198
- Nitric oxide synthase**
- Metabolism of nilutamide in rat lung, 377
- Nitroreduction**
- Metabolism of nilutamide in rat lung, 377
- 1 H NMR**
- Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858
- Nociception**
- Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA $_B$ receptor expression and function in an animal model of neuropathic pain, 196
- Nociceptor**
- Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173
- Noxious heat**
- Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173
- 3-NPA**
- Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- N-reduction**
- Oxygen-insensitive enzymatic reduction of oximes to imines, 354
- NTBC**
- Evaluation of dichloroacetate treatment in a murine model of hereditary tyrosinemia type 1, 1648
- Nuclear factor of activated T cell**
- Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by *N*-lauroyl-D-erythro-sphingosine and *N*-lauroyl-D-erythro-C $_{20}$ -sphingosine, 1229
- Nuclear factor- κ B**
- Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198
- Nuclear receptor**
- Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98
- Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453
- Nucleoside 5'-[β , γ -imido]triphosphate**
- Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- Nucleoside 5'-[γ -thio]triphosphate**
- Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- Nucleoside 5'-triphosphate**
- Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- nucleoside analogs**
- Structure-activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671
- Nucleoside transport inhibitors**
- 6-Benzylthioinosine analogues: Promising anti-toxoplasmic agents as inhibitors of the mammalian nucleoside transporter ENT1 (es), 69
- Nucleoside transport system**
- Contribution of an unidentified sodium-dependent nucleoside transport system to the uptake and cytotoxicity of anthracycline in mouse M5076 ovarian sarcoma cells, 565
- Nucleotides**
- CysLT $_1$ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115
- Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y $_2$ and P2Y $_4$ receptors, 540
- Nystatin**
- Selective lysis of erythrocytes infected with the trophozoite stage of *Plasmodium falciparum* by polyene macrolide antibiotics, 779
- 5-OH-DIPT**
- Oxidative metabolism of 5-methoxy-N,N-diisopropyl tryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes, 1377
- Omeprazole**
- Characterization of the inhibitory activity of tenatoprazole on the gastric H $^+$,K $^+$ -ATPase in vitro and in vivo, 837
- Oncogene**
- Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319
- Oncogenesis**
- JAK/STAT signal transduction: Regulators and implication in hematological malignancies, 713
- μ -, δ - and κ -opioid receptors**
- Synthesis and antinociceptive activity of cyclic endomorphin-2 and morphiceptin analogs, 188

Oral bioavailability

Mobilization of iron from cells by hydroxyquinoline-based chelators, 214

Orexin receptor

Agonist potency differentiates G protein activation and Ca^{2+} signalling by the orexin receptor type 1, 827

Organophosphates

Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497

Organotin

Organotin compounds enhance 17β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAr cells: Potential promotion of 17β -estradiol biosynthesis in human placenta, 1349

Osteoblast

Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818

Osteoporosis

Estrogen receptor-independent inhibition of tumor necrosis factor- α gene expression by phytoestrogen equol is mediated by blocking nuclear factor- κ B activation in mouse macrophages, 136

Ovary

The effects of prostaglandin $\text{F}_{2\alpha}$ treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

Oxaliplatin

The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540

Oxidative stress

Differential toxicity of antimonial compounds and their effects on glutathione homeostasis in a human leukaemia monocyte cell line, 257

Antioxidative function and biodistribution of $[\text{Gd}@\text{C}_{82}(\text{OH})_{22}]_n$ nanoparticles in tumor-bearing mice, 872

Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610

Metabolism of oxidants by blood from different mouse strains, 1753

 α -Oxoaldehydes

Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610

Oxypurinol

Superoxide radical production by allopurinol and xanthine oxidase, 1747

P2Y receptor

Extracellular ATP activates ERK1/ERK2 via a metabotropic P2Y_1 receptor in a Ca^{2+} independent manner in differentiated human skeletal muscle cells, 1497

p38 MAP kinase

Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742

p38 MAPK

Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198

p53

Cell cycle arrest and proapoptotic effects of the anticancer cyclodepsipeptide serratomolide (AT514) are independent of p53 status in breast cancer cells, 32

The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540

Paclitaxel

Methionine dependence of tumours: A biochemical strategy for optimizing paclitaxel chemosensitivity in vitro, 772

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

PAF

Relative effects of phenolic constituents from *Yucca schidigera* Roetzl. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479

Palliative care

The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146

Pantoprazole

Hydroxyl radical scavenging reactivity of proton pump inhibitors, 1337

PAR-1

Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449

Paracellular permeability

Modulation of intestinal barrier properties by miltefosine, 486

PARP inhibitors

Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia-reperfusion, 441

Pathogen response – antibiotics

Antibacterial drug discovery—Then, now and the genomics future, 901

PBP

Targeting the forgotten transglycosylases, 957

PBR

The effects of prostaglandin $\text{F}_{2\alpha}$ treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

PC-Cl3

Differential response of normal, dedifferentiated and transformed thyroid cell lines to cisplatin treatment, 50

PDE

Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219

PDT

Mechanisms involved in δ -aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: Relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin, 42

Penems

Redefining penems, 1085

Pentagalloyl glucose

Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570

PEPT1

The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581

Peptide deformylase

The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042

Peptide deformylase inhibitors

The evolution of peptide deformylase as a target: Contribution of biochemistry, genetics and genomics, 1042

Peptide library

Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319

Peptidoglycan synthesis

Does the cell wall of bacteria remain a viable source of targets for novel antibiotics?, 996

- Peptidomimetic**
Peptide-bond modified glutathione conjugate analogs modulate GST π function in GSH-conjugation, drug sensitivity and JNK signaling, 268
- Peptoids**
Anti-Tat and anti-HIV activities of trimers of *n*-alkylglycines, 596
- Peroxisome proliferator-activated receptor Δ**
Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A₂ receptor gene in human erythroleukemia cells, 1308
- PGF_{2 α}**
The effects of prostaglandin F_{2 α} treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472
- P-glycoprotein**
Modulation of intestinal barrier properties by miltefosine, 486
The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146
Mitochondrial localization and activity of P-glycoprotein in doxorubicin-resistant K562 cells, 1162
- P-gp**
Effects of capsaicin on P-gp function and expression in Caco-2 cells, 1727
- Phage display**
Identification and characterization of surrogate peptide ligand for orphan G protein-coupled receptor mas using phage-displayed peptide library, 319
- Pharmacodynamics**
The role of pharmacodynamic research in the assessment and development of new antibacterial drugs, 1057
- Pharmacokinetics**
Binding kinetics and duration of in vivo action of novel prolyl oligopeptidase inhibitors, 683
Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422
- Pharmacophore model**
Differential interactions of G-proteins and adenylyl cyclase with nucleoside 5'-triphosphates, nucleoside 5'-[γ -thio]triphosphates and nucleoside 5'-[β , γ -imido]triphosphates, 89
- 1,10-Phenanthroline**
Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24), 426
- Phenformin**
AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637
- Phenolic compounds**
Inhibitory effect of dietary phenolic compounds on *Chlamydia pneumoniae* in cell cultures, 735
- Phenyl saligenin phosphate**
Inhibition of neurite outgrowth in differentiating mouse N2a neuroblastoma cells by phenyl saligenin phosphate: Effects on MAP kinase (ERK 1/2) activation, neurofilament heavy chain phosphorylation and neuropathy target esterase activity, 1240
- Phenylsuccinate**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Phosphatidylcholine**
Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114
- Phosphoinositide 3-kinase**
Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735
- Phospholipase A₂**
Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂- α , 144
- Phospholipase C**
Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y₂ and P2Y₄ receptors, 540
- Photo-damage**
Mechanisms involved in δ -aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: Relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin, 42
- PI3-kinase/Akt**
Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia-reperfusion, 441
- Pituitary**
 γ -Aminobutyric acid (GABA)-C receptor stimulation increases prolactin (PRL) secretion in cultured rat anterior pituitary cells, 1705
- [³H]PK 11195**
The effects of prostaglandin F_{2 α} treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472
- PKB/Akt**
Differential response of normal, dedifferentiated and transformed thyroid cell lines to cisplatin treatment, 50
- PKC- δ**
Differential response of normal, dedifferentiated and transformed thyroid cell lines to cisplatin treatment, 50
- Placenta**
Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAR cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349
- Platelet-derived growth factor (PDGF)**
Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806
- Platelets**
Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂- α , 144
- Platinum**
Cytotoxic efficacy of an anthraquinone linked platinum anticancer drug, 1136
- Polychlorinated biphenyls**
Polychlorinated biphenyls induce arachidonic acid release in human platelets in a tamoxifen sensitive manner via activation of group IVA cytosolic phospholipase A₂- α , 144
- Polygodial**
Pharmacological characterisation of the plant sesquiterpenes polygodial and drimaniol as vanilloid receptor agonists, 1248
- Porin mutations and deletions**
The forgotten Gram-negative bacilli: What genetic determinants are telling us about the spread of antibiotic resistance, 1073
- Porins**
Unmet medical needs in antibacterial therapy, 991
- Potassium channel regulation**
The Cdc42 inhibitor secramine B prevents cAMP-induced K⁺ conductance in intestinal epithelial cells, 1720

Potassium channels

Block of hERG channel by ziprasidone: Biophysical properties and molecular determinants, 278

PPAR γ

Niacin induces PPAR γ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646

Pranlukast

CysLT $_1$ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115

Prednisolone

Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858

Prednisone

Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858

Pregnant opiate addict

The effect of methadone and buprenorphine on human placental aromatase, 1255

Pregnenolone xenobiotic receptor

Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693

Privileged structures

Arylpiperazines displaying preferential potency against chloroquine-resistant strains of the malaria parasite *Plasmodium falciparum*, 61

Prodrug

Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683

Progesterone

The effects of prostaglandin F $_{2\alpha}$ treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

Prolactin

γ -Aminobutyric acid (GABA)-C receptor stimulation increases prolactin (PRL) secretion in cultured rat anterior pituitary cells, 1705

Proliferation

ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479

Prolyl oligopeptidase

Binding kinetics and duration of in vivo action of novel prolyl oligopeptidase inhibitors, 683

Promoter

Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A $_2$ receptor gene in human erythroleukemia cells, 1308

Prostaglandin

Niacin induces PPAR γ expression and transcriptional activation in macrophages via HM74 and HM74a-mediated induction of prostaglandin synthesis pathways, 646

Prostate carcinoma

Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791

Protein kinase B

AMP-activated protein kinase (AMPK) activating agents cause dephosphorylation of Akt and glycogen synthase kinase-3, 1637

Protein kinase C

Ceramide-1-phosphate activates cytosolic phospholipase A2 α directly and by PKC pathway, 850

Proteinase activated receptor 2

A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223

Proton pump inhibitors

Characterization of the inhibitory activity of tenatoprazole on the gastric H $^+$,K $^+$ -ATPase in vitro and in vivo, 837

Hydroxyl radical scavenging reactivity of proton pump inhibitors, 1337

Protoporphyrin

Mechanisms involved in δ -aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: Relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin, 42

Pseudomonas

Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910

Pseudopregnancy

The effects of prostaglandin F $_{2\alpha}$ treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

(9) Pulmonary

Effect of culture conditions on the expression and function of Bsep, Mrp2, and Mdr1a/b in sandwich-cultured rat hepatocytes, 1520

9-[2-(Phosphonomethoxy)ethyl]guanine

N 6 -Methyl-AMP aminohydrolase activates N 6 -substituted purine acyclic nucleoside phosphonates, 1370

PUMA

The BH3-only protein, PUMA, is involved in oxaliplatin-induced apoptosis in colon cancer cells, 1540

Purine analogues

The pattern of deoxycytidine- and deoxyguanosine kinase activity in relation to messenger RNA expression in blood cells from untreated patients with B-cell chronic lymphocytic leukemia, 882

Purine receptors

CysLT $_1$ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115

Purine salvage

Structure-activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671

Pyrethroids

Hydrolytic metabolism of pyrethroids by human and other mammalian carboxylesterases, 657

Pyridine N-oxide derivatives

Pyridine N-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122

Pyrimidines

Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y $_2$ and P2Y $_4$ receptors, 540

Quinazoline

Epidermal growth factor receptor inhibitor (PD168393) potentiates cytotoxic effects of paclitaxel against androgen-independent prostate cancer cells, 751
Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422

Quinone reductase 1

NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1

Quinone reductase 2

NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1

- Characterization of the melatonergic MT₃ binding site on the NRH:quinone oxidoreductase 2 enzyme, 74
- QX-222**
The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299
- Rabbits**
The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366
- Radioligand binding**
Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- Radiosensitivity**
Sensitization of breast carcinoma cells to ionizing radiation by small molecule inhibitors of DNA-dependent protein kinase and ataxia telangiectasia mutated, 13
- Raf-1**
14-3-3 ζ interacts with human thromboxane receptors and is involved in the agonist-induced activation of the extracellular-signal-regulated kinase, 624
- RANK ligand**
Differential activity of kaempferol and quercetin in attenuating tumor necrosis factor receptor family signaling in bone cells, 818
- Ras**
Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735
- Rat**
A protective role for proteinase activated receptor 2 in airways of lipopolysaccharide-treated rats, 223
- Rat lung**
Metabolism of nilutamide in rat lung, 377
- Rational drug design**
Mobilization of iron from cells by hydroxyquinoline-based chelators, 214
- Reaction phenotyping**
In vitro-in vivo correlation for drugs and other compounds eliminated by glucuronidation in humans: Pitfalls and promises, 1531
- Reactive oxygen species**
Glyoxal markedly compromises hepatocyte resistance to hydrogen peroxide, 1610
- Receptor**
Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- Receptor antagonists**
The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488
- Receptor overexpression**
Histamine H2 receptor overexpression induces U937 cell differentiation despite triggered mechanisms to attenuate cAMP signalling, 1219
- Receptor theory**
Some implications of receptor theory for in vivo assessment of agonists, antagonists and inverse agonists, 1663
- Redox**
Redox modifications of protein-thiols: Emerging roles in cell signaling, 551
- Redox regulation**
Pyridine N-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122
- Reduced glutathione**
Metabolism of oxidants by blood from different mouse strains, 1753
- Regulation of STAT activity**
JAK/STAT signal transduction: Regulators and implication in hematological malignancies, 713
- Renal and Hepatic Pharmacology**
Effect of culture conditions on the expression and function of Bsep, Mrp2, and Mdr1a/b in sandwich-cultured rat hepatocytes, 1520
- Rengyolone**
Rengyolone inhibits inducible nitric oxide synthase expression and nitric oxide production by down-regulation of NF- κ B and p38 MAP kinase activity in LPS-stimulated RAW 264.7 cells, 1198
- Repressor**
MZFI possesses a repressively regulatory function in ERCC1 expression, 761
- Resistance**
Antibacterial drug discovery—Then, now and the genomics future, 901
Understanding the longevity of the β -lactam antibiotics and of antibiotic/ β -lactamase inhibitor combinations, 930
Glycopeptides: Update on an old successful antibiotic class, 968
Unmet medical needs in antibacterial therapy, 991
When will the genomics investment pay off for antibacterial discovery?, 1096
- Respiratory complexes**
Distribution of tocopheryl quinone in mitochondrial membranes and interference with ubiquinone-mediated electron transfer, 1589
- Resveratrol**
NRH:quinone reductase 2: An enzyme of surprises and mysteries, 1
Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways, 742
- Retinoid X receptor (RXR)**
Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAr cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349
- Reverse transcriptase inhibitors**
3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342
- Rhesus macaque**
Molecular cloning and radioligand binding characterization of the chemokine receptor CCR5 from rhesus macaque and human, 163
- Ribose group**
Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand-receptor interaction, 865
- Ribosomes**
Structure-based drug design meets the ribosome, 1016
- RNAi**
Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum, 702
- RNase A**
Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum, 702
- RND transporters**
Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use, 910

ROS

Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia-reperfusion, 441

Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, 1602

ROS production

Suppression of survival in human SKBR3 breast carcinoma in response to metal-chelator complexes is preferential for copper-dithiocarbamate, 722

Rosiglitazone

Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605

Ruthenium(II) organo-metallic complex

Anti-tumour activity in non-small cell lung cancer models and toxicity profiles for novel ruthenium(II) based organo-metallic compounds, 408

RXR

Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98

Salivary proline-rich protein

Decreased polyphenol transport across cultured intestinal cells by a salivary proline-rich protein, 1570

Sarin

Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure, 497

Schisandrin B

Schisandrin B enhances doxorubicin-induced apoptosis of cancer cells but not normal cells, 584

Schizophrenia

Distinct functional profiles of aripiprazole and olanzapine at RNA edited human 5-HT_{2C} receptor isoforms, 521

Screening

The historical delivery of antibiotics from microbial natural products—Can history repeat?, 981

When will the genomics investment pay off for antibacterial discovery?, 1096

Secramine

The Cdc42 inhibitor secramine B prevents cAMP-induced K⁺ conductance in intestinal epithelial cells, 1720

Serum

Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum, 702

Serum- and glucocorticoid-regulated kinase-1

Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175

Serum response element

Pharmacological characterization of protease activated receptor-1 by a serum responsive element-dependent reporter gene assay: Major role of calmodulin, 1449

ShRNA

Potential role of short hairpin RNA targeting epidermal growth factor receptor in growth and sensitivity to drugs of human lung adenocarcinoma cells, 1265

Signal transduction

Critical role of PI3-kinase/Akt activation in the PARP inhibitor induced heart function recovery during ischemia-reperfusion, 441

14-3-3 ζ interacts with human thromboxane receptors and is involved in the agonist-induced activation of the extra-cellular-signal-regulated kinase, 624

JAK/STAT signal transduction: Regulators and implication in hematological malignancies, 713

The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists, 1488

Long term environmental tobacco smoke activates nuclear transcription factor-kappa B, activator protein-1, and stress responsive kinases in mouse brain, 1602

Signaling cross-talk

Cross-talk between angiotensin II and glucagon receptor signaling mediates phosphorylation of mitogen-activated protein kinases ERK 1/2 in rat glomerular mesangial cells, 1711

siRNA

Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum, 702

Skin inflammation

Involvement of tumor necrosis factor (TNF)- α in phorbol ester 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin edema in mice, 1331

Slow binding inhibition

Binding kinetics and duration of in vivo action of novel prolyl oligopeptidase inhibitors, 683

Smooth muscle cells

Stimulation of serum- and glucocorticoid-regulated kinase-1 gene expression by endothelin-1, 1175

S-nitrosylation

Redox modifications of protein-thiols: Emerging roles in cell signaling, 551

Solid phase peptide synthesis

Synthesis and antinociceptive activity of cyclic endomorphin-2 and morphiceptin analogs, 188

Sp1

The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581

Species difference

Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294

Sphingolipids

Ceramide-1-phosphate activates cytosolic phospholipase A2 α directly and by PKC pathway, 850

Hexadecylphosphocholine disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2 tumour cells, 1114

Spinal cord

Amitriptyline prevents thermal hyperalgesia and modifications in rat spinal cord GABA_B receptor expression and function in an animal model of neuropathic pain, 196

Spin-lattice relaxation rate

Drug-protein recognition processes investigated by NMR relaxation data. A study on corticosteroid-albumin interactions, 858

SR-BI

Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes, 605

Src family kinase

Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735

Stable transfectants

Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta, 1459

Statins

Hydroxymethylglutaryl-coenzyme A reductase inhibitors induce apoptosis in human cardiac myocytes in vitro, 1324

- Staurosporine**
Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562
- Stereospecificity**
Stereospecific recognition of a spirosuccinimide type aldose reductase inhibitor (AS-3201) by plasma proteins: A significant role of specific binding by serum albumin in the improved potency and stability, 338
- Sterol regulatory element-binding protein-1c**
Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453
- S-thiolation**
Redox modifications of protein–thiols: Emerging roles in cell signaling, 551
- Stilbenic derivatives**
Relative effects of phenolic constituents from *Yucca schidigera* Roehl. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479
- Streptococcus pneumoniae***
Surveillance uncovers the smoking gun for resistance emergence, 1036
- Structural genomics**
Crystallizing new approaches for antimicrobial drug discovery, 1048
- Structure**
Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629
- Structure activity relationship**
Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y₂ and P2Y₄ receptors, 540
- Structure guided drug discovery**
Crystallizing new approaches for antimicrobial drug discovery, 1048
- Structure–activity relationship**
Structure–activity relationship for nucleoside analogs as inhibitors or substrates of adenosine kinase from *Mycobacterium tuberculosis*. I. Modifications to the adenine moiety, 1671
- Structure-based drug design**
Structure-based drug design meets the ribosome, 1016
- Sulfation degree**
Interactions of low-molecular-weight semi-synthetic sulfated heparins with human leukocyte elastase and human Cathepsin G, 287
- Superoxide**
CO from enhanced HO activity or from CORM-2 inhibits both O₂^{•−} and NO production and downregulates HO-1 expression in LPS-stimulated macrophages, 307
- Superoxide anion**
Activation of phosphoinositide 3-kinase and Src family kinase is required for respiratory burst in rat neutrophils stimulated with artocarpol A, 1735
- Superoxide radical**
Superoxide radical production by allopurinol and xanthine oxidase, 1747
- Surveillance studies**
Surveillance uncovers the smoking gun for resistance emergence, 1036
- Synaptosomes**
Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools, 399
- Synergism**
Epidermal growth factor receptor inhibitor (PD168393) potentiates cytotoxic effects of paclitaxel against androgen-independent prostate cancer cells, 751
- Systemic and topical glucocorticoids**
Potencies of topical glucocorticoids to mediate genomic and nongenomic effects on human peripheral blood mononuclear cells, 530
- T cells**
Inhibition of interleukin-4 production in activated T cells via the downregulation of AP-1/NF-AT activation by N-lauroyl-D-erythro-sphingosine and N-lauroyl-D-erythro-C₂₀-sphingosine, 1229
- T lymphocytes**
Inhibition of sphingolipid synthesis impairs cellular activation, cytokine production and proliferation in human lymphocytes, 126
- Tamoxifen**
Quaternary ammonium-linked glucuronidation of trans-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358
- Target**
When will the genomics investment pay off for antibacterial discovery?, 1096
- Tat**
Anti-Tat and anti-HIV activities of trimers of n-alkyl-glycines, 596
- Taurine analogues**
Inhibition of rabbit brain 4-aminobutyrate transaminase by some taurine analogues: A kinetic analysis, 1510
- Taxol**
Cellular FLICE-like inhibitory protein (c-FLIP): A novel target for Taxol-induced apoptosis, 1551
- Telavancin**
Glycopeptides: Update on an old successful antibiotic class, 968
- Template**
Natural products — The future scaffolds for novel antibiotics?, 919
- Tenatoprazole**
Characterization of the inhibitory activity of tenatoprazole on the gastric H⁺,K⁺-ATPase in vitro and in vivo, 837
- Tetrachlorobenzoquinone**
Chemical modification at subunit 1 of rat kidney Alpha class glutathione transferase with 2,3,5,6-tetrachloro-1,4-benzoquinone: Close structural connectivity between glutathione conjugation activity and non-substrate ligand binding, 1629
- Δ^9 -Tetrahydrocannabinol**
The effects of cannabinoids on P-glycoprotein transport and expression in multidrug resistant cells, 1146
- TG-domain**
Targeting the forgotten transglycosylases, 957
- The gastric H,K-ATPase**
Characterization of the inhibitory activity of tenatoprazole on the gastric H⁺,K⁺-ATPase in vitro and in vivo, 837
- Thiazolidinedione**
Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A₂ receptor gene in human erythroleukemia cells, 1308
- Thioredoxin**
Redox modifications of protein–thiols: Emerging roles in cell signaling, 551

Thromboxane receptor

14-3-3 ζ interacts with human thromboxane receptors and is involved in the agonist-induced activation of the extra-cellular-signal-regulated kinase, 624

Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A₂ receptor gene in human erythroleukemia cells, 1308

Thymidine

3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342

Thymidine kinase 2

3'-Azido-3'-deoxythymidine (AZT) inhibits thymidine phosphorylation in isolated rat liver mitochondria: A possible mechanism of AZT hepatotoxicity, 1342

Tight binding inhibition

Binding kinetics and duration of in vivo action of novel prolyl oligopeptidase inhibitors, 683

Tight junctions

Modulation of intestinal barrier properties by miltefosine, 486

Tissue inhibitors of matrix metalloproteinases

Fibroblast heterogeneity in collagenolytic response to colchicine, 574

TNF

Involvement of tumor necrosis factor (TNF)- α in phorbol ester 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin edema in mice, 1331

TNF- α

Estrogen receptor-independent inhibition of tumor necrosis factor- α gene expression by phytoestrogen equol is mediated by blocking nuclear factor- κ B activation in mouse macrophages, 136

Inhibition of TNF α -induced activation of nuclear factor κ B by kava (*Piper methysticum*) derivatives, 1206

 α -Tocopherol

Distribution of tocopheryl quinone in mitochondrial membranes and interference with ubiquinone-mediated electron transfer, 1589

 α -Tocopheryl quinone

Distribution of tocopheryl quinone in mitochondrial membranes and interference with ubiquinone-mediated electron transfer, 1589

Topoisomerase I

Circadian regulation of mouse topoisomerase I gene expression by glucocorticoid hormones, 1155

Topoisomerase II inhibitor

Enhancing the antiproliferative effect of topoisomerase II inhibitors using a polypeptide inhibitor of c-Myc, 248

Toxicity

Metabolism of nilutamide in rat lung, 377

Toxoplasma

6-Benzylthioinosine analogues: Promising anti-toxoplasmaic agents as inhibitors of the mammalian nucleoside transporter ENT1 (es), 69

TPA

Involvement of tumor necrosis factor (TNF)- α in phorbol ester 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin edema in mice, 1331

TRAIL

Sensitization to gimatecan-induced apoptosis by tumor necrosis factor-related apoptosis inducing ligand in prostate carcinoma cells, 791

Transactivation

Pyridine N-oxide derivatives inhibit viral transactivation by interfering with NF- κ B binding, 1122

Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta, 1459

Transcription

Selective allosteric ligand activation of the retinoid X receptor heterodimers of NGFI-B and Nurr1, 98

Transcriptional regulation

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

Transepithelial resistance

Investigation of the cytotoxicity of eukaryotic and prokaryotic antimicrobial peptides in intestinal epithelial cells in vitro, 1289

Transglycosylase inhibitors

Targeting the forgotten transglycosylases, 957

Transglycosylation process

Targeting the forgotten transglycosylases, 957

Transmembrane domain

Role of the second transmembrane domain of rat adenosine A₁ receptor in ligand-receptor interaction, 865

Transport protein

Efflux systems in bacterial pathogens: An opportunity for therapeutic intervention? An industry view, 949

Transporter

Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice, 1470

The transcription factor Cdx2 regulates the intestine-specific expression of human peptide transporter 1 through functional interaction with Sp1, 1581

Tributyltin

Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAr cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349

Trigeminal ganglion neuron

Pharmacological characterisation of the plant sesquiterpenes polygodial and drimaniol as vanilloid receptor agonists, 1248

Triglycerides

Different roles of liver X receptor α and β in lipid metabolism: Effects of an α -selective and a dual agonist in mice deficient in each subtype, 453

Triphenyltin

Organotin compounds enhance 17 β -hydroxysteroid dehydrogenase type I activity in human choriocarcinoma JAr cells: Potential promotion of 17 β -estradiol biosynthesis in human placenta, 1349

TrkB

A role for AMPA receptors in mood disorders, 1273

TRPV1

Pharmacological characterisation of the plant sesquiterpenes polygodial and drimaniol as vanilloid receptor agonists, 1248

Tumour hypoxia

Radiolytic and cellular reduction of a novel hypoxia-activated cobalt(III) prodrug of a chloromethylbenzindoline DNA minor groove alkylator, 1683

Tunicamycin

Synergism between staurosporine and drugs inducing endoplasmic reticulum stress, 1562

Turpentine

The 21-aminosteroid U74389G prevents the down-regulation and decrease in activity of CYP1A1, 1A2 and 3A6 induced by an inflammatory reaction, 366

Tyrosine kinase

Blockade of IGF-1 receptor tyrosine kinase has antineoplastic effects in hepatocellular carcinoma cells, 1435

Tyrosine kinase inhibitor

Preclinical analysis of the analinoquinazoline AG1478, a specific small molecule inhibitor of EGF receptor tyrosine kinase, 1422

Ubiquinone

Distribution of tocopheryl quinone in mitochondrial membranes and interference with ubiquinone-mediated electron transfer, 1589

UDP

CysLT₁ leukotriene receptor antagonists inhibit the effects of nucleotides acting at P2Y receptors, 115

UDP-glucuronosyltransferase

Quaternary ammonium-linked glucuronidation of *trans*-4-hydroxytamoxifen, an active metabolite of tamoxifen, by human liver microsomes and UDP-glucuronosyltransferase 1A4, 1358

In vitro-in vivo correlation for drugs and other compounds eliminated by glucuronidation in humans: Pitfalls and promises, 1531

UGT1A1

Proximal HNF1 element is essential for the induction of human UDP-glucuronosyltransferase 1A1 by glucocorticoid receptor, 693

Uncoupling

Generation of adenosine A₃ receptor functionally humanized mice for the evaluation of the human antagonists, 294

Use-dependent block

The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299

Uterus

The effects of prostaglandin F_{2α} treatment on peripheral-type benzodiazepine receptors in the ovary and uterus during pseudopregnancy of rats, 472

Vaccine

Commensals upon us, 893

Vanilloid

Molecular cloning, functional characterization of the porcine transient receptor potential V1 (pTRPV1) and pharmacological comparison with endogenous pTRPV1, 173

Vascular endothelial growth factor receptor

ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479

Vascular smooth muscle cell

Berberine suppresses MEK/ERK-dependent Egr-1 signaling pathway and inhibits vascular smooth muscle cell regrowth after in vitro mechanical injury, 806

VEGF

Relative effects of phenolic constituents from *Yucca schidigera* Roetzl. bark on Kaposi's sarcoma cell proliferation, migration, and PAF synthesis, 1479

Vigabatrin

Inhibition of rabbit brain 4-aminobutyrate transaminase by some taurine analogues: A kinetic analysis, 1510

Vinblastine

Methionine dependence of tumours: A biochemical strategy for optimizing paclitaxel chemosensitivity in vitro, 772

Vincristine

Vincristine transcriptional regulation of efflux drug transporters in carcinoma cell lines, 1695

Vitamin E

Vitamin E inhibits hemolysis induced by hemin as a membrane stabilizer, 799

Voltage dependence of recovery

The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299

Voltage-gated sodium channel

The voltage dependence of recovery from use-dependent block by QX-222 separates mechanisms for drug egress in the cardiac sodium channel, 1299

Whole-cell ligand binding

Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta, 1459

Xanthine oxidase

Superoxide radical production by allopurinol and xanthine oxidase, 1747

Xenopus oocytes

Block of hERG channel by ziprasidone: Biophysical properties and molecular determinants, 278

X-ray structure

Structure-based drug design meets the ribosome, 1016

ZD6474

ZD6474 inhibits proliferation and invasion of human hepatocellular carcinoma cells, 479

Ziprasidone

Block of hERG channel by ziprasidone: Biophysical properties and molecular determinants, 278