

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/biochempharm

Calmodulin potentiates G $\beta\gamma$ activation of phospholipase C- β 3

Jennifer S. McCullar^a, Dean A. Malencik^b, Walter K. Vogel^c, Kristi M. Crofoot^c,
Sonia R. Anderson^b, Theresa M. Filtz^{a,b,c,*}

^a Molecular and Cellular Biology Graduate Program, Oregon State University, Corvallis, OR 97331, United States

^b Department of Biochemistry and Biophysics, Oregon State University, Corvallis, OR 97331, United States

^c Department of Pharmaceutical Sciences, Oregon State University, Corvallis, OR 97331, United States

ARTICLE INFO

Article history:

Received 29 June 2006

Accepted 9 October 2006

Keywords:

Phosphatidylinositol hydrolysis

Phospholipase C- β

Gbetagamma

1321N1 cells

Calmodulin

Fluorescence anisotropy

ABSTRACT

Phospholipase C- β (PLC- β) isozymes (EC 3.1.4.11) hydrolyze the membrane phospholipid phosphatidylinositol-4,5-bisphosphate to generate intracellular second messenger signaling molecules inositol-1,4,5-trisphosphate (IP₃) and diacylglycerol (DAG) in response to receptor activation and other cellular stimuli. PLC β 1 and PLC β 3 isozymes were previously demonstrated to bind the calcium-sensitive molecule calmodulin [McCullar JS, Larsen SA, Millimaki RA, Filtz TM. Calmodulin is a phospholipase C- β interacting protein. *J Biol Chem* 2003;278(36):33708–13]. We have now shown through fluorescence anisotropy that calmodulin/PLC β 3 affinities increase with increasing calcium in a physiologically relevant concentration range. The bimolecular affinity constants for calmodulin interaction with PLC β 1 or PLC β 3 were estimated as 260 and 200 nM, respectively, from fluorescence anisotropy data. There was no effect of calmodulin on basal or G α q-stimulated catalytic activity for either isozyme. However, the interaction between calmodulin and PLC β 3 leads to potentiation of activation by the G-protein $\beta\gamma$ dimer in an in vitro assay. 1321N1 cells treated with calmodulin inhibitors concurrent with and post-stimulation of muscarinic receptors significantly reduced [³H]PIP hydrolysis. Together these data are suggestive of cooperative role for calmodulin in the G-protein $\beta\gamma$ dimer-stimulated activity of PLC β 3.

© 2006 Elsevier Inc. All rights reserved.

1. Introduction

Phosphatidylinositol phospholipid-specific phospholipase C (PLC)¹ is a key intracellular signaling molecule that catalyzes the hydrolysis of PIP₂ into IP₃, a regulator of cytosolic calcium levels, and diacylglycerol, a well-characterized activator of protein kinase C [2]. The identified PLC isozymes have been classified by sequence homology into six families, β , γ , δ , ϵ , ζ ,

and η [3–5]. Each family has unique mechanisms of activation and regulation. The activity of the PLC β family of isozymes is stimulated by membrane G-protein coupled receptors (GPCR) through heterotrimeric guanine nucleotide binding (G) proteins, which are composed of a GTP/GDP-binding α subunit and a $\beta\gamma$ dimer.

There are four identified and characterized isoforms of PLC β numbered 1–4. PLC β 2 and PLC β 4 have limited tissue

* Corresponding author at: Department of Pharmaceutical Sciences, 203 Pharmacy Building Oregon State University, Corvallis, OR 97331, United States. Tel.: +1 541 737 5803; fax: +1 541 737 3999.

E-mail address: Theresa.Filtz@oregonstate.edu (T.M. Filtz).

Abbreviations: PLC, phosphatidylinositol phospholipid-specific phospholipase C; PE, phosphatidylethanolamine; PS, phosphatidylserine; PIP 16:0, synthetic phosphatidylinositol-4-phosphate with symmetric 16:0 saturated fatty acyl chains; PIP, phosphatidylinositol-4-phosphate; PIP₂, phosphatidylinositol-4,5-bisphosphate; PIP₃, phosphatidylinositol-3,4,5-trisphosphate; SDS-PAGE, sodium dodecyl phosphate polyacrylamide gel electrophoresis; W-13, N-(4-aminobutyl)-5-chloro-2-naphthalenesulfonamide; BSA, bovine serum albumin 0006-2952/\$ – see front matter © 2006 Elsevier Inc. All rights reserved.

doi:10.1016/j.bcp.2006.10.004

distributions, whereas PLC β 1 and PLC β 3 are nearly ubiquitous in human tissues; PLC β 1 being dominant in brain and PLC β 3 being dominant in heart and smooth muscle [2]. PLC β 1 and PLC β 3 are both activated by calcium and G α q [6–8], but the PLC β 3 isoform is additionally sensitive to activation by G $\beta\gamma$ [9]. The mechanisms of regulation of PLC β 3 are incompletely understood despite the enzyme's importance in a variety of cellular processes [10–14]. Aberrancies in expression of PLC β 3 can lead to tumorigenesis [13,15,16], and PLC β 3 knockout mice show changes in μ -opioid response [12] or early embryonic lethality [17].

Calmodulin is an established, ubiquitous and abundant calcium-sensitive regulatory protein associated with a vast diversity of cellular functions including signal transduction [18,19]. Calmodulin binds four molecules of calcium cooperatively, and undergoes a significant conformational change upon calcium binding that is important for its many calcium-sensitive regulatory functions. Calmodulin binding sites are nearly as diverse as the number of calmodulin binding proteins, but generally are amphipathic α -helices, typically 20–35 amino acids long, with basic and hydrophobic residues sorting to opposite sides on an α -helical projection. The list of various mechanisms by which calmodulin regulates proteins is growing nearly as fast as the list of calmodulin binding proteins [18,20].

We previously reported that calmodulin directly interacts with PLC β 1 and PLC β 3, and that calmodulin inhibitors attenuate inositol phosphate (IP) accumulation in whole cells [1]. To further understand this interaction, we sought to determine the direct effect of calmodulin on PLC- β activity in vitro and to determine the binding affinity and calcium dependence of the PLC β /calmodulin interaction.

2. Materials and methods

2.1. Reagents

PLC β -selective polyclonal rabbit antisera and alkaline phosphatase-conjugated goat anti-rabbit IgG secondary antibody were obtained from Santa Cruz Biotechnology Inc. (Santa Cruz, CA). SuperSignalTM chemiluminescent substrate was purchased from Pierce (Rockford, IL). Fatty acid-free bovine serum albumin (FAF-BSA), W-13 and carbamylcholine chloride (carbachol) were obtained from Calbiochem (San Diego, California). PIP Strip[®] phospholipid blots were obtained from Echelon Biosciences (Salt Lake City, UT). Synthetic and purified bovine brain PIP, PIP2, PS and PE were purchased from Avanti Polar Lipids (Alabaster, AL). Calmodulin was purified in the presence of calcium from bovine brain as previously described [21]. PLC β 1, PLC β 3, G α q, G $\beta\gamma$, and [3H]PIP2 were purified as previously described [22–24]. Alexa Fluor[®] 488 was acquired from Molecular Probes (Eugene, OR) and was conjugated to calmodulin according to the manufacturer's recommended procedure.

2.2. Fluorescence anisotropy

Fluorescence anisotropy and fluorescence emission spectra were recorded with a Perkin-Elmer LS-50 Luminescence

Spectrophotometer maintaining constant temperature (20 °C). Measurements of fluorescence anisotropy for PLC β 1 and PLC β 3 with Alexa-calmodulin in buffer containing 0.1 M KCl and 30 mM MOPS, pH 7.2, were performed at an excitation wavelength of 490 nm (band-pass 5 nm) using a linear polarizer, and the fluorescence emission intensities at 525 nm (band-pass 20 nm) were monitored through a second linear polarizer. Anisotropy, r , was calculated according to the equation, $r = (I_{VV} - GI_{VH}) / (I_{VV} + 2GI_{VH})$, where I_{VV} and I_{VH} are the intensity of vertically or horizontally polarized emitted light, respectively, obtained with vertically polarized exciting light. Four readings were taken for each measurement and averages calculated and recorded. The value of G , 0.982, which corrects for unequal transmission of vertically and horizontally polarized emitted light, was obtained from tables prepared by the Anderson laboratory specific for their instrument (personal communication).

2.3. Reconstitution assay

The catalytic activity of PLC β 1 and PLC β 3 was quantitated using [3H]PIP2 substrate as described previously [25]. Briefly, 45 ng of purified PLC β 3 or 15 ng of PLC β 1 in 20 μ l of 50 mM HEPES pH 7.2, 3 mM EGTA, 80 mM KCl (Buffer 1) and 1 mg/ml fatty acid-free BSA was added to 20 μ l of Buffer 1 containing 15 μ M PIP2, 135 μ M phosphatidylethanolamine and 6–10,000 cpm [3H]PIP2. Ten microliters of 50 mM HEPES pH 7.2, 1 mM EDTA, 3 mM EGTA, 5 mM MgCl₂, 100 mM NaCl, 1% cholate (Buffer 2) was added to each reaction containing either 30 ng G α q or 50 ng G $\beta\gamma$ for G-protein-stimulated activity of PLC. Basal PLC activity was quantitated with the same buffer with no added G-protein. Buffer 1 with 9 mM CaCl₂ (10 μ l) was added to yield a final assay volume of 60 μ l. The reaction proceeded at 30 °C for 10 min and was terminated by the addition of 375 μ l of ice cold mixture of chloroform, methanol and hydrochloric acid in a ratio of 80:40:1, followed by addition of 125 μ l chloroform and 125 μ l 0.1N hydrochloric acid with vigorous mixing. The aqueous and organic phases were separated by centrifugation for 5 min at 2000 \times g. [3H]IP3 product release was quantitated by scintillation counting of 400 μ l of the upper phase. Triplicate samples were run in three separate experiments.

2.4. Protein lipid overlay assays

Protein lipid overlay assays were conducted using either commercial phospholipid membrane arrays containing synthetic phospholipids (PIP-Strips[®]) or nitrocellulose membranes spotted in our laboratory with mammalian-derived brain phospholipids. Nitrocellulose membrane phospholipid blots were prepared by spotting with chloroform containing 0, 50 or 100 pmol of phosphatidylserine, phosphatidylethanolamine or PIP2 in 1 μ l and allowed to dry. Both purchased and nitrocellulose membranes were blocked with 3% (w/v) fatty acid-free BSA in TBST buffer (150 mM NaCl, 10 mM Tris-HCl, pH 8.0, and 0.1% (v/v) Tween-20) for 1 h at room temperature. Blocked membranes were incubated overnight at 4 °C with 0.5 μ g/ml purified PLC β 3 or PLC β 1 in the presence or absence of 0.5 μ g/ml calmodulin. The membranes were then washed three times for 10 min in TBST with 3% fatty acid-free BSA,

followed by incubation with PLC β -selective polyclonal antibodies (0.1 μ g/ml in TBST with 1% fatty acid-free BSA) overnight at room temperature with gentle agitation. Following three washes over 30 min in TBST with 1% fatty acid-free BSA, the membranes were incubated for 1 h with alkaline phosphatase-conjugated goat anti-rabbit antibodies (0.1 μ g/ml) and visualized with SuperSignalTM alkaline phosphatase chemiluminescent substrate according to manufacturer's specifications.

2.5. Inositol phosphate assay

Whole cell inositol phosphate (IP) assays were performed with 1321N1 cells as previously described [26]. Prior to assay, 1321N1 cells were subcultured in 24-well plates until 80% confluent. Cells were then labeled overnight with [3H]myo-inositol, 1 μ Ci/0.5 ml/well, prepared in sterile inositol-free, bicarbonate-buffered DMEM without additives. Following radiolabeling, cells were pre-treated with 10 mM LiCl for 10 min in 20 mM Hepes-buffered DMEM, pH 7.4 (HDMEM) at 37 °C in room air to halt degradation and allow for accumulation of inositol phosphates. Muscarinic receptors were then stimulated with 1 mM carbachol for 20 min. W-13 (a calmodulin inhibitor) was added concurrently with carbachol or 10 min post-carbachol stimulation. Following 20 min stimulation, cells were lysed and total inositol phosphates (IP, IP2, IP3) purified and quantitated as described [26]. Lipids were collected to quantitate residual [3H]inositol phospholipids. Percent (%) conversion was calculated as $([3H]inositol\ phosphates\ (dpm))/([3H]inositol\ phospholipids\ (dpm) + [3H]inositol\ phosphates\ (dpm)) \times 100$. All assays were performed in triplicate and values reported reflect an average of at least three experiments \pm S.E. Students t-tests were performed to assess statistical significance where indicated.

2.6. Inositol trisphosphate accumulation assay

Prior to assay, 1321N1 cells were plated onto six-well plates at a density of 0.5×10^6 cells/well and allowed to adhere overnight. Cells were then labeled with [3H]myo-inositol, 10 μ Ci/2 ml/well, prepared in inositol-free, bicarbonate-buffered DMEM without additives overnight. Following radiolabeling, cells were pre-treated with 10 mM LiCl for 10 min in 1 ml of 20 mM HDMEM, pH 7.4 at 37 °C in room air in the presence or absence of 100 μ M W-13. Wells were treated with vehicle (water) or 1 mM carbachol to stimulate muscarinic receptors for 30, 60 or 90 s. Following stimulation, the reaction was stopped and cells lysed by the addition of an equal volume of 15% trichloroacetic acid solution directly to the wells. The contents of the wells were transferred to glass tubes and spun at 1500 rpm in a table top centrifuge to pellet cell debris. The supernatant was then transferred to a clean glass tube and the TCA extracted five times with ether. After extraction, the inositol phosphate species were separated as described previously [27]. Briefly, 2 ml dH₂O was added to the extracted cell lysates and poured over Poly-prep columns containing 0.5 ml bed volume of AG1-X8 anion-exchange resin (Formate form, 200–400 mesh, Bio-Rad). Tubes were rinsed with 8 ml dH₂O and applied to the columns. The columns were washed with 5 ml 25 mM sodium borate, 60 mM sodium formate, and

eluted with 5 ml 0.1 M formic acid, 0.2 M ammonium formate (IP containing fraction), 5 ml 0.1 M formic acid, 0.4 M ammonium formate (IP2 containing fraction), and 5 ml 0.1 M formic acid, 1.0 M ammonium formate (IP3 containing fraction). The first 2.5 ml of each elution was collected and analyzed for incorporation of [3H]inositol into IP, IP2 and IP3 by liquid scintillation chromatography. The cell membrane pellets were solubilized with 1 ml of 1 M NaOH, which was used to wash the corresponding wells of the plate for residual labeled cell membrane. The membrane prep was neutralized with 1 ml of 1N HCl. The membrane lipids were collected to quantitate total [3H]inositol phospholipids. Percent IP3 accumulation was calculated as $([3H]inositol\ trisphosphate\ (dpm))/([3H]inositol\ phospholipids\ (dpm) + [3H]inositol\ phosphates\ (dpm)) \times 100$. All assays were performed in triplicate and values reported reflect an average of at least two experiments \pm S.E. Students t-tests were performed to assess statistical significance where indicated.

3. Results

3.1. Fluorescence anisotropy analysis of PLC β 3 and PLC β 1 binding to calmodulin

Having established direct binding between PLC β 1 or PLC β 3 and calmodulin in the context of the full-length proteins [1], we sought to determine the affinity of PLC β for calmodulin. Fluorescence anisotropy depends on the rotational freedom of a fluorophore, which is influenced by its environment. By conjugating a fluorophore to calmodulin, we can estimate binding affinity by recording changes in the rotation of the fluorescent calmodulin in the presence of a potential binding partner, in this case PLC β . Using calmodulin conjugated to Alexa Fluor[®] 488 we measured the fluorescence anisotropy of Alexa-calmodulin titrated with PLC β 1 and PLC β 3 in the presence of 1 mM free calcium. We can determine the degree of complex formation of CaM with the PLC β by measuring the fluorescence anisotropy of the Alexa-calmodulin. PLC β (isoenzyme 1 or 3) was titrated with Alexa-calmodulin in the presence and the absence of calcium, and Alexa-calmodulin fluorescence anisotropy was measured. These anisotropy titration curves are shown in Fig. 1 (panels A and B). In the presence of calcium, dissociation constants of 200 nM for Alexa-calmodulin/PLC β 3 and 260 nM for Alexa-calmodulin/PLC β 1 interaction were determined by non-linear curve-fitting of the experimental titration data shown in Fig. 1 (panels A and B). The fit curve is corrected for a known and quantitated glycerol effect from the purified PLC β storage buffer. In the presence of EDTA, titration of Alexa-calmodulin with PLC β 3 caused only small increases in anisotropy that failed to approach saturation over the concentration range covered, indicative of a non-specific interaction (Fig. 1, Panel A). To further validate the calcium dependence of the interaction between PLC β 3 and calmodulin, we performed a calcium titration curve for Alexa-calmodulin/PLC β 3 interaction using 10 nM Alexa-calmodulin and 30 nM PLC β 3. Calcium stimulated increased Alexa-calmodulin anisotropy in the presence of PLC β 3 (Fig. 1, panel C). The EC₅₀ value for calcium dependence of the Alexa-calmodulin/PLC β 3 interaction can




Fig. 1 – Binding of fluorescently tagged calmodulin to PLCβ1 and PLCβ3 monitored by fluorescence anisotropy. Fluorescence anisotropy monitoring of 10 nM Alexa-calmodulin binding to increasing concentrations of (A) PLCβ3 in the presence of 1 mM calcium (■), or presence of 1 mM EDTA (×). Lines were curvefit as described in Section 2. Fluorescence anisotropy monitoring of 10 nM Alexa-calmodulin binding to increasing concentrations of (B) PLCβ1 in the presence of 1 mM calcium (■). Lines were curvefit as described in Section 2. K_d values calculated are 260 nM for PLCβ1 interaction with calmodulin and 200 nM for PLCβ3 interaction with calmodulin. (C) Effect of increasing calcium concentrations on fluorescence anisotropy with 10 nM Alexa-calmodulin (■) or 10 nM Alexa-calmodulin plus 30 nM PLCβ3 (□). Fluorescence

only be estimated as $<1 \mu\text{M}$, a value that is within a physiologically relevant range. This apparent EC_{50} is necessarily greater than the true EC_{50} of the complex because saturating concentrations of PLCβ3 were not achievable with limited amounts of purified protein available to use. The profile of anisotropy versus Ca^{2+} concentration would be concentration dependent until PLCβ3 concentration was greater than the K_d . However, these limited data suggest that increasing calcium concentrations following initial activation of the enzyme may support further interaction by increasing association of CaM with active PLCβ.

3.2. Calmodulin potentiates $G\beta\gamma$ activation of PLCβ3 in vitro

Pre-treatment with the calmodulin inhibitors W-13 and fluphenazine attenuated muscarinic receptor-stimulated phosphoinositide (PI) hydrolysis in 1321N1 cells [1]. To determine whether the potentiation of PI hydrolysis by calmodulin could occur through direct stimulation of PLCβ1 or PLCβ3, we performed reconstitution assays using purified PLCβ proteins and G-protein subunits with purified substrate in vitro. PLCβ3 can be activated in vitro by either the $G\alpha_q$ or $G\beta\gamma$ dimer of the heterotrimeric G-proteins, whereas PLCβ1 is sensitive to $G\alpha_q$ stimulation but relatively insensitive to $G\beta\gamma$ dimers [28]. Calmodulin did not affect basal or $G\alpha_q$ -stimulated PLCβ1 or PLCβ3 activities at any concentration tested (Fig. 2, panels A and B, respectively). However, calmodulin did potentiate $G\beta\gamma$ stimulation of PLCβ3 activity (Fig. 2, panel A, hatched bars). A titration of increasing $G\beta\gamma$ concentrations in the presence of $10 \mu\text{M}$ calmodulin showed an increase in PLCβ3 activity of almost 30% (Fig. 2, panel C).

3.3. Calmodulin does not affect lipid selectivity of PLCβ isozymes

Calmodulin regulates the activity of other signaling proteins via changes in membrane or lipid interactions [29,30] and calmodulin binding sites share structural similarity to some phospholipid binding domains. We assessed the effect of calmodulin on the lipid selectivity of either PLCβ1 or PLCβ3 isozyme using lipid overlay assay with commercially available PIP Strips®. Calmodulin did not affect the lipid selectivity of PLCβ1 or PLCβ3 (Fig. 3, panels B and C). Although no changes in selectivity of lipid were seen in the presence or absence of calmodulin, we found that PLCβ1 and PLCβ3 bind to PIP3 in addition to PI, PIP, PIP2, PA, and PS (Fig. 3, panels B and C). To investigate the lipid bindingspecificity of PLCβs, we purchased mammalian-derived phosphatidylinositol phospholipids for spotting onto nitrocellulose for additional lipid–protein overlay assays. Surprisingly, we found that PLCβ3 binding to synthetic phospholipids (single phospholipid species with 16:0 side-chains) did not mimic PLCβ3 binding to mammalian-derived phospholipids with a

anisotropy (emission intensity) was measured at 525 nm with excitation at 490 nm at 20°C . Shown is representative data from two separate experiments. An apparent EC_{50} value of $<1 \mu\text{M}$ was calculated for calcium effects on PLCβ3/calmodulin interactions.




Fig. 2 – Effect of calmodulin on basal and G-protein-stimulated PLCβ activity in vitro. (A) Basal (solid bars), 50 ng (90 nM) Gβγ-stimulated (hatched bars), and 30 ng (70 nM) GTPγS-Gαq-stimulated (white bars) PLCβ3 activity was quantitated in the presence of increasing concentrations of calmodulin. (B) Basal (solid bars) and 30 ng GTPγS-Gαq-stimulated (white bars) PLCβ1 activity was quantitated in the presence of increasing concentrations of calmodulin. Basal PLCβ activity was measured using PIP2 substrate in detergent/phospholipid vesicles. G-protein-stimulated PLCβ activity was measured in the presence of purified G-protein subunits using PIP2 substrate in detergent/phospholipid vesicles. PLCβ activity was quantitated as percent conversion of [³H]PIP2 substrate to [³H]IP_n (derivatives of inositol phosphate with the individual phosphate positions defined in accordance with the IUPAC convention) as described in Section 2. (C) PLCβ3 activity was measured in the presence of increasing concentrations of Gβγ subunits with (■) or without (□) 10 μM calmodulin. Data shown are

variety of chain lengths and saturations (Fig. 3, panel D). Side-chain specificity in PLCβ lipid binding has not, to our knowledge, been previously reported. The mammalian PIP2 contains the 16:0 side-chain species in only 7% of the total PIP2 species (manufacturer's analysis).

3.4. Calmodulin inhibitors attenuate phosphatidylinositol phospholipid hydrolysis concurrent with muscarinic receptor activation

To further dissect the mechanism by which calmodulin potentiates phosphatidylinositol phospholipid hydrolysis in whole 1321N1 cells, we attempted to assess whether calmodulin inhibitors have differential abilities to attenuate phosphatidylinositol phospholipid hydrolysis under varying cytosolic calcium conditions. Previously, we pre-treated 1321N1 cells with CaM inhibitors prior to activation by carbachol and showed inhibition of IP₃ accumulation [1]. Here, we show the effect in 1321N1 cells of the calmodulin inhibitors when added concurrent with, and post-stimulation of muscarinic receptors by carbachol to investigate the effect of W-13 attenuation of IP accumulation in actively signaling cells. W-13 attenuates carbachol-stimulated IP accumulation in all cellular activation states tested (Fig. 4A) showing that calmodulin inhibitors affect phosphatidylinositol phospholipid turnover when the GPCR signaling cascade is active. The addition of W-13 concurrent with carbachol showed an attenuation of IP accumulation equal to that of pre-treatment [1]. Furthermore, addition of W-13 10 min post-stimulation by carbachol inhibited any further IP accumulation resulting in values significantly lower than cells treated with vehicle alone. The effect of calmodulin inhibitor on the initial rate of PLCβ activation was investigated by pre-treating 1321N1 cells with LiCl and vehicle or W-13 for 10 min followed by carbachol stimulation for 30, 60 and 90 s. IP₃ was separated from total IP. Accumulation of IP₃ after carbachol stimulation was inhibited by W-13 at all time points (Fig. 4B), supporting a role for calmodulin in PLCβ mediated phosphatidylinositol phospholipid hydrolysis.

4. Discussion

Previously, we demonstrated direct interaction of calmodulin with PLCβ1 and PLCβ3 and the importance of calmodulin in regulating G-protein-stimulated inositol phosphate accumulation by using calmodulin inhibitors in a whole cell assay. Additionally, calmodulin binding homology domains were identified in both PLCβ1 and PLCβ3 protein sequences. In our current study, we have characterized the affinity of the direct interaction between calmodulin and PLCβ3 or PLCβ1, quantitated the effect of calcium on the interaction, defined the effects of calmodulin on PLCβ3 activity in vitro, and further investigated the effects of calmodulin inhibitors in whole cells under varying conditions of activation. Cumulatively, these

mean ± S.E.M. of three to five separate experiments performed in triplicate. Asterisk (*) indicates treatment results significantly different from vehicle at $p < 0.001$.




Fig. 3 – Protein lipid overlay. (A) PIP strips with the following membrane lipids: LPA, lysophosphatidic acid; LPC, lysophosphatidylcholine; PI, phosphatidylinositol; PI(3)P, phosphatidylinositol-3-phosphate; PI(4)P, phosphatidylinositol-4-phosphate; PI(5)P, phosphatidylinositol-5-phosphate; PE, phosphatidylethanolamine; PC, phosphatidylcholine; S(1)P, sphingo-1-phosphate; PI(3,4)P, phosphatidylinositol-3,4-phosphate; PI(3,5)P, phosphatidylinositol-3,5-phosphate; PI(4,5)P, phosphatidylinositol-4,5-diphosphate; PI(3,4,5)P, phosphatidylinositol-3,4,5-trisphosphate; PA, phosphatidic acid; PS, phosphatidylserine, and no lipid blank. PIP strips incubated with 0.5 $\mu\text{g}/\text{ml}$ of (B) PLCβ1 or (C) PLCβ3 in the absence or presence of 0.5 $\mu\text{g}/\text{ml}$ calmodulin. (D) Nitrocellulose membranes spotted with indicated amounts of the listed phospholipids (pmol) and incubated with 0.5 $\mu\text{g}/\text{ml}$ PLCβ3. PE and PS have been isolated from bovine brain. Nitrocellulose membranes and PIP strips were washed, incubated with anti-PLCβ isoenzymes, and detected by Western blotting procedures as described in Section 2. Data shown is representative of three similar experiments.

data support and expand the role for calmodulin in potentiating G-protein-stimulated PLCβ activity, particularly Gβγ-stimulated PLCβ3 activity.

Fluorescence anisotropy data demonstrated that 200 nM concentrations of both PLCβ1 and PLCβ3 bound calmodulin in the presence of 1 μM calcium. Because the number of known calmodulin binding proteins is vast and calmodulin binds proteins under both resting and calcium-activated conditions, the total amount of calmodulin in a cell is limited [31]. Therefore, it is predicted that protein must bind to calmodulin with K_d values in the sub-micromolar range to be expected to form physiologically relevant complexes [20], as found for calmodulin's interaction with PLCβ isozymes. The K_d values for binding of PLCβ1 and PLCβ3 to calmodulin are also consistent with other calmodulin binding signaling proteins such as the G-protein receptor kinases, GRK2, and 5, which have affinities of 40 nM and 2 μM , respectively, for calmodulin [32], and caldesmon which has an affinity for calmodulin of 1 μM [33]. The binding of calmodulin to PLCβ3 was sensitive to calcium concentrations over a physiological range with an apparent EC_{50} value of $<1 \mu\text{M}$ (Fig. 1), calculated using a non-saturating PLCβ3 concentration of 30 nM. This apparent EC_{50} is necessarily greater than the true EC_{50} of the complex

because saturating concentrations of PLCβ3 were not achievable with limited amounts of purified protein available. The profile of Alexa-calmodulin anisotropy versus calcium concentration would be concentration dependent until PLCβ3 concentration was greater than the K_d . However, these data suggest that increasing cytosolic calcium concentrations following initial activation of PLCβ enzyme may support further activation by increasing association with calmodulin and potentiating Gβγ-stimulated PLCβ activity. Additionally the calcium dependence of the PLCβ3 isozyme is within the range of other known calcium/calmodulin dependent proteins with calcium dependent EC_{50} values of 0.3 and 3 μM for GRK1 and GRK5, respectively [34].

Previous work in this lab with fragments from PLCβ3 protein suggested that the amino-terminal portion of the protein was a Ca^{2+} -independent calmodulin binding peptide. However, precipitation of full-length PLCβ from whole cell lysates with calmodulin-sepharose beads at physiologic calcium concentrations demonstrated that calmodulin and PLCβ can bind in the presence of calcium [1]. Clearly, the anisotropy data in this paper demonstrate that full-length PLCβ1 and PLCβ3 bind calmodulin in a Ca^{2+} sensitive manner. The EC_{50} value of 1 μM for calcium dependence of PLCβ3




Fig. 4 – Calmodulin antagonist effects on carbachol-stimulated inositol phosphate accumulation under varying conditions of cell activation. (A) 1321N1 cells were stimulated for 20 min with 1 mM carbachol in the presence of 10 mM LiCl. Additionally, cells were either treated with dH₂O (vehicle) or treated with 100 μ M W-13 for 20 min concurrent with carbachol addition (concurrent), or treated for 10 min with 100 μ M W-13 10 min after carbachol addition (post-Cch). PLC β activity was measured as percent of [3H]PIP_n converted to IP_n in whole cells, collected and quantitated as described in Section 2. Shown is cumulative data from three experiments performed in triplicate. **(B)** 1321N1 cells were pre-treated with dH₂O (vehicle) or with 100 μ M W-13 for 15 min. Cells were stimulated for 30, 60 and 90 s with 1 mM carbachol in the presence of 10 mM LiCl. PLC β activity was measured as percent of [3H]PIP_n converted to IP₃ in whole cells, collected and quantitated as described in Section 2. Shown is cumulative data from two experiments performed in triplicate. Asterisk (*) indicates W-13 pre-treatment results significantly different from vehicle pre-treatment at $p < 0.001$.

interaction with calmodulin, although known to be a high estimate, is within physiological ranges. One limitation of our previous study employing protein fragments is that fragments lack the complete protein context which may confer additional functionalities or structural subtleties to a domain. While the binding data published previously using fragments of the PLC β proteins identified a direct interaction between

PLC β and calmodulin, the anisotropy data presented herein should more accurately depict the calcium dependence of the full-length protein interaction. Calcium dependence is a hallmark of calmodulin binding protein interactions, and thus our current findings of calcium dependence are more expected rather than exceptional.

We have shown a direct effect of calmodulin in potentiating G $\beta\gamma$ activation of PLC β 3 (Fig. 2, panels A and C). One of the putative calmodulin binding sites of PLC β 3 overlaps with a site previously identified as important in G $\beta\gamma$ activation of PLC β [35]. Additionally, the β subunit of G $\beta\gamma$ is a known calmodulin binding protein [36]. Thus, calmodulin potentiation of PLC β 3 activation by G $\beta\gamma$ suggests the existence of a signaling complex that involves all three proteins, G $\beta\gamma$, calmodulin and PLC β 3. The ability of calmodulin to potentiate G $\beta\gamma$ activation of PLC β 3 is consistent with our data in whole cells whereby pre-treatment with calmodulin inhibitors attenuated muscarinic GPCR-stimulated inositol phosphate accumulation [1]. While muscarinic receptor-stimulated phosphatidylinositol hydrolysis in 1321N1 cells is characterized primarily as a G α_q /PLC β 1-mediated signaling event [37–39], the full response may involve G $\beta\gamma$ and PLC β 3. The potential for differential activation of PLC β 1 and PLC β 3 in varying spatial and temporal contexts is not at all well understood. Our data suggests that calmodulin is a co-factor in G $\beta\gamma$ -mediated PLC β activation that may selectively increase PLC β 3 activity concurrently with temporal increases in intracellular calcium concentration. Possible mechanisms for potentiation of G $\beta\gamma$ -stimulated PLC β activity by calmodulin include increased recruitment of G $\beta\gamma$ to PLC β 3 by calmodulin, or calmodulin potentiation of G $\beta\gamma$ -stimulated PLC β 3 hydrolytic activity by supporting the catalytic domain interaction with the membrane interface.

The role of calmodulin in regulating PLC β 1 remains elusive, despite our demonstration of direct binding. Calmodulin binding could potentiate PLC β 1 activation, independent of G $\beta\gamma$, by serving as a temporal scaffold or recruiter of other signaling regulatory proteins. The functions of calmodulin are not limited to direct effects on activity. For example, calmodulin binding of G $\beta\gamma$ sterically interferes with binding to Gai/o subunits [36]. In addition, the effect of calmodulin on GAP activities of the enzymes is unknown [40].

We determined that calmodulin does not alter the lipid selectivity profile of PLC β 3 or PLC β 1 for phospholipids bound to nitrocellulose (PIP strips[®]) in overlay assays. However, the PIP strip[®] overlay assays did reveal a previously unappreciated affinity of PLC β 3 and PLC β 1 for PIP₃. PIP₃ is a membrane phospholipid that has a well-deserved reputation as a regulatory lipid in a variety of cell signaling pathways [41–45], including a reciprocal association with calmodulin [29,46] in other protein contexts. In addition to demonstrating an affinity of PLC β 3 and PLC β 1 for PIP₃, the PIP strip[®] overlay assay also revealed that PLC β isozymes have different affinities for synthetic and wild-type lipids. The single saturated chain species of synthetic phospholipids that are available on the commercial PIP Strip[®] membranes is very different from the wide array of lengths and saturations in side-chains of lipids derived from mammalian sources that more accurately represent the composition of biological cell membranes [47]. The fatty acyl chain specificity of PLC β

isozymes has not yet been investigated, but has been reported in other proteins integral to GPCR signaling, such as RGS4 [29]. It is possible that calmodulin does contribute to lipid selectivity of PLC β s, but it is clear from the above data that chain length is an important factor in determining the role of calmodulin in lipid selectivity of PLC β isozymes.

To further investigate the role of calmodulin in PLC activity in an actively signaling environment, we revisited the whole cell assays. Previously, we determined that pre-treatment of 1321N1 cells with W-13 attenuates inositol phosphate accumulation after stimulation by carbachol. To determine if calmodulin inhibitors were disrupting a pre-activation function integral to PLC β isozymes or eliciting a more direct effect in active signaling, we investigated the effects of W-13 on phosphatidylinositol phospholipid turnover in an actively signaling environment. We observed an effect of W-13 when added concurrently with carbachol equal to that of pre-treatment with W-13. Additionally, W-13 inhibited IP accumulation when added post-activation of muscarinic receptors. These results support a role for calmodulin in active signaling of PLC β s, suggesting that calmodulin does not simply act to pre-couple proteins necessary for activation. Because there are other PLCs in this signaling environment, in addition to other targets of calmodulin, we wanted to investigate early effects on IP₃ accumulation in a time frame relevant to activation of PLC β s. W-13 attenuated IP₃ accumulation at all time points, providing additional support for the hypothesis that the inhibition of calmodulin affects PLC β signaling and not downstream pathways.

In conclusion, these studies further our understanding of the calmodulin/PLC β interaction by showing that calmodulin can potentiate activation by the G $\beta\gamma$ dimer without affecting activation by G α_q . This work begins to detail the separate regulatory systems involving G $\beta\gamma$ and G α_q , and suggests cooperativity between G $\beta\gamma$, calmodulin and PLC β 3; however, temporal and subcellular aspects of GPCR- and calcium-modulated cell signaling will also need to be better understood for a full understanding of calmodulin effects on PLC β 3-mediated signaling. Additionally, further research is required to demonstrate the importance of non-substrate phospholipids and membrane binding on the regulation of PLC β 3 activity.

Acknowledgements

This work was supported by the National Institutes of Health, National Institute of General Medical Sciences, PHS Grant GM61244 and an American Foundation for Pharmaceutical Education Predoctoral Fellowship (JSM).

REFERENCES

- [1] McCullar JS, Larsen SA, Millimaki RA, Filtz TM. Calmodulin is a phospholipase C- β interacting protein. *J Biol Chem* 2003;278(36):33708–13.
- [2] Rebecchi MJ, Pentyala SN. Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol Rev* 2000;80:1291–335.
- [3] Hwang JI, Oh YS, Shin KJ, Kim H, Ryu SH, Suh PG. Molecular cloning and characterization of a novel phospholipase C, PLC- ϵ . *Biochem J* 2005;389(Pt 1):181–6.
- [4] Song C, Hu C-D, Masago M, Kariya K.-I., Yamawaki-Kataoka Y, Shibatohe M, et al. Regulation of a novel human phospholipase C, PLC ϵ , through membrane targeting by Ras. *J Biol Chem* 2001;276(4):2752–7.
- [5] Saunders CM, Larman MG, Parrington J, Cox LJ, Royse J, Blayney LM, et al. PLC ζ : a sperm-specific trigger of Ca²⁺ oscillations in eggs and embryo development. *Development* 2002;129(15):3533–44.
- [6] Lee CH, Park D, Wu D, Rhee SG, Simon MI. Members of the Gq α subunit gene family activate phospholipase C β isozymes. *J Biol Chem* 1992;267(23):16044–7.
- [7] Venkatakrishnan G, Exton JH. Identification of determinants in the α -subunit of Gq required for phospholipase C activation. *J Biol Chem* 1996;271(9):5066–72.
- [8] Wu DQ, Lee CH, Rhee SG, Simon MI. Activation of phospholipase C by the α subunits of the Gq and G11 proteins in transfected Cos-7 cells. *J Biol Chem* 1992;267(3):1811–7.
- [9] Park D, Jhon D-Y, Lee C-W, Lee K-H, Rhee SG. Activation of phospholipase C isozymes by G protein $\beta\gamma$ subunits. *J Biol Chem* 1993;268:4573–6.
- [10] Rhee SG, Bae YS. Regulation of phosphoinositide-specific phospholipase C isozymes. *J Biol Chem* 1997;272(24):15045–8.
- [11] Lian L, Wang Y, Draznin J, Eslin D, Bennett JS, Poncz M, et al. The relative role of PLC β and PI3K γ in platelet activation. *Blood* 2005;106(1):110–7.
- [12] Xie W, Samoriski GM, McLaughlin JP, Romoser VA, Smrcka A, Hinkle PM, et al. Genetic alteration of phospholipase C β 3 expression modulates behavioral and cellular responses to μ opiods. *PNAS* 1999;96(18):10385–90.
- [13] Yang H, Shen F, Herenyiova M, Weber G. Phospholipase C (EC 3.1.4. 11): a malignancy linked signal transduction enzyme. *Anticancer Res* 1998;18(3A):1399–404.
- [14] Yue C, Dodge KL, Weber G, Sanborn BM. Phosphorylation of serine 1105 by protein kinase A inhibits phospholipase C β 3 stimulation by Galphaq. *J Biol Chem* 1998;273(29):18023–7.
- [15] Ståhlberg P, Wang S, Larsson C, Weber G, Öberg K, Gobl A, et al. Suppression of the neoplastic phenotype by transfection of phospholipase C β 3 to neuroendocrine tumor cells. *FEBS Lett* 1999;450(3):210–6.
- [16] Weber G, Friedman E, Grimmond S, Hayward N, Phelan C, Skogseid B, et al. The phospholipase C β 3 gene located in the MEN1 region shows loss of expression in endocrine tumours. *Hum Mol Genet* 1994;3(10):1775–81.
- [17] Wang S, Gebre-Medhin S, Betsholtz C, Ståhlberg P, Zhou Y, Larsson C, et al. Targeted disruption of the mouse phospholipase C β 3 gene results in early embryonic lethality. *FEBS Lett* 1998;441:261–5.
- [18] Chin D, Means AR. Calmodulin: a prototypical calcium sensor. *Trends Cell Biol* 2000;10(8):322–8.
- [19] Vetter SW, Leclerc E. Novel aspects of calmodulin target recognition and activation. *Eur J Biochem* 2003;270(3):404–14.
- [20] Black DJ, Tran QK, Persechini A. Monitoring the total available calmodulin concentration in intact cells over the physiological range in free Ca²⁺. *Cell Calcium* 2004;35(5):415–25.
- [21] Malencik DA, Anderson SR. Dityrosine formation in calmodulin: cross-linking and polymerization catalyzed by arthromyces peroxidase. *Biochemistry* 1996;35(14):4375–86.
- [22] Filtz TM, Paterson A, Harden TK. Purification G protein subunit regulation of phospholipase C- β from *Xenopus laevis* oocytes. *J Biol Chem* 1996;271:31121–6.

- [23] Singer AU, Waldo GL, Harden TK, Sondek J. A unique fold of phospholipase C-mediates dimerization and interaction with Gq. *Nat Struct Biol* 2002;9:32–6.
- [24] Paterson A, Harden TK. Expression, purification and reconstitution of recombinant phospholipase C- β isoenzymes. *Meth Neurosci* 1996;29:246–63.
- [25] Smrcka AV, Hepler JR, Brown KO, Sternweis PC. Regulation of polyphosphoinositide-specific phospholipase C activity by purified G_q. *Science* 1991;251:804–7.
- [26] Filtz TM, Li Q, Boyer JL, Nicholas RA, Harden TK. Expression of a cloned P_{2Y}-purinergic receptor that couples to phospholipase C. *Mol Pharmacol* 1994;46:8–14.
- [27] Bishop CV, Stormshak F. Nongenomic action of progesterone inhibits oxytocin-induced phosphoinositide hydrolysis and prostaglandin F₂[α] secretion in the ovine endometrium. *Endocrinology* 2006;147(2):937–42.
- [28] Runnels LW, Scarlata SF. Determination of the affinities between heterotrimeric G protein subunits and their phospholipase C- β effectors. *J Biochem* 1999;38:1488–96.
- [29] Popov SG, Krishna UM, Falck JR, Wilkie TM. Ca²⁺/calmodulin reverses phosphatidylinositol 3,4,5-trisphosphate-dependent inhibition of regulators of G protein-signaling GTPase-activating protein activity. *J Biol Chem* 2000;275(25):18962–24300.
- [30] Wang J, Arbuzova A, Hangyas-Mihalyne G, McLaughlin S. The effector domain of myristoylated alanine-rich C kinase substrate binds strongly to phosphatidylinositol 4,5-bisphosphate. *J Biol Chem* 2001;276(7):5012–9.
- [31] Persechini A, Stemmer PM. Calmodulin is a limiting factor in the cell. *Trends Cardiovasc Med* 2002;12(1):32–7.
- [32] Chuang TT, Paolucci L, De Blasi A. Inhibition of G protein-coupled receptor kinase subtypes by Ca²⁺/calmodulin. *J Biol Chem* 1996;271(45):28691–6.
- [33] Rhoads AR, Friedberg F. Sequence motifs for calmodulin recognition. *FASEB J* 1997;11(5):331–40.
- [34] Levay K, Satpaev D, Pronin A, Benovic J, Slepak VZ. Localization of the sites for Ca²⁺-binding proteins on G protein-coupled receptor kinases. *Biochemistry* 1998;37(39):13650–9.
- [35] Barr AJ, Ali H, Haribabu B, Snyderman R, Smrcka AV. Identification of a region at the N-terminus of phospholipase C- β 3 that interacts with G protein $\beta\gamma$ subunits. *Biochemistry* 2000;39:1800–6.
- [36] Liu M, Yu B, Nakanishi O, Wieland T, Simon M. The Ca²⁺-dependent binding of calmodulin to an N-terminal motif of the heterotrimeric G protein β subunit. *J Biol Chem* 1997;272:18801–7.
- [37] Lechleiter J, Hellmiss R, Duerson K, Ennulat D, David N, Clapham D, et al. Distinct sequence elements control the specificity of G protein activation by muscarinic acetylcholine receptor subtypes. *EMBO* 1990;9:4381–90.
- [38] Lechleiter J, Peralta E, Clapham D. Diverse functions of muscarinic acetylcholine receptor subtypes. *Trends Pharmacol Sci* 1989;Suppl:34–8.
- [39] Peralta EG, Ashkenazi A, Winslow JW, Ramachandran J, Capon DJ. Differential regulation of PI hydrolysis and adenylyl cyclase by muscarinic receptor subtypes. *Nature* 1988;334(6181):434–7.
- [40] Chidiac P, Ross EM. Phospholipase C- β 1 directly accelerates GTP hydrolysis by G α_q and acceleration is inhibited by G $\beta\gamma$ subunits. *J Biol Chem* 1999;274(28):19639–23699.
- [41] Ching TT, Wang DS, Hsu AL, Lu PJ, Chen CS. Identification of multiple phosphoinositide-specific phospholipases D as new regulatory enzymes for phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem* 1999;274(13):8611–7.
- [42] Pasquet J-M, Quek L, Stevens C, Bobe R, Huber M, Duronio V, et al. Phosphatidylinositol 3,4,5-trisphosphate regulates Ca²⁺ entry via Btk in platelets and megakaryocytes without increasing phospholipase C activity. *EMBO J* 2000;19:2793–802.
- [43] Le Blanc C, Mironneau C, Barbot C, Henaff M, Bondeva T, Wetzker R, et al. Regulation of vascular L-type Ca²⁺ channels by phosphatidylinositol 3,4,5-trisphosphate. *Circ Res* 2004;95(3):300–7.
- [44] Menager C, Arimura N, Fukata Y, Kaibuchi K. PIP₃ is involved in neuronal polarization and axon formation. *J Neurochem* 2004;89(1):109–18.
- [45] Czech MP. PIP₂ and PIP₃: complex roles at the cell surface. *Cell* 2000;100(6):603–6.
- [46] Ishii M, Inanobe A, Kurachi Y. PIP₃ inhibition of RGS protein and its reversal by Ca²⁺/calmodulin mediate voltage-dependent control of the G protein cycle in a cardiac K⁺ channel. *PNAS* 2002;99(7):4325–30.
- [47] Milne SB, Ivanova PT, DeCamp D, Hsueh RC, Brown HA. A targeted mass spectrometric analysis of phosphatidylinositol phosphate species. *J Lipid Res* 2005;46(8):1796–802.