



Renal organic anion transporters OAT1 and OAT3 mediate the cellular accumulation of 5-sulfooxymethylfurfural, a reactive, nephrotoxic metabolite of the Maillard product 5-hydroxymethylfurfural

Nadiya Bakhiya^a, Bernhard Monien^a, Heinz Frank^b, Albrecht Seidel^b, Hansruedi Glatt^{a,*}

^a German Institute of Human Nutrition Potsdam-Rehbrücke, Department of Nutritional Toxicology, 14558 Nuthetal, Germany

^b Biochemical Institute for Environmental Carcinogens, Prof. Dr. Gernot Grimmer Foundation, 22927 Grosshansdorf, Germany

ARTICLE INFO

Article history:

Received 30 January 2009

Accepted 16 April 2009

Keywords:

5-Hydroxymethylfurfural

5-Sulfooxymethylfurfural

Nephrotoxicity

Organic anion transporters

ABSTRACT

5-Hydroxymethylfurfural (HMF) is formed when sugars are acidified or heated. It is present at high levels in numerous foods. HMF is inactive in standard genotoxicity tests, but can be metabolized to a chemically reactive intermediate, 5-sulfooxymethylfurfural (SMF), which is mutagenic and carcinogenic. We recently found that direct parental administration of SMF to mice leads to abundant acute necrosis and proteinaceous casts in the proximal tubules as the dominating toxicological effect. Since proximal tubule cells actively mediate the excretion of many organic anions, we hypothesized that transporter-mediated uptake of SMF into the cells could be the reason for this selective organotoxicity. To test this hypothesis, we used human embryonic kidney (HEK293) cells stably expressing human (h) OAT1 or OAT3. SMF was a competitive inhibitor of *p*-aminohippurate uptake by hOAT1 and estrone sulfate uptake by hOAT3 with K_i values of 225 μ M and 1.5 mM, respectively. Moreover, the initial rates of SMF uptake were 5.2- and 3.1-fold higher in cells expressing hOAT1 and hOAT3, respectively, than in control HEK293 cells. Likewise, the sensitivity of hOAT1- and hOAT3-expressing cells to SMF cytotoxicity was significantly higher than that of control cells, and was reduced by addition of probenecid, an inhibitor of OATs. Taken together, these results indicate that OAT1 and OAT3 mediate the uptake of SMF into proximal tubule cells and thereby may be involved in SMF-induced nephrotoxicity.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

5-(Hydroxymethyl)-2-furfural (HMF) is one of the numerous compounds resulting from the heating of foods. It is formed when reducing hexose moieties are heated in the presence of amino acids or proteins (Maillard reaction) [1] and via direct thermal dehydration of fructose, sucrose and glucose [2]. HMF is present at high levels in many foods, with particularly high concentrations found in caramel products (up to 9.5 g/kg), dried fruit (especially plums), old Port and Madeira wines, and balsamic vinegar [3–5]. However, coffee may be a major source of HMF intake in many subjects due to the high amount consumed [5]. The estimated daily intake is in the range of 30–150 mg per person [6,7]. Although HMF is inactive in standard in vitro genotoxicity tests [7,8], it induced and promoted preneoplastic lesions, aberrant crypt foci, in rat colon [9] and initiated papillomas in mouse skin [10]. Furthermore,

in a long-term carcinogenicity study conducted by National Toxicology Program, HMF was found to increase the incidence of hepatocellular adenomas in female mice [11].

Numerous chemicals are mutagenic or carcinogenic after metabolic activation to chemically reactive intermediates. HMF is converted to a chemically reactive allylic ester, 5-sulfooxymethylfurfural (SMF, structural formula depicted in Fig. 1) by sulfotransferases (SULT), enzymes ignored in standard in vitro genotoxicity tests. HMF showed mutagenic activity when tested in bacterial and mammalian target cells engineered for expression of human SULT1A1 [8]. SMF, unlike HMF, formed DNA adducts in cell-free systems and was mutagenic to bacterial and mammalian cells without requiring an activating system [8,12]. When administered topically to mouse skin, SMF initiated papillomas with higher activity than HMF [10].

In a recent study conducted in our laboratory SMF was found to be strongly nephrotoxic in the mouse in vivo (F. Taugner, S. Florian and H. R. Glatt, manuscript in preparation). Male FVB/N mice which received a single dose of SMF (250 mg SMF/kg body mass, i.p.) died or were moribund 5–11 days after the treatment. Histopathological analyses revealed that SMF induced moderate damage in liver and most impressive damage in the kidneys, particularly in the proximal tubules, where abundant necrosis and proteinaceous

Abbreviations: HMF, 5-hydroxymethylfurfural; hOAT, human organic anion transporter; OAT, organic anion transporter; SMF, 5-sulfooxymethylfurfural; SULT, sulfotransferase; UPLC, ultra-performance liquid chromatography.

* Corresponding author. Tel.: +49 33200 88321; fax: +49 33200 88426.

E-mail address: glatt@dife.de (H. Glatt).




Fig. 1. Structural formula of 5-sulfooxymethylfurfural (SMF).

casts were observed. While nearly all proximal tubules in SMF-exposed animals were destroyed, distal tubules and glomeruli were less affected. The molecular mechanism underlying this selective toxicity of SMF in proximal tubules is unknown.

An important physiological function of the proximal tubule is the secretion of a large variety of organic anions, including endogenous catabolites, drugs and other xenobiotics [13]. The organic anion transporters OAT1 and OAT3 are localized at the basolateral membrane of proximal tubule cells and mediate the concentrative uptake of their substrates from blood into renal proximal tubule cells, which is the rate-limiting step in this excretion system [14]. Genes of both transporters were cloned from several species, including rat and man, and are highly conserved across species [14,15]. Both transporters show overlapping substrate specificities and share the driving force – exchange of organic anions against intracellular α -ketoglutarate. Moreover, the activity of OAT1 and OAT3 has been associated with proximal tubular injury due to the accumulation of toxicants, such as uremic toxins, mercuric species and ochratoxin A [16], as well as genotoxic benzylic sulfates formed from alkylated polycyclic aromatic hydrocarbons [17].

The aim of the present investigation was to determine the possible involvement of OAT1 and OAT3 in the uptake, accumulation and toxicity of SMF in renal cells. To this end, we used human embryonic kidney cells (HEK293) stably expressing human (h) OAT1 or OAT3 to assess both uptake and toxicity of SMF.

2. Materials and methods

2.1. Chemicals

5-Hydroxymethylfurfural, sulfur trioxide–pyridine complex, sodium methoxide, and the anhydrous solvents tetrahydrofuran, diethyl ether, and dimethylformamide were purchased from Sigma–Aldrich (Taufkirchen, Germany) and used as received. All solvents for UPLC were purchased from Carl Roth GmbH (Karlsruhe, Germany). Cell culture reagents were from Gibco Invitrogen (Karlsruhe, Germany), unless specified otherwise. [6,7- ^3H (N)]Estrone sulfate (57.3 Ci/mmol) and [glycyl-2- ^3H]p-aminohippurate (4.18 Ci/mmol) were purchased from Perkin-Elmer (Boston, USA). The specific radioactivity of [^3H]p-aminohippurate was reduced to 0.5 Ci/mmol by adding unlabeled p-aminohippurate. [Methyl- ^3H]thymidine (20 Ci/mmol) was purchased from MP Biomedicals (Illkirch Cedex, France).

NMR spectra were obtained on a Bruker AV400 spectrometer at the University of Hamburg. Melting points (mp) were determined on a Büchi 510 apparatus.

2.2. Synthesis of SMF

5-Hydroxymethylfurfural (0.032 g; 0.25 mmol) was dissolved in 1 ml anhydrous tetrahydrofuran and this solution was cooled to 0 °C. After addition of a solution of 80 mg sulfur trioxide–pyridine complex (0.5 mmol) in 0.65 ml anhydrous *N,N*-dimethylformamide the mixture was stirred at 0 °C for 4 h. Then a solution of 40 mg (0.74 mmol) sodium methoxide in 0.35 ml methanol was added and the resulting mixture was centrifuged. The supernatant was added dropwise to 80 ml of anhydrous diethyl ether. The formed precipitate was filtered off, washed with anhydrous diethyl

ether, and dried under vacuum to give a yield of 0.043 g SMF (75% yield); mp 128 °C (dec); ^1H NMR (400 MHz, DMSO- d_6) δ [ppm] 9.63 (s, 1, CHO), 7.54 (d, 1, H_3 , $J_{3,4} = 3.6$ Hz), 6.75 (d, 1, H_4 , $J_{4,3} = 3.6$ Hz), 4.83 (s, 2, $-\text{CH}_2-\text{O}$). The purity of SMF was greater than 99% as determined by liquid chromatography–mass spectrometry (see Section 2.5).

2.3. Tissue culture

hOAT1 and hOAT3 were stably expressed in the human embryonic kidney cell line HEK293 as described previously [17]. Cells were grown in flasks containing Dulbecco's modified minimum essential medium (high glucose) supplemented with fetal bovine serum (10%, Biochrom AG, Berlin, Germany), penicillin (100 units/ml) and streptomycin (100 $\mu\text{g}/\text{ml}$). The medium for the recombinant cells but not for parental HEK293 cells, additionally contained hygromycin (175 $\mu\text{g}/\text{ml}$). Cultures were maintained in a humidified atmosphere containing 5% CO_2 at 37 °C. Cultures were split in a 1:5 ratio every third to fourth day.

2.4. Inhibition of uptake of standard substrates

Cells were seeded in 24-well plates (2×10^5 cells in 1 ml medium per well) 2 days before the experiment. Uptake of [^3H]p-aminohippurate (for OAT1) and [^3H]estrone sulfate (for hOAT3) was assayed at 37 °C in Ringer's solution (130 mM NaCl, 4 mM KCl, 1 mM CaCl_2 , 1 mM MgSO_4 , 20 mM HEPES, 1 mM NaH_2PO_4 , 18 mM glucose, pH 7.4) for 2 min in the absence or presence of SMF. The uptake was terminated by three washes with 0.5 ml of ice-cold Ringer's solution. Cells were then solubilized in 0.5 ml of 1N NaOH. After neutralization with 0.5 ml of 1N HCl, their ^3H content was assayed by liquid scintillation counting. The results were standardized to the level of protein determined using the bicinchoninic acid assay (Pierce, Rockford, USA) with bovine serum albumin as the standard. For quantifying the inhibitory activity, we used varying concentrations of the substrate (10 and 50 μM [^3H]p-aminohippurate for hOAT1; 50 and 250 nM [^3H]estrone sulfate for hOAT3) and SMF (≥ 5 concentrations, up to 1 mM for hOAT1 and 2 mM for hOAT3). Data were plotted and analyzed according to Dixon [18].

2.5. LC–MS/MS analysis of SMF uptake into cells

Cells were incubated in Ringer's solution containing 250 μM SMF at room temperature for 1–15 min. After three washes with ice-cold Ringer's solution, the cells were lysed with 0.25 ml 1N NaOH and, after neutralization with 0.25 ml of 1N HCl and protein precipitation with 1 ml of ice-cold 2-propanol, mixed thoroughly and centrifuged at $15,000 \times g$ for 15 min. The clear supernatant was analyzed using an Acquity ultra-performance liquid chromatography (UPLC) System (Waters, Milford, MA, USA) with a UPLC BEH Phenyl column (1.7 μm , 2.1 mm \times 100 mm, Waters). Samples of 4 μl were injected, and SMF was eluted with 10 mM ammonium acetate/methanol (95:5) at 0.35 ml/min flow rate. The UPLC was connected to a Quattro Premier XE tandem quadrupole mass spectrometer (Waters Micromass, Manchester, UK) with an electrospray interface operated in the negative ion mode. Two fragmentation reactions of SMF were monitored, yielding the sulfate ion radical ($204.9 \rightarrow 96$) and the protonated sulfonate ion ($204.9 \rightarrow 81$). The tune parameter were as follows: temperature of the electrospray source 120 °C; desolvation temperature 500 °C; desolvation gas: nitrogen (850 l/h); cone gas: nitrogen (100 l/h); collision gas: argon (indicated cell pressure $\sim 5 \times 10^{-3}$ mbar). For the fragmentation of SMF, collision energies were 18 and 22 eV for the transitions $204.9 \rightarrow 81$ and $204.9 \rightarrow 96$, respectively. The dwell time was set to 100 ms, and capillary voltage was set to 0.35 kV.

The cone and RF1 lens voltages were 32 and 0.1 V, respectively. SMF was quantified using an external calibration line prepared in water/2-propanol (1:3). The calibration curves were linear in the range between 500 nM (2 pmol/injection) and 1 nM (4 fmol/injection) SMF ($r^2 \geq 0.993$). Details of the method have been published elsewhere [22].

2.6. [^3H]Thymidine-incorporation proliferation assay

Cells (1×10^5 cells per well) were seeded in 24-well plate 24 h before starting exposure. Then, they were incubated at 37 °C for 1 h in the absence or presence of various concentrations of SMF (0.1–1.6 mM). After the incubation, cells were washed twice with medium. [^3H]Thymidine (0.2 $\mu\text{Ci}/\text{ml}$ in culture medium) was incorporated at 37 °C for 1 h. The cells were fixed with 5% trichloroacetic acid (30 min at 4 °C), washed twice with ice-cold PBS and once with ice-cold 95% ethanol to remove unused [^3H]thymidine. For the solubilization of the genomic DNA, the cells were incubated with 0.5 ml of 1N NaOH overnight. After neutralization with 0.5 ml of 1N HCl, the ^3H content was assayed by liquid scintillation counting.

3. Results

3.1. Inhibition of hOAT1 and hOAT3 by SMF

SMF was tested for its ability to inhibit hOAT-mediated uptake of model substrates, *p*-aminohippurate for hOAT1 and estrone sulfate for hOAT3 in recombinant embryonic kidney cells. As shown in Fig. 2, SMF inhibited substrate uptake by both hOATs. The effect increased with the concentration of SMF. The sensitivities of hOAT1- and hOAT3-mediated substrate uptakes to SMF inhibition were different. At a concentrations of 0.5 and 1 mM, SMF inhibited *p*-aminohippurate (10 μM) uptake by hOAT1-expressing cells by 68% and 79%, respectively (Fig. 2A). In the presence of the same SMF concentrations (0.5 and 1 mM) estrone sulfate (50 nM) uptake by hOAT3-cells was inhibited somewhat less (by 27% and 45%, respectively, Fig. 2B).

Subsequently, we studied the inhibition kinetics of SMF using different concentrations of the model substrates and SMF. The K_i values were determined from Dixon plots. Representative plots are shown in Fig. 3. Similar results were obtained in repeat experiments. SMF competitively inhibited the organic anion

transport mediated by both transporters. The mean K_i value ($\pm\text{S.E.}$) of SMF for hOAT1 was determined at $225 \pm 22 \mu\text{M}$ (245, 248 and 182 μM in three experiments). The corresponding value for hOAT3 was $1.46 \pm 0.04 \text{ mM}$ (1.38, 1.53 and 1.47 mM in three experiments).

3.2. hOAT1- and hOAT3-mediated uptake of SMF into cells

To test whether SMF is transported by hOAT, we exposed parental HEK293 and transporter-expressing cells to SMF (250 μM) for 1–15 min and then analyzed the intracellular SMF content by UPLC–MS/MS. The analytical method used was highly specific and sensitive (limit of detection $\sim 5 \text{ fmol}$ per injection). As shown in Fig. 4A, the uptake was linear with time for at least 4–10 min in all three cell lines. The initial slopes amounted to 1.3, 6.6 and 3.9 nmol/mg protein/min for control, hOAT1- and hOAT3-expressing cells, respectively. Thus, expression of hOAT1 and hOAT3 enhanced the uptake rates by a factor of 5.2 and 3.1, respectively. Moreover, the SMF uptake by hOAT-expressing cells was inhibited by addition of probenecid (a known inhibitor of OAT) to the incubation medium, indicating that this uptake was hOAT-mediated (Fig. 4B).

3.3. Influence of hOAT1 and hOAT3 expression on cellular toxicity of SMF

As shown in Fig. 5, hOAT1- and hOAT3-expressing cells were markedly more sensitive than control cells to toxic effects of SMF. The difference was significant for both transporters starting from 0.2 mM SMF concentration. The enhanced cytotoxicity of hOAT-expressing cells to SMF was directly related to the activity of the expressed transporters, as demonstrated using probenecid, an inhibitor of OATs. Probenecid markedly reduced of the cytotoxicity of SMF (0.4 mM) in the hOAT-expressing cells (Fig. 6). Survival of hOAT1-expressing cells was elevated by probenecid from 49% to 71%. Similarly, survival of hOAT3-expressing cells was increased from 55% in the absence to 74% in the presence of probenecid.

4. Discussion

HMF can be activated to a chemically reactive and genotoxic metabolite, SMF. In a recent in vivo study we detected strong toxicity of SMF to renal proximal tubule cells leading to death in




Fig. 2. Inhibition of human OAT-mediated substrate uptake by SMF. Parental HEK293 cells (open bars) and HEK293-derived cells expressing hOAT1 or hOAT3 (solid bars) were assayed in the presence of SMF (500 μM and 1 mM) and in its absence for uptake of [^3H] *p*-aminohippurate (10 μM ; A) and [^3H] estrone sulfate (50 nM; B). The treatment was performed at 37 °C for 2 min. Data are expressed as a percentage of the basal uptake (in the absence of SMF) of the OAT-expressing cells in the corresponding experiment. Values are means and S.E. of three experiments conducted on separate occasions. The absolute values of the basal uptake ($\pm\text{S.E.}$) amounted to 178 ± 12 and $1.60 \pm 0.14 \text{ pmol}/\text{mg}$ protein in 2 min for hOAT1 (*p*-aminohippurate) and hOAT3 (estrone sulfate), respectively. * $p < 0.05$; ** $p < 0.01$, compared to the same cells in the absence of SMF (paired Student's *t*-test using the absolute uptake as the statistical unit).




Fig. 3. Kinetic analyses of the influence of SMF on organic anion uptake by human OAT. Uptake of [³H]p-aminohippurate (10 μM or 50 μM) and [³H]estrone sulfate (50 or 250 nM) was assayed in HEK293-derived cell lines stably expressing hOAT1 (A) and hOAT3 (B), respectively, in the presence of varying concentrations of SMF for 1 min at 37 °C. Uptake was also determined in control cells. This value was subtracted from that observed in OAT-expressing cells. Data are presented as Dixon plots. Values are means ± S.E. of three wells from a single representative experiment. The following K_i values were calculated from the plots: 248 μM for hOAT1 and 1530 μM for hOAT3. Similar values were obtained in repeat experiments (see main text).

most animals after a single dose of 250 mg/kg. At the next lower dose (125 mg/kg), animals developed an atypical renal hyperplasia. The proximal tubule cells are the primary sites of OAT1 and OAT3 expression, and previous studies suggest that these transporters could participate in accumulation of reactive sulfo conjugates in this tissue.

The data reported here provide molecular evidence that hOAT1 and hOAT3 can mediate the uptake of SMF into renal cells. Both uptake and toxicity of SMF were significantly higher in HEK293 cells expressing hOAT1 or hOAT3 than in control cells lacking these transporters. Furthermore, both uptake and toxicity were reduced by the OAT inhibitor probenecid, indicating that SMF is a substrate of hOAT1 and hOAT3. Therefore, OAT1 and OAT3 seem to provide specific paths for the entry of SMF into renal cells, leading to proximal tubule damage. The affinities of SMF interaction with hOAT1 and hOAT3, measured in the inhibition study, were considerably different. The K_i value of hOAT1 for SMF was in the range of 225 μM, while the affinity of the hOAT3 to SMF was approximately seven times lower (K_i value ~ 1.5 mM), suggesting that this transporter contributes less to cellular SMF uptake than hOAT1.

As shown in rodents, HMF is well absorbed from the gastrointestinal tract, rapidly metabolized, and excreted primarily in urine. The principal urinary metabolites are 5-hydroxymethyl-2-furoic acid and its glycine conjugate and furan-2,5-dicarboxylic

acid [19–21]. SMF was not detected in urine. However, SMF is chemically reactive, making its detection in biological samples difficult. The half-life time of SMF in aqueous solutions at 37 °C amounts to nearly 2 h. Recent studies conducted in our laboratory demonstrate the formation of SMF from HMF in the mouse in vivo [22]. SMF was detected in blood samples of FVB/N mice after intravenous administration of HMF. From a dose of 100 mg HMF/kg body mass, approximately 0.04% HMF was metabolized to SMF and exported into the circulation; additional SMF may have been formed and reacted with cellular structures or hydrolyzed back to HMF at its site of formation not reaching the circulation. Several human SULT forms catalyze the sulfo conjugation of HMF with higher catalytic efficiency than murine enzymes; this is true in particular for SULT1A1, the most abundant SULT form in the human organism [8]. Therefore, it may be speculated that a higher percentage of HMF is converted to SMF in humans than in mice. Taking into consideration that the daily intake of HMF exceeds that of other food-borne pro-genotoxins, such as acrylamide or heterocyclic amines, by factors of 10³ and 10⁵, respectively, conversion of a minute fraction to a reactive metabolite still could be toxicologically relevant.

Recently HMF was orally administered to B6C3F₁ mice and F344/N rats to study its subacute, subchronic and chronic toxicity [11]. In a 3-month study, the incidence of cytoplasmic alterations of the kidney was significantly increased in male mice receiving




Fig. 4. Uptake of SMF by human OAT-expressing cells. (A) Control HEK293 cells and HEK293-derived cells that express hOAT1 or hOAT3 were exposed to 250 μM SMF for varying time periods. After washing and lysing the cells, the intracellular levels of SMF were determined using UPLC–MS/MS. Each value represents the mean ± S.E. of six cell cultures. (B) Control and hOAT-expressing cells were exposed to 250 μM SMF for 10 min in the absence or presence of OAT inhibitor probenecid (1 mM). Data are expressed as a percent of the uptake into control (parental HEK293) cells in the absence of probenecid. Values are means ± S.E. of three experiments conducted on separate occasions. The absolute uptake of SMF into control cells amounted to 18.7 ± 4.1 nmol/mg protein in 10 min. **p* < 0.05 compared to control cells in the absence of probenecid (paired Student's *t*-test).




Fig. 5. Cytotoxicity of SMF in control and human OAT-expressing cells. HEK293-derived cells were treated with varying concentrations of SMF for 1 h. Cell proliferation was assayed by the incorporation of [3 H]thymidine (1 h). Data are expressed as a percentage of [3 H]thymidine incorporation of the corresponding cell line in the absence of SMF. Values are means of three independent experiments \pm S.E. * p < 0.05, ** p < 0.01 (Student's *t*-test) in comparison with control cells exposed to the same level of SMF.

188 or 375 mg/kg HMF. In a 2-year carcinogenicity study, HMF showed evidence of carcinogenic activity, clearly increasing the incidence of hepatocellular adenomas in female mice receiving 188 or 375 mg/kg HMF. Thus, SMF formed in vivo in mice seems to cause the main damage at the site of its formation, the liver; probably only a minor part of SMF is exported to the blood. The situation in humans could be different, taking into consideration the species-dependent differences in tissue distribution of HMF-activating SULTs, which are focused to the liver in mice, but widely expressed in numerous tissues in humans [23,24].

Furfuryl alcohol is another common Maillard product. It only differs from HMF by the absence of the aldehyde group in 2 position. Like HMF, furfuryl alcohol is also activated to a mutagenic sulfo conjugate, furfuryl sulfate, by human SULTs [8]. Two-year carcinogenicity studies in male and female Fischer 344/N rats and B6C3F₁




Fig. 6. Effect of co-incubation of probenecid, an OAT inhibitor, with SMF on the proliferation of control and hOAT-expressing cells. HEK293-derived cells were treated with 400 μ M SMF in absence or the presence of 1 mM probenecid for 1 h. Cell proliferation was assayed by incorporation of [3 H]thymidine (1 h). Data are expressed as a percentage of [3 H]thymidine incorporation of the corresponding cell line in the absence of SMF. Values represent means \pm S.E. of three independent experiments. * p < 0.05 (Student's *t*-test) in comparison with corresponding cells without probenecid.

mice had been conducted with furfuryl alcohol administrated by inhalation [25]. Notably, the incidence of renal tubule neoplasms was increased in male mice exposed to furfuryl alcohol, and the severity of nephropathy (typical for old mice and rats) was enhanced in both sexes of the rat as well as in male mice. We therefore suspect that the renal toxicity and carcinogenicity of furfuryl alcohol was mediated by its metabolite furfuryl sulfate, which is structurally related to SMF and thus could also be accumulated in the renal tubule cells by organic anion transporters.

Findings with some other chemicals indicate that kidneys might be a relatively common target tissue of reactive sulfo conjugates. 1-Sulfooxymethylpyrene is an ultimate mutagen and carcinogen formed from an abundant polycyclic aromatic hydrocarbon, 1-methylpyrene. After treating rats with 1-sulfooxymethylpyrene or its metabolic precursor, 1-hydroxymethylpyrene, the highest level of DNA adducts was detected in kidney [26]. Further in vitro studies showed that 1-sulfooxymethylpyrene is an excellent substrate for OAT1 and OAT3 [17].

Renal cell carcinomas account for approximately 3% of all adult malignancies [27]. The clear cell carcinoma, arising from proximal tubular cells, makes up 70–80% of renal neoplasias. The incidence of renal cell carcinomas varies more than 10-fold between different countries and has been increasing in various countries, suggesting an involvement of exogenous factors. It would be interesting to know whether, and to which extent, HMF, furfuryl alcohol, alkylated polycyclic aromatic hydrocarbons and other SULT-dependent promutagens contribute to the induction of clear cell carcinomas.

In conclusion, in the present study we provide evidence for the involvement of organic anion transporters (OATs) in the renal accumulation of SMF. These transport characteristics could be responsible for the selective damage of renal proximal tubules by this reactive metabolite.

Acknowledgements

We thank Christine Gumz for her skillful technical assistance. This work has been carried out with support from the European Commission, Priority 5 on Food Quality and Safety (Contract no. FOOD-CT-2003-506820 Specific Targeted Project), 'Heat-generated food toxicants-identification, characterisation and risk minimisation'. A.S. and H.F. gratefully acknowledge financial support from ECNIS (Environmental Cancer Risk, Nutrition and Individual Susceptibility), a network of excellence operating within the European Union 6th Framework Program, Priority 5: 'Food Quality and Safety' (Contract No. 513943).

References

- [1] Mauron J. The Maillard reaction in food; a critical review from the nutritional standpoint. *Prog Food Nutr Sci* 1981;5:5–35.
- [2] Antal Jr MJ, Mok WS, Richards GN. Mechanism of formation of 5-(hydroxymethyl)-2-furaldehyde from D-fructose and sucrose. *Carbohydr Res* 1990;199: 91–109.
- [3] Bachmann S, Meier M, Känzig A. 5-Hydroxymethyl-2-furfural (HMF) in Lebensmitteln. *Lebensmittelchemie* 1997;51:49–50.
- [4] Murkovic M, Pichler N. Analysis of 5-hydroxymethylfurfural in coffee, dried fruits and urine. *Mol Nutr Food Res* 2006;50:842–6.
- [5] Husøy T, Haugen M, Murkovic M, Jöbstl D, Stølen LH, Bjellaas T, et al. Dietary exposure to 5-hydroxymethylfurfural from Norwegian food and correlations with urine metabolites of short-term exposure. *Food Chem Toxicol* 2008;46: 3697–702.
- [6] Ulbricht RJ, Northup SJ, Thomas JA. A review of 5-hydroxymethylfurfural (HMF) in parenteral solutions. *Fundam Appl Toxicol* 1984;4:843–53.
- [7] Janzowski C, Glaab V, Samimi E, Schlatter J, Eisenbrand G. 5-Hydroxymethylfurfural: assessment of mutagenicity. DNA-damaging potential and reactivity towards cellular glutathione. *Food Chem Toxicol* 2000;38:801–9.
- [8] Glatt HR, Sommer Y. Health risks by 5-hydroxymethylfurfural (HMF) and related compounds. In: Skog K, Alexander J, editors. *Acrylamide and other health hazardous compounds in heat-treated foods*. Cambridge: Woodhead Publishing; 2006. p. 328–57.

- [9] Zhang XM, Chan CC, Stamp D, Minkin S, Archer MC, Bruce WR. Initiation and promotion of colonic aberrant crypt foci in rats by 5-hydroxymethyl-2-furfuraldehyde in thermolyzed sucrose. *Carcinogenesis* 1993;14:773–5.
- [10] Surh YJ, Liem A, Miller JA, Tannenbaum SR. 5-Sulfooxymethylfurfural as a possible ultimate mutagenic and carcinogenic metabolite of the Maillard reaction product, 5-hydroxymethylfurfural. *Carcinogenesis* 1994;15:2375–7.
- [11] NTP. Toxicology and carcinogenesis studies of 5-(hydroxymethyl)-2-furfural (CAS No. 67-47-0) in F344/N rats and B6C3F₁ mice (gavage studies). Natl Toxicol Program Tech Report Series 2008 (draft available at <http://ntp.niehs.nih.gov/>).
- [12] Surh YJ, Tannenbaum SR. Activation of the Maillard reaction product 5-(hydroxymethyl)furfural to strong mutagens via allylic sulfonation and chlorination. *Chem Res Toxicol* 1994;7:313–8.
- [13] Russel FG, Masereeuw R, van Aubel RA. Molecular aspects of renal anionic drug transport. *Annu Rev Physiol* 2002;64:563–94.
- [14] Burckhardt BC, Burckhardt G. Transport of organic anions across the basolateral membrane of proximal tubule cells. *Rev Physiol Biochem Pharmacol* 2003;146:95–158.
- [15] Robertson EE, Rankin GO. Human renal organic anion transporters: characteristics and contributions to drug and drug metabolite excretion. *Pharmacol Ther* 2006;109:399–412.
- [16] Sweet DH. Organic anion transporter (Slc22a) family members as mediators of toxicity. *Toxicol Appl Pharmacol* 2005;204:198–215.
- [17] Bakhiya N, Stephani M, Bahn A, Ugele B, Seidel A, Burckhardt G, et al. Uptake of chemically reactive, DNA-damaging sulfuric acid esters into renal cells by human organic anion transporters. *J Am Soc Nephrol* 2006;17:1414–21.
- [18] Dixon M. The determination of enzyme inhibitor constants. *Biochem J* 1953;55:170–1.
- [19] Jellum E, Borresen HC, Eldjarn L. The presence of furan derivatives in patients receiving fructose-containing solutions intravenously. *Clin Chim Acta* 1973;47:191–201.
- [20] Germond JE, Philippoussian G, Richli U, Bracco I, Arnaud MJ. Rapid and complete urinary elimination of [¹⁴C]-5-hydroxymethyl-2-furfuraldehyde administered orally or intravenously to rats. *J Toxicol Environ Health* 1987;22:79–89.
- [21] Godfrey VB, Chen LJ, Griffin RJ, Lebetkin EH, Burka LT. Distribution and metabolism of (5-hydroxymethyl)furfural in male F344 rats and B6C3F₁ mice after oral administration. *J Toxicol Environ Health A* 1999;57:199–210.
- [22] Monien BH, Frank H, Seidel A, Glatt HR. Conversion of the common food constituent 5-hydroxymethylfurfural into a mutagenic and carcinogenic sulfuric acid ester in the mouse *in vivo*. *Chem Res Toxicol*. 2009 Apr 22. [Epub ahead of print].
- [23] Glatt HR. Sulphotransferases. In: Ioannides C, editor. *Handbook of enzyme systems that metabolise drugs and other xenobiotics*. Sussex: John Wiley & Sons; 2002. p. 353–439.
- [24] Teubner W, Meinel W, Florian S, Kretzschmar M, Glatt H. Identification and localization of soluble sulfotransferases in the human gastrointestinal tract. *Biochem J* 2007;404:207–15.
- [25] NTP. Toxicology and carcinogenesis studies of furfuryl alcohol (CAS No. 98-00-0) in F344/N rats and B6C3F₁ mice (inhalation studies). Natl Toxicol Program Tech Report Series 1999:1–248.
- [26] Glatt HR, Meinel W, Kuhlow A, Ma L. Metabolic formation, distribution and toxicological effects of reactive sulphuric acid esters. *Nova Acta Leopoldina* 2003;329:151–61.
- [27] Martel CL, Lara PN. Renal cell carcinoma: current status and future directions. *Crit Rev Oncol Hematol* 2003;45:177–90.