



# Genistein and daidzein prevent low potassium-dependent apoptosis of cerebellar granule cells

Anna Atlante<sup>a</sup>, Antonella Bobba<sup>a</sup>, Gianluca Paventi<sup>b</sup>, Roberto Pizzuto<sup>c</sup>, Salvatore Passarella<sup>c,\*</sup>

<sup>a</sup> Istituto di Biomembrane e Bioenergetica, CNR, Via G. Amendola 165/A, 70126 Bari, Italy

<sup>b</sup> Dipartimento di Scienze Animali, Vegetali e dell'Ambiente, Università del Molise, via De Sanctis, 86100 Campobasso, Italy

<sup>c</sup> Dipartimento di Scienze per la Salute, Università del Molise, via De Sanctis, 86100 Campobasso, Italy

## ARTICLE INFO

### Article history:

Received 25 July 2009

Accepted 6 October 2009

### Keywords:

Genistein

Daidzein

Apoptosis

Cerebellar granule cells

Antioxidant

Mitochondria

## ABSTRACT

We have investigated the ability of certain dietary flavonoids, known to exert beneficial effects on the central nervous system, to affect neuronal apoptosis. We used cerebellar granule cells undergoing apoptosis due to potassium deprivation in a serum-free medium in either the absence or presence of the flavonoids genistein and daidzein, which are present in soy, and of catechin and epicatechin, which are present in cocoa. These compounds were used in a blood dietary concentration range. We found that genistein and daidzein, but not catechin and epicatechin, prevented apoptosis, with cell survival measured 24 h after the induction of apoptosis being higher than that of the same cells incubated in flavonoid free medium (80% and 40%, respectively); there was no effect in control cells. A detailed investigation of the effect of these compounds on certain mitochondrial events that occur in cells *en route* to apoptosis showed that genistein and daidzein prevented the impairment of glucose oxidation and mitochondrial coupling, reduced cytochrome c release, and prevented both impairment of the adenine nucleotide translocator and opening of the mitochondrial permeability transition pore. Interestingly, genistein and daidzein were found to reduce the levels of reactive oxygen species, which are elevated in cerebellar granule cell apoptosis. These findings strongly suggest that the prevention of apoptosis depends mainly on the antioxidant properties of genistein and daidzein. This could lead to the development of a flavonoid-based therapy in neuropathies.

© 2009 Elsevier Inc. All rights reserved.

## 1. Introduction

Deregulated apoptotic mechanisms have been implicated in many pathologic conditions, including AIDS, infectious diseases, inflammation, cancer, heart failure, osteoporosis, stroke and trauma. In particular, the process is involved in many human neurological

disorders, including Alzheimer's, Parkinson's and Huntington's diseases and amyotrophic lateral sclerosis [1]. It is intriguing that key events/components in the cellular regulation of apoptosis have been identified and thus may be targeted by therapeutic strategies [2]. In this regard, targeting apoptosis with dietary bioactive agents could be a therapeutic approach to either prevent apoptosis in those diseases that have an apoptotic origin [3] or induce apoptosis, for instance in cancer cells. To better investigate whether and how some dietary compounds can affect apoptosis, the availability of an experimental system is needed in which a dissection of the steps that lead to cell death has already been done.

We have already shown how apoptosis of cerebellar granule cells (CGCs) occurs when they are deprived of extracellular K<sup>+</sup> (SK5 cells) (for Ref., see [4–9]). In particular, we investigated the role of mitochondria in this process. Briefly, in early apoptosis (0–3 h after induction) the rate of glucose oxidation by CGCs decreases [5], mitochondria are subjected to time-dependent uncoupling [5] and elevated production of reactive oxygen species (ROS) occurs [6]. Cytochrome c (cyt c) is released from the mitochondria while still coupled [6,7] and an increase in the ATP level occurs [8]. In late apoptosis (3–8 h after induction), an alteration of the adenine nucleotide translocator (ANT) occurs, with ANT becoming a

**Abbreviations:** Act D, actinomycin D; ANT, adenine nucleotide translocator; Ap5A, P<sub>1</sub>P<sub>5</sub>-di(adenosine-5')penta-phosphate; ASC, ascorbate; ATP D.S., ATP detecting system; ATR, atractyloside; BME, basal medium Eagle; CE, catechin; CGC, cerebellar granule cell; CN<sup>−</sup>, potassium cyanide; CsA, cyclosporine A; Cyt c, cytochrome c; DIV, days *in vitro*; DMSO, dimethyl sulfoxide; DZN, daidzein; DZN-S-K5 cells, S-K5 cells treated with daidzein; EC, epicatechin; Fe<sup>3+</sup>-cyt c, ferricytochrome c; Fe<sup>2+</sup>-cyt c, ferrocyanochrome c; GEN, genistein; GEN-S-K5 cells, S-K5 cells treated with genistein; G6PD, glucose-6-phosphate dehydrogenase; MIM, mitochondrial inner membrane; MOM, mitochondrial outer membrane; mPT, mitochondrial permeability transition; mPTP, mitochondrial permeability transition pore; PTE, pterine; RCR, respiratory control ratio; S.D., standard deviation; S-K25 cells, control cells; S-K5 cells, apoptotic cells; xh S-K5 cells, apoptotic cells xh after the induction of apoptosis; SOD, superoxide dismutase; SUCC, succinate; XO, xanthine oxidase.

\* Corresponding author. Tel.: +39 0874 404243/080 5443364;

fax: +39 080 5443317.

E-mail addresses: [a.atlante@ibbe.cnr.it](mailto:a.atlante@ibbe.cnr.it) (A. Atlante), [passarel@unimol.it](mailto:passarel@unimol.it) (S. Passarella).

component of the mitochondrial permeability transition pore (mPTP); the latter is dispensable for occurrence of apoptosis [9]. On the assumption that neuropathies are a result of neuronal apoptosis, the identification of compounds able to protect neurons against apoptosis is highly desirable.

Given that the soy flavonoids have been claimed to improve human memory and neuro-cognitive performance, there is reason to suspect that this might be due to their ability to protect neurons against stress-induced injury, perhaps preventing apoptosis [10,11]. However, although the biological processes modulated by flavonoids, and especially by the isoflavone genistein (GEN), have been extensively studied, there is no clear understanding of the cellular and molecular mechanisms of action involved [12]. Flavonoids may target mitochondria in apoptosis, given that they have been reported to impair mitochondrial ATPase [13], to modulate the mPTP [14,15] and interact with other mitochondria-associated pro-apoptotic factors such as DIABLO/smac [16,17].

On the other hand, in recent reviews GEN was shown to have a variety of effects, including binding to estrogen receptors, antioxidative activity, the capacity to increase cellular reduced glutathione and effects on other physiological functions (for Ref., see [18]). In particular, GEN was reported to be a potent inhibitor of tyrosine kinase (TK, E.C.2.7.10) in tumor tissues [19,20] and of xanthine oxidase (XO, E.C.1.2.3.2) [21]. Recently, protection by GEN of the rat brain synaptosome and cultured hippocampal neurons from insult induced by  $\beta$ -amyloid peptide A $\beta$ 25–35 (for Ref. see [18]) was found. Indeed, GEN was shown to be either pro-apoptotic or anti-apoptotic depending both on its concentration and the cell system [14,22–24]. Similarly, the GEN analogue daidzein (DZN), which does not inhibit TK [25], can inhibit D-gal-induced apoptosis via the Bcl-2/Bax apoptotic pathway [26] and may be a potential medical candidate for neurodegeneration therapy. DZN exhibits an antioxidant activity [27] and shows other biological properties, including estrogen-like and estrogenin-dependent effects (for Ref. see [28]).

Herein we report investigations on the effect of GEN and DZN on the steps outlined above in the process by which CGCs undergo apoptosis as a result of potassium deprivation. We found that GEN and DZN, the latter with a lower efficiency, when used in the “dietary concentration range” (up to 10  $\mu$ M) (for Refs. see [29–31]), can prevent apoptosis from occurring in a manner consistent with their antioxidant activity. This provides them with a role as potential drugs in neurodegenerative disease therapy. In contrast with [32,33], we found that GEN does not impair xanthine oxidase, at least in CGCs.

## 2. Materials and methods

### 2.1. Reagents

Tissue culture medium and fetal calf serum were purchased from GIBCO (Grand Island, NY) and tissue culture dishes were from NUNC (Taasttrup, Denmark). All enzymes and biochemicals were from Sigma Chemicals Co. (St. Louis, MO, USA).

All procedures involving the use of animals were performed in compliance with relevant laws and institutional guidelines. The animals were anesthetized and insensitive to pain throughout the procedure.

### 2.2. Cell cultures

Primary cultures of CGCs were obtained from dissociated cerebella of 7-day-old Wistar rats as in Levi et al. [34]. Cells were plated in basal medium Eagle (BME) supplemented with 10% fetal calf serum, 25 mM KCl, 2 mM glutamine and 100  $\mu$ g/ml gentamicin on dishes coated with poly-L-lysine. Cells were plated at

$2 \times 10^6$  per 35 mm dish,  $6 \times 10^6$  per 60 mm dish, or  $15 \times 10^6$  per 90 mm dish. Arabinofuranosylcytosine (10  $\mu$ M) was added to the culture medium 18–22 h after plating to prevent proliferation of non-neuronal cells.

### 2.3. Induction of apoptosis

Primary neuronal cultures at 6–7 days in vitro (DIV), which showed typical neuronal morphologies with healthy cell bodies and intact processes, were used for the experiments. Apoptosis was induced as in D'Mello et al. [35]: cells were washed twice and switched to serum-free BME (S-) containing 5 mM KCl supplemented with 2 mM glutamine and 100  $\mu$ g/ml gentamicin. In some experiments, a variety of compounds (including the flavonoids) were also added at the induction time at the indicated concentrations selected to avoid any possible interference with cell viability. The flavonoids were dissolved in dimethyl sulfoxide (DMSO) and stored at  $-20^\circ\text{C}$  in the dark, with the final DMSO concentration kept below 0.1%. The drug exposure was terminated by removing the flavonoid-containing media, followed by washing twice and replacing with fresh media. Sister cultures prepared under the same conditions were used in each experiment. Control cells (without or with DMSO) were treated identically but maintained in serum-free BME medium supplemented with 25 mM KCl for the indicated times; they are referred to as S-K25 cells. Apoptotic cells are referred to as S-K5 cells, and to indicate the different times  $\times$  after apoptosis induction, they are referred to as xtime-S-K5.

The occurrence of apoptosis was checked, as in [6,35], by measuring DNA laddering and prevention of death due to the addition of actinomycin D (Act D). The occurrence of necrosis was assessed, as in [6], by checking the release of L-lactate dehydrogenase and the ability of MK801 to prevent CGC death.

### 2.4. Assessment of CGC survival

As advised in [36,37], cell survival was quantified by counting the number of intact nuclei using a hemacytometer after lysing the cells in detergent-containing solution as in [38,39]. Each measurement was performed in triplicate and reported as the mean  $\pm$  SEM. The data are expressed as the percentage of intact nuclei in the control cultures at each time point.

### 2.5. DNA fragmentation analysis

Fragmentation of DNA was performed as in [6,40]. Briefly, CGCs ( $6 \times 10^6$ ) were plated in poly L-lysine-coated 60 mm tissue culture dishes, collected with cold phosphate-buffered saline (PBS, pH 7.2) and, after removal of the medium and washing once with cold PBS, the cells were centrifuged at  $3500 \times g$  for 5 min. The pellet was lysed in 10 mM Tris-HCl, 10 mM EDTA, and 0.2% Triton X-100 (pH 7.5). After 30 min on ice, the lysates were centrifuged at  $17000 \times g$  for 10 min at  $4^\circ\text{C}$ . The supernatant was digested with proteinase K and then extracted twice with phenol-chloroform/isoamyl alcohol (24:1). The aqueous phase, containing soluble DNA, was recovered and nucleic acids were precipitated with sodium acetate and ethanol overnight. The pellet was washed with 70% ethanol, air-dried and dissolved in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 7.5). After digestion with RNase A (50 ng/mL at  $37^\circ\text{C}$  for 30 min), the sample was subjected to electrophoresis in a 1.8% agarose gel and visualized by ethidium bromide staining. Soluble DNA from equal numbers of cells was loaded in each lane.

### 2.6. Cell suspension and homogenate preparations

Before each experiment, the culture medium was removed and the plated CGCs were washed with phosphate-buffered saline

medium containing 138 mM NaCl, 2.7 mM KCl, 8 mM Na<sub>2</sub>HPO<sub>4</sub>, and 15 mM KH<sub>2</sub>PO<sub>4</sub> at pH 7.4 and then collected by gentle scraping into a volume of medium depending on the particular experiment.

### 2.6.1. Cell suspensions

Cell suspensions showed full viability, even though they lacked the morphological organization observed in culture dishes such as cell–cell and cell–substrate contacts, and had no neurites. Cell integrity, which remained essentially constant for 3–5 h, was quantitatively assessed by confirming the inability of cells to oxidize externally added succinate (SUCC), which cannot enter intact cells [41], and by confirming the ability of ouabain to block glucose transport into the cells [42]. The final cell suspension routinely contained 85–95% intact cells.

### 2.6.2. Cell homogenate

Cell homogenate from plated CGCs was obtained by washing them with PBS and collecting by gentle scraping to obtain a cell suspension that was then homogenized in a Dounce-Potter homogenizer using 10 strokes at 25 °C. The rupture of cell membranes was checked by confirming the failure of ouabain to inhibit glucose-dependent oxygen consumption (for Ref., see [5]).

The intactness of mitochondria in the cell homogenate was checked by measuring the activities of adenylate kinase (ADK, E.C.2.7.4.3) and glutamate dehydrogenase in the post-mitochondrial supernatant (GDH, E.C.1.4.1.3) (for Ref., see [5]), which are marker enzymes of the mitochondrial intermembrane space and matrix, respectively. Mitochondrial coupling was monitored by measuring the respiratory control ratio (RCR), i.e., (oxygen uptake rate after ADP addition)/(oxygen uptake rate before ADP addition), with succinate used as a respiratory substrate.

## 2.7. Polarographic measurements

### 2.7.1. Cellular and mitochondrial respiration

O<sub>2</sub> consumption was measured polarographically using a Gilson 5/6 oxygraph with a Clark electrode, as in [5,7]. Either intact cells or cell homogenates in PBS (about 1 mg protein) were incubated in a thermostatted (25 °C) water-jacketed glass vessel (final volume 1.5 ml). The sensitivity of the instrument was set so as to allow rates of O<sub>2</sub> uptake as low as 0.5 natom min<sup>−1</sup> mg<sup>−1</sup> protein to be measured. Since the incubation chamber required continuous stirring to allow O<sub>2</sub> to diffuse freely, the design included a magnetic stirring system. Oxygen uptake was initiated by adding either glucose or succinate to the cell suspension or homogenate, respectively.

Protein content was determined according to [43] with bovine serum albumin used as a standard.

### 2.7.2. Cytochrome *c* release

To detect the presence of active cyt *c* in the extramitochondrial phase, the ability of cell homogenate to oxidize ascorbate (ASC) was checked [7]. Briefly, because ASC cannot permeate *per se* the outer mitochondrial membrane (for Ref., see [7]), its oxidation must occur as a result of release from mitochondria of a component that can oxidize ASC and then reduce oxygen via cytochrome *c* oxidase, i.e., cyt *c*. The rate of oxygen uptake was measured as the tangent to the initial part of the progress curve and expressed as natom O min<sup>−1</sup> mg<sup>−1</sup> protein.

## 2.8. Superoxide anion detection in CGCs

Superoxide anion (O<sub>2</sub><sup>•−</sup>) was detected using the ferricytochrome *c* (Fe<sup>3+</sup>–cyt *c*) method [44] as an increase in absorbance at 550 (ε<sub>550nm</sub> = 37 mM<sup>−1</sup> cm<sup>−1</sup> at 25 °C) and was measured using a PerkinElmer LAMBDA-5 spectrophotometer equipped with a

thermostatted holder. A calibration curve was made by using xanthine plus XO as an O<sub>2</sub><sup>•−</sup> – producing system, which produces ferrocytochrome *c* (Fe<sup>2+</sup>–cyt *c*) in a 1:1 stoichiometry.

## 2.9. Xanthine oxidase assay

XO activity was assayed both by fluorimetric measurements (PerkinElmer LS50; excitation and emission wavelengths of 345 and 390 nm, respectively) of isoxanthopterin formation from pterine as in [45], and by photometric measurements at 550 nm of XO-dependent O<sub>2</sub><sup>•−</sup> formation, as in [46].

## 2.10. Measurement of ATP levels

ATP levels in CGC extracts were determined by HPLC as in [8].

## 2.11. ANT measurement

Cell homogenate (0.1 mg protein) containing mitochondria was incubated at 25 °C in 2 ml of standard medium consisting of 200 mM sucrose, 10 mM KCl, 1 mM MgCl<sub>2</sub>, and 20 mM HEPES-Tris at pH 7.2. The appearance of ATP in the extramitochondrial phase, due to externally added ADP, was revealed as in [9] by using an ATP detecting system (ATP D.S.). This consisted of glucose (2.5 mM), hexokinase (HK, E.C.2.7.1.1, 0.5 e.u.), glucose-6-phosphate dehydrogenase (G6PD, E.C.1.1.1.49, 0.5 e.u.) and NADP<sup>+</sup> (0.2 mM) in the presence of P<sub>1</sub>P<sub>5</sub>-di(adenosine-5')penta-phosphate (Ap5A), a specific inhibitor of adenylate kinase (for Ref., see [9]). The rate of NADP<sup>+</sup> reduction in the extramitochondrial phase was followed as an absorbance increase at 334 nm, which was measured as the tangent to the initial part of the progress curve and expressed as nmol NADP<sup>+</sup> reduced/min × mg cell protein. Because the rate of ATP appearance measured in cell homogenates could depend on a number of events, including electron flow across the mitochondrial membrane, electrochemical proton gradient generation, ATP synthase and the adenine nucleotide content of the mitochondria, under each experimental condition, we investigated the rate-limiting step of the process leading to ATP efflux from the mitochondria as in [9].

## 2.12. Measurement of the mitochondrial permeability transition pore opening

To monitor the onset of mPTP opening, cell homogenate was suspended in a medium containing 250 mM sucrose, 2 mM HEPES pH 7.4, 0.5 mM K<sub>2</sub>HPO<sub>4</sub>, 1 μM oligomycin, 2 μM rotenone and 5 mM succinate, and mitochondrial swelling was monitored at 25 °C as the absorbance decrease at 546 nm, as in [9].

## 2.13. Statistical analysis and computing

All statistical analyses in this study were performed using SPSS software. The data were representative of at least three independent experiments carried out with different neuronal preparations, and are reported as the mean with the standard deviation (S.D.). Statistical significance of the data was evaluated using the one-way analysis of variance (ANOVA) followed by the post-hoc Bonferroni test. *p* < 0.05 was considered as significant for all analyses.

Experimental plots were obtained using Grafit (Erithacus software).

## 3. Results


### 3.1. Flavonoids and low-K<sup>+</sup> induced CGC apoptosis

7DIV neuron cultures were kept either in high potassium medium (S-K25 cells) or subjected to low potassium shift (S-K5

cells), which is responsible for triggering apoptosis (for Ref., see [6,35]), in the absence or presence of a variety of flavonoids. That neurons died via apoptosis was confirmed by monitoring DNA laddering, which is a specific hallmark of apoptosis (Fig. 1A), and the sensitivity of death to the transcriptional inhibitor Act D (Fig. 1B). DNA laddering was found only in S-K5 cells (Fig. 1A, lane b), but was completely prevented in the presence of the transcriptional inhibitor Act D (1  $\mu$ g/ml) (Fig. 1A, lane d). That necrosis had not occurred was shown by the absence of significant LDH activity, both in S-K5 and S-K25 culture medium up to 15 h (not shown), and by the insensitivity of death to MK801, a selective NMDA receptor antagonist, which prevents glutamate-dependent necrosis (for Ref., see [44,47]).

In Fig. 1B, we further confirmed that CGC death occurred via apoptosis, as shown by its complete prevention by Act D (1  $\mu$ g/ml) (see [35]), but not by MK801 (1  $\mu$ M). In fairly good agreement with [9,35], the survival of S-K5 CGC 24 h after apoptosis induction was 40% with respect to the control ( $p < 0.001$ , 6 expts). Next, we investigated whether the isoflavones GEN and DZN could prevent CGC apoptosis, as observed 24 h after induction. The effect of epicatechin (EC) and catechin (CE) was also investigated (Fig. 1B). In all cases, the compounds were added at concentrations of up to 20  $\mu$ M; they had no effect on S-K25 CGC survival (not shown).

GEN and DZN (0.1–20  $\mu$ M each) proved to prevent death in a dose-dependent manner, with a survival of about 70% ( $p < 0.001$ , 6 experiments) in the presence of 20  $\mu$ M GEN, and about 60%



**Fig. 1.** The effect of some flavonoids on low-K<sup>+</sup> induced CGC apoptosis. Rat CGCs ( $1 \times 10^6$ /well) at 7 DIV were incubated either in high potassium (S-K25) or low potassium (S-K5) serum-free culture medium. (A) DNA fragmentation. Soluble DNA was extracted from neurons switched to serum-free culture medium containing low K<sup>+</sup> (5 mM) in the absence (lane b) or presence of Act D (1  $\mu$ g/ml; lane d) after 15 h. Lane c contains DNA from control cells maintained in high K<sup>+</sup> (25 mM) for 15 h. DNA from equal numbers of plated cells ( $6 \times 10^6$ ) was loaded in each lane. Size marker was HaeIII-digested  $\Phi \times 174$  phage DNA (lane a). (B) Survival: S-K5 cells were incubated in the absence (None) or presence of the following compounds: genistein (GEN), daidzein (DZN), epicatechin (EC) and catechin (CE) at the indicated concentrations. Where indicated, 1  $\mu$ M MK801 and 1  $\mu$ g Act D were present in the culture medium. At 24 h, the viability of S-K25 and S-K5 cells was determined by counting the number of intact nuclei. Cell viability was expressed as the percentage of that of the S-K25 cells. Control values were  $100 \pm 10$ . The compounds used had no effect on the survival of S-K25 cells. Results are means with the ( $\pm$ S.D.) of triplicate measurements and are representative of six different experiments carried out with different cell preparations from different groups of animals. ANOVA and Bonferroni test: \*\*\*, \*\*\*\* statistically significantly different with  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively, when S-K5 cells were compared against the samples indicated with asterisk/s. The lack of asterisk indicates no statistically significant differences. (C) Phase-contrast micrographs showing neurons maintained for 24 h in serum-free medium containing 25 mM KCl (a) and 5 mM KCl in the absence (b) or presence of either GEN (c) or DZN (d) (20  $\mu$ M each). The phase contrast micrograph was taken using a NIKON DIAPHOT TS-100 epifluorescence microscope (equipped with a 40 $\times$  objective). Images were digitized by using a Nikon Digital Video Camera E995 (Coolpix 995).



( $p < 0.01$ , 6 experiments) in the presence of  $20 \mu\text{M}$  DZN. Treatment with Act D in the presence of flavonoid resulted in the complete prevention of death; this confirms that death occurred by apoptosis. In contrast, both EC and CE, used at concentrations of up to  $20 \mu\text{M}$ , were totally ineffective (Fig. 1B). Death prevention was confirmed in a parallel experiment in which phase-contrast micrographs of the cells were used, as in [48]. Fig. 1C shows that both GEN (panel c) and DZN (panel d) ( $20 \mu\text{M}$  each) treatment resulted in a reduction of cell degeneration and death with respect to untreated S-K5 cells (panel b).

Having established that GEN and DZN can prevent apoptosis of CGCs, their effects on some mitochondria-linked steps in the process leading to apoptosis of CGCs were investigated separately.

### 3.2. Glucose oxidation by either GEN- or DZN- S-K5 cells

The ability of GEN and DZN to prevent the impairment of glucose oxidation, which occurs in early apoptosis [5], was investigated. Oxidation of externally added glucose ( $10 \text{ mM}$ ) by intact cells was monitored by measuring the oxygen consumption that results from glucose uptake by CGCs, followed by aerobic

glycolysis, citric acid cycle action and electron flow along the mitochondrial respiratory chain. In a typical experiment (Fig. 2) the oxygen uptake rates were about 55 and 36 natom O/min per mg cell protein for S-K25 and 3 h S-K5 CGCs, respectively. In both cases, cyanide ( $\text{CN}^-$ ), which prevents electron flow by blocking cytochrome oxidase activity, and ouabain, which can prevent glucose uptake by preventing cellular sodium/potassium ATPase (for Ref., see [5]) (not shown), completely inhibited oxygen consumption. In 3 h-S-K5 cells, the decrease in the oxygen consumption rate was largely prevented by either GEN or DZN ( $20 \mu\text{M}$  each); the rates were 49 and 44 natom O/min per mg cell protein, respectively (Fig. 2). Notice that the glucose oxidation rate remained constant in S-K25 CGCs up to 3 h. As a result of treatment of the cells with superoxide dismutase (SOD, E.C.1.15.1.1,  $50 \text{ U/ml}$ ), which was added to S-K5 cell cultures, the impairment of glucose oxidation was prevented, with an oxygen uptake rate by CGCs equal to 51 natom O/min per mg cell protein) (inset to Fig. 2).

In a set of similar experiments with GEN and DZN concentrations ranging from  $0.1$  to  $20 \mu\text{M}$ , we found that isoflavone prevention of the impairment of glucose oxidation was dose-dependent (see panel B of Fig. 2). GEN proved to be more effective than DZN, as judged by a statistical analysis carried out according to the Bonferroni test.

### 3.3. Mitochondrial coupling in either GEN- or DZN- S-K5 cells

In another set of experiments (Fig. 3), mitochondrial coupling in the homogenates of either S-K25 or 3 h-S-K5 CGCs incubated in the absence or presence of either GEN or DZN was investigated. To achieve this, oxygen uptake was measured arising from the addition of succinate to cell homogenate in either the absence (state 4) or presence (state 3) of ADP ( $1 \text{ mM}$ ) used to stimulate oxidation. The RCR values (oxygen uptake in state 3)/(oxygen uptake rate in state 4), which reflect the ability of mitochondria to produce ATP, were about 5.0, 2.6, 4.2 and 3.5 in S-K25, 3 h-S-K5, GEN- and DZN-3h-S-K5 ( $20 \mu\text{M}$  each) cells, respectively. When SOD ( $50 \text{ U/ml}$ ) was added to S-K5 cell culture, the RCR value was 4.5 (inset to Fig. 3).


Notice that the rate of oxygen uptake in state 4 was not changed by treatment with either GEN or DZN, ruling out the possibility that these compounds can uncouple mitochondria. In a control experiment, we found that GEN and DZN were without effect when oxygen uptake by S-K25 cell homogenates was investigated.

In panel B of Fig. 3, the RCR values (calculated from 5 expts) obtained using GEN or DZN concentrations ranging from  $0.1$  to  $20 \mu\text{M}$  are reported. Fig. 3B shows that the prevention of mitochondrial impairment was dose-dependent, with GEN more effective than DZN.


Other mitochondrial steps of CGC apoptosis were investigated separately at 3 h and/or 5 h after apoptosis induction, times which reflect the early and the late phases of apoptosis, respectively (see above).

### 3.4. The release of cytochrome c from either GEN- or DZN-S-K5 cell mitochondria

Since in CGC apoptosis the release of cyt c from mitochondria occurs as a result of extra ROS production (for Refs., see [7,49,50]), the ability of either GEN or DZN to affect cyt c release was investigated (Fig. 4). To do this, as in [46,50], we resorted to polarographic measurement of the activation of cyt c-dependent ASC oxidation, which itself confirmed that cyt c was released in an active form [50] (for details see Section 2). In a typical experiment, S-K25, S-K5, GEN-S-K5 or DZN-S-K5 cells, at 3 h after apoptosis induction, were compared to one another with respect to their ability to oxidize externally added ASC ( $5 \text{ mM}$ ) (Fig. 4A). No oxygen



**Fig. 2.** Glucose oxidation by either GEN- or DZN-S-K5 cells. (A) Rat CGCs ( $2 \times 10^6$ /well) at 7 DIV were switched from high potassium (25 mM) (S-K25) (a) to low potassium (5 mM) serum-free culture medium in the absence (S-K5) or presence of  $20 \mu\text{M}$  genistein (GEN-S-K5) or  $20 \mu\text{M}$  daidzein (DZN-S-K5) for 3 h. Aliquots corresponding to  $0.2 \text{ mg}$  cell protein were incubated at  $25^\circ\text{C}$  in a water-jacketed glass vessel to monitor  $\text{O}_2$  consumption polarographically. Glucose ( $10 \text{ mM}$ ) and subsequent  $\text{CN}^-$  ( $1 \text{ mM}$ ) additions were made at the indicated times. In the inset, S-K5 cells were treated with SOD ( $50 \text{ U/ml}$ ) (SOD-S-K5). Numbers along the curves are rates of oxygen uptake expressed as natom O/min × mg cell protein. (B) The dependence of the rates of glucose-dependent oxygen consumption ( $V_{\text{O}_2 \text{ glucose}}$ ) versus GEN and DZN concentrations is reported. Data are the mean ± S.D. of five measurements and are representative of at least three different experiments carried out with different cell preparations. ANOVA and Bonferroni test: \*\*, statistically significantly different with  $p < 0.05$  and  $p < 0.01$ , respectively, when S-K5 cells were compared against the samples indicated with asterisk/s. The lack of asterisk indicates no statistically significant differences.




**Fig. 3.** Mitochondrial coupling in either GEN- or DZN-S-K5 cells. (A) Rat CGCs ( $2 \times 10^6$ /well) at 7 DIV, switched from high potassium (25 mM) (S-K25) to low potassium (5 mM) serum-free culture medium in the absence (S-K5) or presence of 20 μM genistein (GEN-S-K5) or 20 μM daidzein (DZN-S-K5) for 3 h, were homogenized and incubated (0.2 mg cell protein) at 25 °C in a water-jacketed glass vessel and the consumption of  $O_2$  was monitored polarographically. Succinate (SUCC, 5 mM) and ADP (1 mM) were added at the indicated times. In the inset, S-K5 cells were treated with SOD (50 U/ml) (SOD-S-K5). Numbers along the curves are rates of oxygen uptake expressed as natom  $O/min \times mg$  cell protein. (B) The dependence of Respiratory Control Ratio (RCR) values versus GEN and DZN concentrations is reported. Data are the mean  $\pm$  S.D. of five measurements and are representative of at least three different experiments carried out with different cell preparations. ANOVA and Bonferroni test: \*, \*\*, \*\*\* statistically significantly different with  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively, when S-K5 cells were compared against the samples indicated with asterisk/s. The lack of asterisk indicates no statistically significant differences.

uptake was found in the S-K25 cell sample in the presence of ASC alone (trace a), showing that mitochondria were intact and that no release of cyt c had occurred. ASC addition to homogenates from S-K5, GEN- or DZN-S-K5 CGCs resulted in oxygen uptake at a rate of about 16, 8 and 11 natom  $O/min \times mg$  cell protein, respectively (traces b-d). The results in Fig. 4A show that prevention of cyt c release occurred as a result of the presence of the flavonoids. The prevention was dose-dependent (Fig. 4B), with GEN being more effective than DZN; prevention values were about 50% and 25% in the GEN- and DZN-S-K5 cells (20 μM each), respectively. As a control, the occurrence of 60% prevention of cyt c release in S-K5 cells in the presence of externally added SOD was confirmed [6].

### 3.5. ATP level, ANT activity and mPTP opening in either GEN- or DZN-S-K5 cells


ATP is required for apoptosis to occur. In early apoptosis, ATP is mostly produced in the mitochondria by oxidative phosphorylation [8] and is exported via the ANT into the cytoplasm (for Refs., see [9]). As in [8], we found that the ATP level in 3 h-S-K5 CGCs was higher compared with the control (1.2 vs. 0.7 nmol/mg cell protein); incubation of S-K5 cells with GEN or DZN (20 μM each)



**Fig. 4.** The release of cytochrome c from either GEN- or DZN-S-K5 cell mitochondria. Rat CGCs ( $2 \times 10^6$ /well) at 7 DIV were incubated in either high potassium (S-K25) or low potassium serum-free culture medium in the absence (S-K5) or presence of either genistein (GEN-S-K5), daidzein (DZN-S-K5) or SOD (50 U/ml) (SOD-S-K5). At 3 h after apoptosis induction, the cells were scraped, collected and homogenized. Cell homogenate (about 0.2 mg protein) was incubated at 25 °C in the presence of 3 μM rotenone, 0.8 μM antimycin and 6 μM myxothiazole in a water-jacketed glass vessel. (A) Cyt c release from S-K25 or S-K5 cells, in the absence or presence of GEN and DZN (20 μM each), at 3 h after apoptosis induction, was detected by means of oxygen consumption caused by externally added ASC (5 mM). (B) Cyt c release from S-K25 or S-K5 cells, in the absence or presence of GEN and DZN at the indicated concentrations, at 3 h after apoptosis induction, was detected as in (A). Results, expressed as natoms  $O/min \times mg$  cell protein, are the means  $\pm$  S.D. of triplicate measurements and are representative of at least six different experiments carried out with different cell preparations obtained from different groups of animals. ANOVA and Bonferroni test: \*, \*\*, \*\*\* statistically significantly different with  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively, when S-K5 cells were compared against the samples indicated with asterisk/s. The lack of asterisk indicates no statistically significant differences.

or SOD (50 U/ml) did not result in a change of the ATP level (Fig. 5A).

The effect of GEN and DZN on the process in which alteration of the ANT takes place, triggering opening of mPTP (see above and [9]), was investigated by measuring in parallel both ANT-mediated transport and mPTP opening in S-K25, S-K5, GEN-S-K5 and DZN-S-K5 cell homogenates. This was done both at 3 h, when ANT function is impaired due to ROS, but no mPTP opening occurs, and at 5 h, when further caspase-dependent ANT impairment occurs and mPTP opening initiates [9]. In the inset, GEN and DZN were present at concentrations equal to 20 μM, when ANT function is impaired due to ROS, but no mPTP opening occurs, and at 5 h, when further caspase-dependent ANT impairment occurs and mPTP opening initiates [9]. In the former case, we resorted to a procedure (see [9]) that allows for the continuous monitoring of ATP efflux from mitochondria incubated with ADP; in the latter, mPTP opening was investigated as mitochondrial swelling as in [9] (for details see Section 2). In a typical experiment, cell homogenates were treated with AP5A to inhibit adenylate kinase. In all cases, the ATP concentration outside the mitochondria was negligible, as no increase in the absorbance measured at 334 nm was found in the presence of the ATP D.S. As a result of the addition of ADP (0.04 mM), an increase in the NADPH absorbance was found, indicating the appearance of ATP in the extramitochondrial phase.




**Fig. 5.** ATP level and ANT activity in homogenates from either GEN- or DZN-S-K5 cells. (A) The amounts of ATP were measured by HPLC in neutralized perchloric acid extracts of either S-K25 or S-K5 CGCs in the absence or presence of GEN (20  $\mu$ M), DZN (20  $\mu$ M) or SOD (50 U/ml), as described in Section 2. ANOVA and Bonferroni test: No statistically significant differences were found when S-K5 cells were compared against each sample. (B) The scheme describes the method used to detect ATP appearance in the extramitochondrial phase due to the addition of ADP to CGC homogenate. (C–D) Appearance of ATP due to the addition of ADP (0.04 mM) to CGC homogenate (0.1 mg protein) from S-K25 or S-K5 cells in the absence or presence of GEN and DZN at the indicated concentrations at 3 (A) and 5 h (B) after apoptosis induction was monitored as described in Section 2.

The explanation for this is as follows (see Fig. 5B): ADP enters mitochondria in exchange for endogenous ATP; inside the matrix ATP is synthesized by oxidative phosphorylation, and the newly synthesized ATP exits the mitochondria in exchange for further ADP via the ANT. That the rate of NADPH formation mirrored, in all cases, the rate of ADP/ATP exchange was confirmed by applying control flux analysis as in [9] (not shown). The rates of ATP efflux were 10, 2.5 and 1.6 nmol NADPH<sup>+</sup> reduced/min  $\times$  mg cell protein for S-K25, 3h-S-K5 and 5h-S-K5 CGCs, respectively (Fig. 5C, D). When SOD (50 U/ml) was added to the cell cultures, almost complete prevention of ANT impairment was found at 3 h after apoptosis induction (10 nmol NADPH<sup>+</sup> reduced/min  $\times$  mg cell) (Fig. 5C); this confirmed that the decrease in ANT efficiency depended on extra ROS production [9]. In 5h-S-K5 cells, where a progressive impairment in ANT occurs mainly due to caspase activation, the SOD prevention decreased to 5.3 nmol NADPH<sup>+</sup> reduced/min  $\times$  mg cell (Fig. 5D).

In GEN-3h-S-K5, GEN-5h-S-K5, DZN-3h-S-K5 and DZN-5h-S-K5 CGCs (20  $\mu$ M each), the rates of ATP efflux were 9.3, 4.0, 5.9 and 3.6 nmol NADPH<sup>+</sup> reduced/min  $\times$  mg cell protein, respectively (Fig. 5C,D). This shows that both GEN and DZN can prevent ANT impairment.


In a parallel experiment (Fig. 6), mPTP opening was measured by monitoring mitochondrial swelling as a change in absorbance at 546 nm of the CGC homogenate (see Section 2). It should be noted that the decrease in light absorbance is a spontaneous process due to apoptosis and is not calcium-dependent (see [51]). As in [9], no mPTP opening was found at 3 h after induction of apoptosis, either in the absence or presence of GEN or DZN (not shown), confirming that ROS themselves cannot cause mPTP opening. In 5h-S-K5 CGCs, mPTP opening occurred. In either GEN- or DZN-S-K5 cells, mPTP opening was prevented in a manner dependent on the compound concentration, with 70% and 50% prevention found for GEN and DZN, respectively, at

a concentration of 20  $\mu$ M. In SOD (50 U/ml) treated cells, 80% prevention was found. As expected, no mitochondrial swelling occurred in the presence of 1  $\mu$ M cyclosporine A (CsA), an inhibitor of mPTP [9,52].



**Fig. 6.** mPTP opening in homogenates from either GEN- or DZN-S-K5 cells. Rat CGCs ( $15 \times 10^6$ /well) at 7 DIV were incubated either in high potassium (S-K25) or low potassium (S-K5) serum-free culture medium in the absence or presence of either GEN or DZN at the indicated concentrations, or SOD (50 U/ml) or cyclosporine A (CsA, 1  $\mu$ M) for 8 h. mPTP opening was monitored by mitochondrial swelling (see Section 2).






**Fig. 7.** Superoxide anion production in either GEN- or DZN-S-K5 cells. Rat CGCs ( $0.5 \times 10^6$ /well) at 7 DIV were incubated in either high potassium (S-K25) or low potassium (S-K5) serum-free culture medium containing  $\text{Fe}^{3+}$ -cyt c ( $10 \mu\text{M}$ ) in the absence or presence of either GEN or DZN at the indicated concentrations or SOD ( $50 \text{ U/ml}$ ). At 3 h after apoptosis induction, the culture solution was taken and the increase in absorbance at  $550 \text{ nm}$  due to  $\text{Fe}^{3+}$ -cyt c reduction was determined. The experimental data are reported as nmol ROS formed  $\pm$  S.D./ $10^6$  cells, calculated on the basis of the stoichiometry of the reaction using the extinction coefficient determined under our experimental conditions (see Section 2). The experiment was repeated 4 times with different cell preparations. ANOVA and Bonferroni test: \*\*\*, \*\*\*\* statistically significantly different with  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively, when S-K5 cells were compared against the samples indicated with asterisk/s. The lack of asterisk indicates no statistically significant differences.

### 3.6. Superoxide anion production in either GEN- or DZN- S-K5 cells

In CGC apoptosis, extra ROS production takes place ([6,53] and Refs. therein) with all of the above steps shown to be mostly dependent on ROS production. Since both GEN and DZN are antioxidants, we investigated whether they, used up to  $20 \mu\text{M}$ , could decrease  $\text{O}_2^{\bullet-}$  levels at 3 h into apoptosis, when maximum  $\text{O}_2^{\bullet-}$  production takes place [6] (Fig. 7). In agreement with [6], the superoxide level ( $4.0 \pm 0.1 \text{ nmol O}_2^{\bullet-}/10^6$  cells) was substantially decreased by externally added SOD ( $p < 0.001$ ). A dose-dependent decrease in  $\text{O}_2^{\bullet-}$  levels was found in both GEN- or DZN-S-K5 cells; in GEN ( $10 \mu\text{M}$ ) and DZN ( $20 \mu\text{M}$ )-S-K5 cells, the level of  $\text{O}_2^{\bullet-}$  was reduced by about 50%.

Since in S-K5 cells XO can contribute to ROS production [49], and since GEN was reported to be a XO inhibitor [32,33], we investigated whether GEN could inhibit XO in S-K5 cells. To do this we resorted to a procedure in which XO was assayed fluorimetrically by monitoring the increase in fluorescence arising from the formation of isoxanthopterin from pterine (PTE) [45]. The rates of isoxanthopterin generation were about 14 and  $6 \text{ nmol/min} \times \text{mg cell protein}$  for 3 h S-K5 and S-K25 cells, respectively (Fig. 8A(a,b)). In GEN-3h-S-K5 cells, no inhibition of the fluorescence increase (Fig. 8A(a)) due to pterine addition to the cell homogenate occurred, showing that GEN itself had no effect on XO activity. This was also confirmed when GEN was added to the assay mixture before starting the reaction (Fig. 8A(b)). As a control, we confirmed that the reaction was strongly inhibited by allopurinol, a specific XO inhibitor (see [44,45]), either incubated with cultured cells (a) or added during the reaction (b). As expected, no change in the XO reaction was found in SOD-treated CGCs (not shown).

In parallel, we assayed XO by adding xanthine ( $10 \mu\text{M}$ ) plus  $\text{Fe}^{3+}$ -cyt c to the homogenates. In this system,  $\text{O}_2^{\bullet-}$  formation due to endogenous XO was measured as an absorbance increase at  $550 \text{ nm}$  resulting from reduction of  $\text{Fe}^{3+}$ -cyt c (Fig. 8B). The rates of  $\text{O}_2^{\bullet-}$  formation, equal to 10 and  $2.5 \text{ nmol/min} \times \text{mg cell protein}$  in 3h-S-K5 and S-K25, respectively, were decreased by 60% by GEN ( $20 \mu\text{M}$ ) incubated with 3h-S-K5-cells (a) or added before the



**Fig. 8.** Xanthine oxidase activity in GEN-S-K5 cells. Rat CGCs ( $3 \times 10^6$ /well) at 7 DIV were switched from high potassium ( $25 \text{ mM}$ ) (S-K25) to low potassium ( $5 \text{ mM}$ ) serum-free culture medium (S-K5). After 3 h of incubation, CGCs were collected and homogenized in PBS. The protein concentration was adjusted to  $25 \mu\text{g/ml}$  and aliquots were then assayed for XO activity. (A) The fluorimetric increase due to isoxanthopterin formation ( $345/390 \text{ nm}$ ) was initiated by the addition of pterine ( $10 \mu\text{M}$ ) (see Section 2). In (a) either  $20 \mu\text{M}$  GEN (GEN-S-K5) or  $10 \mu\text{M}$  allopurinol (ALLO-S-K5) was added to the cell well at the same time as the switch in  $\text{K}^+$  concentration. In (b)  $20 \mu\text{M}$  GEN or  $10 \mu\text{M}$  ALLO was added to the S-K5 suspension before pterine addition or during the reaction, respectively. Numbers along the traces are values of XO activity expressed as nmol of isoxanthopterin formed/ $\text{min} \times \text{mg cell protein}$ . (B) Both S-K5 and S-K25-CGCs were incubated with  $\text{Fe}^{3+}$ -Cyt c ( $10 \mu\text{M}$ ). The absorbance increase due to  $\text{Fe}^{3+}$ -Cyt c ( $550 \text{ nm}$ ) reduction was initiated by the addition of xanthine (XANT,  $10 \mu\text{M}$ ) (see Section 2). In (a) and (b) either  $20 \mu\text{M}$  GEN (GEN-S-K5),  $10 \mu\text{M}$  allopurinol (ALLO-S-K5) or SOD ( $50 \text{ U/ml}$ ) (SOD-S-K5) was added to the cell well at the same time as the switch in  $\text{K}^+$  concentration. In (c)  $20 \mu\text{M}$  GEN and  $10 \mu\text{M}$  ALLO was added to the S-K5 suspension before XANT addition. Numbers along the traces are values of XO activity expressed as nmol  $\text{O}_2^{\bullet-}$  formed/ $\text{min} \times \text{mg cell protein}$ .

reaction was started (b). The rates were decreased to zero (a, c) or partially decreased (b) by allopurinol ( $10 \mu\text{M}$ ) and SOD ( $50 \text{ U/ml}$ ), respectively.

These data suggest that GEN cannot inhibit XO itself, even if it can scavenge ROS. Surprisingly enough with respect to the above experiment, but in agreement with [32,33], we found that GEN can inhibit commercial XO activity as assayed either fluorimetrically or photometrically (not shown).

## 4. Discussion

We show here that both GEN and its analogue DZN, used at dietary concentrations, can prevent low potassium-dependent apoptosis in CGCs, and may perhaps be of use in neuroprotection. Indeed, the flavonoid effect proved to be rather specific; in spite of their shared antioxidant capability, GEN and DZN, but not CE and



EC, prevented CGC death (Fig. 1). On the other hand, a comparison made between GEN and CE showed that GEN could prevent cell death both in CGCs and in rat mesencephalic cultures, whereas CE was effective only in the latter case [54]. Hence, the flavonoid effect depends on the experimental system investigated.

Since prevention of apoptosis occurred with a parallel prevention of the impairment of certain steps of the mitochondrial pathway to apoptosis, we conclude that the effects of both GEN and DZN involve CGC mitochondria. In fact, we found prevention of the impairment of both aerobic glucose metabolism and, more importantly, of the mitochondrial uncoupling that occurs in apoptosis (Figs. 2 and 3) in the presence of these compounds. We also found the prevention of cyt c release (Fig. 4), ANT alteration (Fig. 5) and mPTP opening (Fig. 6). Finally, because the above steps of the mitochondrial pathway to apoptosis are somehow related to ROS production, which takes place during apoptosis, and since both GEN and DZN proved to decrease ROS levels (Fig. 7), we strongly suggest that the prevention of apoptosis is essentially due to the antioxidant properties of the flavonoids. This is supported by the evidence that the effects of GEN and DZN parallel those produced by the antioxidant SOD (see [6]). Thus, although other effects of GEN as a dietary compound inducing hormonal, metabolic and expression changes [55,56] cannot be ruled out, we maintain that its role as an antioxidant is crucial in accounting for these experimental findings. Accordingly, we first showed that these compounds could remove the extra ROS produced in cells undergoing apoptosis (Fig. 7). Whether the mechanism by which GEN decreases ROS levels in CGC apoptosis includes other effects remains a matter of speculation. For instance, proteosomal inhibition by GEN could cause an increase in the scavenging efficiency of the cell antioxidant system targeted by proteosoma *en route* to apoptosis (see [6]). In addition, the antioxidant properties of GEN and DZN could result from enhancement of the activities of other antioxidant molecules and enzymes (superoxide dismutase, glutathione peroxidase and reductase) both *in vitro* and *in vivo* [30,57]. In particular, DZN was shown to decrease the GSH/GSSG ratio, suggesting that it could have pro-oxidant effects in the brain [58].

Since GEN is relatively more effective at inhibiting hydroxyl radical-induced oxidation compared with DZN, which is the stronger agent against superoxide radical-induced oxidation [59], we might speculate that the greater protection against apoptosis provided by GEN suggests a special role for hydroxyl radicals in apoptosis. GEN and DZN differ from one another with respect to other biological activities. For example, in distinction from DZN, GEN is a strong inhibitor of TK [19]. Since DZN is also neuroprotective, we assume that the inhibition of TK has no/minimal role in preventing apoptosis. In this regard, it should be noted that in normal cells TK activity is rather poor, whereas in tumor cells, TK activity can increase 10–20 times [60]. Hence, the activity of GEN in cancer chemoprevention could arise from the inhibition of TK [61].

In CGCs, the ATP level proved to be insensitive to GEN and DZN (Fig. 5A). On the other hand, GEN was shown to enhance mitochondrial energy metabolism by increasing ATP synthesis in the neurodegeneration of ovariectomized rats, as reported by Shi et al. [62]. Indeed, several flavonoids proved to increase cellular ATP levels, possibly by increasing ATP production via oxidative phosphorylation [63]. In our case, because mitochondrial impairment in apoptosis is somehow prevented by these isoflavones, we suggest that oxidative phosphorylation contributes more significantly to ATP production, which is normally counterbalanced by anaerobic glycolysis, leading to extra production of L-lactate in apoptosis [4,8].

As expected, because of their role as antioxidants, GEN and DZN were found to almost completely prevent the alteration of ANT,

which occurs in apoptosis (Fig. 5), as investigated in 3h-S-K5 CGCs when ROS alone affects the molecule. The effect was only partial at 5 h when a further progressive decrease in the ANT function occurs arising from caspase-dependent proteolysis. Conversely, at 5 h after apoptosis induction, GEN and DZN partially prevented mPTP opening, as occurs with SOD (Fig. 6). Since neither GEN nor DZN proved to have any effect on ATP levels in S-K25 cells, we conclude that their effects are a result of the prevention of oxidative damage to mitochondria rather than to broad activation.

We have also shown that GEN can reduce ROS levels in S-K5 cells. This does not depend on the inhibition by GEN of XO, which in distinction with [21], does not occur in CGCs. The reason for this anomaly is under investigation; it may be possible that the XO domain where GEN binds, thus causing competitive inhibition (see [21]) is protected by a high xanthine concentration, perhaps due to microcompartmentation or by other unknown cell component/s.

Comparison of the results reported in this paper with those of others shows that any effect due to either GEN or DZN depends on their concentrations. Cell treatment with GEN at high non-dietary concentrations results in a variety of different effects, including ATP synthase inhibition [13], mPTP opening [15] and apoptosis induction [14]. The clear lesson is that simple generalizations of the roles of these compounds cannot be made [22]. Their effects depend on a variety of factors that are related to the experimental system investigated and the flavonoid concentrations used.

## Acknowledgements

The authors thank Prof. Shawn Doonan for his critical reading of the manuscript and Mr. Vito Petragallo (IBBE, CNR) for his skilful technical assistance with tissue culture.

This work was partially financed by FIRB RBNE03B8KK\_003, Fondi di Ricerca di Ateneo del Molise (to S.P.) and MIUR – Contributi straordinari di ricerca/aree obiettivo 1 (to E.M.).

## References

- [1] Alenzi FQ. Apoptosis and diseases: regulation and clinical relevance. *Saudi Med J* 2005;26:1679–90.
- [2] Martin KR. Targeting apoptosis with dietary bioactive agents. *Exp Biol Med* 2006;231:117–29.
- [3] Zhao B. Natural antioxidants protect neurons in Alzheimer's disease and Parkinson's disease. *Neurochem Res* 2009;34:630–8.
- [4] Atlante A, de Bari L, Bobba A, Marra E, Passarella S. Transport and metabolism of L-lactate occur in mitochondria from cerebellar granule cells and are modified in cells undergoing low potassium dependent apoptosis. *Biochim Biophys Acta* 2007;1767:1285–99.
- [5] Atlante A, Gagliardi S, Marra E, Calissano P. Neuronal apoptosis in rats is accompanied by rapid impairment of cellular respiration and is prevented by scavengers of reactive oxygen species. *Neurosci Lett* 1998;245:127–30.
- [6] Atlante A, Bobba A, Calissano P, Passarella S, Marra E. The apoptosis/necrosis transition in cerebellar granule cells depends on the mutual relationship of the antioxidant and the proteolytic systems which regulate ROS production and cytochrome c release *en route* to death. *J Neurochem* 2003;84:960–71.
- [7] Bobba A, Atlante A, Giannattasio S, Sgarrella G, Calissano P, Marra E. Early release and subsequent caspase-mediated degradation of cytochrome c in apoptotic cerebellar granule cells. *FEBS Lett* 1999;457:126–30.
- [8] Atlante A, Giannattasio S, Bobba A, Gagliardi S, Petragallo V, Calissano P, et al. An increase in the ATP levels occurs in cerebellar granule cells *en route* to apoptosis in which ATP derives from both oxidative phosphorylation and anaerobic glycolysis. *Biochim Biophys Acta* 2005;1708:50–62.
- [9] Atlante A, Bobba, de Bari L, Fontana F, Calissano P, Marra E, et al. Caspase-dependent alteration of the ADP/ATP translocator triggers the mitochondrial permeability transition which is not required for the low-potassium-dependent apoptosis of cerebellar granule cells. *J Neurochem* 2006;97:1166–81.
- [10] Spencer JPE. Food for thought: the role of dietary flavonoids in enhancing human memory, learning and neuro-cognitive performance. *Proc Nutr Soc* 2008;67:238–52.
- [11] Spencer JPE. Flavonoids: modulators of brain function? *Br J Nutr* 2008;99E (Suppl 1):ES60–77.
- [12] Barnes S, Boersma B, Patel R, Kirk M, Darley-Usmar VM, Kim H, et al. Isoflavonoids and chronic disease: mechanisms of action. *Biofactors* 2000;12:209–15.
- [13] Zheng J, Ramirez VD. Inhibition of mitochondrial proton F0F1-ATPase/ATP synthase by polyphenolic phytochemicals. *Br J Pharm* 2000;130:1115–23.

- [14] Yoon HS, Moon SC, Kim ND, Park BS, Jeong MH, Yoo YH. Genistein induces apoptosis of RPE-J cells by opening mitochondrial PTP. *Biochem Biophys Res Commun* 2000;276:151–6.
- [15] Salvi M, Brunati AM, Clari G, Toninello A. Interaction of genistein with the mitochondrial electron transport chain results in opening of the membrane transition pore. *Biochim Biophys Acta* 2002;1556:187–96.
- [16] Martin KR. Targeting apoptosis with dietary bioactive agents. *Exp Biol Med* (Maywood) 2006;231:117–29.
- [17] Khan N, Afaq F, Mukhtar H. Apoptosis by dietary factors: the suicide solution for delaying cancer growth. *Carcinogenesis* 2007;28:233–9.
- [18] Zeng H, Chen Q, Zhao B. Genistein ameliorates beta-amyloid peptide (25–35)-induced hippocampal neuronal apoptosis. *Free Radic Biol Med* 2004;36:180–8.
- [19] Akiyama T, Ishida J, Nakagawa S, Ogawara H, Watanabe S, Ito N, et al. Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J Biol Chem* 1987;262:5592–5.
- [20] Ohigashi T, Ueno M, Nonnaka S, Nakanoma T, Furukawa Y, Deguchi N, et al. Tyrosin kinase inhibitors reduce bel-2 expression and induced apoptosis in androgen-depent cells. *Am J Physiol* 2000;278:C66–72.
- [21] Lin C-M, Chen C-S, Chen C-T, Liang Y-C, Lin J-K. Molecular modeling of flavonoids that inhibits xanthine oxidase. *Biochem Biophys Res Commun* 2004;324:167–72.
- [22] Das A, Banik NL, Ray SK. Mechanism of apoptosis with the involvement of calpain and caspase cascades in human malignant neuroblastoma SH-SY5Y cells exposed to flavonoids. *Int J Cancer* 2006;119:2575–85.
- [23] Linford NJ, Dorsa DM. 17 $\beta$ -Estradiol and the phytoestrogen genistein attenuate neuronal apoptosis induced by the endoplasmic reticulum calcium-ATP inhibitor thapsigargin. *Steroids* 2002;67:1029–40.
- [24] Hee K, Oh YB, Min WJ, Sang DH, Hyun SH, Kyoon H, et al. Neuroprotective effects of estrogen against beta-amyloid toxicity are mediated by estrogen receptors in cultured neuronal cells. *Neurosci Lett* 2001;302:58–62.
- [25] Shuba LM, Asai T, Pelzer S, McDonald TF. Activation of cardiac chloride conductance by the tyrosine kinase inhibitor, genistein. *Br J Pharmacol* 1996;119:335–45.
- [26] Mao Z, Zheng YL, Zhang YQ, Han BP, Zhu XW, Chang Q, et al. The anti-apoptosis effects of daidzein in the brain of D-galactose treated mice. *Molecules* 2007;12:1455–70.
- [27] Lai HH, Yen GC. Inhibitory effect of isoflavones on peroxynitrite-mediated low-density lipoprotein oxidation. *Biosci Biotechnol Biochem* 2002;66:22–8.
- [28] Guo JM, Xiao BX, Liu DH, Grant M, Zhang S, Lai YF, et al. Biphasic effect of daidzein on cell growth of human colon cancer cells. *Food Chem Toxicol* 2004;42:1641–6.
- [29] Klein CB, King AA. Genistein genotoxicity: critical considerations of in vitro exposure dose. *Toxicol Appl Pharmacol* 2007;224:1–11.
- [30] Foti P, Erba D, Riso P, Spadafranca A, Crisculi F, Testolin G. Comparison between daidzein and genistein antioxidant activity in primary and cancer lymphocytes. *Arch Biochem Biophys* 2005;433:421–7.
- [31] King RA, Bursill DB. Plasma and urinary kinetics of the isoflavones daidzein and genistein after a single soy meal in humans. *Am J Clin Nutr* 1998;67:867–72.
- [32] Sumbayev VV. Genistein effect on xanthine oxidase activity. *Ukr Biokhim Zh* 2001;73:39–43.
- [33] Lin CM, Chen CS, Chen CT, Liang YC, Lin JK. Molecular modeling of flavonoids that inhibits xanthine oxidase. *Biochem Biophys Res Commun* 2002;294:167–72.
- [34] Levi G, Aloisi F, Ciotti MT, Gallo V. Autoradiographic localization and depolarization induced release of acidic amino acids in differentiating cerebellar granule cell cultures. *Brain Res* 1984;290:77–86.
- [35] D'Mello SR, Galli C, Ciotti T, Calissano P. Induction of apoptosis in cerebellar granule neurons by low potassium: inhibition of death by insulin-like growth factor I and cAMP. *Proc Natl Acad Sci USA* 1993;90:10989–93.
- [36] Stefanis L, Troy CM, Qi H, Greene LA. Inhibitors of trypsin-like serine proteases inhibit processing of the caspase Nedd-2 and protect PC12 cells and sympathetic neurons from death evoked by withdrawal of trophic support. *J Neurochem* 1997;69:1425–37.
- [37] Stefanis L, Park DS, Friedman WJ, Greene LA. Caspase-dependent and -independent death of camptothecin-treated embryonic cortical neurons. *J Neurosci* 1999;19:6235–47.
- [38] Soto AM, Sonnenschein C. The role of estrogen on the proliferation of human breast tumor cells (MCF-7). *J Steroid Biochem* 1985;23:87–94.
- [39] Volontè C, Ciotti MT, Battistini L. Development of a method for measuring cell number: application to CNS primary neuronal cultures. *Cytometry* 1994;17:274–6.
- [40] Hockenbery D, Nunez G, Millman C, Schreiber RD, Korsmeyer SJ. Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature* 1990;348:334–6.
- [41] Berry MN, Edwards AM, Barritt GJ. Assessment of integrity of isolated hepatocytes. In: Burdon RH, Van Knippenberg PH, editors. *Laboratory techniques in chemistry and molecular biology*. 21. Amsterdam-New York-Oxford: Elsevier, Sc. Publ.; 1991. p. 83–98.
- [42] Sjödin RA. Measurement of Na<sup>+</sup>-K<sup>+</sup> pump in muscle. *Methods Enzymol* 1989;173:695–714.
- [43] Waddel WJ, Hill C. A simple ultraviolet spectrophotometer method for the determination of protein. *J Lab Clin Med* 1956;48:311–4.
- [44] Atlante A, Gagliardi S, Minervini GM, Ciotti MT, Marra E, Calissano P. Glutamate neurotoxicity in rat cerebellar granule cells: a major role for xanthine oxidase in oxygen radical formation. *J Neurochem* 1997;68:2038–45.
- [45] Atlante A, Valenti D, Gagliardi S, Passarella S. A sensitive method to assay the xanthine oxidase activity in primary cultures of cerebellar granule cells. *Brain Res Brain Res Protoc* 2000;6:1–5.
- [46] Atlante A, Calissano P, Bobba A, Azzariti A, Marra E, Passarella S. Cytochrome c is released from mitochondria in a reactive oxygen species (ROS)-dependent fashion and can operate as a ROS scavenger and as a respiratory substrate in cerebellar neurons undergoing excitotoxic death. *J Biol Chem* 2000;275:37159–66.
- [47] Bobba A, Atlante A, Petragallo VA, Marra E. Different sources of reactive oxygen species contribute to low potassium-induced apoptosis in cerebellar granule cells. *Int J Mol Med* 2008;21:737–45.
- [48] Bobba A, Atlante A, Moro L, Calissano P, Marra E. Nitric oxide has dual opposite roles during early and late phases of apoptosis in cerebellar granule neurons. *Apoptosis* 2007;12:1597–610.
- [49] Bobba A, Atlante A, de Bari L, Passarella S, Marra E. Apoptosis and cytochrome c release in cerebellar granule cells. *In Vivo* 2004;18:335–44.
- [50] Atlante A, de Bari L, Bobba A, Marra E, Calissano P, Passarella S. Cytochrome c, released from cerebellar granule cells undergoing apoptosis or excitotoxic death, can generate protonmotive force and drive ATP synthesis in isolated mitochondria. *J Neurochem* 2003;86:591–604.
- [51] Kristal BS, Brown AM. Apoptogenic ganglioside GD3 directly induces the mitochondrial permeability transition. *J Biol Chem* 1999;274:23169–75.
- [52] Basso E, Fante L, Fowlkes J, Petronilli V, Forte MA, Bernardi P. Properties of the permeability transition pore in mitochondria devoid of Cyclophilin D. *J Biol Chem* 2005;280:18558–61.
- [53] Valencia A, Moran J. Reactive oxygen species induce different cell death mechanisms in cultured neurons. *Free Rad Biol Med* 2004;36:1112–25.
- [54] Mercer LD, Kelly BL, Horne MK, Beart PM. Dietary polyphenols protect dopamine neurons from oxidative insults and apoptosis: investigations in primary rat mesencephalic cultures. *Biochem Pharmacol* 2005;69:339–45.
- [55] Szkudelska K, Nogowski L. Genistein – a dietary compound inducing hormonal and metabolic changes. *J Steroid Biochem Mol Biol* 2007;105:37–45.
- [56] Takahashi Y, Odbayar TO, Ide T. A Comparative Analysis of Genistein and Daidzein in Affecting Lipid Metabolism in Rat Liver. *J Clin Biochem Nutr* 2009;44:223–30.
- [57] Cai Q, Wei H. Effect of dietary genistein on antioxidant enzyme activities in SENCAR mice. *Nutr Cancer* 1996;25:1–7.
- [58] Choi EJ. The prooxidant, rather than antioxidant, acts of daidzein in vivo and in vitro: daidzein suppresses glutathione metabolism. *Eur J Pharmacol* 2006;542:162–9.
- [59] Wijeratne SS, Cuppett SL. Soy isoflavones protect the intestine from lipid hydroperoxide mediated oxidative damage. *J Agric Food Chem* 2007;55:9811–6.
- [60] Paul MK, Mukhopadhyay AK. Tyrosine kinase - Role and significance in Cancer. *Int J Med Sci* 2004;1:101–15.
- [61] Sarkar FH, Li Y. Mechanisms of cancer chemoprevention by soy isoflavone genistein. *Cancer Metastasis Rev* 2002;21:265–80.
- [62] Shi C, Xu J. Increased vulnerability of brain to estrogen withdrawal-induced mitochondrial dysfunction with aging. *J Bioenerg Biomembr* 2008;40:625–30.
- [63] Santos AC, Uyemura SA, Lopes JL, Bazon JN, Mingatto FE, Curti C. Effect of naturally occurring flavonoids on lipid peroxidation and membrane permeability transition in mitochondria. *Free Radic Biol Med* 1998;24:1455–61.