



Biochemical Pharmacology, Volume 79, issue 12, 15 June 2010


Contents

COMMENTARY

RLIP76: A versatile transporter and an emerging target for cancer therapy

1699–1705


Rit Vatsyayan, Poorna Chandra Rao Lelsani, Sanjay Awasthi, Sharad S. Singhal



Progenitor cell-derived smooth muscle cells in vascular disease

1706–1713

Augusto Orlandi, Martin Bennett




ANTIBIOTICS AND CHEMOTHERAPEUTICS

Suppression of PMA-induced tumor cell invasion by dihydroartemisinin via inhibition of PKC α /Raf/MAPKs and NF- κ B/AP-1-dependent mechanisms

1714–1726


Yong Pil Hwang, Hyo Jeong Yun, Hyung Gyun Kim, Eun Hee Han, Gye Won Lee, Hye Gwang Jeong

Proposed signal transduction pathways by which PMA induced and DHA-inhibited invasion and migration of HT-1080 cells.




Inhibition of calcium-independent phospholipase A₂ activates p38 MAPK signaling pathways during cytostasis in prostate cancer cells1727–1735

Bin Sun, Xiaoling Zhang, Christopher Yonz, Brian S. Cummings
Role of iPLA₂ in prostate cancer cell signaling and growth.




Mode of action of Nifurtimox and N-oxide-containing heterocycles against Trypanosoma cruzi: Is oxidative stress involved?1736–1745

Mariana Boiani, Lucia Piacenza, Paola Hernández, Lucia Boiani, Hugo Cerecetto, Mercedes González, Ana Denicola
No production of ROS/RNS was observed by *Trypanosoma cruzi* epimastigotes treated with several N-oxide-containing heterocycles and Nifurtimox at 25 µM.




Different contribution of BH3-only proteins and caspases to doxorubicin-induced apoptosis in p53-deficient leukemia cells1746–1758

Nuria López-Royuela, Patricia Pérez-Galán, Patricia Galán-Malo, Victor J. Yuste, Alberto Anel, Santos A. Susín, Javier Naval, Isabel Marzo



Modulation of stress genes expression profile by nitric oxide-releasing aspirin in Jurkat T leukemia cells1759–1771


Niharika Nath, Mitali Chattopadhyay, Ravinder Kodela, Song Tian, Peter Vlismas, Daniel Boring, James A. Crowell, Khosrow Kashfi



An RNA-directed nucleoside anti-metabolite, 1-(3-C-ethynyl-beta-D-ribo-pentofuranosyl)cytosine (ECyd), elicits antitumor effect via *TP53*-induced Glycolysis and Apoptosis Regulator (TIGAR) downregulation

1772–1780


Vivian Wai Yan Lui, Cecilia Pik Yuk Lau, Crystal Sao Fong Cheung, Kakiu Ho, Margaret Heung Ling Ng, Suk Hang Cheng, Bo Hong, Sai-Wah Tsao, Chi Man Tsang, Kenny Ieng Kit Lei, Yasundo Yamasaki, Akira Mita, Anthony T.C. Chan



Remarkable interference with telomeric function by a G-quadruplex selective bisantrene regioisomer

1781–1790

Marco Folini, Claudia Pivetta, Giuseppe Zagotto, Cinzia De Marco, Manlio Palumbo, Nadia Zaffaroni, Claudia Sissi




CARDIOVASCULAR PHARMACOLOGY

D-Serine modulates neurogenic relaxation in rat corpus cavernosum

1791–1796

Mehdi Ghasemi, Fatemeh Rezania, Jackie Lewin, Kevin P. Moore, Ali R. Mani




INFLAMMATION AND IMMUNOPHARMACOLOGY

The intestinal antiinflammatory agent glycomacropeptide has immunomodulatory actions on rat splenocytes


1797–1804

Pilar Requena, Raquel González, Rocío López-Posadas, Ana Abadía-Molina, María Dolores Suárez, Antonio Zarzuelo, Fermín Sánchez de Medina, Olga Martínez-Augustín



Endogenous *N*-acyl-dopamines induce COX-2 expression in brain endothelial cells by stabilizing mRNA through a p38 dependent pathway 1805–1814

Carmen M. Navarrete, Moisés Pérez, Amaya García de Vinuesa, Juan A. Collado, Bernd L. Fiebich, Marco A. Calzado, Eduardo Muñoz



Schematic diagram describing the effect of endogenous *N*-acyl-dopamines on COX-2 expression in brain endothelial cells.

Falcarinol is a covalent cannabinoid CB₁ receptor antagonist and induces pro-allergic effects in skin 1815–1826


Marco Leonti, Laura Casu, Stefan Raduner, Filippo Cottiglia, Costantino Floris, Karl-Heinz Altmann, Jürg Gertsch



METABOLIC DISORDERS AND ENDOCRINOLOGY

The dopamine receptor D2 agonist bromocriptine inhibits glucose-stimulated insulin secretion by direct activation of the α 2-adrenergic receptors in beta cells 1827–1836

J.E. de Leeuw van Weenen, E.T. Parlevliet, P. Maechler, L.M. Havekes, J.A. Romijn, D.M. Ouwens, H. Pijl, B. Guigas



CORRIGENDUM

Corrigendum to “Heteronemin, a spongean sesterterpene, inhibits TNF α -induced NF- κ B activation through proteasome inhibition and induces apoptotic cell death” [Biochem. Pharmacol. 79 (2010) 610–622] 1837

Marc Schumacher, Claudia Cerella, Serge Eifes, Sébastien Chateaufvieux, Franck Morceau, Marcel Jaspars, Mario Dicato, Marc Diederich

INDEXED/ABSTRACTED IN: Curr. Cont. ASCA, Biosis Data, CAB Inter., Chemical Abstracts Service, Curr. Cont./Life Sci., CABS, EMBASE/Excerpt. Med., Curr. Cont. ISI/BIOMED Database, MEDLINE, PASCAL-CNRS Data, Curr. Cont. Sci. Cit. Ind., Curr. Cont. SCISEARCH Data, Ind. Med., Reference Update. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®.



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

www.elsevier.com/locate/biochempharm