



An autoradiographic survey of mouse brain nicotinic acetylcholine receptors defined by null mutants

Christopher G. Baddick, Michael J. Marks *

Institute for Behavioral Genetics, University of Colorado, Boulder, CO, USA

ARTICLE INFO

Article history:

Received 24 March 2011

Accepted 29 April 2011

Available online 10 May 2011

Keywords:

Nicotinic acetylcholine receptor

Null mutant mice

Epibatidine

A-85380

α -Conotoxin MII

α -Bungarotoxin

ABSTRACT

Nine nicotinic receptor subunits are expressed in the central nervous system indicating that a variety of nicotinic acetylcholine receptors (nAChR) may be assembled. A useful method with which to identify putative nAChR is radioligand binding. In the current study the binding of [125 I] α -bungarotoxin, [125 I] α -conotoxin MII, 5[125 I]-3-((2S)-azetidinylmethoxy)pyridine (A-85380), and [125 I]epibatidine has been measured autoradiographically to provide data on many nAChR binding sites. Each binding site was evaluated semi-quantitatively for samples prepared from wild-type and α 2, α 4, α 6, α 7, β 2, β 4, α 5 and β 3 null mutant mice. Deletion of the α 7 subunit completely and selectively eliminated [125 I] α -bungarotoxin binding. The binding of [125 I] α -conotoxin MII was eliminated in most brain regions by deletion of either the α 6 or β 2 subunit and is reduced by deletion of either the α 4 or β 3 subunit. The binding of 5[125 I]A-85380 was completely eliminated by deletion of the β 2 subunit and significantly reduced by deletion of the α 4 subunit. Most, but not all, α 4-independent sites require expression of the α 6 subunit. The effect of gene deletion on total [125 I]epibatidine binding was very similar to that on [125 I]A-85380 binding. [125 I]Epibatidine also labels β 4* nAChR, which was readily apparent for incubations conducted in the presence of 100 nM cytosine. The effects of α 3 gene deletion could not be evaluated, but persistence of residual sites implies the expression of α 3* nAChR. Taken together these results confirm and extend previously published evaluations of the effect of nAChR gene deletion and help to define the nAChR subtypes measurable by ligand binding.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

1.1. Historical ligands

Nicotinic acetylcholine receptors (nAChRs) in brain have been studied using ligand binding assays for many years. Early studies of [125 I] α -bungarotoxin (α Bgt) [1] and [3 H]-nicotine [2] binding to rat brain membrane provided some of the first evidence that nAChRs might be expressed in brain. Subsequently, the demonstration that α Bgt and nicotine binding sites had different anatomical distributions [3] and biochemical properties [4] provided the first evidence that more than one putative nAChR subtype is expressed in brain. When the nine nAChR subunit genes expressed in mammalian brain (α 2– α 7, β 2– β 4) were cloned and sequenced nearly 20 years ago, the number of potential subtypes

expanded dramatically [5,6]. Much of the recent research has attempted to identify the subunit compositions of those nAChR subtypes that are actually expressed (i.e. native receptors) in brain as well as other tissues [7].

1.2. Epibatidine

Although [125 I] α Bgt and [3 H]nicotine (as well as [3 H]acetylcholine [8], [3 H]cytosine [9] and [3 H]methylcarbaryl [10]) have been useful in identifying and characterizing putative nAChR binding sites in brain, these ligands do not label a wide array of binding sites. This deficiency was overcome with the discovery and characterization of epibatidine [11]. Epibatidine binds with extraordinarily high affinity to rat brain membranes [11–13]. Saturation binding and differential inhibition studies in rat brain [13,14] and inhibition experiments in mouse brain [15,16] demonstrated that epibatidine binds to multiple nAChR subtypes. Furthermore, epibatidine potentially activates α 3 β 2, α 3 β 4, α 4 β 2, α 7, and α 8 nAChRs expressed in *Xenopus laevis* oocytes [17]. Epibatidine also binds with very high affinity to heteromeric receptors expressed in *Xenopus* oocytes [18] and HEK cells [19].

Abbreviations: nAChR, nicotinic cholinergic receptor; α Bgt, α -bungarotoxin; A-85380, 3-((2S)-azetidinylmethoxy)pyridine; α Ctx MII, α -conotoxin MII.

* Corresponding author at: Institute for Behavioral Genetics, University of Colorado, 1480 30th Street, Campus Box 447, Boulder, CO, USA.

Tel.: +1.303.492.9677, fax: +1 303 492 8063.

E-mail address: marksm@colorado.edu (M.J. Marks).

1.3. α ConotoxinMII

The ability to measure other nAChR subtypes was expanded by the discovery and initial characterization of α ConotoxinMII (α CtxMII) [20] and the demonstration that this ligand labels a unique population of nAChR binding sites in mouse brain that is concentrated in catecholaminergic cells and their terminals and in visual pathways [21,22]. This distribution represents a subset of epibatidine binding sites that can also be visualized as a subset of the sites observed when epibatidine binding is conducted in the presence of a low concentration of cytosine (50 nM) [22].

1.4. A-85380

As part of the nicotinic research program at Abbott Laboratories a potent, high-affinity ligand 3-((2S)-azetidinylmethoxy)-pyridine (A-85380) has been developed [23]. Radiolabeled 5- 125 I-A-85380 labels $\beta 2^*$ nAChR selectively [24] and its analogs are extremely useful ligands for positron emission tomography [25,26].

1.5. Knock-out mice

Nicotinic receptor knock-out mice have proven to be valuable tools to identify native nAChR expression and function [27–29]. Knock-out mice have been used to define populations of ligand sites that identify natively expressed nAChR subtypes [22,30–48]. The results described in this study have extended the information by examining the effect of nAChR subunit gene deletion on the binding of [125 I]epibatidine, [125 I]A-85380, [125 I] α CtxMII and [125 I] α Bgt in order to provide a comprehensive overview of the expression of native nAChR in mouse brain. Although some compensation is likely to occur following deletion of a nAChR subunit, current evidence indicates that no nAChR subtypes that are not normally present are expressed following deletion of major nAChR subunits such as $\alpha 3$ [49], $\alpha 4$ [30,46,50], $\alpha 6$ [22,51–53], $\alpha 7$ [33,44], $\beta 2$ [32,39,42,44,45,54] and $\beta 4$ [44,54]. However, deletion of the auxiliary subunits $\alpha 5$ [51,55,56] or $\beta 3$ [51,55] appears to change the relative ratio of nAChR that mediated synaptosomal dopamine release with differential sensitivity to inhibition by α CtxMII.

2. Materials and methods

2.1. Materials

The radioligands [125 I]epibatidine (specific activity 2200 Ci/mmol), 5[125 I]-A-85380 (2200 Ci/mmol) and [125 I] α bungatotoxin (α Bgt) (250 mCi/mmol) and Kodak MR film were obtained from PerkinElmer New England Nuclear, Shelton, CT, USA. α ConotoxinMII (α CtxMII) and [125 I] α Ctx MII (2200 Ci/mmol) were prepared as described previously ([20,21], respectively) were obtained from J. Michael McIntosh, University of Utah, Salt Lake City, UT, USA. Unlabeled I-epibatidine was a gift from Kenneth Kellar, Georgetown University, Washington, DC. Unlabeled 5I-A-85380 was purchased from Tocris Bioscience, Ellisville, MO, USA. HEPES (free acid and Na salt) are products of BDH and were purchased from VWR, Chester, PA, USA. The following chemicals were purchased from Sigma Chemical Co., St. Louis, MO, USA: 2-methylbutane, NaCl, KCl, CaCl_2 , MgSO_4 , bovine serum albumin, leupeptin, pepstatin, aprotinin, EDTA, EGTA, and phenylmethylsulfonyl fluoride (PMSF). M-1 Embedding Matrix was purchased from Anatomical Pathology USA, Pittsburgh, PA, USA. Superfrost Plus Microscope Slides were obtained from Fisher Scientific, Fair Lawn, NJ, USA.

2.2. Mice

All procedures involving mice were reviewed and approved by the Animal Care and Utilization Committee of the University of Colorado, Boulder, CO, USA. Mice were bred in the Specific Pathogen Free Colony at the Institute for Behavioral Genetics, University of Colorado, Boulder, CO, USA weaned at 25 days of age and housed with like-sexed littermates. Animals were maintained on a 12 h light/12 h dark cycle (lights on 7 AM–7 PM) and allowed free access to food and water. The following nicotinic knock-out mice were used in this study: $\alpha 2$ [57]; $\alpha 4$ [50]; $\alpha 6$ [22]; $\alpha 7$ [33]; $\beta 2$ [32]; $\beta 4$ [54]; $\alpha 5$ [36] and $\beta 3$ [35]. Mice differing in nAChR genotype were derived by mating heterozygotes that had been backcrossed to C57BL/6J for at least 10 generations. Tail clippings were obtained from mice about 40 days of age and genotype determined as described previously [51].

2.3. Autoradiography

Mice were sacrificed by cervical dislocation, the brains rapidly removed and quickly frozen by immersion in isopentane (-35°C). Brains were stored at -70°C until sectioning. Brains were sectioned (14 μ thickness) using an IEC or Leica cryostat. Sections were thaw mounted on Fisher Superfrost Plus microscope slides. Slides containing the sections were stored at -70°C until use. For all binding conditions samples were removed from the -70°C freezer and warmed to room temperature under vacuum in a desiccator. After warming the slides were distributed to plastic holders modified to contain 50 slides.

The binding of 6-[125 I] epibatidine was conducted as follows [58]. Samples were incubated in binding buffer (NaCl, 140 mM; KCl, 1.5 mM; CaCl_2 , 2 mM; MgSO_4 , 1 mM; bovine serum albumin, 1 g/L; HEPES buffer, pH 7.5, 25 mM) containing 500 nM [125 I]epibatidine with a final specific activity of 110 Ci/mmol (attained by diluting the commercial [125 I]epibatidine (2200 Ci/mmol) with unlabeled 6I-epibatidine) for 2 h at 22° . Three different incubation conditions were used. The first condition measured total [125 I]epibatidine binding and contained no further additions to the buffer. The second condition included 100 nM cytosine. The third condition included 100 nM cytosine and 100 nM α -conotoxin MII. Following the incubation, the slides were washed by immersion in ice-cold protein free binding buffer (2×30 s), ice-cold $0.1 \times$ protein-free binding buffer (2×10 s) and ice-cold 5 mM HEPES, pH 7.5 (2×5 s each). Samples were then dried under a gentle stream of air and desiccated overnight before exposure to Kodak MR film for 17 days.

The binding of [125 I]A-85380 was conducted as follows [59]. Samples were incubated in binding buffer containing 200 pM [125 I]A-85380 with a final specific activity of 110 Ci/mmol (attained by diluting the commercial [125 I]A-85380 (2200 Ci/mmol with unlabeled 5I-A-85380) for 2 h at 22° . Following the incubation, the slides were washed by immersion in ice-cold protein free binding buffer (2×30 s), ice-cold $0.1 \times$ protein-free binding buffer (2×10 s) and ice-cold 5 mM HEPES, pH 7.5 (2×5 s each). Samples were then dried under a gentle stream of air and desiccated overnight before exposure to Kodak MR film for 17 days.

The binding of [125 I] α -CtxMII was conducted as follows [21]. Samples were incubated for 10 min in binding buffer without BSA but including 1 mM PMSF for 10 min. Samples were then incubated with 0.5 nM [125 I] α CtxMII in binding buffer that also contained 5 mM EDTA and 5 mM EGTA as well as leupeptin, pepstatin and aprotinin (each 10 $\mu\text{g}/\text{ml}$) for 2 h at 22° . Following the incubation, samples were incubated twice for 15 min at 22° in binding buffer. Subsequently, the slides were washed by immersion in ice-cold protein free binding buffer (2×30 s), ice-cold $0.1 \times$

protein-free binding buffer (2×10 s) and ice-cold 5 mM HEPES, pH 7.5 (2×5 s each). Samples were then dried under a gentle stream of air and desiccated overnight before exposure to Kodak MR film for 5 days.

The binding of [125 I] α Bgt was conducted as follows [60]. Samples were incubated with 0.25 nM [125 I] α Bgt in binding buffer for 2 h at 22°. Following the incubation, samples were incubated twice for 30 min at 22° in binding buffer. Subsequently, the slides were washed by immersion in ice-cold protein free binding buffer (2×30 s), ice-cold 0.1 \times protein-free binding buffer (2×10 s) and ice-cold 5 mM HEPES, pH 7.5 (2×5 s each). Samples were then dried under a gentle stream of air and desiccated overnight before exposure to Kodak MR film for 5 days.

2.4. Photography

Films were placed on a Northern Light Precision Illuminator set to full intensity and images were captured using a Nikon D40 camera equipped with a 60 mm microimaging extension lens. Ambient light was kept constant. Subsequently images were transferred to Photoshop for further processing.

2.5. Estimate of signal intensity

Visual assessment was used to obtain semi-quantitative estimates of signal intensity. Relative binding among regions was determined and graded as follows: ■■■■, highest labeling; ■■■, substantial labeling; ■■, modest labeling; ■, weak labeling; □, no significant labeling. The effect of gene deletion for each brain region relative to that of wild-type mice was graded as follows: +++, 75–100% of maximum (virtually no effect); ++, 50–75%; +, 25–50%; <25%; and –, no detectable signal remaining.

3. Results

3.1. Autoradiographic Illustration of the effect of nAChR gene deletion on ligand binding

The autoradiograms in Fig. 1 illustrate pictorially the effect of deletion of $\alpha 2$, $\alpha 4$, $\alpha 6$, $\alpha 7$, $\beta 2$, $\beta 4$, $\alpha 5$ and $\beta 3$ on the binding of [125 I]epibatidine, [125 I]A-85380, [125 I] α CtxMII and [125 I] α Bgt at approximately -3.5 mm Bregma. This figure illustrates several features of the data. The binding of each ligand to tissue from wild-




Fig. 1. Autoradiographic images of coronal mouse brain sections at approximately -3.5 mm Bregma. Autoradiograms for each of the ligands [125 I]epibatidine (three conditions: total binding, binding in the presence of 100 nM cytosine and binding in the presence of 100 nM cytosine and 100 nM α -conotoxinMII), [125 I]A-85380, [125 I] α -conotoxinMII and [125 I] α -bungarotoxin are shown for wild-type mice and for each of the nAChR null mutants for sections at a level approximately -3.5 mm Bregma.

type mice is shown in the top row of the figure. The effect of each nAChR gene deletion is shown for each binding condition below that of the wild-type. Similar figures at different anatomical levels (approximate Bregma: –0.1 mm, –0.6 mm, –2.1 mm, –2.5 mm and –5.2 mm) are found in the [Supplementary data, Figs. 1–5](#). These images can also be accessed and downloaded at <http://ibgwww.colorado.edu> (under the animal research heading). The descriptions that follow have also been applied to the results summarized in [Tables 1–6](#) in the following sections.

3.1.1. [¹²⁵I]αBungarotoxin binding

The simplest response to nAChR gene deletion is observed for [¹²⁵I]αBgt binding. Highest binding is observed in superior colliculus and hippocampus with less intense labeling of cortical and midbrain regions. Specific binding is eliminated by deletion of the α7 gene and is virtually unaffected by the other null mutations.

3.1.2. [¹²⁵I]α-ConotoxinMII binding

At this level only the superior colliculus shows intense [¹²⁵I]αCtXMII binding, although light labeling of the interpeduncular nucleus is also observed. Deletion of α2, α7, β4 or α5 subunits has no detectable effect on these binding sites. In contrast, deletion of the β2 and α6 subunits completely eliminated [¹²⁵I]αCtXMII binding in superior colliculus. A significant, but not quite complete, reduction was observed in β3 knock-out mice, while a more modest reduction occurred in α4 knock-out mice.

3.1.3. [¹²⁵I]A-85380 Binding

Significantly more sites are labeled by [¹²⁵I]A-85380 than by either [¹²⁵I]αBgt or [¹²⁵I]αCtXMII. The binding of this ligand is completely eliminated by deletion of the β2 subunit. Deletion of the α4 subunit also eliminates many [¹²⁵I]A-85380 binding sites. However, significant labeling persists in the superficial gray region of the superior colliculus and in the interpeduncular nucleus. A small amount of binding is also seen in the substantia nigra. In contrast to the widespread effects observed following β2 and α4 gene deletion, the effects of deletion of the α6 or β3 gene are restricted to the superficial gray area of the superior colliculus. No significant reductions were noted for α2, α7, β4 and α5 null mutants.

3.1.4. [¹²⁵I]Epibatidine binding

The general pattern of [¹²⁵I]epibatidine binding is similar to that observed with [¹²⁵I]A-85380 and the effects of β2, α4, α6 and β3 gene deletion are also similar. However, intense labeling in the interpeduncular nucleus persists in the β2 knock-out mice.

It is known that [¹²⁵I]epibatidine binding is heterogeneous [15,16,58] with a subset of the sites resistant to inhibition by cytosine. The effect of the null mutations demonstrates the heterogeneity of the cytosine-resistant [¹²⁵I]epibatidine binding sites. Deletion of β2 eliminates binding in the superior colliculus and deletion of either α6 or β3 significantly reduces this binding. Deletion of α4 has less effect, while no detectable reductions were observed in α2, α7, β4 or α5 null mutants. However, deletion of β4 significantly reduces labeling in the interpeduncular nucleus, while no significant effects were noted for the other null mutants.

A subset of the cytosine-resistant [¹²⁵I]epibatidine binding sites can be inhibited by αCtXMII [39]. Deletion of the β4 subunit has the most noticeable effect on those residual sites that are resistant to both cytosine and αCtXMII inhibition in interpeduncular nucleus.

3.2. Semi-quantitative analysis of nAChR gene deletion on nicotinic binding sites

All nAChR in the CNS must include either the α7, β2 and/or β4 subunit [31,44,61]. The heteromeric nAChR include α subunits and

Table 1

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –0.1 mm Bregma.

	Cpu	GP	S Cx	M Cx	OT
0.5 nM [¹²⁵ I]-Epibatidine					
WT	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++
α4	+	–	–	–	–
α6	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++
β2	–	–	–	–	–
β4	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM cytosine					
WT	■	■	□	□	□
α2	++++	++++	n/a	n/a	n/a
α4	–	–	n/a	n/a	n/a
α6	++++	++++	n/a	n/a	n/a
α7	++++	++++	n/a	n/a	n/a
β2	–	–	n/a	n/a	n/a
β4	++++	++++	n/a	n/a	n/a
α5	++++	++++	n/a	n/a	n/a
β3	++++	++++	n/a	n/a	n/a
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytosine + 100 nM α-CtXMII					
WT	■	■	□	□	□
α2	++++	++++	n/a	n/a	n/a
α4	–	–	n/a	n/a	n/a
α6	++++	++++	n/a	n/a	n/a
α7	++++	++++	n/a	n/a	n/a
β2	–	–	n/a	n/a	n/a
β4	++++	++++	n/a	n/a	n/a
α5	++++	++++	n/a	n/a	n/a
β3	++++	++++	n/a	n/a	n/a
200 pM 5-[¹²⁵ I]A-85380					
WT	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++
α4	+	+	–	–	–
α6	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++
β2	–	–	–	–	–
β4	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I] α-ConotoxinMII					
WT	■ ■ ■	□	□	□	□
α2	++++	n/a	n/a	n/a	n/a
α4	+++	n/a	n/a	n/a	n/a
α6	–	n/a	n/a	n/a	n/a
α7	++++	n/a	n/a	n/a	n/a
β2	–	n/a	n/a	n/a	n/a
β4	++++	n/a	n/a	n/a	n/a
α5	++++	n/a	n/a	n/a	n/a
β3	+	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I] α-Bungarotoxin					
WT	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++
α4	++++	++++	++++	++++	++++
α6	++++	++++	++++	++++	++++
α7	–	–	–	–	–
β2	++++	++++	++++	++++	++++
β4	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At Bregma = –0.1 mm, the following brain regions are visible. Caudate Putamen, CPU; Globus Pallidus, GP; Somatosensory Cortex, S Cx; Motor Cortex, M Cx; Olfactory Tubercles, OT. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

Table 2

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –0.6 mm Bregma.

	Cpu	GP	Th	MHb	S Cx	M Cx
0.5 nM [¹²⁵ I]-Epibatidine						
WT	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++
α4	+	—	—	++++	—	—
α6	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++
β2	—	—	—	++++	—	—
β4	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine						
WT	■ ■	□	■ ■	■ ■ ■ ■	■	■
α2	+++	n/a	+++	++++	+++	+++
α4	—	n/a	—	++++	—	—
α6	—	n/a	++++	++++	++++	++++
α7	+++	n/a	++++	++++	++++	++++
β2	—	n/a	—	+++	—	—
β4	—	n/a	++++	+	++++	++++
α5	+++	n/a	++++	++++	++++	++++
β3	—	n/a	+++	++++	+++	+++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine + 100 nM α-CtxMII						
WT	■	□	■ ■	■ ■ ■ ■	■	■
α2	+++	n/a	+++	++++	+++	+++
α4	—	n/a	—	++++	—	—
α6	—	n/a	++++	++++	++++	++++
α7	+++	n/a	++++	++++	++++	++++
β2	—	n/a	—	+++	—	—
β4	—	n/a	++++	+	++++	++++
α5	+++	n/a	++++	++++	++++	++++
β3	—	n/a	+++	++++	+++	+++
200 pM 5-[¹²⁵ I]-A85380						
WT	■ ■ ■	■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++
α4	+	—	—	++	—	—
α6	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++
β2	—	—	—	—	—	—
β4	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I]α-ConotoxinMII						
WT	■ ■ ■ ■	□	□	□	□	□
α2	++++	n/a	n/a	n/a	n/a	n/a
α4	+	n/a	n/a	n/a	n/a	n/a
α6	—	n/a	n/a	n/a	n/a	n/a
α7	++++	n/a	n/a	n/a	n/a	n/a
β2	—	n/a	n/a	n/a	n/a	n/a
β4	++++	n/a	n/a	n/a	n/a	n/a
α5	++++	n/a	n/a	n/a	n/a	n/a
β3	+	n/a	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I]α-Bungarotoxin						
WT	■ ■	■ ■	■ ■	■ ■	■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++
α4	++++	++++	++++	++++	++++	++++
α6	++++	++++	++++	++++	++++	++++
α7	—	—	—	—	—	—
β2	++++	++++	++++	++++	++++	++++
β4	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At bregma = –0.6 mm the following brain regions are visible: Caudate Putamen, CPU; Globus Pallidus, GP; Thalamus, Th; Medial Habenular, MHb; Somatosensory Cortex, S Cx; Motor Cortex, M Cx. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

may also include the auxiliary α5 and β3 subunits. The results that follow will discuss the effect of the deletion of each nAChR subunit on the various binding sites. The semi-quantitative analyses are summarized in Tables 1–6 for analyses of binding at the levels of approximately –0.1 mm, –0.6 mm, –2.1 mm, –2.5 mm, –3.5 and –5.2 mm Bregma, respectively. The symbols for each ligand in the

rows for the wild-type mice (□, ■, ■ ■, ■ ■ ■, and ■ ■ ■ ■) illustrate the expression level from undetectable (□) to most intense (■ ■ ■ ■). The symbols for each null mutant (++++, +++, ++, + and –) illustrate the relative effect of that particular gene deletion from no significant change from control (++++) to complete elimination (–).

Table 3

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –2.1 mm Bregma.

	Th	opt	DLG	MHb	fr	HP	Au1	RSG
0.5 nM [¹²⁵ I]-Epibatidine								
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	—	++	++	++++	++++	—	—	—
α6	++++	++++	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++	++++
β2	—	—	—	++++	++++	—	—	—
β4	++++	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine								
WT	■	■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	□	□	□
α2	++++	+++	++++	++++	++++	n/a	n/a	n/a
α4	++++	—	+++	++	++	n/a	n/a	n/a
α6	++++	+++	+++	++++	++++	n/a	n/a	n/a
α7	++++	+++	++++	++++	++++	n/a	n/a	n/a
β2	++++	—	—	+++	+++	n/a	n/a	n/a
β4	++++	—	+++	+	+	n/a	n/a	n/a
α5	++++	+++	+++	++++	++++	n/a	n/a	n/a
β3	++++	—	+++	++++	++++	n/a	n/a	n/a
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine + 100 nM α-CtxMII								
WT	■	■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	□	□	□
α2	++++	+++	++++	++++	++++	n/a	n/a	n/a
α4	++++	—	+++	++	++	n/a	n/a	n/a
α6	++++	+++	+++	++++	++++	n/a	n/a	n/a
α7	++++	+++	++++	++++	++++	n/a	n/a	n/a
β2	++++	—	—	+++	+++	n/a	n/a	n/a
β4	++++	—	+++	+	+	n/a	n/a	n/a
α5	++++	+++	+++	++++	++++	n/a	n/a	n/a
β3	++++	—	+++	++++	++++	n/a	n/a	n/a
200 pM 5-[¹²⁵ I]-A85380								
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	—	+	+	+++	++	—	—	—
α6	++++	++++	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++	++++
β2	—	—	—	—	—	—	—	—
β4	++++	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I] α-ConotoxinMII								
WT	□	■ ■ ■ ■	■ ■	■ ■	□	□	□	□
α2	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
α4	n/a	++	+	++	n/a	n/a	n/a	n/a
α6	n/a	++	—	—	n/a	n/a	n/a	n/a
α7	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
β2	n/a	++	—	+	n/a	n/a	n/a	n/a
β4	n/a	++++	++++	+++	n/a	n/a	n/a	n/a
α5	n/a	++	++++	++++	n/a	n/a	n/a	n/a
β3	n/a	++	+	++	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I] α-Bungarotoxin								
WT	■ ■ ■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■	■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	++++	++++	++++	++++	++++	++++	++++	++++
α6	++++	++++	++++	++++	++++	++++	++++	++++
α7	—	—	—	—	—	—	—	—
β2	++++	++++	++++	++++	++++	++++	++++	++++
β4	++++	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At bregma = –2.1 mm the following brain regions are visible: Thalamus, Th; Optic Tract, opt; Dorsal Lateral Geniculates, DLG; Medial Habenular, MHb; Fasciculus Retroflexus, fr; Hippocampus, HP; Primary Auditory Cortex, Au1; Retrosplenial Granular Cortex, RSG. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

3.2.1. α7 nAChR gene deletion

Deletion of the α7 nAChR gene gives the simplest pattern: complete elimination of [¹²⁵I]αBgt binding in every brain region with no significant effect on the binding of [¹²⁵I]epibatidine, [¹²⁵I]A-85380, or [¹²⁵I]αCtxMII.

3.2.2. β2 nAChR gene deletion

Deletion of the β2 nAChR gene has the most effect on the ligand binding sites, although it had no noticeable effect on [¹²⁵I]αBgt binding. Deletion of the β2 nAChR completely eliminates both [¹²⁵I]A-85380 and high-affinity [¹²⁵I]αCtxMII in every brain region

Table 4

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –2.5 mm Bregma.

	DS	VLG	DLG	OPT	fr	HP	V1	RSG
0.5 nM [¹²⁵ I]-Epibatidine								
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	—	++	++	++++	++++	—	—	—
α6	++++	+	+	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++	++++
β2	—	—	—	—	++++	—	—	—
β4	++++	++++	++++	++++	+	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	+	+	++++	+++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine								
WT	□	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	□	□	□
α2	n/a	+++	+++	+++	++++	n/a	n/a	n/a
α4	n/a	++	++	++	+++	n/a	n/a	n/a
α6	n/a	++	++	+	++++	n/a	n/a	n/a
α7	n/a	+++	+++	+++	++++	n/a	n/a	n/a
β2	n/a	—	—	—	++++	n/a	n/a	n/a
β4	n/a	++	++	++	+	n/a	n/a	n/a
α5	n/a	++++	++++	++++	+++	n/a	n/a	n/a
β3	n/a	++	++	+	+++	n/a	n/a	n/a
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine + 100 nM α-CtxMII								
WT	Υ	■ ■ ■	■ ■ ■	■ ■ ■	■ ■ ■ ■	□	□	□
α2	n/a	++++	++++	+++	++++	n/a	n/a	n/a
α4	n/a	+++	+++	++	++++	n/a	n/a	n/a
α6	n/a	+++	+++	+	++++	n/a	n/a	n/a
α7	n/a	++++	++++	+++	++++	n/a	n/a	n/a
β2	n/a	—	—	—	++++	n/a	n/a	n/a
β4	n/a	+++	+++	++	+++	n/a	n/a	n/a
α5	n/a	++++	++++	++++	++++	n/a	n/a	n/a
β3	n/a	++	++	+	++++	n/a	n/a	n/a
200 pM 5-[¹²⁵ I]-A85380								
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	—	++	++	++	++++	—	—	—
α6	++++	++	+++	++	—	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++	++++
β2	—	—	—	—	—	—	—	—
β4	++++	++++	++++	++	++++	++++	++++	++++
α5	++++	++++	++	++++	++++	++++	++++	++++
β3	++++	++	+++	++	—	++++	++++	++++
0.5 nM [¹²⁵ I] α-ConotoxinMII								
WT	Υ	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	□	□	□	□
α2	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
α4	n/a	++	++	++	n/a	n/a	n/a	n/a
α6	n/a	—	—	—	n/a	n/a	n/a	n/a
α7	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
β2	n/a	—	—	—	n/a	n/a	n/a	n/a
β4	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
α5	n/a	++++	++++	++++	n/a	n/a	n/a	n/a
β3	n/a	+	+	+	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I] α-Bungarotoxin								
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++
α4	++++	++++	++++	++++	++++	++++	++++	++++
α6	++++	++++	++++	++++	++++	++++	++++	++++
α7	—	—	—	—	—	—	—	—
β2	++++	++++	++++	++++	++++	++++	++++	++++
β4	++++	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At bregma = –2.5 mm the following brain regions are visible: Dorsal Subiculum, DS; Ventral Lateral Geniculates, VLG; Dorsal Lateral Geniculates, DLG; Olivary Pretectal Nucleus, OPT; Fasciculus Retroflexus, fr; Hippocampus, HP; Primary Visual Cortex, V1; Retrosplenial Granular Cortex, RSG. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

as well as a significant fraction of high-affinity [¹²⁵I]epibatidine binding. Significant [¹²⁵I]epibatidine binding persists in the medial habenula, fasciculus retroflexus, interpeduncular nucleus and inferior colliculus of β2 knock-out mice.

Inclusion of 100 nM cytosine in the binding assays selectively inhibits primarily α4β2* nAChR [14,15,46] and facilitates identifi-

cation of less widely expressed nAChR subtypes. With the exception of the binding in the medial habenula, fasciculus retroflexus, and the interpeduncular nucleus and a subset of the signal in the optic tracts and inferior colliculus, deletion of the β2 subunit eliminated the cytosine-resistant [¹²⁵I]epibatidine binding sites.

Table 5

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –3.5 mm Bregma.

	IPN	MG	SuG	DpG	SNpr	SNpc	VTA	V1	HP	DS
0.5 nM [¹²⁵ I]-Epibatidine										
WT	■ ■ ■ ■	■ ■ ■	■ ■ ■ ■	■	■	■ ■ ■	■ ■ ■	■ ■ ■	■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++	++++	++++
α4	++++	+	++	—	—	+	+	—	—	—
α6	++++	++	++	++++	++++	++	++	++++	++++	++++
α7	++++	++++	++++	++++	++++	+++	+++	++++	++++	++++
β2	++++	—	—	—	—	—	—	—	—	—
β4	++	+++	++++	++++	++++	+++	+++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++	++	++++	++++	++	++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine										
WT	■ ■ ■ ■	■	■ ■ ■	□	□	□	□	□	□	□
α2	++++	++++	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α4	++++	++++	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α6	++++	++++	++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α7	++++	++++	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β2	++++	—	—	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β4	+	—	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α5	++++	++++	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β3	++++	++++	++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine + 100 nM α-CtxMII										
WT	■ ■ ■ ■	■	■ ■ ■	□	□	□	□	□	□	□
α2	++++	++++	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α4	++++	++	++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α6	++++	++++	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α7	++++	—	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β2	++++	—	—	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β4	+	—	++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α5	++++	++++	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β3	++++	++++	++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
200 pM 5-[¹²⁵ I]-A85380										
WT	■ ■ ■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■	■ ■ ■	■ ■ ■	■ ■	■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++	++++	++++	++++
α4	++++	+	++	—	++	+	+	—	—	—
α6	++++	++++	++++	++++	++++	++	++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++	++++	++++	++++
β2	—	—	—	—	—	—	—	—	—	—
β4	+++	++++	++++	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++	++	++++	++++	++++
0.5 nM [¹²⁵ I] α-ConotoxinMII										
WT	■ ■	□	■ ■ ■ ■	□	□	□	□	□	□	□
α2	++++	n/a	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α4	++	n/a	+++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α6	++	n/a	—	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α7	+++	n/a	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β2	—	n/a	—	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β4	+++	n/a	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α5	+++	n/a	++++	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β3	+++	n/a	+	n/a	n/a	n/a	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I] α-Bungarotoxin										
WT	■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■	■ ■	□	□	■	■	■
α2	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
α4	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
α6	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
α7	—	—	—	—	—	n/a	n/a	—	—	—
β2	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
β4	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
α5	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++
β3	++++	++++	++++	++++	++++	n/a	n/a	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At bregma = –3.52 mm the following brain regions are visible: Interpeduncular Nucleus, IPN; Medial Geniculates, MG; Superior Colliculus Superficial Gray layer, SuG; Superior Colliculus deep gray layer, DpG; Substantia Nigra pars reticulata, SNpr; Substantia Nigra pars compacta, SNpc; Ventral Tegmental Area, VTA; Hippocampus, HP; Primary Visual Cortex, V1; Dorsal Subiculum, DS. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

3.2.3. β4 nAChR gene deletion

Deletion of the β4 nAChR gene had no detectable effect on [¹²⁵I]αBgt, [¹²⁵I]αCtxMII or [¹²⁵I]A-85380 binding. In general, deletion of β4 had little effect on total [¹²⁵I]epibatidine binding with the exception of partial reductions in the signal intensity in

the fasciculus retroflexus, interpeduncular nucleus and the inferior colliculus. The effect of β4 gene deletion was more readily observable when 100 nM cytosine was included in the incubation with [¹²⁵I]epibatidine. Under these experimental conditions, when binding to α4β2* nAChR sites is removed, robust effects

Table 6

Semi-quantitative visual analysis of coronal mouse brain sections at approximately –5.2 mm Bregma.

	CIC	DCIC	ECIC	PAG	DT	PN	CEnt
0.5 nM [¹²⁵ I]-Epibatidine							
WT	■ ■	■ ■ ■	■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++
α4	++++	+	+	++	+++	—	—
α6	++++	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++
β2	+	++++	—	—	+	—	—
β4	++	++	++	+++	++++	+++	+++
α5	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine							
WT	■	■ ■ ■	□	■	□	□	□
α2	++++	++++	n/a	++++	n/a	n/a	n/a
α4	++	++++	n/a	++++	n/a	n/a	n/a
α6	—	—	n/a	++++	n/a	n/a	n/a
α7	++++	++++	n/a	++++	n/a	n/a	n/a
β2	++++	++++	n/a	++++	n/a	n/a	n/a
β4	—	—	n/a	—	n/a	n/a	n/a
α5	++++	++++	n/a	++++	n/a	n/a	n/a
β3	++++	++++	n/a	++++	n/a	n/a	n/a
0.5 nM [¹²⁵ I]-Epibatidine + 100 nM Cytisine + 100 nM α-CtxMII							
WT	■	■ ■ ■	□	■	□	□	□
α2	++++	++++	n/a	++++	n/a	n/a	n/a
α4	++	++++	n/a	++++	n/a	n/a	n/a
α6	—	—	n/a	++++	n/a	n/a	n/a
α7	++++	++++	n/a	++++	n/a	n/a	n/a
β2	++++	++++	n/a	++++	n/a	n/a	n/a
β4	—	—	n/a	—	n/a	n/a	n/a
α5	++++	++++	n/a	++++	n/a	n/a	n/a
β3	++++	++++	n/a	++++	n/a	n/a	n/a
200 pM 5-[¹²⁵ I]-A85380							
WT	■ ■	■ ■ ■	■ ■	■ ■ ■	■ ■ ■ ■	■ ■ ■	■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++
α4	++++	+	+	++	+++	—	—
α6	++++	++++	++++	++++	++++	++++	++++
α7	++++	++++	++++	++++	++++	++++	++++
β2	+	++++	—	—	+	—	—
β4	++	++	++	+++	++++	+++	+++
α5	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++
0.5 nM [¹²⁵ I] α-CtxMII							
WT	□	□	□	□	□	□	□
α2	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α4	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α6	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α7	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β2	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β4	n/a	n/a	n/a	n/a	n/a	n/a	n/a
α5	n/a	n/a	n/a	n/a	n/a	n/a	n/a
β3	n/a	n/a	n/a	n/a	n/a	n/a	n/a
250 pM [¹²⁵ I] α-Bungarotoxin							
WT	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■	■ ■ ■ ■
α2	++++	++++	++++	++++	++++	++++	++++
α4	++++	++++	++++	++++	++++	++++	++++
α6	++++	++++	++++	++++	++++	++++	++++
α7	—	—	—	—	—	—	—
β2	++++	++++	++++	++++	++++	++++	++++
β4	++++	++++	++++	++++	++++	++++	++++
α5	++++	++++	++++	++++	++++	++++	++++
β3	++++	++++	++++	++++	++++	++++	++++

Autoradiography was performed as described in Section 2.3. At bregma = –5.2 mm the following brain regions are visible: CIC—Central nucleus of the Inferior Colliculus; DCIC—Dorsal cortex of the inferior Colliculus; ECIC—External Cortex of the inferior colliculus; DT—Dorsal Tegmental nucleus; PN—Pontine Nucleus; Cent—Caudomedial entorhinal cortex. Regions with no detectable signal above background are designated with □ and relative quantitation was not attempted (n/a).

of β4 gene deletion could be observed in medial habenula, fasciculus retroflexus, interpeduncular nucleus and inferior colliculus, regions that express high levels of cytosine-resistant [¹²⁵I]epibatidine binding. In addition, detectable reductions in cytosine-resistant [¹²⁵I]epibatidine binding were also noted in the dorsolateral, ventrolateral and medial geniculate nuclei, olivary

pretectal nucleus, superficial gray level of the superior colliculus and the inferior colliculus.

3.2.4. α2 nAChR gene deletion

Little effect of the deletion of the α2 nAChR subunit could be noted for any binding sites. However, modest, relatively subtle

reductions may have occurred for cytosine-resistant [125 I]epibatidine binding sites in several cortical layers, and some regions of the visual tract including the dorsolateral and ventrolateral geniculate nuclei, the olivary pretectal nucleus, the optic tract and the superficial gray area of the superior colliculus.

3.2.5. $\alpha 4$ nAChR gene deletion

Deletion of the $\alpha 4$ nAChR gene had no effect on [125 I] α Bgt binding. All other binding sites were at least partially reduced in $\alpha 4$ knock-out mice.

Significantly less [125 I]A-85380 binding was observed in the brain of $\alpha 4$ knock-out mice than wild-type mice. Indeed, deletion of $\alpha 4$ eliminated [125 I]A-85380 binding in many brain regions. However, in contrast to the complete elimination of [125 I]A-85380 binding following deletion of the $\beta 2$ subunit, low, but detectable binding, was noted in the caudate putamen, globus pallidus, optic tracts, dorsolateral, ventrolateral and medial geniculate nuclei, substantia nigra pars compacta and ventral tegmental area. In addition, higher levels of [125 I]A-85380 binding persisted in medial habenula, olivary pretectal nucleus, substantia nigra pars reticulata and superficial gray region of the superior colliculus in $\alpha 4$ knock-outs. Deletion of $\alpha 4$ had little effect on [125 I]A-85380 binding in the fasciculus retroflexus and the interpeduncular nucleus.

Deletion of the $\alpha 4$ nAChR subunit generally reduced, but did not eliminate, [125 I] α CtxMII binding sites throughout the brain.

Given the complexity of [125 I]epibatidine binding sites, the pattern of response to deletion of $\alpha 4$ on these sites was more complex than that for [125 I]A-85380 binding. However, those sites that remained in $\beta 2$ knock-out mice also remained in $\alpha 4$ knock-outs. Furthermore, the additional sites that differed between $\beta 2$ and $\alpha 4$ knock-outs for [125 I]A-85380 binding were also noted for total [125 I]epibatidine binding. As was the case for the $\beta 2$ knock-outs, significant signal remained in the medial habenula, fasciculus retroflexus, interpeduncular nucleus and inferior colliculus of $\alpha 4$ knock-out mice. In addition, as was the case for the [125 I]A-85380 and [125 I] α CtxMII binding, deletion of $\alpha 4$ reduced, but did not eliminate $\beta 2$ -dependent, cytosine-resistant [125 I]epibatidine binding sites particularly in the visual system.

3.2.6. $\alpha 6$ nAChR gene deletion

Deletion of the $\alpha 6$ nAChR gene had very specific effects on nAChR binding sites. Virtually all [125 I] α CtxMII binding was eliminated by $\alpha 6$ gene deletion with the exception of partial reductions in the labeling in the interpeduncular nucleus and the optic tracts. While effects of $\alpha 6$ gene deletion for both [125 I]A-85380 and total [125 I]epibatidine binding were not generally obvious, partial reductions were noted for the dorsolateral and ventrolateral geniculates, the superficial gray layer of the superior colliculus, substantia nigra pars compacta and ventral tegmental areas. Addition of 100 nM cytosine to the [125 I]epibatidine binding assay helped to illustrate the effect of $\alpha 6$ gene deletion in the regions listed above as well as in the caudate putamen, the optic tracts and olivary pretectal nucleus.

3.2.7. $\alpha 5$ nAChR gene deletion

Deletion of the $\alpha 5$ nAChR gene had little detectable effect on the expression of any of the ligand binding sites.

3.2.8. $\beta 3$ nAChR gene deletion

The patterns for the effect of deletion of the $\beta 3$ nAChR subunit were very similar to those following deletion of the $\alpha 6$ nAChR subunit: significant reductions in high-affinity [125 I] α CtxMII binding, selective reductions [125 I]A-85380 binding and selective reductions in [125 I]epibatidine binding that were particularly evident in samples including 100 nM cytosine. However, the

reduction in signal following deletion of $\beta 3$ was less complete than that following deletion of $\alpha 6$.

4. Discussion

4.1. Null mutants and nicotinic binding sites

The examination of the effects of nAChR gene mutation on various nicotinic binding sites has helped to define the nature of the complex nAChR subtypes measured with these ligands.

4.1.1. [125 I] α -Bungarotoxin binding sites

The binding of [125 I] α Bgt shows the simplest response to nAChR gene deletion. Deletion of the $\alpha 7$ gene completely eliminates [125 I] α Bgt binding. No noticeable effect of deletion of any of the other nAChR genes on [125 I] α Bgt binding was observed. This result is completely consistent with the original cloning of the $\alpha 7$ gene as a α Bgt binding protein [62] as well as a functional receptor inhibited by α Bgt [63,64]. The original report describing the $\alpha 7$ knock-out mouse confirmed that deletion of $\alpha 7$ eliminated [125 I] α Bgt binding as well as α Bgt-sensitive function [33]. Deletion of $\alpha 7$ had no effect on the other binding sites measured here, although a lower affinity, α Bgt-sensitive epibatidine binding site, which measures the same subtype as [125 I] α Bgt is eliminated by deletion of the $\alpha 7$ gene [39,44]. Thus, all [125 I] α Bgt binding sites in mouse brain are $\alpha 7^*$ nAChR.

4.1.2. [125 I]A-85380 binding sites

A-85380 was originally identified as a potent $\alpha 4\beta 2^*$ nAChR agonist [23]. It is now recognized as a more general $\beta 2^*$ nAChR agonist [24]. Compounds modified by incorporation of a substitution in the 5 position of the pyridine moiety are extremely useful ligands for positron emission tomography [25,26]. A-85380 also serves as the progenitor for the potent agonist, saztidine that rapidly desensitizes the $\alpha 4\beta 2$ nAChR [65]. Confirming earlier results [24], deletion of the $\beta 2$ nAChR subunit completely eliminated [125 I]A-85380 binding throughout the mouse brain. The results summarized in this paper also illustrate why A-85380 was originally considered to be an $\alpha 4\beta 2$ nAChR selective ligand: Deletion of the $\alpha 4$ nAChR subunit eliminates most [125 I]A-85380 binding sites. However, residual [125 I]A-85380 binding sites persist in a few distinct brain areas in the $\alpha 4$ knock-out mice. The brain regions remaining in $\alpha 4$ knock-outs include the dopaminergic pathways and visual tracts all of which show partial reductions in signal following deletion of either the $\alpha 6$ or $\beta 3$ genes. In addition, the medial habenula and interpeduncular nucleus express residual [125 I]A-85380 binding sites in $\alpha 4$ knock-out mice which are not obviously reduced by deletion of any of the genes investigated here. These are likely to correspond to $\alpha 3\beta 2^*$ nAChR that will be discussed in more detail below.

4.1.3. [125 I] α -ConotoxinMII binding sites

α CtxMII was originally isolated from the cone snail, *Conus magus*, and characterized as an inhibitor of $\alpha 3\beta 2$ nAChR expressed in *Xenopus* oocytes [20]. However, with the exception of the medial habenula, fasciculus retroflexus and interpeduncular nucleus the binding of [125 I] α CtxMII is unaffected by deletion of the $\alpha 3$ nAChR subunit [38]. The deletion of the $\beta 2$ nAChR subunit completely eliminates [125 I] α CtxMII binding [39,52,55] and that result has been confirmed in the current study. In what was initially a surprise, the binding of [125 I] α CtxMII in catecholaminergic and visual pathways was eliminated by deletion of the $\alpha 6$ nAChR subunit [22], an observation that has also been confirmed in the present study. In yet another result that was initially a surprise, deletion of the $\beta 3$ nAChR subunit also substantially reduced [125 I] α CtxMII binding [35] in catecholaminergic and visual path-

ways. This result has been confirmed with immunochemical experiments [43] and by the results of the current study. In contrast to the effect of $\alpha 6$ nAChR gene deletion, deletion of the $\beta 3$ nAChR subunit did not completely eliminate [125 I] α CtxMII binding or α CtxMII-sensitive function [43,53], indicating either that a small population of $\alpha 6\beta 2^*$ nAChR normally exist or that these receptors are assembled in the absence of the $\beta 3$ subunit. The complexity of [125 I] α CtxMII binding sites is further revealed by the demonstration that the $\alpha 4$ nAChR subunit is included in an $\alpha 4\alpha 6\beta 2\beta 3$ nAChR [52,53,66,67] that is also illustrated in the current paper by the partial reduction of [125 I] α CtxMII binding in $\alpha 4$ knock-out mice. Thus, [125 I] α CtxMII binding sites represent a complex mixture of nAChR subtypes. A relatively small proportion of the [125 I] α CtxMII binding sites, confined primarily to the medial habenula/interpeduncular nucleus pathway, are $\alpha 3\beta 2^*$ nAChR. Most of the [125 I] α CtxMII binding sites are $\alpha 6\beta 2^*$ nAChR including $\alpha 4\beta 6\beta 2\beta 3$ nAChR, $\alpha 6\beta 2\beta 3$ nAChR and, perhaps a small native population of $\alpha 6\beta 2$ nAChR, the relative proportions of which vary among dopaminergic terminal regions [67].

4.1.4. [125 I]Epibatidine binding sites

By far the most complex collection of nAChR binding sites are those that bind epibatidine. Epibatidine binds with very high affinity to heteromeric nAChR ($\alpha 2$, $\alpha 3$ and $\alpha 4$ assembled with $\beta 2$ or $\beta 4$ subunits) expressed in *Xenopus* oocytes [19] and HEK cells [18]. Epibatidine has also been used to label receptors analyzed by immunoprecipitation [40,41,43,47,48,67–76]. It is an excellent ligand to identify a diverse number of receptor subtypes using this approach. Pharmacological differences among diverse epibatidine binding sites have been used to identify subsets of these sites [14–16]. The combination of the pharmacological approaches with the use of null mutant mice has been used to investigate the diversity of natively expressed receptors analogous to the experiments described here [16,39,42,44,46,77,78].

Most [125 I]epibatidine binding sites respond to nAChR gene deletion in a manner very similar to, if not quite identical to, that observed for [125 I]A-85380: Deletion of either the $\beta 2$ or the $\alpha 4$ nAChR subunits eliminates most high-affinity [125 I]epibatidine binding. In as much as the cytosine-sensitive [125 I]epibatidine binding sites are virtually identical to the sites measured by either [3 H]nicotine (or [3 H]cytosine), the loss of cytosine-sensitive [125 I]epibatidine binding by deletion of either the $\alpha 4$ or $\beta 2$ subunits corresponds to the elimination of [3 H]nicotine binding following deletion of these subunits [30,32].

Inhibition of high-affinity [125 I]epibatidine binding sites by cytosine is a useful experimental approach to separate subsets of cytosine-sensitive and cytosine-resistant [125 I]epibatidine binding sites [14,15,39,45,46,78–80]. Most epibatidine binding sites in brain are sensitive to inhibition by cytosine and represent primarily, if not exclusively, $\alpha 4\beta 2^*$ nAChR sites [14,15,46]. Inhibition of the predominant $\alpha 4\beta 2^*$ nAChR allows detection of effects that may be masked by the large number of cytosine-sensitive sites. The cytosine-resistant epibatidine binding sites represent a diverse population and that is illustrated by the different effects occurring following deletion of the various nAChR genes. Deletion of the $\alpha 4$ reduced the cytosine-resistant [125 I]epibatidine binding in several brain regions, particularly those affected by $\alpha 6$ or $\beta 3$ deletion. Elimination of additional sites following deletion of the $\beta 2$ subunit demonstrates the existence of $\beta 2^*$ nAChR that do not also include the $\alpha 4$ subunit. $\beta 2^*$ nAChR that do not include $\alpha 4$ subunits are particularly noticeable in dopaminergic terminal regions and visual tracts. These [125 I]epibatidine binding sites are substantially reduced by deletion of either the $\alpha 6$ or $\beta 3$ gene implying the existence of $\alpha 6\beta 2\beta 3^*$ nAChR in this subset of sites. This assignment parallels that for [125 I] α CtxMII binding sites and is consistent with results reported previously for

$\alpha 6$ null mutant mice [22]. An effect of $\beta 4$ gene deletion is also clearly visible for the cytosine-resistant [125 I]epibatidine binding sites: A dramatic reduction in signal was also noted in the medial habenula, fasciculus retroflexus, interpeduncular nucleus and inferior colliculus.

Further refinement of the cytosine-resistant [125 I]epibatidine binding sites was accomplished by including 100 nM α CtxMII in the incubation. These cytosine-resistant, α CtxMII-sensitive sites detect nAChR that interact with α CtxMII including lower affinity sites that are not measured directly by the binding of [125 I] α CtxMII, which are predominantly $\alpha 6\beta 2\beta 3^*$ nAChR. Indeed, little further inhibition by 100 nM α CtxMII was noted for $\alpha 6$ and $\beta 3$ knock-out mice in most regions. However, a further reduction in [125 I]epibatidine binding was observed in the superior colliculus of $\alpha 6$ and $\beta 3$ knock-out mice with the addition of 100 nM α CtxMII. The existence of additional α CtxMII sensitive sites, likely corresponding to $\alpha 3\beta 2^*$ nAChR [81], is indicated (see also below).

4.2. Potential role of subunits the deletion of which has little effect on ligand binding

While deletion of most of the nAChR subunits had detectable effects on one or more of the ligand binding sites measured in the current study, deletion of either the $\alpha 2$ or $\alpha 5$ subunit did not have obvious effects.

4.2.1. $\alpha 3$ subunit

Deletion of the $\alpha 3$ nAChR results in early postnatal death such that null mutants generally do not survive for more than a few weeks [49]. Consequently, $\alpha 3$ knock-out mice were not included in the experiments described here. However, it is evident from the sites that were eliminated by deletion of $\alpha 2$, $\alpha 4$ or $\alpha 6$ nAChR subunits that, by process of elimination, $\alpha 3^*$ nAChR account for these residual sites. Many of these residual regions including medial habenula, fasciculus retroflexus, interpeduncular nucleus and inferior colliculus are found in the cytosine-resistant component of high-affinity [125 I]epibatidine binding sites, regions that are reduced in $\beta 4$ knock-out mice. This strongly indicates the presence of $\alpha 3\beta 4^*$ nAChR in these regions and is consistent with the observations made on the effect of $\alpha 3$ gene deletion in 8-day-old $\alpha 3$ knock-out mice [38]. The existence of potential $\alpha 3\beta 2^*$ nAChR is suggested by the observation that 100 nM α CtxMII decreases [125 I]epibatidine binding in the dorso- and ventrolateral geniculate nuclei and in the superficial gray area of the superior colliculus which is evident particularly in sections from $\alpha 6$ knock-out mice. This observation is reasonable given that α CtxMII is known to interact with $\alpha 3\beta 2$ nAChR, albeit with lower affinity than with $\alpha 6\beta 2^*$ nAChR [20,82] and is consistent with the demonstration that $\alpha 3^*$ nAChR are detectable in visual pathways [81,83,84].

4.2.2. $\alpha 2$ subunit

In situ hybridization studies in either rat [85] or mouse [86] brain indicate that the $\alpha 2$ subunit is sparsely expressed. The experiments describing the binding of [3 H]epibatidine to both $\alpha 2\beta 2$ nAChR and $\alpha 2\beta 4$ nAChR expressed in *Xenopus* oocytes [19] suggest that such sites should be detectable in brain. Indeed, by the use of ligand binding and immunoprecipitation, [125 I]epibatidine binding sites including the $\alpha 2$ subunit have been identified in mouse brain [47]; however, even in regions of highest expression (olfactory bulbs and interpeduncular nucleus) the $\alpha 2^*$ nAChR are sparsely expressed. There is also evidence that some $\alpha 2^*$ nAChR are present in visual pathways [81,83,84]. Given the semi-quantitative nature of these studies relatively small effects of $\alpha 2$ gene deletion on any of the binding sites may have escaped detection. Development of an $\alpha 2^*$ nAChR selective ligand would facilitate the identification of these receptors.

4.2.3. $\alpha 5$ subunit

The $\alpha 5$ subunit is an accessory subunit the deletion of which has significant effects on nicotine-mediated responses in vivo [36,87,88] as well as on nAChR-mediated function [48,51,89]. Furthermore, immunoprecipitation studies have established that a subset of nAChR measured by epibatidine binding include the $\alpha 5$ subunit [31,40,48,72,81,90,91]. Nevertheless, no study, including the current one, has been able to detect a change in the total number of epibatidine binding sites [36,48], despite the significant changes in function, pharmacology and physiology resulting from $\alpha 5$ gene deletion [48,51,89]. Among the possible reasons for the failure to detect changes in binding sites include their relative rarity [48,72,90] or the possibility that an alternate subunit such as $\alpha 4$ or $\beta 2$ is now incorporated in nAChR which normally include the $\alpha 5$ subunit thereby preserving measurable binding sites.

4.3. Summary

The results presented here provide a survey of the effects of nAChR gene deletion of nicotinic binding sites measured with [125 I] α Bgt, [125 I] α CtxMII, [125 I]A-85380 and [125 I]epibatidine under several conditions. Since the experiments were done in one laboratory under standard conditions the effects are easily comparable. The data presented in this paper are completely consistent with the findings of similar experiments with null mutant mice. It should be noted that all comparisons made in the current study are semi-quantitative and therefore provide a general picture of receptor expression and composition. More detailed comparisons will require a quantitative autoradiographic approach with a larger number of replicates than were included here.

Autoradiographic studies have their limitations:

- The effects of deletion of genes with relatively low expression (such as the $\alpha 2$ nAChR subunit) are difficult to detect autoradiographically and often require alternative approaches such as immunoprecipitation [47].
- Deletion of an auxiliary subunit (such as $\alpha 5$) may have no detectable effect on traditional ligand binding. Consequently its role can escape detection. The lack of a detectable effect on binding could result because the $\alpha 5^*$ nAChR represent a relatively small fraction of the total heteromeric receptors. However, immunoprecipitation and functional assays have demonstrated an important role for this subunit [48,89,90], including behavioral consequences [36,92,93]. The $\alpha 5$ nAChR has been repeatedly shown to have an important role in human tobacco use [92]. Functional assays are essential to define fully the role of nAChR subunits.
- Autoradiographic analyses are inadequate to examine effects of subunit deletion, and therefore nAChR composition, when alternate subunits can substitute for the deleted gene. This caveat applies for auxiliary subunits such as $\alpha 5$ as discussed above. It could also apply for alternately assembled nAChR such as an $\alpha 7^*$ nAChR that includes a $\beta 2$ subunit [94], which in the absence of $\beta 2$ could assemble as an $\alpha 7$ nAChR homopentamer.
- Autoradiographic analyses also require sufficient signal:noise for detection of binding sites. This requirement can be problematic when ligands with relatively high non-specific binding especially peptide probe such as [125 I] α Bgt and [125 I] α CtxMII. Thus, in ventral tegmental area little [125 I] α Bgt binding was observed in the current study, consistent with similar observations for mouse [95] or rat [96]. However, functional $\alpha 7$ -nAChR have been identified in the ventral tegmental area [94,97] indicating that the resolution and/or sensitivity achieved with autoradiography is inadequate to identify these binding sites. Indeed, an electron

micrographic analysis identified $\alpha 7$ nAChR binding sites in the rat ventral tegmental area [98].

Therefore, failure to detect changes in binding site density following gene deletion, particularly for sparsely expressed or auxiliary subunits, does not necessarily establish that nAChR containing this subunits are absent. Additional studies using functional, pharmacological and immunochemical assays, in addition to more quantitative autoradiographic experiments, particularly in concert with null mutant mice, will provide more definitive description of the native nAChR populations to expand on the general picture provided here.

Conflict of interest

None.

Acknowledgements

The authors thank Dr. Sharon R. Grady and Dr. Heidi C. O'Neill for critical reading of the manuscript and helpful comments. The authors also thank the original providers of the null mutant mice: Dr. Jim Boulter, UCLA— $\alpha 2$; Dr. Richard Paylor, Baylor University School of Medicine— $\alpha 5$, $\alpha 7$ and $\beta 4$; Dr. John Drago, University of Melbourne— $\alpha 4$; Dr. Marina Picciotto, Yale University— $\beta 2$; Dr. Stephan Heinemann, Salk Institute— $\beta 3$ and Dr. Uwe Maskos, Pasteur Institute— $\alpha 6$. This work was supported by the following grants from the National Institutes on Drug Abuse of the National Institutes of Health of the United States of America: R01 DA003194, R01 DA012242 and P30 DA015663.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bcp.2011.04.019.

References

- [1] Patrick J, Stallcup B. Alpha-bungarotoxin binding and cholinergic receptor function on a rat sympathetic nerve line. *J Biol Chem* 1977;252:8629–33.
- [2] Romano C, Goldstein A. Stereospecific nicotine receptors on rat brain membranes. *Science* 1980;210:647–50.
- [3] Clarke PB, Schwartz RD, Paul SM, Pert CB, Pert A. Nicotinic binding in rat brain: autoradiographic comparison of [3 H]acetylcholine, [3 H]nicotine, and [125 I]-alpha-bungarotoxin. *J Neurosci* 1985;5:1307–15.
- [4] Marks MJ, Collins AC. Characterization of nicotine binding in mouse brain and comparison with the binding of alpha-bungarotoxin and quinuclidinyl benzilate. *Mol Pharmacol* 1982;22:554–64.
- [5] Sargent PB. The diversity of neuronal nicotinic acetylcholine receptors. *Annu Rev Neurosci* 1993;16:403–43.
- [6] Lindstrom J, Schoepfer R, Conroy W, Whiting P, Das M, Saedi M, et al. The nicotinic acetylcholine receptor gene family: structure of nicotinic receptors from muscle and neurons and neuronal alpha-bungarotoxin-binding proteins. *Adv Exp Med Biol* 1991;287:255–78.
- [7] Gotti C, Clementi F, Fornari A, Gaimarri A, Guiducci S, Manfredi I, et al. Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem Pharmacol* 2009;78:703–11.
- [8] Schwartz RD, McGee Jr R, Kellar KJ. Nicotinic cholinergic receptors labeled by [3 H]acetylcholine in rat brain. *Mol Pharmacol* 1982;22:56–62.
- [9] Pabreza LA, Dhawan S, Kellar KJ. [3 H]cytisine binding to nicotinic cholinergic receptors in brain. *Mol Pharmacol* 1991;39:9–12.
- [10] Aboud LG, Grassi S. [3 H]methylcarbamylcholine, a new radioligand for studying brain nicotinic receptors. *Biochem Pharmacol* 1986;35:4199–202.
- [11] Badio B, Daly JW. Epibatidine. A potent analgesic and nicotinic agonist. *Mol Pharmacol* 1994;45:563–9.
- [12] Davila-Garcia MI, Musachio JL, Perry DC, Xiao Y, Horti A, London ED, et al. [125 I]IPH, an epibatidine analog, binds with high affinity to neuronal nicotinic cholinergic receptors. *J Pharmacol Exp Ther* 1997;282:445–51.
- [13] Houghtling RA, Davila-Garcia MI, Kellar KJ. Characterization of (+/–)(–)[3 H]epibatidine binding to nicotinic cholinergic receptors in rat and human brain. *Mol Pharmacol* 1995;48:280–7.
- [14] Perry DC, Xiao Y, Nguyen HN, Musachio JL, Davila-Garcia MI, Kellar KJ. Measuring nicotinic receptors with characteristics of alpha4beta2, alpha3beta2

- and $\alpha 3\beta 4$ subtypes in rat tissues by autoradiography. *J Neurochem* 2002;82:468–81.
- [15] Marks MJ, Smith KW, Collins AC. Differential agonist inhibition identifies multiple epibatidine binding sites in mouse brain. *J Pharmacol Exp Ther* 1998;285:377–86.
 - [16] Zoli M, Lena C, Picciotto MR, Changeux JP. Identification of four classes of brain nicotinic receptors using beta2 mutant mice. *J Neurosci* 1998;18:4461–72.
 - [17] Gerzanich V, Peng X, Wang F, Wells G, Anand R, Fletcher S, et al. Comparative pharmacology of epibatidine: a potent agonist for neuronal nicotinic acetylcholine receptors. *Mol Pharmacol* 1995;48:774–82.
 - [18] Xiao Y, Baydyuk M, Wang HP, Davis HE, Kellar KJ. Pharmacology of the agonist binding sites of rat neuronal nicotinic receptor subtypes expressed in HEK 293 cells. *Bioorg Med Chem Lett* 2004;14:1845–8.
 - [19] Parker MJ, Beck A, Luetje CW. Neuronal nicotinic receptor beta2 and beta4 subunits confer large differences in agonist binding affinity. *Mol Pharmacol* 1998;54:1132–9.
 - [20] Cartier GE, Yoshikami D, Gray WR, Luo S, Olivera BM, McIntosh JM. A new alpha-conotoxin which targets $\alpha 3\beta 2$ nicotinic acetylcholine receptors. *J Biol Chem* 1996;271:7522–8.
 - [21] Whiteaker P, McIntosh JM, Luo S, Collins AC, Marks MJ. 125I-alpha-conotoxin MII identifies a novel nicotinic acetylcholine receptor population in mouse brain. *Mol Pharmacol* 2000;57:913–25.
 - [22] Champiaux N, Han ZY, Bessis A, Rossi FM, Zoli M, Marubio L, et al. Distribution and pharmacology of alpha 6-containing nicotinic acetylcholine receptors analyzed with mutant mice. *J Neurosci* 2002;22:1208–17.
 - [23] Sullivan JP, Donnelly-Roberts D, Briggs CA, Anderson DJ, Gopalakrishnan M, Piattoni-Kaplan M, et al. A-85380 [3-(2(S)-azetidinylmethoxy) pyridine]: in vitro pharmacological properties of a novel, high affinity alpha 4 beta 2 nicotinic acetylcholine receptor ligand. *Neuropharmacology* 1996;35:725–34.
 - [24] Mukhin AG, Gundisch D, Horti AG, Koren AO, Tamagnan G, Kimes AS, et al. 5-lodo-A-85380, an $\alpha 4\beta 2$ subtype-selective ligand for nicotinic acetylcholine receptors. *Mol Pharmacol* 2000;57:642–9.
 - [25] Rueter LE, Donnelly-Roberts DL, Curzon P, Briggs CA, Anderson DJ, Bitner RS. A-85380: a pharmacological probe for the preclinical and clinical investigation of the $\alpha 4\beta 2$ neuronal nicotinic acetylcholine receptor. *CNS Drug Rev* 2006;12:100–12.
 - [26] Sihver W, Nordberg A, Langstrom B, Mukhin AG, Koren AO, Kimes AS, et al. Development of ligands for in vivo imaging of cerebral nicotinic receptors. *Behav Brain Res* 2000;113:143–57.
 - [27] Picciotto MR. Nicotine as a modulator of behavior: beyond the inverted U. *Trends Pharmacol Sci* 2003;24:493–9.
 - [28] Cordero-Erausquin M, Marubio LM, Klink R, Changeux JP. Nicotinic receptor function: new perspectives from knockout mice. *Trends Pharmacol Sci* 2000;21:211–7.
 - [29] Champiaux N, Changeux JP. Knockout and knockin mice to investigate the role of nicotinic receptors in the central nervous system. *Prog Brain Res* 2004;145:235–51.
 - [30] Marubio LM, del Mar Arroyo-Jimenez M, Cordero-Erausquin M, Lena C, Le Novère N, de Kerchove d'Exaerde A, et al. Reduced antinociception in mice lacking neuronal nicotinic receptor subunits. *Nature* 1999;398:805–10.
 - [31] Millar NS, Gotti C. Diversity of vertebrate nicotinic acetylcholine receptors. *Neuropharmacology* 2009;56:237–46.
 - [32] Picciotto MR, Zoli M, Lena C, Bessis A, Lallemand Y, Le Novère N, et al. Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature* 1995;374:65–7.
 - [33] Orr-Urtreger A, Goldner FM, Saeki M, Lorenzo I, Goldberg L, De Biasi M, et al. Mice deficient in the $\alpha 7$ neuronal nicotinic acetylcholine receptor lack α -bungarotoxin binding sites and hippocampal fast nicotinic currents. *J Neurosci* 1997;17:9165–71.
 - [34] Champiaux N, Changeux JP. Knock-out and knock-in mice to investigate the role of nicotinic receptors in the central nervous system. *Curr Drug Targets CNS Neurol Disord* 2002;1:319–30.
 - [35] Cui C, Booker TK, Allen RS, Grady SR, Whiteaker P, Marks MJ, et al. The beta3 nicotinic receptor subunit: a component of alpha-conotoxin MII-binding nicotinic acetylcholine receptors that modulate dopamine release and related behaviors. *J Neurosci* 2003;23:11045–53.
 - [36] Salas R, Orr-Urtreger A, Broide RS, Beaudet A, Paylor R, De Biasi M. The nicotinic acetylcholine receptor subunit alpha 5 mediates short-term effects of nicotine in vivo. *Mol Pharmacol* 2003;63:1059–66.
 - [37] Salas R, Pieri F, De Biasi M. Decreased signs of nicotine withdrawal in mice null for the beta4 nicotinic acetylcholine receptor subunit. *J Neurosci* 2004;24:10035–9.
 - [38] Whiteaker P, Peterson CG, Xu W, McIntosh JM, Paylor R, Beaudet AL, et al. Involvement of the $\alpha 3$ subunit in central nicotinic binding populations. *J Neurosci* 2002;22:2522–9.
 - [39] Marks MJ, Laverty DS, Whiteaker P, Salminen O, Grady SR, McIntosh JM, et al. John Daly's compound, epibatidine, facilitates identification of nicotinic receptor subtypes. *J Mol Neurosci* 2010;40:96–104.
 - [40] Grady SR, Moretti M, Zoli M, Marks MJ, Zanardi A, Pucci L, et al. Rodent habenulo-interpeduncular pathway expresses a large variety of uncommon nAChR subtypes, but only the $\alpha 3\beta 4$ and $\alpha 3\beta 2\alpha 4$ subtypes mediate acetylcholine release. *J Neurosci* 2009;29:2272–82.
 - [41] Gotti C, Moretti M, Meinerz NM, Clementi F, Gaimarri A, Collins AC, et al. Partial deletion of the nicotinic cholinergic receptor alpha 4 or beta 2 subunit genes changes the acetylcholine sensitivity of receptor-mediated 86Rb⁺ efflux in cortex and thalamus and alters relative expression of alpha 4 and beta 2 subunits. *Mol Pharmacol* 2008;73:1796–807.
 - [42] Marks MJ, Whiteaker P, Grady SR, Picciotto MR, McIntosh JM, Collins AC. Characterization of [(125)I]epibatidine binding and nicotinic agonist-mediated (86) Rb⁺ efflux in interpeduncular nucleus and inferior colliculus of beta2 null mutant mice. *J Neurochem* 2002;81:1102–15.
 - [43] Gotti C, Moretti M, Clementi F, Riganti L, McIntosh JM, Collins AC, et al. Expression of nigrostriatal alpha 6-containing nicotinic acetylcholine receptors is selectively reduced, but not eliminated, by beta 3 subunit gene deletion. *Mol Pharmacol* 2005;67:2007–15.
 - [44] Marks MJ, Whiteaker P, Collins AC. Deletion of the $\alpha 7$, $\beta 2$, or $\beta 4$ nicotinic receptor subunit genes identifies highly expressed subtypes with relatively low affinity for [3H]epibatidine. *Mol Pharmacol* 2006;70:947–59.
 - [45] Marks MJ, Stitzel JA, Grady SR, Picciotto MR, Changeux JP, Collins AC. Nicotinic-agonist stimulated (86)Rb⁺ efflux and [(3)H]epibatidine binding of mice differing in beta2 genotype. *Neuropharmacology* 2000;39:2632–45.
 - [46] Marks MJ, Meinerz NM, Drago J, Collins AC. Gene targeting demonstrates that $\alpha 4$ nicotinic acetylcholine receptor subunits contribute to expression of diverse [3H]epibatidine binding sites and components of biphasic 86Rb⁺ efflux with high and low sensitivity to stimulation by acetylcholine. *Neuropharmacology* 2007;53:390–405.
 - [47] Whiteaker P, Wilking JA, Brown RW, Brennan RJ, Collins AC, Lindstrom JM, et al. Pharmacological and immunohistochemical characterization of $\alpha 2^*$ nicotinic acetylcholine receptors (nAChRs) in mouse brain. *Acta Pharmacol Sin* 2009;30:795–804.
 - [48] Brown RW, Collins AC, Lindstrom JM, Whiteaker P. Nicotinic $\alpha 5$ subunit deletion locally reduces high-affinity agonist activation without altering nicotinic receptor numbers. *J Neurochem* 2007;103:204–15.
 - [49] Xu W, Gelber S, Orr-Urtreger A, Armstrong D, Lewis RA, Ou CN, et al. Megacystis, mydriasis, and ion channel defect in mice lacking the $\alpha 3$ neuronal nicotinic acetylcholine receptor. *Proc Natl Acad Sci U S A* 1999;96:5746–51.
 - [50] Ross SA, Wong JY, Clifford JJ, Kinsella A, Massalas JS, Horne MK, et al. Phenotypic characterization of an $\alpha 4$ neuronal nicotinic acetylcholine receptor subunit knock-out mouse. *J Neurosci* 2000;20:6431–41.
 - [51] Salminen O, Murphy KL, McIntosh JM, Drago J, Marks MJ, Collins AC, et al. Subunit composition and pharmacology of two classes of striatal presynaptic nicotinic acetylcholine receptors mediating dopamine release in mice. *Mol Pharmacol* 2004;65:1526–35.
 - [52] Salminen O, Whiteaker P, Grady SR, Collins AC, McIntosh JM, Marks MJ. The subunit composition and pharmacology of alpha-Conotoxin MII-binding nicotinic acetylcholine receptors studied by a novel membrane-binding assay. *Neuropharmacology* 2005;48:696–705.
 - [53] Salminen O, Drapeau JA, McIntosh JM, Collins AC, Marks MJ, Grady SR. Pharmacology of alpha-conotoxin MII-sensitive subtypes of nicotinic acetylcholine receptors isolated by breeding of null mutant mice. *Mol Pharmacol* 2007;71:1563–71.
 - [54] Xu W, Orr-Urtreger A, Nigro F, Gelber S, Sutcliffe CB, Armstrong D, et al. Multiorgan autonomic dysfunction in mice lacking the beta2 and the beta4 subunits of neuronal nicotinic acetylcholine receptors. *J Neurosci* 1999;19:9298–305.
 - [55] Grady SR, Salminen O, Laverty DC, Whiteaker P, McIntosh JM, Collins AC, et al. The subtypes of nicotinic acetylcholine receptors on dopaminergic terminals of mouse striatum. *Biochem Pharmacol* 2007;74:1235–46.
 - [56] Grady SR, Salminen O, McIntosh JM, Marks MJ, Collins AC. Mouse striatal dopamine nerve terminals express $\alpha 4\alpha 5\beta 2$ and two stoichiometric forms of $\alpha 4\beta 2^*$ -nicotinic acetylcholine receptors. *J Mol Neurosci* 2010;40:91–5.
 - [57] Nakauchi S, Brennan RJ, Boulter J, Sumikawa K. Nicotine gates long-term potentiation in the hippocampal CA1 region via the activation of $\alpha 2^*$ nicotinic ACh receptors. *Eur J Neurosci* 2007;25:2666–81.
 - [58] Whiteaker P, Jimenez M, McIntosh JM, Collins AC, Marks MJ. Identification of a novel nicotinic binding site in mouse brain using [(125)I]-epibatidine. *Br J Pharmacol* 2000;131:729–39.
 - [59] Mineur YS, Brunzell DH, Grady SR, Lindstrom JM, McIntosh JM, Marks MJ, et al. Localized low-level re-expression of high-affinity mesolimbic nicotinic acetylcholine receptors restores nicotine-induced locomotion but not place conditioning. *Genes Brain Behav* 2009;8:257–66.
 - [60] Pauly JR, Marks MJ, Gross SD, Collins AC. An autoradiographic analysis of cholinergic receptors in mouse brain after chronic nicotine treatment. *J Pharmacol Exp Ther* 1991;258:1127–36.
 - [61] Millar NS. Assembly and subunit diversity of nicotinic acetylcholine receptors. *Biochem Soc Trans* 2003;31:869–74.
 - [62] Schoepfer R, Conroy WG, Whiting P, Gore M, Lindstrom J. Brain alpha-bungarotoxin binding protein cDNAs and MABs reveal subtypes of this branch of the ligand-gated ion channel gene superfamily. *Neuron* 1990;5:35–48.
 - [63] Couturier S, Bertrand D, Matter JM, Hernandez MC, Bertrand S, Millar N, et al. A neuronal nicotinic acetylcholine receptor subunit ($\alpha 7$) is developmentally regulated and forms a homo-oligomeric channel blocked by alpha-BTX. *Neuron* 1990;5:847–56.
 - [64] Seguela P, Wadiche J, Dineley-Miller K, Dani JA, Patrick JW. Molecular cloning, functional properties, and distribution of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J Neurosci* 1993;13:596–604.
 - [65] Xiao Y, Fan H, Musachio JL, Wei ZL, Chellappan SK, Kozikowski AP, et al. Sazetidine-A, a novel ligand that desensitizes $\alpha 4\beta 2$ nicotinic acetylcholine receptors without activating them. *Mol Pharmacol* 2006;70:1454–60.
 - [66] Champiaux N, Gotti C, Cordero-Erausquin M, David DJ, Przybylski C, Lena C, et al. Subunit composition of functional nicotinic receptors in dopaminergic neurons investigated with knock-out mice. *J Neurosci* 2003;23:7820–9.

- [67] Gotti C, Guiducci S, Tedesco V, Corbioli S, Zanetti L, Moretti M, et al. Nicotinic acetylcholine receptors in the mesolimbic pathway: primary role of ventral tegmental area $\alpha 6\beta 2^*$ receptors in mediating systemic nicotine effects on dopamine release, locomotion, and reinforcement. *J Neurosci* 2010;30:5311–25.
- [68] Yeh JJ, Yasuda RP, Davila-Garcia MI, Xiao Y, Ebert S, Gupta T, et al. Neuronal nicotinic acetylcholine receptor $\alpha 3$ subunit protein in rat brain and sympathetic ganglion measured using a subunit-specific antibody: regional and ontogenic expression. *J Neurochem* 2001;77:336–46.
- [69] Hernandez SC, Vicini S, Xiao Y, Davila-Garcia MI, Yasuda RP, Wolfe BB, et al. The nicotinic receptor in the rat pineal gland is an $\alpha 3\beta 4$ subtype. *Mol Pharmacol* 2004;66:978–87.
- [70] Turner JR, Kellar KJ. Nicotinic cholinergic receptors in the rat cerebellum: multiple heteromeric subtypes. *J Neurosci* 2005;25:9258–65.
- [71] Marritt AM, Cox BC, Yasuda RP, McIntosh JM, Xiao Y, Wolfe BB, et al. Nicotinic cholinergic receptors in the rat retina: simple and mixed heteromeric subtypes. *Mol Pharmacol* 2005;68:1656–68.
- [72] Mao D, Perry DC, Yasuda RP, Wolfe BB, Kellar KJ. The $\alpha 4\beta 2\alpha 5$ nicotinic cholinergic receptor in rat brain is resistant to up-regulation by nicotine in vivo. *J Neurochem* 2008;104:446–56.
- [73] Martin-Ruiz CM, Court JA, Molnar E, Lee M, Gotti C, Mamalaki A, et al. $\alpha 4$ but not $\alpha 3$ and $\alpha 7$ nicotinic acetylcholine receptor subunits are lost from the temporal cortex in Alzheimer's disease. *J Neurochem* 1999;73:1635–40.
- [74] Balestra B, Vailati S, Moretti M, Hanke W, Clementi F, Gotti C. Chick optic lobe contains a developmentally regulated $\alpha 2\alpha 5\beta 2$ nicotinic receptor subtype. *Mol Pharmacol* 2000;58:300–11.
- [75] Zoli M, Moretti M, Zanardi A, McIntosh JM, Clementi F, Gotti C. Identification of the nicotinic receptor subtypes expressed on dopaminergic terminals in the rat striatum. *J Neurosci* 2002;22:8785–9.
- [76] Moretti M, Mugnaini M, Tessari M, Zoli M, Gaimarri A, Manfredi I, et al. A comparative study of the effects of the intravenous self-administration or subcutaneous minipump infusion of nicotine on the expression of brain neuronal nicotinic receptor subtypes. *Mol Pharmacol* 2010;78:287–96.
- [77] Marks MJ, Whiteaker P, Calcaterra J, Stitzel JA, Bullock AE, Grady SR, et al. Two pharmacologically distinct components of nicotinic receptor-mediated rubidium efflux in mouse brain require the $\beta 2$ subunit. *J Pharmacol Exp Ther* 1999;289:1090–103.
- [78] Whiteaker P, Marks MJ, Grady SR, Lu Y, Picciotto MR, Changeux JP, et al. Pharmacological null mutation approaches reveal nicotinic receptor diversity. *Eur J Pharmacol* 2000;393:123–35.
- [79] Nguyen HN, Rasmussen BA, Perry DC. Subtype-selective up-regulation by chronic nicotine of high-affinity nicotinic receptors in rat brain demonstrated by receptor autoradiography. *J Pharmacol Exp Ther* 2003;307:1090–7.
- [80] Salas R, Pieri F, Fung B, Dani JA, De Biasi M. Altered anxiety-related responses in mutant mice lacking the $\beta 4$ subunit of the nicotinic receptor. *J Neurosci* 2003;23:6255–63.
- [81] Gotti C, Moretti M, Zanardi A, Gaimarri A, Champiaux N, Changeux JP, et al. Heterogeneity and selective targeting of neuronal nicotinic acetylcholine receptor (nAChR) subtypes expressed on retinal afferents of the superior colliculus and lateral geniculate nucleus: identification of a new native nAChR subtype $\alpha 3\beta 2(\alpha 5 \text{ or } \beta 3)$ enriched in retinocollicular afferents. *Mol Pharmacol* 2005;68:1162–71.
- [82] McIntosh JM, Olivera BM, Cruz LJ. Conus peptides as probes for ion channels. *Methods Enzymol* 1999;294:605–24.
- [83] Moretti M, Vailati S, Zoli M, Lippi G, Riganti L, Longhi R, et al. Nicotinic acetylcholine receptor subtypes expression during rat retina development and their regulation by visual experience. *Mol Pharmacol* 2004;66:85–96.
- [84] Cox BC, Marritt AM, Perry DC, Kellar KJ. Transport of multiple nicotinic acetylcholine receptors in the rat optic nerve: high densities of receptors containing $\alpha 6$ and $\beta 3$ subunits. *J Neurochem* 2008;105:1924–38.
- [85] Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, et al. Distribution of $\alpha 2$, $\alpha 3$, $\alpha 4$, and $\beta 2$ neuronal nicotinic receptor subunit mRNAs in the central nervous system: a hybridization histochemical study in the rat. *J Comp Neurol* 1989;284:314–35.
- [86] Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, et al. Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. *J Neurosci* 1992;12:2765–84.
- [87] Salas R, Sturm R, Boulter J, De Biasi M. Nicotinic receptors in the habenulo-interpeduncular system are necessary for nicotine withdrawal in mice. *J Neurosci* 2009;29:3014–8.
- [88] Jackson KJ, Marks MJ, Vann RE, Chen X, Gamage TF, Warner JA, et al. Role of $\alpha 5$ nicotinic acetylcholine receptors in pharmacological and behavioral effects of nicotine in mice. *J Pharmacol Exp Ther* 2010;334:137–46.
- [89] Bailey CD, De Biasi M, Fletcher PJ, Lambe EK. The nicotinic acetylcholine receptor $\alpha 5$ subunit plays a key role in attention circuitry and accuracy. *J Neurosci* 2010;30:9241–52.
- [90] Heath CJ, King SL, Gotti C, Marks MJ, Picciotto MR. Cortico-thalamic connectivity is vulnerable to nicotine exposure during early postnatal development through $\alpha 4\beta 2/\alpha 5$ nicotinic acetylcholine receptors. *Neuropsychopharmacology* 2010;35:2324–38.
- [91] Gotti C, Moretti M, Gaimarri A, Zanardi A, Clementi F, Zoli M. Heterogeneity and complexity of native brain nicotinic receptors. *Biochem Pharmacol* 2007;74:1102–11.
- [92] Gangitano D, Salas R, Teng Y, Perez E, De Biasi M. Progesterone modulation of $\alpha 5$ nAChR subunits influences anxiety-related behavior during estrus cycle. *Genes Brain Behav* 2009;8:398–406.
- [93] Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ. Habenular $\alpha 5$ nicotinic receptor subunit signalling controls nicotine intake. *Nature* 2011;471:597–601.
- [94] Liu Q, Huang Y, Xue F, Simard A, DeChon J, Li G, et al. A novel nicotinic acetylcholine receptor subtype in basal forebrain cholinergic neurons with high sensitivity to amyloid peptides. *J Neurosci* 2009;29:918–29.
- [95] Pauly JR, Stitzel JA, Marks MJ, Collins AC. An autoradiographic analysis of cholinergic receptors in mouse brain. *Brain Res Bull* 1989;22:453–9.
- [96] Clarke PB, Pert CB, Pert A. Autoradiographic distribution of nicotine receptors in rat brain. *Brain Res* 1984;323:390–5.
- [97] Mansvelder HD, McGehee DS. Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 2000;27:349–57.
- [98] Jones IW, Wonnacott S. Precise localization of $\alpha 7$ nicotinic acetylcholine receptors on glutamatergic axon terminals in the rat ventral tegmental area. *J Neurosci* 2004;24:11244–52.