



Review

Positive allosteric modulators as an approach to nicotinic acetylcholine receptor-targeted therapeutics: Advantages and limitations

Dustin K. Williams^a, Jingyi Wang^b, Roger L. Papke^{a,*}^a Department of Pharmacology and Therapeutics, University of Florida College of Medicine, Gainesville, FL, United States^b Department of Chemistry, University of Florida, Gainesville, FL, United States

ARTICLE INFO

Article history:

Received 21 March 2011

Accepted 2 May 2011

Available online 7 May 2011

Keywords:

Alzheimer's disease

Schizophrenia

Drug development

Electrophysiology

Modeling

ABSTRACT

Neuronal nicotinic acetylcholine receptors (nAChR), recognized targets for drug development in cognitive and neuro-degenerative disorders, are allosteric proteins with dynamic interconversions between multiple functional states. Activation of the nAChR ion channel is primarily controlled by the binding of ligands (agonists, partial agonists, competitive antagonists) at conventional agonist binding sites, but is also regulated in either negative or positive ways by the binding of ligands to other modulatory sites. In this review, we discuss models for the activation and desensitization of nAChR, and the discovery of multiple types of ligands that influence those processes in both heteromeric nAChR, such as the high-affinity nicotine receptors of the brain, and homomeric $\alpha 7$ -type receptors. In recent years, $\alpha 7$ nAChRs have been identified as a potential target for therapeutic indications leading to the development of $\alpha 7$ -selective agonists and partial agonists. However, unique properties of $\alpha 7$ nAChR, including low probability of channel opening and rapid desensitization, may limit the therapeutic usefulness of ligands binding exclusively to conventional agonist binding sites. New enthusiasm for the therapeutic targeting of $\alpha 7$ has come from the identification of $\alpha 7$ -selective positive allosteric modulators (PAMs) that work effectively on the intrinsic factors that limit $\alpha 7$ ion channel activation. While these new drugs appear promising for therapeutic development, we also consider potential caveats and possible limitations for their use, including PAM-insensitive forms of desensitization and cytotoxicity issues.

© 2011 Elsevier Inc. All rights reserved.

Contents

1. Introduction	916
2. Basic principles of allosteric proteins and modulation	916
2.1. Models of heteromeric nAChR activation and desensitization	917
2.2. Modeling potential mechanisms of positive allosteric modulators	918
2.3. Models for the unique properties of homomeric $\alpha 7$ nAChR and potential mechanisms for $\alpha 7$ PAM activity	919
3. Known allosteric modulators of heteromeric nAChR	919
4. Positive allosteric modulation of $\alpha 7$ nAChR	920
4.1. Structures, binding sites, and mechanisms of $\alpha 7$ PAMs	921
4.2. Demonstrated effects of PAMs with relevance to biological systems	924
4.3. Limiting factors of allosteric modulation as a therapeutic approach	925
5. Summary	927
Acknowledgements	927
References	927

Abbreviations: A-585539, (1S,4S)-2,2-dimethyl-5-(6-phenylpyridazin-3-yl)-5-aza-2-azoniabicyclo[2.2.1]heptane; A-867744, 4-(5-(4-chlorophenyl)-2-methyl-3-propionyl-1H-pyrrol-1-yl)benzenesulfonamide; 5-HI, 5-hydroxyindole; 5-HT, 5-hydroxytryptamine; α -btx, α -bungarotoxin; ACh, acetylcholine; CCMI, N-(4-chlorophenyl)- α -[[(4-chloro-phenyl)amino]methylene]-3-methyl-5-isoxazoleacetamide; CNS, central nervous system; dFBr, desformylflustrabromine; D_s, type II modulator-sensitive desensitization; ERK1/2, extracellular signal-related kinase-1 and -2; GTS-21, 3-(2,4-dimethoxybenzylidene)-anabaseine; JNJ-1930942, 2-[[[4-fluoro-3-(trifluoromethyl)phenyl]amino]-4-(4-pyridinyl)-5-thiazolemethanol; LY-2087101, [2-(4-fluoro-phenylamino)-4-methyl-thiazol-5-yl]-thio-phen-3-yl-methanone; nAChR, nicotinic acetylcholine receptor; MWC, Monod, Wyman, Changeux; NS-1738, 1-(5-chloro-2-hydroxy-phenyl)-3-(2-chloro-5-trifluoromethyl-phenyl)-urea; PAM, positive allosteric modulator; PNU-120596, 1-(5-chloro-2,4-dimethoxy-phenyl)-3-(5-methyl-isoxazol-3-yl)-urea; PNU-282987, N-[[[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-chlorobenzamide]; SB-206553, 3,5-dihydro-5-methyl-N-3-pyridinylbenzo [1,2-b:4,5-b']-di pyrrole-1(2H)-carboxamide; SLURP-1, secreted mammalian Ly-6/uPAR related protein 1; TM, transmembrane; TQS, 4-naphthalene-1-yl-3a,4,5,9b-tetrahydro-3-H-cyclopenta[c]quinoline-8-sulfonic acid amide.

* Corresponding author at: Department of Pharmacology and Therapeutics, University of Florida, P.O. Box 100267, Gainesville, FL 32610-0267, United States.
Tel.: +1 352 392 4712; fax: +1 352 392 9696.

E-mail address: rlpapke@ufl.edu (R.L. Papke).

1. Introduction

Many drugs, like nicotine, were used socially or clinically long before their mechanisms of action were identified and understood. Now that we begin to understand the molecular substrates that mediate the effects of nicotine in the brain and on human behavior, we recognize that there is tremendous therapeutic potential for new agents targeting neuronal and somatic nicotinic-type receptors, all of which are ligand-gated ion channels activated by the neurotransmitter acetylcholine (ACh). In order to advance nicotinic receptor-based therapeutics, we now face the traditional challenges of drug development: optimizing drugs for receptor subtype specificity, potency, pharmacokinetic properties, and, most importantly, for every hypothetical indication we need to consider the optimal mechanism of action.

Numerous distinct nicotinic acetylcholine receptor (nAChR) subtypes can be identified based on subunit composition and stoichiometry [1], and for particular indications we identify broad classes of nAChR that can be classified as distinct pharmacological subtypes to be either targeted or avoided for the sake of reducing side effects. The first two classes of nAChR to be identified and pharmacologically characterized were the primary mediators of synaptic transmission outside the central nervous system (CNS), at neuromuscular junctions and autonomic ganglia. As drugs are being developed for CNS indications ranging from analgesia to schizophrenia, both of these peripheral nicotinic systems are potential sources for off-target effects. Most nAChR function in the CNS is dichotomized between two broad classes of nAChR: heteromeric (primarily $\alpha 4$ -containing) and homomeric $\alpha 7$ receptors. Both of these receptor subtypes are most likely to be activated by ACh in the micromolar range. However, while $\alpha 4^*$ receptors initially open with high probability [2], they relax into high-affinity desensitized states that resist further activation. In contrast, $\alpha 7^*$ nAChR open with very low probability, and, although they are profoundly desensitized by high concentrations of agonist, their desensitization is readily reversible [3].

One question at the core of therapeutic drug development is how a drug will affect the endogenous signaling mediated by the natural activator. The naïve assumption is that an agonist will replace or augment the stimulation provided by natural activator. However, for most nAChR subtypes a primary effect of the prolonged presence of an agonist is to produce desensitization, decreasing the effect of the normal fluctuations in ACh signaling [4]. In this regard, the effect of an agonist may be similar to that of an antagonist, as is the case with two drugs commonly used to paralyze the neuromuscular junction, succinylcholine (a potent neuromuscular receptor agonist), and tubocurarine (a competitive antagonist). Likewise, treatments for nicotine addiction include both partial agonists of $\alpha 4^*$ receptors, such as varenicline and the putative $\alpha 4^*$ receptor antagonist bupropion [5]. However, for this indication it is not the natural ACh-mediated signaling the drugs are intended to oppose, but the pulsatile stimulation of nAChRs delivered by cigarette smoking.

An important alternative to the therapeutic development of agonists has been the development of positive allosteric modulators (PAMs), which can synergize and augment natural signals rather than oppose or attempt to replace them, while potentially preserving the topological distribution and rhythm of those natural neurotransmitter-mediated signals. Since PAMs bind to sites away from well-conserved agonist binding domains, increased receptor subtype selectivity may be an additional advantage of allosteric modulators. A proven area of success for the therapeutic development of PAMs is in regard to sedative-hypnotic drugs, such as barbiturates and benzodiazepines targeting GABA_A receptors, which belong to the same superfamily of ligand-gated ion channels as nAChRs [6]. Despite GABA_A PAMs

having been known and used clinically for decades, nAChR PAMs have only relatively recently been identified.

In this article we review new advances in the development of nAChR PAMs, potential therapeutic applications, and possible limitations for their use. We will discuss the fundamental differences between heteromeric and homomeric nAChRs in the context of receptor models and energy “landscapes”, discussing how PAMs may alter receptor function to promote either short-term or sustained levels of increased ion channel opening.

2. Basic principles of allosteric proteins and modulation

The term “allosteric” was first introduced following observations that bacterial enzymes were inhibited by the end product of synthetic pathways, even though the end product had limited structural similarities with the enzyme active site substrate [7]. The inhibition appeared to be non-competitive with substrate, which led to hypotheses that the non-competitive inhibitor produced conformational alterations in the protein [8] and the formulation of the well-known Monod, Wyman, Changeux (MWC) model of protein allostery [9–11]. The basic concept proposed that proteins are dynamic structures existing in multiple discrete functional states or conformations, all of which are accessible to the protein under resting conditions, and that binding of a ligand alters the resting equilibrium by reversibly stabilizing the protein in the conformation to which the ligand has greatest affinity. The conformation of each subunit was proposed to be constrained by the other subunits, and protein symmetry always conserved. Therefore, the binding of ligand at one site was predicted to alter the affinity of the other binding sites within the oligomer for the ligands. The model was soon applied to hemoglobin to describe the cooperative nature of oxygen binding, and to proteins involved in signal transduction, including membrane receptors as diverse as G-protein coupled receptors and ligand-gated ion channels, including nAChRs [10,11].

The MWC concept of allosteric proteins has enjoyed widespread acceptance. Unfortunately, the use of the term “allosteric” to describe ligands and sites of ligand interaction has become somewhat ambiguous. The term “allosteric ligand” is commonly used to refer to any ligand that binds to any site on the receptor and has the effect of promoting a specific conformation [12]. Under this definition, with the exception of true channel blockers all ligands including agonists, antagonists, and modulators are probably “allosteric”. However, the term “allosteric ligand” is also commonly used to describe ligands that modulate or regulate the protein through sites that are distinct from traditional agonist binding sites in protein receptors [10,11]. The terms “orthosteric” and “allosteric” are sometimes used to differentiate between canonical agonist binding domains and sites located elsewhere from which ligands positively or negatively modulate the efficacy or potency of ligands binding to the primary agonist binding sites on the receptor. These terms are basically inaccurate, but can be useful to distinguish agonists from modulators so long as it is appreciated that both types of ligand may produce global changes in protein conformation that influence receptor function.

In the case of nAChR, the distinction can be made between the primary allosteric sites where the binding of agonists or partial agonists is both necessary and sufficient to produce ion conducting conformations of the receptors. In contrast, ligands binding to modulatory allosteric sites are unable to produce receptor activation on their own, but facilitate agonist-evoked responses by modifying energy barriers associated with transitions between functional conformations (see below). However, observations that

some positive allosteric modulators may produce activation of GABA_A [13] and $\alpha 7$ receptors [14] independent of traditional agonist, blur the functional distinction between agonist and modulator. Nonetheless, such observations are entirely consistent with the MWC model since an allosteric modulator would alter the resting distribution of functional states of the receptor, perhaps by enhancing the small probability of spontaneous openings in the absence of orthosteric agonist.

2.1. Models of heteromeric nAChR activation and desensitization

The functions of allosteric proteins, including nAChRs and other ligand-gated channels, are typically modeled under the theoretical framework of MWC allostery. Discrete functional states (i.e. protein conformations) are defined, with basic models of nAChR required to account for resting closed states (C), open states (O), and desensitized states (D). There are sufficient data, from both




Fig. 1. The energy landscapes for nAChR state transitions. A) Hypothetical energy wells and barriers for the conformational states of heteromeric nAChR (e.g. muscle-type or neuronal $\alpha 4$ nAChR) as functions of the level of agonist occupancy (see description in text). Under equilibrium conditions, the distributions of receptors into the resting closed (C), brief open (O^*), long-lived open (O'), and desensitized (D) states will be determined by the relative free energy of the states (represented by vertical displacements). Dynamically, the transition rates between the states will be inversely related to the log of the energy barriers between the states. B) Two ways in which PAMs may operate on the energy profile of receptors in the doubly liganded state to increase the probability of channel opening: either transiently, after a jump in agonist concentration (left schematic) or under steady-state conditions (right schematic). C) Hypothetical scheme representing state transitions of $\alpha 7$ nAChR under various levels of agonist occupancy. In the absence of PAMs, $\alpha 7$ nAChR do not exhibit a long-lived open state, but do show a unique desensitized state (D_s) that is preferentially favored at higher levels of agonist occupancy. They also have an intrinsic desensitized state (D_i), which is analogous to the D state of heteromeric receptors (A). Very efficacious type II PAMs, such as PNU-120596, appear to either destabilize the D_s state, or hypothetically may convert it into a conducting state similar to the O' state of heteromeric receptors. This hypothesis is represented by (O') symbols inserted into the panel.

heteromeric muscle-type and neuronal nAChR [15–17], to indicate the existence of two distinct open states which have similar ionic conductance, but different mean channel open times. In Fig. 1 we represent these two open states as O^* , short-lived openings that occur in isolation, and O' , longer openings that sometimes occur in bursts. Because the behavior of individual receptor molecules are stochastic in nature, typical Markov assumptions are made to describe the interconnection between these several states and may, with sufficient data, allow assignment of rate constants for the interstate conversions. A complementary perspective is provided by Eyring rate theory, which describes the relationships among the receptor states in regard to their respective free energy levels and intervening energy barriers. The heights of the energy barriers are inversely proportionate to the logs of the transition rate constants in standard presentations of Markov models.

In Fig. 1A we show hypothetical energy landscapes that provide a qualitative description of key features associated with heteromeric nAChR, such as $\alpha 4$ -containing neuronal nAChR. Heteromeric nAChR have two agonist binding sites and, therefore, can have three different levels of agonist occupancy, with 0, 1, or 2 agonist molecules bound. The relative stabilities and interconversion rates between the states depend on the level of agonist occupancy [10,18]. Therefore, to clarify the presentation of these landscapes, we have separated the distinct profiles that would be associated with the different levels of agonist occupancy by a given agonist. In reality, the overall free energy of the unliganded condition is probably higher (more positive) than the average free energy corresponding to the singly liganded condition, which in turn is higher than the doubly liganded condition. For simplicity, the C state corresponding to each level of agonist occupancy is used to align the energy landscapes, and barriers between states are meant to represent the net transition, including any intermediate states such as the recently proposed flip state between resting closed and open states [19]. Recent work with tethered agonists has provided the additional perspective and complexity that within the orthosteric site a single ligand can bind in multiple orientations, each of which may be associated with a unique energy landscapes [20].

With no agonist molecules bound (left-most profile in Fig. 1A), receptors are most stable in the resting closed state, with some equilibration between the resting closed and the desensitized states, and vanishingly low probability of channel opening (P_{open}). With a single agonist binding site occupied (middle scheme, Fig. 1A), there is an increased probability of channel opening, particularly to the brief O^* state. Early studies that first identified the two distinct open states associated the short-lived events with singly liganded receptors [17], and it was unclear from those studies whether or not singly liganded receptors could also open to the longer-lived state. When experiments were conducted at increasing concentrations of agonist, the frequency of long-lived openings increased, essentially as the probability of having doubly bound receptors increased. Using conditional elimination of single agonist binding sites in muscle-type receptors, we have recently shown [21] that while singly bound receptors are most likely to open to the O^* state, they can also open to the O' state. The probability of opening to the longer-lived O' state from activation at one binding site increases under conditions of saturated agonist binding to that site. In addition, those experiments demonstrated that even singly bound receptors were ultimately most stable in the D state, as shown schematically.

Receptors with two ligands bound have a high probability of converting to the long-lived O' open state, as represented in the right-most scheme in the Fig. 1A. It has been reported [2] that following the rapid application of a high concentration of agonist, sufficient to saturate the binding sites and fully populate the resting closed state, $\alpha 4\beta 2$ nAChRs have approximately an 80%

probability of synchronously passing into open states prior to reaching an equilibration which will ultimately favor the D state. Receptors with both agonist sites bound may also open to the O^* state, but isolated brief openings are only about 10% as common as longer openings or bursts [17,22]. Conversions occurring between the O^* and O' would be undetectable as unique events in single channel recordings and would add to the apparent open duration of O' events.

The model as presented in Fig. 1 assumes that the agonist binding steps proceed from a rapid change in agonist concentration and all receptors are initially in the resting closed state. Therefore, the affinity associated with the association and dissociation steps shown below the state-transition landscapes would correspond with the micromolar ACh potency for channel activation. However, it is known that agonists bind with much higher apparent affinity once the receptors have relaxed into the low energy D state sometime after the initial presentation of agonist. This could be modeled by assigning different association and dissociation rates to each of the states in the model. However, if dissociation rates from the O and D states are so slow that they approach zero, then the apparent high affinity of the receptors at equilibrium might be related to the slow conversion between the D and C states, with dissociation still occurring primarily from receptors in the C state.

2.2. Modeling potential mechanisms of positive allosteric modulators

Allosteric modulators may affect receptor function in any of several ways that could have the net effect of increasing channel activation under particular sets of conditions. One requirement for “true” positive allosteric modulators is that they are distinct from agonists, having no intrinsic channel activation properties. Therefore, we can consider PAM mechanisms in terms of modifying the normal receptor mechanisms discussed above.

One potential effect of a PAM would be to enhance agonist binding to the resting state of the receptor, increasing potency. This was proposed to be the mechanism for benzodiazepine enhancement of GABA_A receptor function [23]. This particular form of modulation would be most effective at increasing responses to weak stimuli, with little or no effect on the response to strong stimuli, since the energy landscape under the conditions of receptor saturation would be unchanged. It is generally believed that the relatively large therapeutic index associated with benzodiazepines, compared to other GABA_A modulators such as barbiturates, may be due to the fact that they do not produce more GABA_A receptor activation than would be created by high concentration of GABA alone.

Positive allosteric modulators are commonly observed to increase agonist efficacy, at least in regard to the peak current evoked by an agonist application. Two ways they might potentially do so are illustrated in Fig. 1B as alternative energy landscapes for the doubly liganded condition. In the scheme on the left, the energy barrier from C to O' is reduced, allowing for faster opening after a rapid increase in agonist concentration. The energy barrier from O' to D is increased, which would have the effect of increasing mean channel open time and the probability of multiple openings prior to desensitization. These alterations in the energy profile would produce a transient increase in P_{open} , conditional on all receptors initially being in the C state. Note, however, that in this scheme the absolute disposition of the four states is unchanged from the initial condition in Fig. 1A. Therefore, even with these changes to the energy profile, the receptors would ultimately relax to the same equilibrium conditions, strongly favoring occupancy of the D state.

The scheme on the right in Fig. 1B allows for increased P_{open} by destabilizing the D state relative to the other states in the model. As drawn, this would primarily be associated with increasing the exit

rates out of the D state. A PAM working through this mechanism might have a relatively small effect on the peak transient current evoked by a rapid agonist application, but would have a large effect under equilibrium conditions, producing a significant amount of steady-state current. However, since, as mentioned above, $\alpha 4$ -containing nAChRs naturally open with high probability upon stimulation with ACh, the potential range for enhanced activation of these receptors may be limited.

The two hypothetical energy landscapes in Fig. 1B represent mechanisms that could produce either transient or steady-state potentiation, and both assume that the PAM effects are on the same population of receptors that are functional in the absence of the PAM. However, it has been reported in some systems that there are two pools of surface nAChRs and that only a small percentage (10–20%) are capable of being activated at all [2,24]. Hypothetically, PAMs might also work by switching receptors from the non-activatable pool into the activatable population. It may be especially important to consider this possible mechanism for agents that are known to have potential effects on cellular signal transduction, such as genistein, a tyrosine kinase inhibitor and an $\alpha 7$ potentiator [25].

2.3. Models for the unique properties of homomeric $\alpha 7$ nAChR and potential mechanisms for $\alpha 7$ PAM activity

The hypothetical energy profiles illustrated for heteromeric nAChR in Fig. 1A are unlikely to be applicable to homomeric $\alpha 7$ nAChR. Since $\alpha 7$ nAChR do not require specialized subunits to provide the complementary surface of the ligand-binding domain, there are five putatively equivalent binding sites per receptor rather than the two likely non-equivalent sites in heteromeric nAChR [26]. Additionally, $\alpha 7$ nAChR manifest a unique “fast” desensitized state, entry into which appears to be concentration-dependent, suggesting that the state is stabilized by high fractional occupancy of the agonist binding sites [27]. Fig. 1C illustrates a series of hypothetical energy landscapes for $\alpha 7$ nAChR as related to the level of binding site occupancy by a full agonist.

Key features of $\alpha 7$ activation and desensitization are consistent with the model in Fig. 1C. The P_{open} of $\alpha 7$ receptors is very low under all conditions, and open times are very brief, usually less than 100 μs [3,28]. These features are associated with the steep barrier for entry into the open state, which is located in a very shallow trough. In this model, we present the rapid form of desensitization unique to $\alpha 7$ as the D_s state and propose that it is analogous to the O^* state of heteromeric nAChR. The D_s state is more stable than the short-lived O^* state, and it is most readily entered into with high levels of agonist occupancy. The true intrinsic desensitized state (D_i in Fig. 1C) of $\alpha 7$ receptors is less stable than that of heteromeric nAChR, and so the absolute free energy difference between the D_i and C states at high levels of agonist occupancy is less for $\alpha 7$ than for the heteromeric nAChR modeled in Fig. 1.

All of the general mechanisms for the potentiation of nAChR discussed above and illustrated in Fig. 1B might be applied to $\alpha 7$ -selective PAMs. Additionally, as will be discussed below, the intrinsically low P_{open} of $\alpha 7$ greatly broadens the range for possible potentiation of this important therapeutic target and suggests that these PAMs may draw on unique mechanisms. However, since $\alpha 7$ is a receptor with high calcium permeability, it must also be considered whether extreme increases in $\alpha 7$ channel activation may lead to unexpected, and potentially undesirable, effects. For this reason, drug development in this area benefits from the generation of compounds with widely different activity profiles and depends on the testing of such compounds in a variety of appropriate model systems.

3. Known allosteric modulators of heteromeric nAChR

Several ions and molecules have been identified to potentiate agonist-evoked responses of heteromeric nAChRs. However, the development of PAMs with activity on heteromeric nAChRs has developed slowly relative to the identification and development of $\alpha 7$ PAMs. In general, the known heteromeric receptor PAMs produce only modest potentiation and are poorly selective for specific heteromeric receptor subtypes. In addition, several PAMs of $\alpha 7$ nAChR actually have negative effects on $\alpha 4\beta 2$ and $\alpha 3\beta 4$ mediated responses (see below).

Calcium ions were shown to allosterically potentiate nAChR-mediated responses from heteromeric $\alpha 2$ -, $\alpha 3$ -, and $\alpha 4$ -containing receptors expressed heterologously and in native tissues including the medial habenula, chromaffin cells, and ganglion neurons [29–31]. Responses were potentiated by low millimolar concentrations (<10 mM) of calcium, which are considered to be physiologically relevant, up to about 4-fold in a voltage-insensitive manner that depended on agonist concentration and extracellular, not intracellular, calcium [30]. Although at millimolar concentrations of external calcium single-channel conductance was decreased, increased frequency of single-channel openings, with little to no effect on open durations, was found to account for the potentiation [29,31]. In habenular neurons, similar potentiation was observed with barium and strontium ions, but not magnesium ions [29], while in $\alpha 3\beta 4$ receptors expressed in *Xenopus* oocytes, neither barium nor magnesium ions potentiated responses [30].

Zinc ions, which are found in high concentrations in many brain regions and are key functional components of many proteins, were shown to potentiate $\alpha 2$ -, $\alpha 3$ -, $\alpha 4$ -containing nAChR in a biphasic nature, with potentiation occurring between 1 and 100 μM and inhibition occurring at higher concentrations. Responses of $\alpha 4\beta 4$ receptors were potentiated about 5-fold with more modest potentiation observed on $\alpha 2\beta 4$, $\alpha 2\beta 4$, $\alpha 3\beta 4$, and $\alpha 4\beta 2$ receptors. Zinc did not potentiate but rather inhibited $\alpha 3\beta 2$ receptors with an IC_{50} of $\sim 100 \mu\text{M}$. Both potentiation and inhibition of responses by zinc were voltage independent [32]. Similar to its effects on $\alpha 3\beta 2$ receptors, ionic zinc inhibited $\alpha 7$ -mediated responses with an IC_{50} of 27 μM in a voltage-insensitive manner [33]. With site-directed cysteine mutations and reactive sulfhydryl reagents, residues E59 and H162 on the $\alpha 4$ subunit were found to be critical for potentiation by zinc, potentially forming part of the zinc binding site at subunit interfaces not containing the orthosteric site [34]. Potentiating effects of lead ions were observed on $\alpha 3\beta 2$ receptors at concentrations between 1 and 250 μM , while similar concentrations inhibited responses by $\alpha 3\beta 4$ and $\alpha 4\beta 2$ receptors [35].

Naturally synthesized steroids (i.e. progesterone, testosterone, estradiol, corticosterone) were shown to exert allosteric effects on heteromeric $\alpha 4$ -containing and $\alpha 3$ -containing nAChR, the effects however, are generally inhibitory at low μM concentrations [36–39]. One exception is 17 β -estradiol, which has been found to potentiate human, but not rat, $\alpha 4\beta 2$ and $\alpha 4\beta 4$ responses about 3–4-fold with no potentiation of responses by $\alpha 3\beta 2$ or $\alpha 3\beta 4$ receptors [39–41]. Single-channel recordings from potentiated $\alpha 4\beta 2$ receptors suggest that 17 β -estradiol increases the frequency of channel openings, with little effect on unitary conductance [41]. Chimeric $\alpha 4$ subunits and site-directed mutagenesis were used to identify a C-terminal region of the $\alpha 4$ subunit as critical for potentiation by estradiols [40,41].

The protein lynx-1, which contains remarkable structural similarity to snake neurotoxins, is expressed by cerebellar, cortical, hippocampal neurons and was found to co-localize with neuronal nAChR and modestly increase peak responses of $\alpha 4\beta 2$ receptors when applied in soluble form [42], but was shown to increase macroscopic current decay rates, reduce rate of recovery from

desensitization, and decrease agonist potency when lynx-1 was co-expressed with $\alpha 4\beta 2$ receptors in *Xenopus* oocytes and mammalian cells [43], and appears to be a mechanism used to negatively modulate $\alpha 4\beta 2$ -containing receptors [44]. In general, the modulatory properties of divalent cations, proteins, and steroids for nAChRs support the hypothesis that allosteric modulation is a natural mechanism controlling the function of endogenous nAChRs [45].

The anthelmintic agents levamisole and morantel are full agonists at nematode nAChR, but have been found to allosterically potentiate human $\alpha 3$ -containing receptors expressed in *Xenopus* oocytes [46,47]. Micromolar concentrations of levamisole were found to potentiate agonist-evoked responses of $\alpha 3\beta 2$ and $\alpha 3\beta 4$ receptors about 4-fold, while millimolar concentrations significantly inhibited responses, likely due to open channel block [46]. Low micromolar concentrations ($\leq 10 \mu\text{M}$) of morantel potentiated ACh-evoked responses on $\alpha 3\beta 2$ up to 8-fold primarily by increasing the frequency of single-channel openings and the duration of bursts, rather than increasing the unitary conductance or altering macroscopic response decay kinetics [47]. Although both levamisole and morantel produce weak activation of human $\alpha 3$ -containing receptors, they appear to bind away from the orthosteric site since inhibition by DH β E is non-competitive [46,47]. Evidence that morantel binds in the N-terminal extracellular region at subunit interfaces not containing orthosteric sites was provided by experiments utilizing site-directed cysteine mutations and reactive sulfhydryl reagents [48].

The acetylcholinesterase inhibitor galantamine has been shown to potentiate $\alpha 4\beta 2$, $\alpha 3\beta 4$, and $\alpha 6\beta 4$ nAChRs at concentrations between 0.1 and 1 μM , and at concentrations $>10 \mu\text{M}$ galantamine was found to inhibit responses [49]. However, it has recently been reported that galantamine potentiation of $\alpha 4\beta 2^*$ nAChR required the presence of $\alpha 5$ subunits [50]. At low concentrations, galantamine was shown to increase the frequency of single-channel openings of nAChR receptors expressed in PC12 cells [51]. The maximal potentiation produced by galantamine on heteromeric receptors is modest. Nonetheless, some have speculated that the therapeutic actions of galantamine are primarily produced through the compound's action on nAChRs, since the cholinesterase activity of galantamine is also weak compared to other known cholinesterase inhibitors [49].

Three (2-amino-5-keto)thiazole compounds (LY-2087101, LY-1078733, LY-2087133) were identified in a high-throughput screen as PAMs of nAChRs [52]. Although non-selective for a single nAChR subtype, the compounds do show good selectivity against other ligand- and voltage-gated ion channels. Agonist-evoked peak responses of $\alpha 2\beta 4$, $\alpha 4\beta 2$, $\alpha 4\beta 4$, and $\alpha 7$, but not $\alpha 1\beta 1\gamma\delta$, $\alpha 3\beta 2$, or $\alpha 3\beta 4$, were potentiated from ~ 2 -fold up to ~ 20 -fold with concentrations $<10 \mu\text{M}$, and inhibited with concentrations $>10 \mu\text{M}$. These compounds showed the greatest potentiation of $\alpha 4\beta 2$ -mediated responses from receptors expressed in *Xenopus* oocytes. All three compounds were unable to displace [^3H]epibatidine from $\alpha 4\beta 2$, $\alpha 4\beta 4$, and $\alpha 3\beta 4$ receptors, and potentiation was unaffected by blockade of a number of intracellular second messenger systems, suggesting these compounds work directly on the receptors through allosteric mechanisms [52]. In the ventral tegmental area, LY-2087101 enhanced nicotine-dependent increases of dopamine neuron firing in a DH β E sensitive manner, suggesting the effect was mediated through potentiation of $\alpha 4\beta 2$ nAChR [53].

Desformylflustrabromine (dFBr) was identified as a PAM of $\alpha 4\beta 2$ nAChR from screens of compounds isolated from *Flustra foliacea*, a bryozoan commonly found in the North Sea. Low micromolar ($<10 \mu\text{M}$) concentrations of dFBr modestly potentiate (~ 2 – 3 -fold) agonist-evoked currents from human $\alpha 4\beta 2$ expressed in *Xenopus* oocytes, while higher concentrations inhibit responses,

likely through open-channel block [54,55]. No potentiation was observed for $\alpha 7$ or $\alpha 3\beta 4$ receptors [56], and in fact dFBr was shown to inhibit $\alpha 7$ responses with an IC_{50} of 44 μM [55]. Recordings of single-channel currents reveal that dFBr potentiates responses of $\alpha 4\beta 2$ by increasing the frequency of openings and modestly increasing the open durations. In addition, dFBr was reported to activate desensitized receptors [54]. dFBr is one of the few known compounds capable of discriminating $\alpha 4\beta 2$ from other nAChR subtypes to produce selective potentiation of responses. Initial screens of a series of carbamates [57] and a series of piperidines [58] have suggested the discovery of other novel molecules that selectively potentiate $\alpha 4\beta 2$ -mediated responses with good potency, but these compounds have not yet been fully characterized.

4. Positive allosteric modulation of $\alpha 7$ nAChR

Much of the preclinical data with nicotinic ligands provide positive support for selective stimulation of $\alpha 7$ nAChR as a therapeutic strategy to treat cognitive deficits in Alzheimer's disease and schizophrenia [59–61]. In addition, the $\alpha 7$ gene has been linked to sensory gating deficits observed in schizophrenia [62,63], and $\alpha 7$ agonists have been shown to normalize sensory gating deficits in animal models (for example, [64,65]), as well as in schizophrenic humans [66].

Positive allosteric modulators selective for $\alpha 7$ share some of the putative therapeutic activities ascribed to $\alpha 7$ agonists, including enhancement of memory-related behavioral performance in the eight-arm radial maze, social recognition, and Morris water maze tasks [67,68], and normalization of sensory gating deficits [67,69–72]. As noted above, PAMs of $\alpha 7$ have special appeal because they overcome the limitations associated with the intrinsically low P_{open} and rapid desensitization of $\alpha 7$ nAChR, potentially with significant benefit for processes that depend on channel activation [73].

Calcium ions were the first known allosteric potentiators of $\alpha 7$ responses [74], followed by the discovery of the antiparasitic agent ivermectin [75], the serotonin metabolite 5-hydroxyindole (5-HI; [76,77]), the acetylcholinesterase inhibitor galantamine [78], serum albumin peptides [79], and secreted mammalian Ly-6/uPAR related protein 1 (SLURP-1 [80]). In addition, the tyrosine kinase inhibitor genistein augments $\alpha 7$ responses, and some evidence suggests it functions directly on $\alpha 7$ as a PAM [25,81], but other evidence suggests an indirect mechanism through kinase inhibition [82]. Unfortunately, these “first generation” $\alpha 7$ PAMs lack desirable potency, efficacy, and/or selectivity profiles. For example, ivermectin is non-selective [83–86]; galantamine exhibits poor efficacy as a PAM, is non-selective [49,87], and works through a dual mechanism as an acetylcholinesterase inhibitor; and 5-HI has low potency and also potentiates 5-hydroxytryptamine $_3$ (5-HT $_3$) receptors [88].

Since the discovery of the first $\alpha 7$ modulators, small molecule PAMs with diverse potency, efficacy and/or selectivity profiles have emerged (Fig. 2, Table 1). These include 1-(5-chloro-2,4-dimethoxy-phenyl)-3-(5-methyl-isoxazol-3-yl)-urea (PNU-120596) [69,89], 2-(amino-5-keto)thiazole compounds (e.g. LY-2087101) [52], 1-(5-chloro-2-hydroxy-phenyl)-3-(2-chloro-5-trifluoromethyl-phenyl)-urea (NS-1738) [68], N-(4-chlorophenyl)-alpha-[[[4-chloro-phenyl]amino]methylene]-3-methyl-5-isoxazoleacetamide (compound 6, also known as CCMI) [67], 4-naphthalene-1-yl-3a,4,5,9b-tetrahydro-3-H-cyclopenta[c]quinoline-8-sulfonic acid amide (TQS) [81], 3,5-dihydro-5-methyl-N-3-pyridinylbenzo [1,2-b:4,5-b']-di pyrrole-1(2H)-carboxamide (SB-206553) [71], 4-(5-(4-chlorophenyl)-2-methyl-3-propionyl-1H-pyrrol-1-yl)benzenesulfonamide (A-867744) [70,90], 2-[[4-fluoro-3-(trifluoromethyl)phenyl]amino]-4-(4-pyridinyl)-5-thia-




Fig. 2. Structures of $\alpha 7$ -active PAMs. The PAMs classified as type I, which increase the amplitude but do not strongly alter the kinetics of $\alpha 7$ -mediated agonist-evoked responses, are shown on the left. They include: CCM1, N-(4-chlorophenyl)- α -[[(4-chloro-phenyl)amino]methylene]-3-methyl-5-isoxazoleacetamide; NS-1738, 1-(5-chloro-2-hydroxy-phenyl)-3-(2-chloro-5-trifluoromethyl-phenyl)-urea; Galantamine; 5-HI, 5-hydroxyindole; LY-2087101, [2-(4-fluoro-phenylamino)-4-methyl-thiazol-5-yl]-thiophen-3-yl-methanone; Genistein; and Ivermectin. Type II PAMs, which appear to slow or reverse $\alpha 7$ desensitization are shown on the right and include: PNU-120596, 1-(5-chloro-2,4-dimethoxy-phenyl)-3-(5-methyl-isoxazol-3-yl)-urea; TQS, 4-naphthalene-1-yl-3a,4,5,9b-tetrahydro-3-H-cyclopenta[c]quinoline-8-sulfonic acid amide; and A-867744, 4-(5-(4-chlorophenyl)-2-methyl-3-propionyl-1H-pyrrol-1-yl)benzenesulfonamide. PAMs proposed to be intermediate in their activity are shown in the center and include: JNJ-1930942, 2-[[4-fluoro-3-(trifluoromethyl)phenyl]amino]-4-(4-pyridinyl)-5-thiazolemethanol; and SB-206553, 3,5-dihydro-5-methyl-N-3-pyridinylbenzo [1,2-b:4,5-b']-di pyrrole-1(2H)-carboxamide.

zolemethanol (JNJ-1930942) [72], and a series of amide derivatives (compounds 2–4) [91]. While these “second generation” PAMs generally provide a range of potencies and efficacies useful to consider for therapeutic development, some still suffer from lack of pure $\alpha 7$ -selectivity. For example, CCM1 was discovered from a library of GABA_A PAMs [67], SB-206553 is a 5-HT_{2B/C} receptor antagonist [71], and LY-2087101 (and analogues) potentiates responses at several nAChR subtypes [52]. In addition, NS-1738 [68], TQS [81], SB-206553 [71], A-867744 [90], and compounds 2–4 [91] have been shown to inhibit responses of $\alpha 3\beta 4$ and $\alpha 4\beta 2$ nAChR.

The identified $\alpha 7$ PAMs have considerable diversity, ranging from proteins to small molecules. Even the small molecule PAMs vary significantly in structure, as well as in properties and probably mechanisms of the modulation [92]. Gronlien et al. [81] proposed that $\alpha 7$ PAMs be divided into two classes, type I and type II, based on the functional properties of modulation. Following the rapid application of agonists, all PAMs appear to increase receptor sensitivity to agonists, current magnitudes, and empirical Hill coefficients; the type I PAMs (for example 5-HI, NS-1738, or CCM1) do so with little or no effect on the basic onset and decay kinetics, or shape, of the response, while the type II PAMs (for example PNU-120596, TQS, or A-867744) markedly slow response decay kinetics and can even activate receptors that have been desensitized by applications of high agonist concentrations or by application of

agonists like GTS-21 (3-(2,4-dimethoxybenzylidene)-anabaseine) that produce residual inhibition or desensitization [14,67,69,81,90,93]. An observation yet to be explained that is sometimes seen with responses potentiated by type II PAMs is an initial biphasic current which rapidly rises and decays, followed by the emergence of a distinct secondary component. Low concentrations of PNU-120596 and TQS produce such responses, but at higher PAM concentrations the two components merge [81,90,94]. Responses potentiated by A-867744 apparently do not show the secondary component at low or high concentrations [90]. Since the two classes of $\alpha 7$ modulators were proposed, additional PAMs (Fig. 2) have been identified that display properties intermediate to the type I and type II classes [71,72,94].

4.1. Structures, binding sites, and mechanisms of $\alpha 7$ PAMs

As noted above, the potential mechanisms for $\alpha 7$ PAMs include the same mechanisms that might apply to the activation of heteromeric nAChR (Fig. 1B) and, additionally, unique effects associated with the special (PAM-sensitive) D_s state of $\alpha 7$ nAChR. Specifically, modulation of the $\alpha 7$ D_s state appears to underlie the extremely effective modulation produced by type II PAMs such as PNU-120596. One question that arises is whether type II PAMs stabilize functional states intrinsic to the receptor or create new conducting states. PNU-120596 has no effect on ion selectivity and

Table 1
Summary of published $\alpha 7$ PAM EC₅₀ and maximal potentiation values*.

PAM	EC ₅₀ , μ M	Max. % or fold-increase	Cell-type (species of $\alpha 7$)	Stimulating agonist	PAM preapplication	Measurement	Reference
Ivermectin	6.8	3.5-fold	<i>Xenopus laevis</i> oocytes (rat)	100 μ M ACh	60 s	Peak current	Collins and Millar [114]
BSA	10	6-fold	Chick ciliary ganglion	20 μ M nicotine	Not specified	Peak current	Conroy et al. [79]
Genestein	20	267%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Gronlien et al. [81]
	40	227%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	No	Peak current	Gronlien et al. [81]
5-HI	2500	1208%	<i>X. laevis</i> oocytes (human)	60 μ M ACh	No	Peak current	Zwart et al. [77]
	631	541%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Gronlien et al. [81]
	1585	350%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	No	Peak current	Gronlien et al. [81]
CCMI	0.7	45% of [EC ₁₀₀] ACh response	<i>X. laevis</i> oocytes (human)	[EC ₅] ACh	30 s	Peak current	Ng et al. [67]
NS-1738	3.4	322%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Timmermann et al. [68]
	1.6	1170%	GH4C1 (human)	300 μ M ACh	60 s	Peak current	Timmerman et al. [68]
	12.5	12-fold	GH4C1 (rat)	300 μ M ACh	150 s	Current; area	Friis et al. [89]
PNU-120596	0.2	400%	SH-EP1 (human)	100 μ M ACh	120 s	Calcium transient	Hurst et al. [69]
	1.6	455%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Gronlien et al. [81]
	1.5	37-fold	<i>X. laevis</i> oocytes (rat)	50 μ M ACh	Duration not specified	Peak current	Young et al. [113]
	0.1	203% of 10 μ M nicotine response	GH4C1 (rat)	0.5 μ M nicotine	10 min	Calcium transient	Dunlop et al. [71]
	4.9	2500-fold	GH4C1 (rat)	300 μ M ACh	150 s	Current; area	Friis et al. [89]
	5	Not reported	IMR-32 (human)	10 nM PNU-282987	60 s	Calcium transient	Gopalakrishnan et al. [99]
	1	Not reported	IMR-32 (human)	100 nM PNU-282987	60 s	Calcium transient	Gopalakrishnan et al. [99]
TQS	3.2	418%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Gronlien et al. [81]
	6.2	65-fold	<i>X. laevis</i> oocytes (human)	100 μ M ACh	Not specified	Peak current	Gill et al. [14]
A-867744	1	52% of max potentiation by PNU-120596	IMR-32 (human)	10 μ M PNU-282987	3–5 min	Calcium transient	Faghih et al. [70]
	1.1	733%	<i>X. laevis</i> oocytes (human)	100 μ M ACh	60 s	Peak current	Malysz et al. [90,94]
SB-206553	1.5	82% of 10 μ M nicotine response	GH4C1 (rat)	0.5 μ M nicotine	10 min	Calcium transient	Dunlop et al. [71]
JNJ-1930942	1.9	21-fold	GH4C1 (human)	100 μ M choline	Duration not specified	Calcium transient	Dinklo et al. [72]

* EC₅₀ values for potentiation of $\alpha 7$ -mediated responses by SLURP-1, galantamine, and LY-2087101 not found in literature.

little, if any, effect on channel conductance [69], supporting the interpretation that PNU-120596 stabilizes intrinsic states of the channel.

In the absence of modulators, it may be the case that wild-type $\alpha 7$ receptors can only activate in a manner similar to heteromeric receptors with a single occupied binding site, that is, primarily to the short-lived O^* state. We hypothesize that an effect of type II PAMs is to convert the D_s state of $\alpha 7$ nAChR to a functional homolog of the O' state of heteromeric nAChR (Fig. 1C). While the cooperative effects of a second binding event help to promote additional conformational change associated with the long-lived O' state in heteromeric receptors, the analogous cooperative effects may be negative rather than positive in $\alpha 7$, effectively promoting the unique D_s desensitized state [27]. It is interesting to note that the $\alpha 7$ L248T (also known as L9T) gain-of-function mutation has also been reported to slow or eliminate the unique form of $\alpha 7$ desensitization. The change in structure related to the binding of a type II PAM or to the L248T mutation in the transmembrane (TM) domain, may have the effect of reversing the negative cooperativity related to the effect of multiple ligand binding in $\alpha 7$.

As shown in Fig. 2, the $\alpha 7$ PAMs are constituted of several aromatic rings, hydrogen-bonding donors or hydrogen-bonding acceptors, which are oriented differently in space. All of the PAMs are neutral (uncharged) under physiological pH except galantamine, consistent with the hypothesis that this molecule has a unique binding site near the orthosteric site for agonists [95]. Several of the known $\alpha 7$ PAMs contain unsymmetrical aryl ureas or amides (CCMI, NS-1738, PNU-120596, and SB-206553), a commonality which has been applied in pharmacophore *in silico* screening methods to identify novel PAMs [96].

Although the specific orientation of the pharmacophore is likely a key, deciding element of a modulator's potentiation profile, there appear to be no strict governing rules. Even the type I modulators CCMI, 5-HI, and LY-2087101 may interact with the receptor differently, solely because of the different hydrogen bonding and hydrophobic group orientation, as shown in a solvent density guided docking study [97]. However, this study evaluated PAM binding in the ligand-binding domain, which is an unlikely binding site for most allosteric modulators. Although pharmacophore orientations among individual PAMs appear variable, there are some common elements shared between the two classes of PAMs on the electrostatic potential surface. For example, the type I modulator LY-2087101 and the type I/type II modulator JNJ-1930942 share the 2-(4-fluorophenylamino)-thiozoly group. Perhaps, the differences in their effects on the $\alpha 7$ desensitized states may arise from the dissimilar orientations of the misaligned aromatic rings (shown in Fig. 3A). As shown in Fig. 3B, the type I PAM NS-1738 has a unique electron deficit on the hydroxyl group (typical of hydrogen bond donors), while PNU-120596 features the electron-rich isoxazole ring (suggesting function as a hydrogen bond acceptor or having a role in dipole–dipole interactions). We speculate that type I PAMs prefer hydrogen-bonding donors on aromatic rings, while type II PAMs favor electron-rich aromatic rings without hydrogen-bonding donors.

While the mechanisms of action of PAMs remain unknown, some progress is being made through introduction of point mutations, use of $\alpha 7$ -5HT₃ chimeras, and molecular docking simulations to identifying locations on the receptor that are important for PAM function (Fig. 3C–D). It is important to appreciate that alteration of PAM function by point mutations




Fig. 3. $\alpha 7$ PAM structures and putative binding sites. A) Pharmacophore orientation comparison between LY-2087101 and JNJ-1930942. The structures of the two compounds were optimized using Chem 3D Pro v.12.0 (CambridgeSoft, Cambridge, MA): their energies were first minimized by molecular mechanics to minimum RMS gradient of 0.001 separately. Then LY-2087101 and JNJ-1930942 were overlaid on top of each other and shown in green and magenta, respectively. B) Molecular electrostatic potential surfaces of NS-1738 (left) and PNU-120596 (right). The structures of the two compounds were optimized using Gaussian09 (Gaussian, Inc., Wallingford, CT). The electrostatic potential on a total electron density isosurface ($0.002 \text{ Bohr}/\text{\AA}^3$) of the two compounds is displayed in GaussView (Gaussian, Inc., Wallingford, CT) with red color indicating the most negative electrostatic potential and blue color indicating the most positive electrostatic potential, respectively. The structures of the two compounds are displayed on top of each electrostatic potential surface. C) Structural domains of nAChR regulating PAM activity and/or binding. The key elements of the $\alpha 7$ receptor were modeled using the 2BG9 template for the transmembrane domain fused to the N-terminal extracellular domain made from the 2PGZ template. The C-loop is shown in yellow, and the M2 and M3 linker loop is shown in green. The view was made from the outside of the channel pore, and the four transmembrane helices (M1–M4) were lined clockwise as shown, putting the M2 helix toward the channel pore. Residues involved for different $\alpha 7$ PAMs potentiation are displayed as spheres of different colors as indicated: calcium, tan; galantamine, cyan; common residues for type I PAMs Ivermectin and LY-2087101 and the type II PAM PNU-120596, blue; residues only for Ivermectin, orange; residues only for PNU-120596, red. Leucine 248 is also shown in magenta in both subunits. D) A close-up view of the transmembrane domain, made by rotating the view in part C 90° vertically away from the plane of the paper.

does not necessarily indicate the site of ligand interaction. The use of the *Caenorhabditis elegans* $\alpha 7$ homolog ACR-16, which is insensitive to potentiation by ivermectin, 5-HI, genistein, NS-1738, and PNU-120596, has been suggested as a complementary approach to use of $\alpha 7/5$ -HT₃ chimeras to identify $\alpha 7$ domains that are critical for allosteric potentiation [98].

In general, PAMs have little to no effect on equilibrium binding of radiolabeled $\alpha 7$ competitive antagonists or high-affinity agonists in membrane preparations from rodent brain or heterologous expression systems [67,68,71,72,90,91,99], supporting the concept that PAMs bind in a non-competitive manner away from the traditional agonist binding site. However, while A-867744 was unable to displace [³H]methylcaconitine (MLA) binding, it displaced the high-affinity $\alpha 7$ agonist [³H]A-585539 (1S,4S)-2,2-dimethyl-5-(6-phenylpyridazin-3-yl)-5-aza-2-azoniabicyclo[2.2.1]heptane from rat brain membranes ($K_i \sim 23$ nM), while PNU-120596, TQS, and 5-HI did not displace either ligand under the same conditions [90]. The ability of A-867744 to displace [³H]A-585539 binding was lost in an $\alpha 7/5$ -HT₃ chimera, suggesting A-867744 may interact away from the extracellular regions of $\alpha 7$ and has the ability to induce or stabilize significant conformational changes in the $\alpha 7$ orthosteric site from a distance [90]. Further support that modulators influence protein conformation is provided in a study which compared effects of ACh and PNU-120596 on thiol modification rates at cysteine residues introduced in the inner beta sheet, transition zone, and agonist binding site [100]. Within the (non-conservative) L248T mutant background, ACh and PNU-120596 were found to produce similar changes in cysteine accessibility, suggesting that PNU-120596 may induce or stabilize conformational changes that are similar to those of the channel gating process. This study also exploited a mutant receptor, $\alpha 7$ C116A/L248T/W149C, which responded to PNU-120596 alone, to show that sulfhydryl modification at W149C greatly reduced ACh-evoked currents with little effect on PNU-120596 evoked currents, providing further support that PNU-120596 binds at a non-orthosteric site [100]. A similar study provides evidence that divalent cation potentiators also produce conformational changes via modulatory allosteric sites [101].

Putative allosteric modulator binding sites of GABA_A receptors have been identified in the N-terminal extracellular domains at subunit interfaces not forming orthosteric agonist binding sites (benzodiazepine binding site) and in the transmembrane region (barbiturate, alcohol, neurosteroid, general anesthetic binding sites [102–106]). This information, along with the differing functional profiles of type I and type II PAMs may be seen as consistent with the hypothesis that multiple allosteric sites also exist on $\alpha 7$. The site where calcium binds to potentiate responses on $\alpha 7$ was localized to the N-terminal extracellular domain [74,107]. In addition, the crystal structure of galantamine bound to acetylcholine binding protein [95], mutagenesis studies [108], and computer docking simulations [97,109] provide further evidence for nAChR allosteric sites in the extracellular domain, some of which may be related to benzodiazepine binding sites on GABA_A receptors [95,110]. Using $\alpha 7/5$ -HT₃ chimeras, the N-terminal extracellular domain of $\alpha 7$ has been shown to be sufficient for potentiation by the type I PAMs NS-1738 [111] and 5-HI [112], but not the type II PAM PNU-120596 [111]. In addition, PNU-120596, but not NS-1738, potentiated currents (ACh-evoked) in a reverse 5-HT₃/ $\alpha 7$ chimera containing $\alpha 7$ TM regions [111]. These studies also provide evidence that the extracellular $\alpha 7$ M2–M3 loop is implicated in potentiation by genistein and NS-1738 [111,112].

Other studies using $\alpha 7/5$ -HT₃ chimeras suggest that TM1–TM3 regions are critical for potentiation by PNU-120596 and the type I modulators LY-2087101 and ivermectin [113,114]. Furthermore, mutations at several $\alpha 7$ amino acid residues, which are hypothesized to contribute to an intrasubunit cavity within the four transmembrane domains, significantly reduced potentiation by PNU-120596,

LY-2087101 [113], and ivermectin [114]. Through site-directed mutagenesis, these studies together identify potentially important differences in modulator interactions with the receptors. For example, the A225D mutation reduced potentiation by PNU-120596 significantly more than it reduced potentiation by LY-2087101 [113] and the S276V mutation had no effect on potentiation by PNU-120596 [113], but conferred inhibitory properties to ivermectin [114]. Based on their work with $\alpha 7/5$ -HT₃ chimeras, mutagenesis studies, blind computer docking simulations, and evidence that an intrasubunit transmembrane site appears to be important for potentiation of glycine and GABA receptors [103,115–118], Young et al. proposed the intrasubunit cavity is a highly conserved modulatory site of Cys-loop ion channels [113], a hypothesis which has been supported by others [98,103]. The fact that CCMI was discovered from a library of GABA_A PAMs (that bind away from the benzodiazepine site), provides further support for the existence of conserved allosteric sites and/or mechanisms [67]. The findings that the extracellular domain appears to be sufficient for potentiation by some type I PAMs (NS-1738 and 5-HI), but not others (ivermectin and LY-2087101) suggest that more than one mechanism may produce the type I potentiation profile, and perhaps the same applies to the type II profile.

Kinetic models of ligand-gated ion channels and MWC allosteric theory predict that binding of agonist and channel conformational state (i.e. gating) are tightly coupled and exercise mutual influence through the principle of reciprocity [12]. Although by definition, PAMs generally have no intrinsic ability to open the ion channel, they do alter the distribution of receptors across conformational states [90,100,101,103,112], and basic allosteric theory predicts that binding of agonists at orthosteric sites facilitates binding of PAMs at allosteric sites, and vice versa [103,119]. Some experimental evidence is beginning to emerge that is consistent with this prediction. For example, the potency of the agonist PNU-282987 (N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-chlorobenzamide) to evoke calcium responses was increased with higher concentrations of PNU-120596, and, vice versa, the potency of PNU-120596 to potentiate responses was increased with higher PNU-282987 concentrations in IMR-32 cells [99]. In our own studies, we have observed that the onset of potentiation by PNU-120596 is dependent on agonist concentration for both $\alpha 7$ nAChR-expressing *Xenopus* oocytes and outside-out patches from cultured mammalian cells (manuscript in preparation). In addition, computer docking simulations predicted that binding of either PNU-120596 or LY-2087101 to the proposed TM intrasubunit cavity has lower free energy (higher affinity) in an “open channel” model than in a “closed channel” model [113].

However, 10 μ M JNJ-1930942 had no effect on the equilibrium binding of the high-affinity $\alpha 7$ agonist [³H]A-585539 to membranes from GH4C1 cells expressing human $\alpha 7$ [72]. In addition, JNJ-1930942 does not appear to significantly increase choline affinity, as measured by shifts in the IC₅₀ of MLA in the absence (~ 1 nM) or presence (~ 2.3 nM) of 5 μ M JNJ-1930942 [72]. Likewise, there were no significant effects of PNU-120596 on the binding of the $\alpha 7$ agonist ABT-107 or the antagonist MLA as revealed by the displacement of [³H]A-585539 in rat cortical homogenates [120]. Conclusions regarding changes in agonist affinity are hard to interpret in these experiments since binding is determined only at equilibrium. The degree to which antagonist affinity is affected by modulator is unknown, and the degree to which antagonists are purely competitive is unknown.

4.2. Demonstrated effects of PAMs with relevance to biological systems

Positive allosteric modulators have been demonstrated to potentiate many types of $\alpha 7$ -mediated responses that may be

important within biological systems. For example, extracellular signal-related kinase-1 and -2 (ERK1/2) phosphorylation was enhanced in PC12 cells by A-867744 [120] and PNU-120596 [120–122] with several structurally diverse agonists. Several studies have provided evidence that activation of ERK1/2, the prototypical mitogen activated kinase, is important for $\alpha 7$ -mediated protection from death in PC12 cells [121–125] and have implicated ERK1/2 in cognitive functions [125–129].

Several PAMs, including 5-HI, PNU-120596, SB-206553, and A-867744, enhance agonist-evoked $\alpha 7$ responses from hippocampal CA1 stratum radiatum interneurons [69,71,90,130,131] and glial cells [132] in acute brain slices. Spontaneous and choline- or ACh-induced increases in GABAergic inhibitory post-synaptic currents were enhanced by 5-HI [133,134], PNU-120596 [69], LY-2087101 [53], and A-867744 [90] in hippocampal neurons. JNJ-1930942 increased the amplitude of excitatory post-synaptic potentials in hippocampal dentate gyrus [72]. In cerebellar slices, 5-HI enhanced ACh-induced frequency increases of excitatory post-synaptic currents mediated by glutamate [77]. PNU-120596 (10 μ M) potentiated $\alpha 7$ induced increases in [3 H]-D-aspartate release from prefrontal cortex synaptosomes, as well as [3 H]-dopamine release from prefrontal cortex *in vitro* and *in vivo* [135]. The nicotinic facilitation of long-term potentiation was enhanced by PNU-120596 [136] and JNJ-1930942 [72] in rat dentate gyrus. Physiological concentrations of choline (~10 μ M) and 1–5 μ M PNU-120596 activated $\alpha 7$ -containing receptors in tuberomammillary neurons and hippocampal CA1 pyramidal neurons sufficiently to depolarize a cell and facilitate the firing of action potentials [137,138].

Supporting the potential therapeutic significance of these *in vitro* assays, NS-1738 [68] and CCMI [67] have been shown to enhance performance in behavioral measures of cognitive function; and CCMI [67], PNU-120596 [69,70], SB-206553 [71], A-867744 [70], and JNJ-1930942 [72] have been shown to reverse auditory gating deficits in drug-induced or DBA/2 models with systemic administration to rodents, indicating that these PAMs have sufficient pharmacokinetic properties to modulate brain $\alpha 7$ *in vivo* and that a sufficient level of endogenous agonist is present in relevant brain regions. Pharmacokinetic studies generally suggest that brain concentrations of PAMs are low compared to the EC₅₀ for potentiation determined *in vitro* [67,68,70,72]. However, since $\alpha 7$ has an intrinsically low P_{open} , even modest potentiation of $\alpha 7$ -mediated signals may be sufficient to produce significant *in vivo* effects.

A recent study has provided evidence that simultaneous modulation of $\alpha 5$ GABA_A and $\alpha 7$ nicotinic receptors may function to synergistically enhance long-term potentiation in rodent brain slices and memory-related behavior *in vivo* [139]. This study provides proof-of-concept evidence that molecules engineered to modulate multiple targets might provide an optimized approach for specific therapeutic purposes.

4.3. Limiting factors of allosteric modulation as a therapeutic approach

Despite having identified and initially characterized several $\alpha 7$ PAMs, our current knowledge of the limiting factors for PAM-based therapy and the most desirable functional characteristics of PAMs as therapeutic agents is incomplete and many questions persist unanswered. For example, are there different conditions when either a type I or type II PAM might be most advantageous? To what degree might a type II PAM disturb native temporal characteristics of a neuronal circuit? Can a PAM alter the native channel kinetics and still provide an acceptable therapeutic index? Are the most efficacious PAMs most desirable? Can $\alpha 7$ PAMs induce cytotoxicity due to high levels of channel-mediated calcium flux? What are the

limiting factors of allosteric potentiation, and under what condition is the potentiation optimized? Does the ability of a PAM to modulate responses change over significant amounts of time? Are the known $\alpha 7$ PAMs sufficiently selective to avoid undesired effects? Some of these questions have begun to be answered, but clearly, our search for understanding and identifying characteristics of the ideal $\alpha 7$ PAM continues.

If PAM therapy is going to work, obviously the endogenous agonist must be present in sufficient concentrations and able to evoke responses that the allosteric modulator can potentiate. Under conditions of degeneration in the cholinergic system, as in Alzheimer's disease, the question of whether sufficient agonist levels are present for effective modulation arises. Alternatively, under conditions of trauma, choline and possibly ACh concentrations may become unregulated [140,141] potentially resulting in over-stimulation if PAMs were present. There are some instances when stimulation of $\alpha 7$, and especially potentiation, may be undesirable. For example, activation of $\alpha 7$, particularly by nicotine, has been associated with cell proliferation, angiogenesis, and inhibition of apoptosis in some cancers, including non-small cell lung cancer and malignant pleural mesothelioma [142–144]. In addition, $\alpha 7$ has been shown to modulate aspects of immune system function, [145–147], and in immuno-compromised individuals potentiated activation of $\alpha 7$ may be more harmful than beneficial. Another cautionary perspective may come from the consideration that *in vitro* characterizations of PAMs performed at room temperature may not be entirely predictive of the behavior of the drugs *in vivo*. Despite several observations that $\alpha 7$ PAMs are effective *in vivo* [67–72] it has been suggested that the modulatory effects of NS-1738, PNU-120596, and SB-206553 may actually be reduced at physiological temperatures (John Dunlop, personal communication, also published in abstract form [148]).

Numerous articles in the literature state that type II PAMs reverse or eliminate desensitization, commonly citing references that show modulators both activate previously desensitized receptors and produce non-decaying currents in the presence of relatively high agonist throughout relatively prolonged applications [67,69,81,90]. In addition, we have used large non-decaying steady-state currents generated in *Xenopus* oocytes with 60 μ M choline and 10 μ M PNU-120596 to demonstrate channel block by agonists [93] and to study properties of antagonists [149]. Interestingly, applications of the type II PAM TQS [81] or A-867744 [90] were indeed shown to re-activate receptors that had been desensitized by a high concentration of agonist and produced a non-decaying current on the time scale of the experiments. However, the peak of the potentiated current in the presence of either TQS or A-867744 was only about 50% of the peak response recorded in the absence of PAM, indicating that only a fraction of the receptor population was actually conducting current at any given moment. These and a few other published observations, along with our own unpublished observations (manuscript in preparation) indicate the existence of desensitized states which are non-conducting in the presence of type II PAMs, meaning that at least two forms of desensitized states exist: type II modulator sensitive (D_s) and insensitive (D_i). On one hand, the existence of D_i states may limit the efficacy of PAMs to potentiate receptor-mediated responses, but on the other hand might actually provide an intrinsic safety mechanism against the dramatic potentiation produced by some modulators (see below).

The extent to which type II modulators are capable of allowing otherwise desensitized receptors to conduct current appears to depend on both agonist and modulator concentrations as well as duration of applications, with high concentrations and prolonged applications producing more D_i . When agonist concentrations are increased in the presence of a fixed PAM concentration, responses are generally enhanced over the full agonist concentration ranges,

but the magnitude of potentiation often tends to peak at intermediate concentrations and actually decrease at the higher agonist concentrations [69,72,81], meaning that in these cases, the maximal increases in efficacy do not appear to reach a true plateau. A similar phenomenon is sometimes observed when modulator concentrations are increased in the presence of a constant agonist concentration [68,71,72,77,94]. To date, published studies involving $\alpha 7$ PAMs generally have utilized protocols that make short duration applications with long interstimulus intervals, a procedure which does not strongly induce D_i states [69], and may account for the lack of articles reporting type II modulator-insensitive desensitization. The only published example to our knowledge of agonist applications made with short interstimulus intervals [67] in the continued presence of PAMs does indeed show decreased responses when several agonist pulses are delivered in succession, indicative of the induction of modulator-insensitive states. In this study, D_i states appeared to accumulate in the presence of the type I PAM CCMI, as well as the type II PAM PNU-120596 [67]. Even though a 7 s application is not particularly long in duration, Dinklo et al. (2010) provide an example of a slowly decaying current despite the continued presence of agonist and modulator, indicating a conversion of receptors to JNJ-1930942-resistant states [72]. When 60 μM ACh was applied every 3 min on top of 60 μM choline and 10 μM PNU-120596 in *Xenopus* oocytes expressing $\alpha 7$, only the initial ACh response was greatly potentiated (~ 130 -fold), with much less potentiation observed with subsequent ACh applications (< 20 -fold), relative to the response to 60 μM ACh alone. After several minutes of no 60 μM ACh stimulation (application of choline and PNU-120596 continued), subsequent responses to 60 μM ACh (immediately after choline and PNU-120596 application stopped) were again greatly potentiated (~ 80 -fold), indicating receptor recovery from the D_i states induced by ACh, choline, and PNU-120596 stimulation [93].

Additional evidence of D_i states is found in a recent study measuring calcium currents evoked by nicotine, PNU-282987, and PNU-120596 in bovine chromaffin cells [150]. In this study, the authors were only able to detect an α -bungarotoxin (α -btX)-sensitive and PNU-120596-dependent increase in calcium response when low (1 μM) concentrations of nicotine were used to stimulate the cells. At concentrations of ≥ 3 μM nicotine, neither PNU-120596 nor α -btX had an effect on the response, indicating that the $\alpha 7$ component was lost by the higher nicotine concentrations [150]. To isolate and more closely examine the disappearing $\alpha 7$ component, varying concentrations of the $\alpha 7$ agonist PNU-282987 with 1 μM PNU-120596 were applied for 5 min to evoke a calcium response. Any fluorescence increases that were insensitive to blockade by α -btX were subtracted from the total response. Interestingly, as stimulating agonist (in this case PNU-282987) concentrations increased, the $\alpha 7$ -dependent calcium responses decreased, indicating greater D_i as agonist concentrations increased in the presence of a fixed concentration of PNU-120596 [150].

Two modes of potentiation, analogous to the two modes of activation proposed for $\alpha 7$ [27], may exist. For example, relatively low occupancy of agonist and PAM binding sites may generate an equilibrium condition between open, D_s , and D_i states that allows for large prolonged steady-state currents, whereas higher occupancy of agonist and PAM binding sites promote D_i states, which ultimately result in biphasic and reduced steady-state currents. The potentiation of $\alpha 7$ responses likely depends on a complex interplay of many factors, including the effective agonist and modulator concentrations, nature of agonist/modulator combinations, and duration of applications [67,81,131].

Activation of $\alpha 7$ can have either protective or toxic effects depending on the mode of stimulation [151]. Positive allosteric modulators have been identified which increase the open

probability of $\alpha 7$ by several orders of magnitude, inviting the question of whether PAMs may take the activation of this receptor subtype, which has high permeability to the natural catalytic ion calcium, to dangerously high levels. Calcium-mediated toxicity has been reported in SH-SY5Y cells and mice expressing mutant forms of $\alpha 7$ that display dramatically prolonged responses after stimulation [152–154]. *In vivo* toxicity profiles for PAMs are lacking, but there are some *in vitro* experiments that suggest type I and type II PAMs may have different profiles. Ng et al. showed that 24-h exposure to the type I PAM CCMI did not reduce the viability of SH-SY5Y cells, whereas the type II PAM PNU-120596 did under the same conditions and in a MLA-sensitive manner [67]. Prior to the toxicity assays the authors tested the cells to verify the presence of functional receptors and successful modulation by PNU-120596 and CCMI. A similar experiment with stable $\alpha 7$ -expressing GH4C1 cells showed that the type II PAM PNU-120596 was toxic, while JNJ-1930942 was not toxic in the same paradigm [72]. However, Hu et al. failed to detect toxic effects in undifferentiated PC12 cells and cortical neurons after treatment with the type I PAM CCMI, but they also did not detect any toxic effects with the type II PAMs PNU-120596 or A-867744 [155]. The reason for the contradictory results is unclear, but two shortcomings of this study are that functional $\alpha 7$ expression and potentiation were indirectly evaluated, and only one agonist, PNU-282987, was tested at only one concentration, 10 μM . The result of this study, in light of the existence of D_i states discussed above, invites consideration of the interesting possibility that D_i states provide an intrinsic safety mechanism that shuts the channel down with strong activation. In IMR-32 cells, PNU-282987 was reported as the most potent of several agonists tested in the presence of 10 μM PNU-120596 to induce calcium responses, with an EC_{50} of 0.1 ± 0.1 μM [99]. In *Xenopus* oocytes, the EC_{50} of PNU-282987 was reported as approximately 200 nM in the presence of 3 μM PNU [81]. Given that 24-h applications of 10 μM PNU-282987 and PAM is expected to be a strong stimulus, the possibility that sufficient amounts of D_i were induced to protect cells from toxicity seems plausible. In addition, del Barrio et al. demonstrated that D_i states are induced with responses evoked by PNU-282987 in the presence of 1 μM PNU-120596 in adrenal chromaffin cells, which are related to PC12 cells [150]. In contrast, the toxicity studies of both Ng et al. [67] and Dinklo et al. [72] used the weakly potent agonist choline at concentrations of ~ 100 μM .

In many cases, non-potentiated $\alpha 7$ -mediated responses that are below or at the limits of experimental detection become easily measurable with the application of type II PAMs. For example, increases in ERK1/2 phosphorylation in PC12 cells that were evoked by agonists were detectable only in the presence of PNU-120596 [120–122,155]. Similarly, agonist-evoked calcium responses were undetectable in PC12 cells [122,155], IMR-32 cells [70,90,99,120,156], primary cortical neurons [99], chromaffin cells [150], and CHO-K1 cells [157] unless PAMs were co-applied. In these experiments, type I PAMs were generally unable to increase either ERK1/2 phosphorylation or calcium responses to detectable levels [99,155,157], however, calcium responses were significantly potentiated with 5-HI to detectable levels in GH4C1 cells stably expressing $\alpha 7$ [158]. In addition, in hippocampal CA1 pyramidal cells, PNU-120596 was shown to enhance $\alpha 7$ -mediated responses evoked by physiologically relevant concentrations of choline, which are difficult to detect under normal conditions, to the point of depolarizing the cells and facilitating action potential firing [138]. Together these observations raise questions of whether PAMs (primarily type II) may alter the $\alpha 7$ component of neuronal systems sufficiently to interrupt natural synchrony and regulation of important signaling events. These results also bring to light technical challenges in high-throughput screening methods to identify type I PAMs: potentiated responses may still

fall below experimental detection limits, and the sampling rate of data collection in commonly used assays is too slow to detect the rise and decay of an $\alpha 7$ response potentiated by a type I PAM.

5. Summary

The recent programs of drug development to discover and characterize nAChR PAMs, particularly ones which are selective for $\alpha 7$ nAChR, have created both new opportunities and challenges. We are at a crucial point in the testing of PAMs with unique attributes, and we must identify the correct model systems to determine whether they should be moved forward into clinical trials. We must also consider whether they will be safe from serious side effects, as they will reach into all of the many areas of $\alpha 7$ nAChR activity. An additional consideration is that evidence would suggest that PAMs would work primarily, if not exclusively, through increasing signal transduction mediated by $\alpha 7$ ion channel activation. Some studies have reported PAM enhancement of intracellular signal transduction pathways [120–122], and the tacit assumption is that in these cases, a PAM-sensitive, channel-mediated calcium influx is the essential connection between receptor activation and signal transduction. However, there are numerous cases, especially in non-neuronal cells, where there is important $\alpha 7$ -mediated signal transduction under conditions when no ion channel activation can be detected [146,159]. Likewise, we have learned from the development of $\alpha 7$ agonists and other drugs that the most efficacious compounds are not always the best leads to follow. DMXBA (GTS-21), one of the first $\alpha 7$ -selective drugs to be identified [160], has been shown to have good activity in many model systems [161], including human trials [162], and yet it has very low efficacy as an ion channel activator [163,164] and strongly stabilizes receptor desensitization [93]. Considering the extremely low P_{open} of $\alpha 7$ nAChR, especially in the presence of high concentrations of agonists, the common condition for these receptors is for them to be in non-conducting states. Experiments with PNU-120596 have shown that $\alpha 7$ nAChR have multiple non-conducting states that are invisibly and dynamically interconvertible. It has been shown that many intracellular proteins interact with $\alpha 7$ receptors [165], and it seems reasonable to hypothesize that the dynamics of the conformational changes associated with conversions among multiple non-conducting states may extend into the largely mysterious intracellular domain and thereby have channel-independent signal transducing effects.

The identification of $\alpha 7$ PAMs has opened up new avenues for therapeutic approaches, and taught us much about the mysteries of these important receptors. They also show that there are likely to be more mysteries to uncover and understand about $\alpha 7$ nAChR.

Acknowledgements

This work was supported by the National Institutes of Health grants, R01 GM057481 and T32-AG000196. We thank Dr. Nicole A. Horenstein for many helpful comments and Clare Stokes for editorial assistance.

References

- [1] Gotti C, Clementi F, Fornari A, Gaimarri A, Guiducci S, Manfredi I, et al. Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem Pharmacol* 2009;78:703–11.
- [2] Li P, Steinbach JH. The neuronal nicotinic $\alpha 4\beta 2$ receptor has a high maximal probability of being open. *Br J Pharm* 2010;160:1906–15.
- [3] Uteshev VV, Meyer EM, Papke RL. Activation and inhibition of native neuronal $\alpha 4\beta 2$ nicotinic ACh receptors. *Brain Res* 2002;948:33–46.
- [4] Picciotto MR, Addy NA, Mineur YS, Brunzell DH. It is not “either/or”: activation and desensitization of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and mood. *Prog Neurobiol* 2008;84:329–42.
- [5] Papke RL, Trocme-Thibierge C, Guendisch D, Abbas Al Rubaiy SA, Bloom SA. Electrophysiological perspectives on the therapeutic use of nicotinic acetylcholine receptor partial agonists. *J Pharmacol Exp Ther* 2011;337:367–79.
- [6] Ortells MO, Lunt GG. Evolutionary history of the ligand-gated ion-channel superfamily of receptors. *Trends Neurosci* 1995;18:121–7.
- [7] Monod J, Jacob F. Teleonomic mechanisms in cellular metabolism, growth, and differentiation. *Cold Spring Harb Symp Quant Biol* 1961;26:389–401.
- [8] Monod J, Changeux JP, Jacob F. Allosteric proteins and cellular control systems. *J Mol Biol* 1963;6:306–29.
- [9] Monod J, Wyman J, Changeux JP. On the nature of allosteric transitions: a plausible model. *J Mol Biol* 1965;12:88–118.
- [10] Changeux JP, Edelstein SJ. Allosteric receptors after 30 years. *Neuron* 1998;21:959–80.
- [11] Changeux JP, Edelstein SJ. Allosteric mechanisms of signal transduction. *Science* 2005;308:1424–8.
- [12] Colquhoun D. Binding gating, affinity and efficacy: the interpretation of structure-activity relationships for agonists and of the effects of mutating receptors. *Br J Pharmacol* 1998;125:924–47.
- [13] Campo-Soria C, Chang Y, Weiss DS. Mechanism of action of benzodiazepines on GABAA receptors. *Br J Pharmacol* 2006;148:984–90.
- [14] Gill JK, Savolainen M, Young GT, Zwart R, Sher E, Millar NS. Agonist activation of $\alpha 7$ nicotinic acetylcholine receptors via an allosteric transmembrane site. *Proc Natl Acad Sci USA* 2011;108:5867–72.
- [15] Papke RL, Boulter J, Patrick J, Heinemann S. Single-channel currents of rat neuronal nicotinic acetylcholine receptors expressed in *Xenopus laevis* oocytes. *Neuron* 1989;3:589–96.
- [16] Papke RL, Millhauser G, Lieberman Z, Oswald RE. Relationships of agonist properties to the single channel kinetics of nicotinic acetylcholine receptors. *Biophys J* 1988;53:1–10.
- [17] Colquhoun D, Sakmann B. Fast events in single-channel currents activated by acetylcholine and its analogues at the frog muscle end-plate. *J Physiol (Lond)* 1985;369:501–57.
- [18] Auerbach A. The gating isomerization of neuromuscular acetylcholine receptors. *J Physiol* 2010;588:573–86.
- [19] Lape R, Colquhoun D, Sivilotti LG. On the nature of partial agonism in the nicotinic receptor superfamily. *Nature* 2008;454:722–7.
- [20] Wang J, Horenstein NA, Stokes C, Papke RL. Tethered agonist analogs as site-specific probes for domains of the human $\alpha 7$ nicotinic acetylcholine receptor that differentially regulate activation and desensitization. *Mol Pharmacol* 2010;78:1012–25.
- [21] Williams DK, Stokes C, Horenstein NA, Papke RL. The effective opening of nicotinic acetylcholine receptors with single agonist binding sites. *J Gen Physiol* 2011;137:369–84.
- [22] Colquhoun D, Ogden DC. Activation of ion channels in the frog end-plate by high concentrations of acetylcholine. *J Physiol (Lond)* 1988;395:131–59.
- [23] Wyman RE, Rogers CJ, Macdonald RL. Differential regulation of gamma-aminobutyric acid receptor channels by diazepam and phenobarbital. *Ann Neurol* 1989;25:213–20.
- [24] McNeerney ME, Pardi D, Pugh PC, Nai Q, Margiotta JF. Expression and channel properties of $\alpha 4\beta 2$ nicotinic acetylcholine receptors on chick ciliary and choroid neurons. *J Neurophysiol* 2000;84:1314–29.
- [25] Charpentier E, Wiesner A, Huh KH, Ogier R, Hoda JC, Allaman G, et al. $\alpha 7$ neuronal nicotinic acetylcholine receptors are negatively regulated by tyrosine phosphorylation and Src-family kinases. *J Neurosci* 2005;25:9836–49.
- [26] Palma E, Bertrand S, Binzoni T, Bertrand D. Neuronal nicotinic $\alpha 7$ receptor expressed in *Xenopus* oocytes presents five putative binding sites for methyllycaconitine. *J Physiol* 1996;491:151–61.
- [27] Papke RL, Meyer E, Nutter T, Uteshev VV. $\alpha 7$ -selective agonists and modes of $\alpha 7$ receptor activation. *Eur J Pharmacol* 2000;393:179–95.
- [28] Mike A, Castro NG, Albuquerque EX. Choline and acetylcholine have similar kinetic properties of activation and desensitization on the $\alpha 7$ nicotinic receptors in rat hippocampal neurons. *Brain Res* 2000;882:155–68.
- [29] Mulle C, Léna C, Changeux JP. Potentiation of nicotinic receptor response by external calcium in rat central neurons. *Neuron* 1992;8:937–45.
- [30] Vernino S, Amador M, Luetje CW, Patrick J, Dani JA. Calcium modulation and high calcium permeability of neuronal nicotinic acetylcholine receptors. *Neuron* 1992;8:127–34.
- [31] Amador M, Dani JA. Mechanism for modulation of nicotinic acetylcholine receptors that can influence synaptic transmission. *J Neurosci* 1995;15:4525–32.
- [32] Hsiao B, Dweck D, Luetje CW. Subunit-dependent modulation of neuronal nicotinic receptors by zinc. *J Neurosci* 2001;21:1848–56.
- [33] Palma E, Maggi L, Miledi R, Eusebi F. Effects of Zn^{2+} on wild and mutant neuronal $\alpha 7$ nicotinic receptors. *Proc Natl Acad Sci USA* 1998;95:10246–50.
- [34] Hsiao B, Mihalak KB, Repicky SE, Everhart D, Mederos AH, Malhotra A, et al. Determinants of zinc potentiation on the $\alpha 4\beta 2$ subunit of neuronal nicotinic receptors. *Mol Pharmacol* 2006;69:27–36.
- [35] Zwart R, Van Kleef RG, Milikan JM, Oortgiesen M, Vijverberg HP. Potentiation and inhibition of subtypes of neuronal nicotinic acetylcholine receptors by Pb^{2+} . *Eur J Pharmacol* 1995;291:399–406.
- [36] Bertrand D, Valera S, Bertrand S, Ballivet M, Rungger D. Steroids inhibit nicotinic acetylcholine receptors. *Neuroreport* 1991;2:277–80.
- [37] Valera S, Ballivet M, Bertrand D. Progesterone modulates a neuronal nicotinic acetylcholine receptor. *Proc Natl Acad Sci USA* 1992;89:9949–53.
- [38] Ke L, Lukas RJ. Effects of steroid exposure on ligand binding and functional activities of diverse nicotinic acetylcholine receptor subtypes. *J Neurochem* 1996;67:1100–12.

- [39] Paradiso K, Sabey K, Evers AS, Zorumski CF, Covey DF, Steinbach JH. Steroid inhibition of rat neuronal nicotinic $\alpha 4 \beta 2$ receptors expressed in HEK 293 cells. *Mol Pharmacol* 2000;58:341–51.
- [40] Paradiso K, Zhang J, Steinbach JH. The C terminus of the human nicotinic $\alpha 4 \beta 2$ receptor forms a binding site required for potentiation by an estrogenic steroid. *J Neurosci* 2001;21:6561–8.
- [41] Curtis L, Buisson B, Bertrand S, Bertrand D. Potentiation of human $\alpha 4 \beta 2$ neuronal nicotinic acetylcholine receptor by estradiol. *Mol Pharmacol* 2002;61:127–35.
- [42] Miwa JM, Ibanez-Tallon I, Crabtree GW, Sanchez R, Sali A, Role LW, et al. lynx1 , an endogenous toxin-like modulator of nicotinic acetylcholine receptors in the mammalian CNS. *Neuron* 1999;23:105–14.
- [43] Ibanez-Tallon I, Miwa JM, Wang HL, Adams NC, Crabtree GW, Sine SM, et al. Novel modulation of neuronal nicotinic acetylcholine receptors by association with the endogenous prototoxin lynx1 . *Neuron* 2002;33:893–903.
- [44] Miwa JM, Freedman R, Lester HA. Neural systems governed by nicotinic acetylcholine receptors: emerging hypotheses. *Neuron* 2011;70:20–33.
- [45] Pereira EF, Hilmas C, Santos MD, Alkondon M, Maelicke A, Albuquerque EX. Unconventional ligands and modulators of nicotinic receptors. *J Neurobiol* 2002;53:479–500.
- [46] Levandoski MM, Piket B, Chang J. The anthelmintic levamisole is an allosteric modulator of human neuronal nicotinic acetylcholine receptors. *Eur J Pharmacol* 2003;471:9–20.
- [47] Wu TY, Smith CM, Sine SM, Levandoski MM. Morantel allosterically enhances channel gating of neuronal nicotinic acetylcholine $\alpha 3 \beta 2$ receptors. *Mol Pharmacol* 2008;74:466–75.
- [48] Seo S, Henry JT, Lewis AH, Wang N, Levandoski MM. The positive allosteric modulator morantel binds at noncanonical subunit interfaces of neuronal nicotinic acetylcholine receptors. *J Neurosci* 2009;29:8734–42.
- [49] Samochocki M, Höffle A, Fehrenbacher A, Jostock R, Ludwig J, Christner C, et al. Galantamine is an allosterically potentiating ligand of neuronal nicotinic but not of muscarinic acetylcholine receptors. *J Pharmacol Exp Ther* 2003;305:1024–36.
- [50] Kuryatov A, Berrettini W, Lindstrom J. Acetylcholine receptor (AChR) $\alpha 5$ subunit variant associated with risk for nicotine dependence and lung cancer reduces ($\alpha 4 \beta 2$) $\alpha 5$ AChR function. *Mol Pharmacol* 2011;79:119–25.
- [51] Schrattenholz A, Pereira EF, Roth U, Weber KH, Albuquerque EX, Maelicke A. Agonist responses of neuronal nicotinic acetylcholine receptors are potentiated by a novel class of allosterically acting ligands. *Mol Pharmacol* 1996;49:1–6.
- [52] Broad LM, Zwart R, Pearson KH, Lee M, Wallace L, McPhie GI, et al. Identification and pharmacological profile of a new class of selective nicotinic acetylcholine receptor potentiators. *J Pharmacol Exp Ther* 2006;318:1108–17.
- [53] de Filippi G, Mogg AJ, Phillips KG, Zwart R, Sher E, Chen Y. The subtype-selective nicotinic acetylcholine receptor positive allosteric potentiator 2087101 differentially facilitates neurotransmission in the brain. *Eur J Pharmacol* 2010;643:218–24.
- [54] Weltzin MM, Schulte MK. Pharmacological characterization of the allosteric modulator desformylflustrabromine and its interaction with $\alpha 4 \beta 2$ neuronal nicotinic acetylcholine receptor orthosteric ligands. *J Pharmacol Exp Ther* 2010;334:917–26.
- [55] Kim JS, Padnya A, Weltzin M, Edmonds BW, Schulte MK, Glennon RA. Synthesis of desformylflustrabromine and its evaluation as an $\alpha 4 \beta 2$ and $\alpha 7$ nACh receptor modulator. *Bioorg Med Chem Lett* 2007;17:4855–60.
- [56] Sala F, Mulet J, Reddy KP, Bernal JA, Wikman P, Valor LM, et al. Potentiation of human $\alpha 4 \beta 2$ neuronal nicotinic receptors by a *Flustra foliacea* metabolite. *Neurosci Lett* 2005;373:144–9.
- [57] Springer SK, Woodin KS, Berry V, Boezio AA, Cao L, Clarkin K, et al. Synthesis and activity of substituted carbamates as potentiators of the $\alpha 4 \beta 2$ nicotinic acetylcholine receptor. *Bioorg Med Chem Lett* 2008;18:5643–7.
- [58] Albrecht BK, Berry V, Boezio AA, Cao L, Clarkin K, Guo W, et al. Discovery and optimization of substituted piperidines as potent, selective, CNS-penetrant $\alpha 4 \beta 2$ nicotinic acetylcholine receptor potentiators. *Bioorg Med Chem Lett* 2008;18:5209–12.
- [59] Haydar SN, Dunlop J. Neuronal nicotinic acetylcholine receptors – targets for the development of drugs to treat cognitive impairment associated with schizophrenia and Alzheimer's disease. *Curr Top Med Chem* 2010;10:144–52.
- [60] Thomsen MS, Hansen HH, Timmerman DB, Mikkelsen JD. Cognitive improvement by activation of $\alpha 7$ nicotinic acetylcholine receptors: from animal models to human pathophysiology. *Curr Pharm Des* 2010;16:323–43.
- [61] Levin ED, McClernon FJ, Rezvani AH. Nicotinic effects on cognitive function: behavioral characterization, pharmacological specification, and anatomic localization. *Psychopharmacology (Berl)* 2006;184:523–39.
- [62] Freedman R, Leonard S, Gault JM, Hopkins J, Cloninger CR, Kaufmann CA, et al. Linkage disequilibrium for schizophrenia at the chromosome 15q13–14 locus of the $\alpha 7$ -nicotinic acetylcholine receptor subunit gene (CHRNA7). *Am J Med Genet* 2001;105:20–2.
- [63] Hajos M, Rogers BN. Targeting $\alpha 7$ nicotinic acetylcholine receptors in the treatment of schizophrenia. *Curr Pharm Des* 2010;16:538–54.
- [64] Stevens KE, Kem WR, Mahnir VM, Freedman R. Selective $\alpha 7$ nicotinic agonists normalize inhibition of auditory response in DBA mice. *Psychopharmacology* 1998;136:320–7.
- [65] Simosky JK, Stevens KE, Kem WR, Freedman R. Intragastric DMXB-A, an $\alpha 7$ nicotinic agonist, improves deficient sensory inhibition in DBA/2 mice. *Biol Psychiatry* 2001;50:493–500.
- [66] Olincy A, Harris JG, Johnson LL, Pender V, Kongs S, Allensworth D, et al. Proof-of-concept trial of an $\alpha 7$ nicotinic agonist in schizophrenia. *Arch Gen Psychiatry* 2006;63:630–8.
- [67] Ng HJ, Whittemore ER, Tran MB, Hogenkamp DJ, Broide RS, Johnstone TB, et al. Nootropic $\alpha 7$ nicotinic receptor allosteric modulator derived from GABAA receptor modulators. *Proc Natl Acad Sci USA* 2007;104:8059–64.
- [68] Timmermann DB, Gronlien JH, Kohlhaas KL, Nielsen EO, Dam E, Dyhring T, et al. An allosteric modulator of the ($\alpha 7$) nicotinic acetylcholine receptor possessing cognition enhancing properties in vivo. *J Pharmacol Exp Ther* 2007;323:294–307.
- [69] Hurst RS, Hajos M, Raggenbass M, Wall TM, Higdon NR, Lawson JA, et al. A novel positive allosteric modulator of the $\alpha 7$ neuronal nicotinic acetylcholine receptor: in vitro and in vivo characterization. *J Neurosci* 2005;25:4396–405.
- [70] Faghieh R, Gopalakrishnan SM, Gronlien JH, Malysz J, Briggs CA, Wetterstrand C, et al. Discovery of 4-(5-(4-chlorophenyl)-2-methyl-3-propionyl-1H-pyrrol-1-yl)benzenesulfonamide (A-867744) as a novel positive allosteric modulator of the $\alpha 7$ nicotinic acetylcholine receptor. *J Med Chem* 2009;52:3377–84.
- [71] Dunlop J, Lock T, Jow B, Sitzia F, Grauer S, Jow F, et al. Old and new pharmacology: positive allosteric modulation of the $\alpha 7$ nicotinic acetylcholine receptor by the 5-hydroxytryptamine(2B/C) receptor antagonist SB-206553 (3,5-dihydro-5-methyl-N-3-pyridinylbenzo[1,2-b:4,5-b']dipyrrole-1(2H)-carboxamide). *J Pharmacol Exp Ther* 2009;328:766–76.
- [72] Dinklo T, Shaban H, Thuring JW, Lavreysen H, Stevens KE, Zheng L, et al. Characterization of 2-[[4-fluoro-3-(trifluoromethyl)phenyl]amino]-4-(4-pyridinyl)-5-thiazolemethanol (JNJ-1930942), a novel positive allosteric modulator of the ($\alpha 7$) nicotinic acetylcholine receptor. *J Pharmacol Exp Ther* 2011;336:560–74.
- [73] Briggs CA, Gronlien JH, Curzon P, Timmermann DB, Ween H, Thorin-Hagene K, et al. Role of channel activation in cognitive enhancement mediated by $\alpha 7$ nicotinic acetylcholine receptors. *Br J Pharmacol* 2009;158:1486–94.
- [74] Galzi JL, Bertrand S, Corringer PJ, Changeux JP, Bertrand D. Identification of calcium binding sites that regulate potentiation of a neuronal nicotinic acetylcholine receptor. *EMBO J* 1996;15:5824–32.
- [75] Krause RM, Buisson B, Bertrand S, Corringer PJ, Galzi JL, Changeux JP, et al. Ivermectin: a positive allosteric effector of the $\alpha 7$ neuronal nicotinic acetylcholine receptor. *Mol Pharmacol* 1998;53:283–94.
- [76] Gurley D, Harris EW, Li C, Johnson EC, Lanthorn T. 5-hydroxyindole potentiates the nicotinic acetylcholine receptor $\alpha 7$ subtype. *Soc Neurosci Abs* 2000;716:15.
- [77] Zwart R, De Filippi G, Broad LM, McPhie GI, Pearson KH, Baldwinson T, et al. 5-Hydroxyindole potentiates human $\alpha 7$ nicotinic receptor-mediated responses and enhances acetylcholine-induced glutamate release in cerebellar slices. *Neuropharm* 2002;43:374–84.
- [78] Lopes C, Pereira EF, Wu HQ, Purushottamachar P, Njar V, Schwarcz R, et al. Competitive antagonism between the nicotinic allosteric potentiating ligand galantamine and kynurenic acid at $\alpha 7$ nicotinic receptors. *J Pharmacol Exp Ther* 2007;322:48–58.
- [79] Conroy WG, Liu QS, Nai Q, Margiotta JF, Berg DK. Potentiation of $\alpha 7$ -containing nicotinic acetylcholine receptors by select albumins. *Mol Pharmacol* 2003;63:419–28.
- [80] Chimienti F, Hogg RC, Plantard L, Lehmann C, Brackh N, Fischer J, et al. Identification of SLURP-1 as an epidermal neuromodulator explains the clinical phenotype of Mal de Meleda. *Hum Mol Genet* 2003;12:3017–24.
- [81] Gronlien JH, Haakerud M, Ween H, Thorin-Hagene K, Briggs CA, Gopalakrishnan M, et al. Distinct profiles of ($\alpha 7$) nAChR positive allosteric modulation revealed by structurally diverse chemotypes. *Mol Pharmacol* 2007;72:715–24.
- [82] Cho CH, Song W, Leitzell K, Teo E, Meleth AD, Quick MW, et al. Rapid upregulation of $\alpha 7$ nicotinic acetylcholine receptors by tyrosine dephosphorylation. *J Neurosci* 2005;25:3712–23.
- [83] Sigel E, Baur R. Effect of ivermectin B1a on chick neuronal gamma-aminobutyrate receptor channels expressed in *Xenopus* oocytes. *Mol Pharmacol* 1987;32:749–52.
- [84] Krusek J, Zemkova H. Effect of ivermectin on gamma-aminobutyric acid-induced chloride currents in mouse hippocampal embryonic neurones. *Eur J Pharmacol* 1994;259:121–8.
- [85] Shan Q, Haddrill JL, Lynch JW. Ivermectin, an unconventional agonist of the glycine receptor chloride channel. *J Biol Chem* 2001;276:12556–64.
- [86] Khakh BS, Proctor WR, Dunwiddie TV, Labarca C, Lester HA. Allosteric control of gating and kinetics at P2X(4) receptor channels. *J Neurosci* 1999;19:7289–99.
- [87] Moriguchi S, Marszalec W, Zhao X, Yeh JZ, Narahashi T. Mechanism of action of galantamine on N-methyl-D-aspartate receptors in rat cortical neurons. *J Pharmacol Exp Ther* 2004;310:933–42.
- [88] van Hooft JA, van der Haar E, Vijverberg HP. Allosteric potentiation of the 5-HT3 receptor-mediated ion current in N1E-115 neuroblastoma cells by 5-hydroxyindole and analogues. *Neuropharmacology* 1997;36:649–53.
- [89] Friis S, Mathes C, Sunesen M, Bowlby MR, Dunlop J. Characterization of compounds on nicotinic acetylcholine receptor $\alpha 7$ channels using higher throughput electrophysiology. *J Neurosci Methods* 2009;177:142–8.
- [90] Malysz J, Gronlien JH, Anderson DJ, Hakerud M, Thorin-Hagene K, Ween H, et al. In vitro pharmacological characterization of a novel allosteric modula-

- tor of alpha 7 neuronal acetylcholine receptor, 4-(5-(4-chlorophenyl)-2-methyl-3-propionyl-1H-pyrrol-1-yl)benzenesulfonamide (A-867744), exhibiting unique pharmacological profile. *J Pharmacol Exp Ther* 2009;330:257–67.
- [91] Arias HR, Gu RX, Feuerbach D, Guo BB, Ye Y, Wei DQ. Novel positive allosteric modulators of the human alpha7 nicotinic acetylcholine receptor. *Biochemistry* 2011.
 - [92] Faghih R, Gopalakrishnan M, Briggs CA. Allosteric modulators of the alpha7 nicotinic acetylcholine receptor. *J Med Chem* 2008;51:701–12.
 - [93] Papke RL, Kem WR, Soti F, López-Hernández GY, Horenstein NA. Activation and desensitization of nicotinic alpha7-type acetylcholine receptors by benzyldiene anabaseines and nicotine. *J Pharmacol Exp Ther* 2009;329:791–807.
 - [94] Malysz J, Gronlien JH, Timmermann DB, Hakerud M, Thorin-Hagene K, Ween H, et al. Evaluation of alpha7 nicotinic acetylcholine receptor agonists and positive allosteric modulators using the parallel oocyte electrophysiology test station. *Assay Drug Dev Technol* 2009;7:374–90.
 - [95] Hansen SB, Taylor P. Galanthamine and non-competitive inhibitor binding to ACh-binding protein: evidence for a binding site on non-alpha-subunit interfaces of heteromeric neuronal nicotinic receptors. *J Mol Biol* 2007;369:895–901.
 - [96] Capelli AM, Castelletti L, Salvagno C, Oliosi B, Di Lenarda E, Virginio C, et al. Identification of novel alpha7 nAChR positive allosteric modulators with the use of pharmacophore in silico screening methods. *Bioorg Med Chem Lett* 2010;20:4561–5.
 - [97] Dey R, Chen L. In search of allosteric modulators of alpha7-nAChR by solvent density guided virtual screening. *J Biomol Struct Dyn* 2011;28:695–715.
 - [98] Sattelle DB, Buckingham SD, Akamatsu M, Matsuda K, Pienaar IS, Jones AK, et al. Comparative pharmacology and computational modelling yield insights into allosteric modulation of human alpha7 nicotinic acetylcholine receptors. *Biochem Pharmacol* 2009;78:836–43.
 - [99] Gopalakrishnan SM, Philip BM, Gronlien JH, Malysz J, Anderson DJ, Gopalakrishnan M, et al. Functional characterization and high-throughput screening of positive allosteric modulators of alpha7 nicotinic acetylcholine receptors in IMR-32 neuroblastoma cells. *Assay Drug Dev Technol* 2011.
 - [100] Barron SC, McLaughlin JT, See JA, Richards VL, Rosenberg RL. An allosteric modulator of alpha7 nicotinic receptors, N-(5-Chloro-2,4-dimethoxyphenyl)-N'-(5-methyl-3-isoxazolyl)-urea (PNU-120596), causes conformational changes in the extracellular ligand binding domain similar to those caused by acetylcholine. *Mol Pharmacol* 2009;76:253–63.
 - [101] McLaughlin JT, Barron SC, See JA, Rosenberg RL. Conformational changes in alpha7 acetylcholine receptors underlying allosteric modulation by divalent cations. *BMC Pharmacol* 2009;9:1.
 - [102] Olsen RW, Chang CS, Li G, Hancher HJ, Wallner M. Fishing for allosteric sites on GABA(A) receptors. *Biochem Pharmacol* 2004;68:1675–84.
 - [103] Chang Y, Huang Y, Whiteaker P. Mechanism of allosteric modulation of the Cys-loop receptors. *Pharmaceuticals* 2010;3:2592–609.
 - [104] Forman SA, Miller KW. Anesthetic sites and allosteric mechanisms of action on Cys-loop ligand-gated ion channels. *Can J Anaesth* 2011;58:191–205.
 - [105] Baenziger JE, Corringer PJ. 3D structure and allosteric modulation of the transmembrane domain of pentameric ligand-gated ion channels. *Neuropharmacology* 2011;60:116–25.
 - [106] Nury H, Van Renterghem C, Weng Y, Tran A, Baaden M, Dufresne V, et al. X-ray structures of general anaesthetics bound to a pentameric ligand-gated ion channel. *Nature* 2011;469:428–31.
 - [107] Eisele JL, Bertrand S, Galzi JL, Devillers-Thierry A, Changeux JP, Bertrand D. Chimeric nicotinic-serotonergic receptor combines distinct ligand binding and channel specificities. *Nature* 1993;366:479–83.
 - [108] Ludwig J, Hoffle-Maas A, Samochocki M, Luttmann E, Albuquerque EX, Fels G, et al. Localization by site-directed mutagenesis of a galantamine binding site on alpha7 nicotinic acetylcholine receptor extracellular domain. *J Recept Signal Transduct Res* 2010;30:469–83.
 - [109] Luttmann E, Ludwig J, Hoffle-Maas A, Samochocki M, Maelicke A, Fels G. Structural model for the binding sites of allosterically potentiating ligands on nicotinic acetylcholine receptors. *ChemMedChem* 2009;4:1874–82.
 - [110] Ernst M, Brauchart D, Boresch S, Sieghart W. Comparative modeling of GABA(A) receptors: limits, insights, future developments. *Neuroscience* 2003;119:933–43.
 - [111] Bertrand D, Bertrand S, Cassar S, Gubbins E, Li J, Gopalakrishnan M. Positive allosteric modulation of the alpha7 nicotinic acetylcholine receptor: ligand interactions with distinct binding sites and evidence for a prominent role of the M2-M3 segment. *Mol Pharmacol* 2008;74:1407–16.
 - [112] Gronlien JH, Ween H, Thorin-Hagene K, Cassar S, Li J, Briggs CA, et al. Importance of M2-M3 loop in governing properties of genistein at the alpha7 nicotinic acetylcholine receptor inferred from alpha7/5-HT3A chimera. *Eur J Pharmacol* 2010;647:37–47.
 - [113] Young GT, Zwart R, Walker AS, Sher E, Millar NS. Potentiation of alpha7 nicotinic acetylcholine receptors via an allosteric transmembrane site. *Proc Natl Acad Sci USA* 2008;105:14686–91.
 - [114] Collins T, Millar NS. Nicotinic acetylcholine receptor transmembrane mutations convert ivermectin from a positive to a negative allosteric modulator. *Mol Pharmacol* 2010;78:198–204.
 - [115] Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, et al. Sites of alcohol and volatile anaesthetic action on GABA(A) and glycine receptors. *Nature* 1997;389:385–9.
 - [116] Hosie AM, Wilkins ME, da Silva HM, Smart TG. Endogenous neurosteroids regulate GABAA receptors through two discrete transmembrane sites. *Nature* 2006;444:486–9.
 - [117] Ye Q, Koltchine VV, Mihic SJ, Mascia MP, Wick MJ, Finn SE, et al. Enhancement of glycine receptor function by ethanol is inversely correlated with molecular volume at position alpha267. *J Biol Chem* 1998;273:3314–9.
 - [118] Lobo IA, Trudell JR, Harris RA. Accessibility to residues in transmembrane segment four of the glycine receptor. *Neuropharmacology* 2006;50:174–81.
 - [119] Kenakin T. Allosteric theory: taking therapeutic advantage of the malleable nature of GPCRs. *Curr Neuropharmacol* 2007;5:149–56.
 - [120] Malysz J, Anderson DJ, Gronlien JH, Ji J, Bunnelle WH, Hakerud M, et al. In vitro pharmacological characterization of a novel selective alpha7 neuronal nicotinic acetylcholine receptor agonist ABT-107. *J Pharmacol Exp Ther* 2010;334:863–74.
 - [121] Gubbins EJ, Gopalakrishnan M, Li J. Alpha7 nAChR-mediated activation of MAP kinase pathways in PC12 cells. *Brain Res* 2010;1328:1–11.
 - [122] El Kouhen R, Hu M, Anderson DJ, Li J, Gopalakrishnan M. Pharmacology of alpha7 nicotinic acetylcholine receptor mediated extracellular signal-regulated kinase signalling in PC12 cells. *Br J Pharmacol* 2009;156:638–48.
 - [123] Utugisawa K, Nagane Y, Obara D, Tohogi H. Over-expression of alpha7 nicotinic acetylcholine receptor induces sustained ERK phosphorylation and N-cadherin expression in PC12 cells. *Brain Res Mol Brain Res* 2002;106:88–93.
 - [124] Ren K, Puig V, Papke RL, Itoh Y, Hughes JA, Meyer EM. Multiple calcium channels and kinases mediate alpha7 nicotinic receptor neuroprotection in PC12 cells. *J Neurochem* 2005;94:926–33.
 - [125] Bitner RS, Bunnelle WH, Anderson DJ, Briggs CA, Buccafusco J, Curzon P, et al. Broad-spectrum efficacy across cognitive domains by alpha7 nicotinic acetylcholine receptor agonism correlates with activation of ERK1/2 and CREB phosphorylation pathways. *J Neurosci* 2007;27:10578–87.
 - [126] Atkins CM, Selcher JC, Petraitis JJ, Trzaskos JM, Sweatt JD. The MAPK cascade is required for mammalian associative learning. *Nat Neurosci* 1998;1:602–9.
 - [127] Selcher JC, Atkins CM, Trzaskos JM, Paylor R, Sweatt JD. A necessity for MAP kinase activation in mammalian spatial learning. *Learn Mem* 1999;6:478–90.
 - [128] Sweatt JD. The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J Neurochem* 2001;76:1–10.
 - [129] Adams JP, Sweatt JD. Molecular psychology: roles for the ERK MAP kinase cascade in memory. *Annu Rev Pharmacol Toxicol* 2002;42:135–63.
 - [130] Lopez-Hernandez G, Placzek AN, Thinschmidt JS, Lestage P, Trocme-Thibierge C, Morain P, et al. Partial agonist and neuromodulatory activity of S 24795 for alpha7 nAChR responses of hippocampal interneurons. *Neuropharmacology* 2007;53:134–44.
 - [131] López-Hernández GY, Thinschmidt JS, Morain P, Trocme-Thibierge C, Kem WR, Soti F, et al. Positive modulation of alpha7 nAChR responses in rat hippocampal interneurons to full agonists and the alpha7-selective partial agonists, 4OH-GTS-21 and S 24795. *Neuropharm* 2009;56:821–30.
 - [132] Velez-Fort M, Audinat E, Angulo MC. Functional alpha 7-containing nicotinic receptors of NG2-expressing cells in the hippocampus. *Glia* 2009;57:1104–14.
 - [133] Arnaiz-Cot JJ, Gonzalez JC, Sobrado M, Baldelli P, Carbone E, Gandia L, et al. Allosteric modulation of alpha 7 nicotinic receptors selectively depolarizes hippocampal interneurons, enhancing spontaneous GABAergic transmission. *Eur J Neurosci* 2008;27:1097–110.
 - [134] Mok MHS, Kew JN. Excitation of rat hippocampal interneurons via modulation of endogenous agonist activity at the alpha7 nicotinic ACh receptor. *J Physiol* 2006;574:699–710.
 - [135] Livingstone PD, Dickinson JA, Srinivasan J, Kew JN, Wonnacott S. Glutamate-dopamine crosstalk in the rat prefrontal cortex is modulated by Alpha7 nicotinic receptors and potentiated by PNU-120596. *J Mol Neurosci* 2010;40:172–6.
 - [136] Welsby PJ, Rowan MJ, Anwyl R. Intracellular mechanisms underlying the nicotinic enhancement of LTP in the rat dentate gyrus. *Eur J Neurosci* 2009;29:65–75.
 - [137] Gusev AG, Uteshev VV. Physiological concentrations of choline activate native alpha7-containing nicotinic acetylcholine receptors in the presence of PNU-120596 [1-(5-chloro-2,4-dimethoxyphenyl)-3-(5-methylisoxazol-3-yl)-urea]. *J Pharmacol Exp Ther* 2010;332:588–98.
 - [138] Kalappa BI, Gusev AG, Uteshev VV. Activation of functional alpha7-containing nAChRs in hippocampal CA1 pyramidal neurons by physiological levels of choline in the presence of PNU-120596. *PLoS One* 2010;5:e13964.
 - [139] Johnstone TB, Gu Z, Yoshimura RF, Villegier AS, Hogenkamp DJ, Whitemore ER, et al. Allosteric modulation of related ligand-gated ion channels synergistically induces long-term potentiation in the hippocampus and enhances cognition. *J Pharmacol Exp Ther* 2011;336:908–15.
 - [140] Jope RS, Gu X. Seizures increase acetylcholine and choline concentrations in rat brain regions. *Neurochem Res* 1991;16:1219–26.
 - [141] Scremin OU, Jenden DJ. Time-dependent changes in cerebral choline and acetylcholine induced by transient global ischemia in rats. *Stroke* 1991;22:643–7.
 - [142] Egleton RD, Brown KC, Dasgupta P. Nicotinic acetylcholine receptors in cancer: multiple roles in proliferation and inhibition of apoptosis. *Trends Pharmacol Sci* 2008;29:151–8.
 - [143] Paleari L, Cesario A, Fini M, Russo P. Alpha7-nicotinic receptor antagonists at the beginning of a clinical era for NSCLC and mesothelioma? *Drug Discov Today* 2009;14:822–36.

- [144] Tournier JM, Birembaut P. Nicotinic acetylcholine receptors and predisposition to lung cancer. *Curr Opin Oncol* 2011;23:83–7.
- [145] Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, et al. Nicotinic acetylcholine receptor $\alpha 7$ subunit is an essential regulator of inflammation. *Nature* 2003;421:384–8.
- [146] de Jonge WJ, Ulloa L. The $\alpha 7$ nicotinic acetylcholine receptor as a pharmacological target for inflammation. *Br J Pharmacol* 2007;151:915–29.
- [147] Bencherif M, Lippello PM, Lucas R, Marrero MB. $\alpha 7$ nicotinic receptors as novel therapeutic targets for inflammation-based diseases. *Cell Mol Life Sci* 2011;68:931–49.
- [148] Peri R, Randall A, Bowlby M, Yeola S, Jow F, Lock Y, et al. Functional signatures of $\alpha 7$ nicotinic acetylcholine receptor positive allosteric modulators: effects on channel biophysics and pharmacology. *FASEB J Abstr* 2009;9467.
- [149] Lopez-Hernandez GY, Thinschmidt JS, Zheng G, Zhang Z, Crooks PA, Dwojskin LP, et al. Selective inhibition of acetylcholine-evoked responses of $\alpha 7$ neuronal nicotinic acetylcholine receptors by novel tris- and tetrakis-azaaromatic quaternary ammonium antagonists. *Mol Pharmacol* 2009;76:652–66.
- [150] del Barrio L, Egea J, Leon R, Romero A, Ruiz A, Montero M, et al. Calcium signalling mediated through $\alpha 7$ and non- $\alpha 7$ nAChR stimulation is differentially regulated in bovine chromaffin cells to induce catecholamine release. *Br J Pharmacol* 2011;162:94–110.
- [151] Li Y, Papke RL, He Y-J, Millard B, Meyer EM. Characterization of the neuroprotective and toxic effects of $\alpha 7$ nicotinic receptor activation in PC12 cells. *Brain Res* 1999;81:218–25.
- [152] Lukas RJ, Lucero L, Buisson B, Galzi JL, Puchacz E, Fryer JD, et al. Neurotoxicity of channel mutations in heterologously expressed $\alpha 7$ -nicotinic acetylcholine receptors. *Eur J Neurosci* 2001;13:1849–60.
- [153] Orrenius S, Zhivotovsky B, Nicotera P. Regulation of cell death: the calcium-apoptosis link. *Nat Rev Mol Cell Biol* 2003;4:552–65.
- [154] Orr-Urtreger A, Broide RS, Kasten MR, Dang H, Dani JA, Beaudet AL, et al. Mice homozygous for the L250T mutation in the $\alpha 7$ nicotinic acetylcholine receptor show increased neuronal apoptosis and die within 1 day of birth. *J Neurochem* 2000;74:2154–66.
- [155] Hu M, Gopalakrishnan M, Li J. Positive allosteric modulation of $\alpha 7$ neuronal nicotinic acetylcholine receptors: lack of cytotoxicity in PC12 cells and rat primary cortical neurons. *Br J Pharmacol* 2009;158:1857–64.
- [156] Ween H, Thorin-Hagene K, Andersen E, Gronlien JH, Lee CH, Gopalakrishnan M, et al. $\alpha 3^*$ and $\alpha 7$ nAChR-mediated Ca^{2+} transient generation in IMR-32 neuroblastoma cells. *Neurochem Int* 2010;57:269–77.
- [157] Roncarati R, Seredenina T, Jow B, Jow F, Papini S, Kramer A, et al. Functional properties of $\alpha 7$ nicotinic acetylcholine receptors co-expressed with RIC-3 in a stable recombinant CHO-K1 cell line. *Assay Drug Dev Technol* 2008;6:181–93.
- [158] Dunlop J, Roncarati R, Jow B, Bothmann H, Lock T, Kowal D, et al. In vitro screening strategies for nicotinic receptor ligands. *Biochem Pharmacol* 2007;74:1172–81.
- [159] Skok MV. To channel or not to channel? Functioning of nicotinic acetylcholine receptors in leukocytes. *J Leukoc Biol* 2009;86:1–3.
- [160] de Fiebre CM, Meyer EM, Zoltewicz J, Henry JC, Muraskin S, Kem WR, et al. Characterization of a family of anabaseine-derived compounds reveals that the 3-(4)-dimethylaminocinnamylidene derivative (DMAC) is a selective agonist at neuronal nicotinic $\alpha 7/[^{125}\text{I}]\alpha$ -bungarotoxin receptor subtypes. *Mol Pharm* 1995;47:164–71.
- [161] Meyer EM, Tay ET, Papke RL, Meyers C, Huang G, de Fiebre CM. Effects of 3-[2,4-dimethoxybenzylidene]anabaseine (DMXB) on rat nicotinic receptors and memory-related behaviors. *Brain Res* 1997;768:49–56.
- [162] Freedman R, Olincy A, Buchanan RW, Harris JG, Gold JM, Johnson L, et al. Initial phase 2 trial of a nicotinic agonist in schizophrenia. *Am J Psychiatry* 2008;165:1040–7.
- [163] Papke RL, Meyer EM, de Fiebre CM. Differential discrimination between human and rat $\alpha 7$ nAChR by GTS-21 and its primary metabolite, 4-OH, 2-methoxybenzylidene anabaseine. 26th Annual Meeting of the Society for Neuroscience; 1996. p. 602.4.
- [164] Briggs CA, Anderson DJ, Brioni JD, Buccafusco JJ, Buckley MJ, Campbell JE, et al. Functional characterization of the novel nicotinic receptor ligand GTS-21 in vitro and in vivo. *Pharm Biochem Behav* 1997;57:231–41.
- [165] Paulo J, Brucker W, Hawrot E. Proteomic analysis of an $\alpha 7$ nicotinic acetylcholine receptor interactome. *J Proteome Res* 2009;8:1849–58.