

REPETITIONS IN THE POLYPEPTIDE  
SEQUENCE OF CYTOCHROMES

Charles R. Cantor and Thomas H. Jukes

Department of Chemistry and Space Sciences Laboratory  
University of California,  
Berkeley, California\*

Received April 1, 1966

It seems likely that many proteins have evolved from comparatively short primordial peptides by processes of duplication, deletions, and amino acid substitutions due to point mutations (Eck and Dayhoff, 1966; Jukes, 1966a). In most cases, the evidence for this may have vanished due to the masking effect of evolutionary changes in the amino acid sequences. In a few cases, however, remnants of the early peptides are still discernible in the forms of partially repetitive sequences. Proteins from "primitive" organisms may be the best source in which to search for such manifestations.

The phenomena of gene duplications and deletions are perceptible as translations of the genetic message in the polypeptide sequences of certain proteins. Presumably this indicates an evolutionary origin and significance for the occurrence of these phenomena in these instances. Examples are to be found in the  $\alpha$  and  $\beta$  chains of hemoglobin A, the genes for which occupy separate chromosomal loci. Their separation is thought to result from duplication and translocation (Ingram, 1963). The two chains also show the occurrence of deletions, indicated by the presence of gaps in the homology of the polypeptides, for instance -

---

\*Supported by a National Science Foundation Cooperative Graduate Fellowship and by grant NsG 479 by the National Aeronautics and Space Administration to the University of California.



An examination of the primary structure of the cytochromes c indicates a region of partial duplication in the cytochrome c of Neurospora crassa (Heller and Smith, 1965). Using throughout the numbering system employed for the vertebrate cytochromes c, residues 5 to 19 are homologous with residues 20 to 34 as follows:

Residue No.	5															19
	lys	gly	ala	asn	leu	phe	lys	thr	arg	cys	ala	glu	cys	his	gly	
	20															34
	glu	gly	gly	asn	leu	thr	gln	lys	ile	gly	pro	ala	leu	his	gly	
Base																
Changes	1	0	1	0	0	2	1	1	1	1	1	1	2	0	0	

This internal homology is not perceptible in the cytochromes c of other species, including those of certain vertebrates (Margoliash and Smith, 1965), two yeasts (bakers yeast and Candida krusei) (Narita, et al, 1963; Narita and Titani, 1965) and the moth Samia cynthia (Chan and Margoliash, 1966). As an example, the corresponding sequences for human cytochrome c show no evidence of homology beyond what might be interpreted as being coincidental, as follows

Residue No.	5															19
	lys	gly	lys	lys	ile	phe	ile	met	lys	cys	ser	gln	cys	his	thr	
	20															34
	val	glu	lys	gly	gly	lys	his	lys	thr	gly	pro	asn	leu	his	gly	
Base																
Changes	2	1	0	2	2	3	2	1	1	1	1	2	2	0	2	

The only two identical pairs are the lysines at positions 7 and 22 and the histidines at 18 and 33. In other cytochromes c, this last vestige of a common origin of the two sequences in this region has disappeared; residues 22 and 33 are occupied by asparagine and tryptophan in tuna fish cytochrome c (Kreil, 1963) as follows:

Residue No.	5															19
	lys	gly	lys	lys	thr	phe	val	gln	lys	cys	ala	gln	cys	his	thr	
	20															34
	val	glu	asn	gly	gly	lys	his	lys	val	gly	pro	asn	leu	trp	gly	
Base																
Changes	2	1	1	2	2	3	2	1	2	1	1	2	2	3	2	

```
i Tuna                                10                               19
cytochrome c:                         phe val gln lys cys ala gln cys his thr -
                                     20          24
                                   val glu asn gly gly
ii Chromatium                          1                              10
                                       phe ala gly lys cys ser gln cys his thr leu
                                      16           20
                                   val ala asp glu gly - - - ser ala lys cys his thr phe
                                      27
                                   - - asp glu gly ser
```

Base changes,  
i vs. ii:      0   1   1   1   0   0   1   2   0   0   1   0   0   0   0   -

Base changes,  
internal in ii: - - 0 0 0 1 - - - 1 1 1 0 0 0 1

322

## REFERENCES

- Brimacombe, R., J. Trupin, M. Nirenberg, P. Leder, M. Bernfield, and T. Jaouni. 1965. *Proc. Natl. Acad. Sci., U.S.*, 54:954.
- Chan, S.K. and E. Margoliash. 1966. *J. Biol. Chem.* 241:335.
- Dus, K., R. G. Bartsch, and M. D. Kamen. 1962. *J. Biol. Chem.* 237:3083.
- Fitch, W. M. 1966. *Fed. Proc.* 25:775; *J. Mol. Biol.* 16:000.
- Ingram, V.M. 1963. *The Hemoglobins in Genetics and Evolution*. Columbia University Press, N.Y.
- Jukes, T.H. 1966a. *Molecules and Evolution*. Columbia University Press, N.Y. (in the press).
- Jukes, T.H. 1966b. *Science* (in the press).
- Kreil, G. 1963. *Z. Physiol. Chem.* 334:154.
- Margoliash, E. and E. L. Smith. 1965. In V. Bryson and H. J. Vogel (Eds.), *Evolving Genes and Proteins*, p. 221. Academic Press, N.Y.
- Narita, K., K. Titani, Y. Yaoi, and H. Murakami. 1963. *Biochem. Biophys. Acta* 77:688.
- Narita, K. and K. Titani. 1965. *Proc. Japan Acad.* 41:831.
- Smithies, O. 1964. *Cold Spring Harbor Symp. Quant. Biol.* 29:309.