

REFERENCES

1. Hayaishi, O., and Stanier, R. Y., J. Bacteriol. 62, 691 (1951).
2. Jacoby, W. B., and Bonner, D. M., J. Biol. Chem. 205, 699, 709 (1953).
3. Knox, W. E., Biochem. J. 53, 379 (1953).
4. Braunstein, A. E., Goryachenkova, E. V., and Pashkina, T. S., Biokhimiya. 14, 163 (1949).
5. Wiss, O., Helv. Chim. Acta, 32, 1694 (1949).
6. Turner, J. R., and Drucker, H., Biochem. Biophys. Res. Commun., 42, 698 (1971).
7. Van Holde, K. E., and Baldwin, R. L., J. Phys. Chem., 62, 734 (1958).
8. Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J., J. Biol. Chem. 193, 265 (1951).
9. Wada, H., and Snell, E. E., J. Biol. Chem., 237, 127 (1962).
10. Soda, K., Yorifuji, T., Misono, H., and Moriguchi, M., 114, 629 (1969).

ERRATUM

Volume 43, Number 1, April 2, 1971, in the Communication,
"Effect of Indole-3-Acetic Acid on the Synthesis of
Cyclic 3'-5'-Adenosine Phosphate by Bengal Gram Seeds"
by S. Azhar and C. R. Krishna Murti, page 59, line 25:

the concentration of IAA should read 0.18 μ M instead
of 0.18 M.