

Errata

Volume 91, Number 1, November 14, 1979

In "Effects of Cholera Toxin and 5'-Guanylylimidodiphosphate on Human Spermatozoal Adenylate Cyclase Activity," by C. Y. Cheng and B. Boettcher, pp. 1-9, on page 5, Table 1a and Table 2, the unit of human sperm adenylate cyclase activity should read "pmoles of cAMP formed/ 10^8 cells/20 min" instead of "pmoles of cAMP formed/ 10^9 cells/20 min."

Volume 91, Number 1, November 14, 1979

In "Purification and Characterization of Dicyclohexylcarbodiimide Binding Protein from Mouse Liver Mitochondria," by Vivek A. Aroskar and Narayan G. Avadhani, pp. 17-23, through a printer's error, figure 1 on p. 19 was reproduced incorrectly. For the convenience of our readers, figure 1 and its legend appear below as they should.




Fig. 1. Electrophoretic Analysis of the Proteolipid: Electrophoresis on 10% polyacrylamide-8M urea gels was carried out as described in the Materials and Methods. A, 7.5 μg proteolipid was electrophoresed and the gel was stained with Amido black; B, purified mitoplasts (25 mg each) were labeled with ^{14}C amino acid mixture with and without added chloramphenicol. After 60 min of labeling, proteolipid fractions were isolated and electrophoresed as described in A; C, photograph of a gel stained with Sudan black B.

March 13, 1980

Errata-continued

Volume 91, Number 1, November 14, 1979

In "Membrane Transport in Resealed Haemoglobin-containing Human Erythrocyte 'Ghosts' Prepared by a Dialysis Procedure," by U. Sprandel, A. R. Hubbard, and R. A. Chalmers, pp. 79-85, through an unfortunate error, figures 4 and 5 were reversed. For the convenience of our readers, Figures 4 and 5 and their legends are reproduced below.




Fig. 4a: L-Phenylalanine (2mM) transport across the membranes of normal erythrocytes (open circles) and resealed erythrocyte 'ghosts' (closed circles).

Fig. 4b: Effect of increasing L-phenylalanine concentrations on initial rate (first minute) of influx of normal erythrocytes (open circles) and resealed erythrocyte 'ghosts' (closed circles).




Fig. 5a: Uric acid (0.5mM) transport across the membranes of normal erythrocytes (open circles) and resealed erythrocyte 'ghosts' (closed circles).

Fig. 5b: Effect of increasing urate concentrations on initial rate (first minute) of influx of normal erythrocytes (open circles) and resealed erythrocyte 'ghosts' (closed circles).

Errata-continued

Volume 91, Number 1, November 14, 1979

In "Increased Paracetamol-Induced Hepatotoxicity after Chronic Alcohol Consumption," by Rolf Teschke, Gerd Stutz, and Georg Strohmeyer, pp. 368-374, on p. 368, in line 8 of the Summary, "GOT and GPT activities" should read GOT and GDH activities."

Volume 91, Number 3, December 14, 1978

In "The Monosaccharide Transporter from Human Erythrocytes is Heterogeneously Glycosylated, " by F. R. Gorga, S. A. Baldwin, and G. E. Lienhard, pp. 955-961, the legend of Figure 2 on page 958 should read "Panel B shows the radioactivity of treated (O) and untreated transporter (●)," rather than "Panel B shows the radioactivity of treated (●) and untreated (O) transporter."