

Volume 100, Number 1, May 15, 1981

In "Increase in Histidine Decarboxylase Activity in Mouse Skin after Application of the Tumor Promoter Tetradecanoylphorbol Acetate," by Takehiko Watanabe, Yoshitaka Taguchi, Kazuyuki Sasaki, Kenichiro Tsuyama, and Yukihiro Kitamura, pp. 427-432, Ref. (9) was omitted from the reference list. It should read:

9. Watanabe, T., Nakamura, H., Liang, L. Y., Yamatodani, A.,
and Wada, H. (1979) Biochem. Pharmacol. 28, 1149-1155.

Volume 100, Number 2, May 29, 1981

In "Reaction of Woodward's Reagent K with Pancreatic Porcine Phospholipase A₂: Modification of an Essential Carboxylate Residue," by Dorit Dinur, Evan R. Kantrowitz, and Joseph Hajdu, pp. 785-792: On pages 788 and 789, Figs. 1 and 2 were transposed. For the readers' convenience, the two figures are reprinted below with their respective legends.




Figure 1. Reaction of N-ethyl-5-isoxazolium-3'-sulfonate (WRK) with phospholipase A₂. The reaction was monitored by the loss of enzymatic activity and the data are presented in a semilogarithmic manner. The modification reaction was carried out at 25°C in 0.01 phosphate, pH 4.75. The enzyme concentration in each reaction was 0.17 mg/ml. The residual enzymatic activity was determined by the pH-stat assay at pH 8.0. The WRK concentrations in the various runs were (A) 2.5×10^{-3} M (●) and (B) 5.0×10^{-3} M (○). Curve (C) corresponds to the reaction using 5×10^{-3} M WRK in presence of 30 mM hexadecylphosphorylcholine.