

ERRATUM

Volume 102, Number 2, September 30, 1981

In "Specific Binding of Transforming Growth Factor Correlates with Promotion of Anchorage Independence in EGF Receptorless Mouse JB6 Cells," by Nancy H. Colburn and Thomas D. Gindhart, pages 799-807, on page 801, in Table 1, the concentration of the last agent, TGF, should read "5  $\mu$ g/ml" rather than "5 g/ml." For the readers' convenience, the table is printed below.

TABLE 1  
INDUCTION OF ANCHORAGE INDEPENDENCE BY GROWTH FACTORS

Agent	Conc.	Anchorage Independence Response (Colonies/10 <sup>4</sup> cells)
NONE		<1
TPA	10 ng/ml (1.6 X 10 <sup>-8</sup> M)	2739 $\pm$ 147
EGF	20 ng/ml <sup>a</sup> 2 ng/ml <sup>b</sup>	1547 $\pm$ 50 1194 $\pm$ 42
TGF	5 $\mu$ g/ml	1517 $\pm$ 68

The JB6 Cl 41 mouse epidermal cell line was derived as described (7) and assayed for induction of anchorage independence after exposure of cells to inducer in 0.33% agar (5,7) containing 10% fetal calf serum. The TGF preparation used was the 20-23,000 m.w. peak material obtained from A673 cells after separation by Biogel P-100 chromatography and supplied by C. Fryling and Dr. G. J. Todaro (3).

<sup>a</sup>EGF from Collaborative Research. This partially purified EGF at 2 ng/ml produced little or no colony formation.

<sup>b</sup>Purified EGF supplied by Dr. Bruce Magun.

Results are expressed as the mean value for 2 experiments run in duplicate and the variability as one-half the range.

The acknowledgment section was inadvertently omitted. It is given below.

ACKNOWLEDGMENTS

We thank Dr. Anita Roberts of NCI for comments on the manuscript and Ed Wendel and Ray Sims for technical assistance.