

Volume 104, Number 2, January 29, 1982

In "Evidence for Post-transcriptional Effects of T_3 on Hepatic Ferritin Synthesis," by Donald B. Jump, Cary N. Mariash, and J. H. Oppenheimer, pages 701-707, on page 703, in Table I, the units for non-heme iron and ferritin apoprotein should read " μ g Iron" and " μ g Protein," respectively, instead of "mg." For the reader's convenience, the corrected table appears below.

Table I. Non-Heme Iron and Ferritin Levels in Hypothyroid and Euthyroid Rat Liver^a

Thyroidal Status	Non-heme Iron		Ferritin Apoprotein	% of Total Tissue Protein as Ferritin ^c	Shell Composition ^d Subunit Ratio
	Whole Homogenate	Ferritin ^b			
	μ g Iron/g liver	μ g Protein/g liver			
Euthyroid	59.8 \pm 3.4	48.7 \pm 10.6	247.5 \pm 47.0	0.10	0.50
Hypothyroid	200.1 \pm 20.8*	163.0 \pm 6.0*	929.2 \pm 79.0*	0.35	0.24

^aThe results are the mean \pm S.D. of eight determinations

^bFerritin iron was measured in ferritin immunoprecipitates

^cHypothyroid and euthyroid liver contained 263 \pm 82 mg protein/g liver

^dAssuming 24 subunits per apoprotein shell, heavy subunit (H) M_r 21,000; light subunit (L) M_r 19,000

* $P < .001$