

CUMULATIVE SUBJECT INDEX FOR VOLUMES 134-141

A

- 2-5A, *see* 2',5'-Oligoadenylate
- AA861
antiproliferative effects on leukemia cells (human), **140**, 832
- Acetaldehyde
-protein conjugates, detection in urine of alcoholics, **140**, 924
- Acetoacetyl-CoA thiolase, *see* Acetyl-CoA acetyltransferase
- Acetone
-water mixture, chlorophyll *a* photogeneration of superoxide ion *in vitro*, ESR study (spinach), **139**, 1124
- Acetonitrile
solubilization of iron-molybdenum cofactor of *Azotobacter vinelandii* nitrogenase, **139**, 740
- 2-Acetylaminofluorene, *see* N-2-Fluorenylacetylamide
- Acetylcholine
-activated channels
and α -bungarotoxin receptor, localization on different molecules (human neuroblastoma cells), **137**, 1141
life time in cultured myotubes, regulation by cAMP (chicken embryo), **140**, 243
vasodilating effect, effect of bovine serum albumin (rat aortic rings), **138**, 1211
- Acetylcholine receptors
dimer stabilization by extracellular disulfide bonding (*Torpedo californica*), **139**, 830
localization with voltage-dependent Na⁺ channel at neuromuscular junction (mouse), **139**, 196
-rich membranes, isolation of p60^{v-src}-related tyrosine kinase: association and dissociation of phosphatidylinositol kinase activity (*Narke japonica*), **139**, 473
 α -subunit, detection of highly immunogenic region (*Torpedo californica*), **135**, 82
- Acetylcholinesterase
and thyroglobulin, sequence and structure homologies: functional and clinical significance, **137**, 142
- Acetyl-CoA acetyltransferase
fibroblast activity, effect of mevinolin (CHO-K1 cells), **138**, 953
- N-Acetylcysteine
prevention of doxorubicin-induced killing of breast cancer cells (human), **135**, 330
- N-Acetylgalactosamine
occurrence in linkage region of polygalactosamine-containing glycoproteins (*Cordyceps ophioglossoides*), **140**, 350
- N-Acetylglucosamine transferase I
detection in mung bean seedling extracts, **134**, 1395
- N-Acetylneuraminic acid
accumulation in buoyant lysosomal fractions of fibroblasts from patients with infantile generalized N-acetylneuraminic acid storage disease, **139**, 838
- N-Acetylneuraminic acid storage disease
infantile variant, N-acetylneuraminic acid accumulation in fibroblast buoyant lysosomal fractions (human), **139**, 838
- Acetylsalicylic acid, *see* Aspirin
- Acid endopeptidase
erythrocyte levels and subcellular distribution, Ca²⁺-induced alterations (human), **138**, 87
- α_1 -Acid glycoprotein
without or with high-mannose, hybrid, or complex-type oligosaccharide side chains, clearance by isolated perfused liver (rat), **141**, 949
- Acidosis
induced changes in myocardial tissues, procaine effects (dog), **134**, 900
- Acidotropic compounds
effects on processing and secretion of complement components C3 and C4 (HepG2 cells), **136**, 603
- Acid phosphatase
identification as phosphotyrosine protein phosphatase: role in dephosphorylation of membrane protein band 3 (human erythrocytes), **134**, 557
purple form, interaction with phosphate, associated oxidation of iron chromophore (bovine spleen), **136**, 490
repressible, distinct specificities toward phosphoserine and phosphotyrosyl phosphopeptides (yeast), **139**, 1202
- Acid protease
lysosomal, estrogen-induced, secretion by breast cancer cells (human), **138**, 102
- Acquired immune deficiency syndrome
activity of HTLV-III/LAV, potent and selective inhibition by 2',3'-dideoxycytidine *in vitro*, **140**, 735
cytopathogenicity of HTLV-III/LAV, selective inhibition by aurointricarboxylic acid and Evans blue *in vitro*, **136**, 64
expression of LAV/HTLV-III glycoproteins, suppression by glycosylation inhibitors, **141**, 33
- 4'-(9-Acridinylamino)methanesulfon-*m*-aniside, *see* Am-sacrine
- Acrosin
and precursor form, biochemical characterization (porcine spermatozoa), **138**, 139
- Acrosome reaction
promotion by α -L-fucosidase (bovine seminal plasma), **137**, 1061
- ACTH, *see* Adrenocorticotropin
- Actin
antibodies, effect on IgM secretion by lymphoblastoid B-cell line (human), **140**, 399
fibrous form

Subject Index

- interaction with caldesmon, role of erythrocyte protein 4.1-related factor (bovine brain), **140**, 1051
- 34-kDa binding protein, isolation and characterization (chicken gizzard smooth muscle), **141**, 20
- filament length, evaluation from depolymerization rate, **139**, 109
- gene expression in fibroblasts, effect of serial divisions (human), **141**, 112
- gene promoter, application in expression of hepatitis B surface antigen (*Saccharomyces cerevisiae*), **136**, 1063
- interaction with heavy meromyosin, analysis, **141**, 541
- monomers
- high-affinity binding of Ca^{2+} and Mg^{2+} , **135**, 607
 - sequential binding to plasma gelsolin, inhibition by vitamin D-binding protein, **136**, 72
- nonmuscle, ADP-ribosylation by component I of C_2 toxin from *Clostridium botulinum* type C (chicken), **136**, 802
- nucleating activity, increase in tumorigenic cells, **134**, 1284
- polymerization and nucleotide exchange, effect of erythro-9-[3-(2-hydroxypropyl)]adenine, **137**, 1181
- synthesis, effect of profilin, fluorescence photobleaching recovery study (*Acanthamoeba castellanii*), **140**, 308
- Actomyosin ATPase, *see* Myosin ATPase
- Acyl CoA
- long-chain, role in antagonistic effects of halothane and L-carnitine on mitochondrial energy-linked processes (rat liver), **139**, 303
- Acyl-CoA dehydrogenase
- synthesis *in vitro* (porcine kidney), **137**, 1049
- Acyl groups
- position in oligosaccharide sequences, mass spectrometric determination, **135**, 1126
- Acylpeptide hydrolase
- cleavage of blocked amino-terminal residues, substrate specificity, **139**, 244
- Acyl transferase
- identification in bovine rod outer segments, **138**, 988
- 1-N-Acyl-3'-N-(trifluoroacetyl)kanamycin
- inhibition of simian virus-40 DNA replication *in vitro* (Cos1 cells), **136**, 322
- Adenine phosphoribosyltransferase
- deficient individuals, B-cell lines from, establishment and characterization, **137**, 998
- Adenosine
- and formyl-methionyl-leucyl-phenylalanine, synergistic induction of cAMP levels in monocytes (human), **138**, 1376
 - inhibition of adenylate cyclase catalytic subunit (ram sperm), **137**, 970
 - N^6 -functionalized congeners, potency at A_2 -adenosine receptors: application as potential ligands for affinity chromatography (dog), **136**, 1097; *erratum*, **139**, 375
- Adenosine receptors
- A_1 subtype, carbohydrate content analysis (hamster, rat brain), **140**, 406
 - A_2 subtype, response to adenosine N^6 -functionalized congeners: application as potential ligands for affinity chromatography (dog), **136**, 1097; *erratum*, **139**, 375
- Adenosylhomocysteinase
- inhibition by 3-deazaneplanocin: characterization and effects on promyelocytic leukemia cells (human), **135**, 688
- S-Adenosylhomocysteinase, *see* Adenosylhomocysteinase
- Adenovirus type 2
- core protein VII and DNA chromatin-like complex, transcription *in vitro*, **136**, 86
- Adenylate cyclase
- activity in brown adipose tissue, α_2 -adrenergic effects (fetal rat), **138**, 1390
 - alterations in cell division cycle mutant 35-1 (yeast), **140**, 723
 - basal and stimulated activity in 235-1 pituitary cells, enhancement by phorbol esters, **135**, 276
 - catalytic subunit
 - inhibition by adenosine (ram sperm), **137**, 970
 - interaction with guanine nucleotide-binding stimulatory protein, forskolin stabilizing effect (rat erythrocytes), **137**, 190 - desensitization to β -adrenergic agonist stimulation, induction by tetradecanoyl phorbol acetate (rat reticulocytes), **138**, 125
 - effect of chronic doxorubicin administration (murine heart), **136**, 745
 - GTP-activated forms, as cholera toxin substrates, comparison, **134**, 928
 - induction of cAMP accumulation and hormone release in rat pituitary tumor cells (*Bordetella pertussis*), **136**, 463
 - inhibition by
 - inorganic orthophosphate, dependence on III^{Glc} protein of phosphoenolpyruvate:glycose phosphotransferase system (*Escherichia coli*), **141**, 1138
 - pentobarbital, relationship to functional coupling of guanine nucleotide-binding stimulatory protein and catalytic unit (rat brain), **140**, 237
 - transglutaminase-catalyzed reactions in erythrocyte ghosts (pigeon), **138**, 596 - negative coupling to atrial natriuretic factor receptors in atrial and ventricular cardiocytes (rat), **138**, 427
 - phosphodiesterase system in thrombopathic and normal platelets, regulatory control (dog), **140**, 589
 - sensitivity to stimulatory agonists, increase by fasting, mechanism (rat fat cells), **138**, 1411
- Adenylate kinase
- intracellular localization (*Escherichia coli*), **134**, 527
- Adipocytes
- brown
 - α -adrenergic responses, mediatory role of guanyl nucleotide-binding protein, evaluation with pertussis toxin (hamster), **135**, 823
 - and white, metabolic activity of key enzymes of carbohydrate ketone body, adenosine, and glutamine, comparison (rat), **138**, 687 - glucagon binding *in vitro*, insulin effects (rat epididymis), **141**, 812
 - insulin-like growth factor binding, stimulation by vanadate (rat), **138**, 174
 - plasma membranes, identification of calmodulin-dependent Ca^{2+} -activated K^+ channel (rat), **135**, 934
 - thyrotropin receptor, physicochemical properties (guinea pig), **135**, 208
- Adipokinetic hormone, *see* Lipotropic hormone

Subject Index

- ADP (adenosine 5'-diphosphate)
 -ATP carrier, substrate and inhibitor binding, analysis with 6'-O-dansyl- γ -aminobutyryl atractyloside (bovine cardiac mitochondria), **134**, 266
 induced Ca^{2+} flux in fura-2-loaded platelets, kinetics (human), **136**, 1124
 vasodilating effect, effect of bovine serum albumin (rat aortic rings), **138**, 1211
- ADP-ribose
 monomeric and polymeric, degradation rates in permeabilized HeLa cells, **139**, 37
 protein-bound, hydrolyzing enzyme, identification (rat skeletal muscle), **139**, 932
- (ADP-ribose)_n glycohydrolase
 forms I and II, detection in HeLa S3 cells, **135**, 979
 occurrence in erythrocytes (human), **136**, 1110
- ADP-ribosylation
 associated activity of *Clostridium difficile* toxin B, analysis in fibroblasts (human), **139**, 64
 avian nonmuscle actin by component I of C₂ toxin (*Clostridium botulinum* type C), **136**, 802
 chemoattractant receptors for mediation by pertussis/cholera toxin-sensitive N protein (human), **138**, 887
 cholera toxin-catalyzed
 efficiency improvement by NADP (canine cardiac, rat hepatic membranes), **137**, 328
 and pertussis toxin-catalyzed, substrate identification on cardiac cell membranes (rat), **138**, 1355
- ADP-ribosyltransferase
 cellular, specific for elongation factor 2, characterization (rat liver), **139**, 1210
- Adrenalectomy
 effect on corticotropin-releasing factor binding to peripheral tissues (rat), **134**, 255
- Adrenal gland
 α -atrial natriuretic factor receptor distribution (bovine), **137**, 657
 atrial natriuretic polypeptide molecular size, comparison with kidney and aorta by uv-photoaffinity labeling (rat), **136**, 200
 cortex, inner and outer zones, cAMP-dependent protein kinase activity and protein phosphorylation (guinea pig), **138**, 895
- β -Adrenergic agonists
 stimulation in adenylate cyclase system, desensitization induction by tetradecanoyl phorbol acetate (rat reticulocytes), **138**, 125
- α_1 -Adrenergic receptors
 electrical stimulation, effect on phosphoinositide metabolism (rat muscle), **136**, 863
 expression in myocardium, age-dependent changes (rat), **138**, 387
 mediated activation of membrane phospholipases and iodide fluxes, role of pertussis toxin-sensitive G proteins (FRTL-5 cells), **141**, 1000
 regulation of atriopeptin release from isolated heart (rat), **137**, 94
 stimulation, effect on ³²P-labeling of membrane phospholipids (rabbit, rat aorta), **141**, 1213
 stimulation of phosphoinositide breakdown and respiration in brown adipocytes, mediatory role of guanyl nucleotide-binding protein, evaluation with pertussis toxin (hamster), **135**, 823
- α_2 -Adrenergic receptors
 activation, subsequent insulin release, effect of glucose (murine pancreas), **139**, 557
 photoaffinity labeling with [³H]SKF 102229 (human platelets, rabbit kidney), **137**, 606
 regulation by sodium, effect of 4,4'-diisothiocyano-2,2'-stilbene disulfonic acid (human platelets), **141**, 222
- β -Adrenergic receptors
 detection along renal proximal tubules (rat), **141**, 334
 pulmonary, binding properties (human fetus), **135**, 816
 76-kDa form, identification (human placental basal membranes), **141**, 411
- Adrenocortical cells
 differentiated, steroidogenic activities, effect of transforming growth factor type β (bovine), **139**, 693
 steroidogenesis *in vitro*, effect of human α -atrial natriuretic factor (bovine), **137**, 657
- Adrenocorticotropin hormone, *see* Adrenocorticotropin
- Adrenocorticotropin
 and cAMP, comparative effects on epinephrine-induced melanosome aggregation (goldfish), **138**, 1081
 dual activation of cAMP and inositol-trisphosphate- Ca^{2+} signaling systems (rat adrenal cells), **135**, 742
 12-O-tetradecanoylphorbol 13-acetate-induced secretion by pituitary cells, role of Ca^{2+} : effect of dexamethasone (rat), **138**, 323
- Adriamycin, *see* Doxorubicin
- Affinity chromatography
 with adenosine N⁶-functionalized congeners, potency at A₂-adenosine receptors (dog), **136**, 1097; *erratum*, **139**, 375
 on immobilized Rolipram, purification of cAMP-specific phosphodiesterase (rat heart), **138**, 205
 insulin-like growth factor I with amniotic fluid somatomedin-binding protein (guinea pig), **136**, 253
 in isolation of transcriptionally active DNA (chicken erythrocytes), **137**, 474
- Aflatoxin B₁
 induced hepatocellular carcinomas, expression of *c-myc*, *c-Ha-ras* and *N-ras* genes (rat), **138**, 858
 -specific form of cytochrome P-450 from 3-methylcholanthrene-treated liver, purification and characterization (hamster), **139**, 466
- Age
 effect on
 expression of α_1 -adrenergic receptors in myocardium (rat), **138**, 387
 glycolysis and lipid synthesis (rat brown adipose tissue), **140**, 419
 methylation of *c-myc* proto-oncogene (mouse), **139**, 1299
 related differences in interferon induction of 2-5A synthetase and 2-5A binding protein (guinea pig macrophages), **140**, 455
 senescence-associated changes in
c-myc oncogene expression (human fibroblasts), **135**, 105
c-ras oncogene expression (human fibroblasts), **137**, 917
- Agglutinin
 ketosugar-specific, isolation (rat uterus), **136**, 116
- AIDS, *see* Acquired immune deficiency syndrome
- Ajoene
 inhibition of fibrinogen binding to platelets (human), **141**, 145

Subject Index

- L-Alanine
 hydration state in water/dimethyl sulfoxide mixtures, ¹⁷O NMR, **135**, 363
 stimulation of phosphoinositide turnover in olfactory cilia (catfish), **137**, 36
- L-Alanine:4,5-dioxovalerate transaminase, *see* Aminolevulinate aminotransferase
- D-Alanine²-D-leucine⁵-enkephalin
 and bradykinin, effects on inositol trisphosphate release from NCB-20 cells, **135**, 507
- D-Alanine²-D-leucine⁵-enkephalin receptors
 effects of swainsonine and tunicamycin, comparison (NCB-20 cells), **134**, 1387
- Albumin
 -copper complexes in bloodstream, functional role, **135**, 1064
 formaldehyde-treated and galactosylated, cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
 mRNA, hepatocyte levels, regulation by dexamethasone and estrogen *in vitro* (*Xenopus laevis*), **136**, 8
 pre-, intron V, complementary cytoplasmic RNA, detection (rat liver), **136**, 989
 serum, *see* Serum albumin
- Alcohol
 -specific product, detection in urine of alcoholics, **140**, 924
- Alcohol dehydrogenase
 class I and II isozymes, role in serotonin metabolism (human liver), **139**, 1009
 subunits α , β , and γ , mRNA size heterogeneity and developmental changes (human), **140**, 1020
 testicular isozymes, isolation and characterization (human), **134**, 1056
- Aldehyde oxidase
 inhibition by *N*-hydroxy-2-acetylaminofluorene (rabbit liver), **138**, 312
 oxidation of tropylium tetrafluoroborate (rabbit liver), **140**, 609
 reduction of chromate (rabbit liver), **137**, 8
- Aldehydes
 peptide, and α_1 -antitrypsin, as elastase inhibitors in treatment of emphysema, comparison (human), **139**, 896
- Aldolase, *see* Fructose-bisphosphate aldolase
- Aldosterone
 angiotensin-stimulated production, effect of osmolality (bovine adrenal glomerulosa cells), **139**, 46
 plasma levels, effects of α -melanocyte-stimulating hormone *in vivo* (rat), **141**, 1279
 secretion by glomerulosa cells, K⁺-induced, inhibition by diazepam (bovine adrenal), **135**, 994
- Aldosteronism
 primary, associated ouabain-like activity in plasma (human), **135**, 41
- Aldrin
 epoxidation, lipid peroxidation-dependent, analysis in hepatic microsomes, hepatocytes, and granuloma cells (rat), **138**, 24
- Alizarin
 inhibition of glutathione *S*-transferase isoenzymes from human and rat tissues, comparison, **141**, 1170
- Alkaline elution
 gravity-flow, detection of carcinogen-induced DNA strand breaks (MDBK cells), **137**, 1006
- Alkaline phosphatase
 catalytic rate enhancement by diffusion of adsorbed substrate on membrane surface, **138**, 803
 expression in subclones of neoplastic salivary duct cells (human), **138**, 625
 inhibition of cholesterol 7 α -hydroxylase (rat), **134**, 797
 proliferation and expression in murine osteoblast-like cells, inhibition by porcine type- β transforming growth factor, **140**, 56
- N*-Alkanes
 induction of cytochrome *P*-450 mRNA synthesis (*Candida maltosa*), **136**, 1148
- Allergenicity
 quaternary ammonium salts (guinea pig), **135**, 221
- Allopurinol
 beneficial action during myocardial reperfusion injury, mechanism, **137**, 76
- Alloxan
 induced diabetes, associated decreased expression of epidermal growth factor receptors (rat), **137**, 258
- Allyl chloride
 epoxidation by chloroperoxidase (*Caldariomyces fumago*), **136**, 778
- Alveolyn
 -like protein associated with murine type 2 pneumocytes, immunochemical characterization, **139**, 1135
- Alzheimer's disease
 associated filaments, self-assembly of microtubule-associated protein τ from porcine brain into similar filaments, **141**, 790
 brain temporal region homogenates, quantitation of phosphorylcholine hydrolysis (human), **139**, 315
 β -protein, synthetic peptides corresponding to 17- and 28-residue sequences, *in vitro* formation of amyloid fibrils (human), **141**, 782
- α -Amanitin
 effect on phosphorylation of proteins crosslinked to poly (A)⁺ RNA from free polyribosomes (rat), **136**, 356
- Amidase
 Ca²⁺-dependent, identification and isolation (human heart), **136**, 773
- α -Amidation
 peptide C-terminal
 analysis in neonatal pancreas (rat), **138**, 179
 catalyzing enzyme, purification and characterization (*Xenopus laevis* skin), **137**, 984
 tissue distribution and characterization (rat), **140**, 230
- Amiloride
 effect on platelet serotonin release and clot retraction (human), **140**, 329
 -sensitive sodium transport, inhibition by atrial natriuretic factor and cGMP (LLC-PK₁ cells), **134**, 852
- Aminacrine
 induced frameshift mutations, effect of mutation in *uvrB* gene (*Salmonella typhimurium*), **138**, 505
- Amine oxidase (flavin-containing)
 inactivation by 1-phenylcyclopropylamine: identification of cysteine residue involved in labile adduct formation, **135**, 154
- Amino acid composition
 aromatase (human placenta), **135**, 713

Subject Index

- basic fibroblast growth factor amino-terminal extended form (bovine pituitary), **138**, 580
- crystalline serine hydroxymethyltransferase (*Hyphomicrobium methylovorans*), **139**, 71
- delta toxin and active fragment (*Bacillus thuringiensis*), **141**, 106
- plastocyanin (*Ceratophyllum demersum*), **136**, 610
- prostatropin multiple forms (bovine brain), **136**, 1155
- pulmonary surfactant-associated proteins (bovine), **135**, 527
- soluble hydrogenase from *Desulfovibrio africanus*, **139**, 658
- thylakoid 9-kDa phosphoprotein (pea), **138**, 146
- D-Amino acid oxidase
- biosynthesis *in vitro* and *in vivo*, comparison (porcine kidney), **141**, 1222
- Amino acids
- blocked, N-terminal, from blocked peptides, cleavage by acylpeptide hydrolase, substrate specificity, **139**, 244
- branched-chain, plasma levels, effect of sustained exercise (rat), **137**, 149
- interferon α_2 , functionally important, identification with analogs produced from synthetic genes (human), **134**, 1404
- nonproteinic, identification in xylem sap (*Pisum sativum*), **139**, 383
- ribonuclease T₁ L-glutamic acid-58, modification to L-glutamine and L-aspartic acid (*Escherichia coli*), **138**, 789
- Amino acid sequence
- acetylcholinesterase and thyroglobulin, homology: functional and clinical significance, **137**, 142
- acidic fibroblast growth factor
- bovine brain, **133**, 554; *erratum*, **134**, 1419
- human brain, **138**, 611; **140**, 874
- adipokinetic hormone
- from corpora cardiaca (*Heliothis zea*), **135**, 622
- type II from corpora cardiaca, determination by fast atom bombardment mass spectrometry (locust), **134**, 723
- amino-terminal
- acidic and basic forms of fibroblast growth factor (human brain), **135**, 541
- amyloid fibril protein from insulin-producing pancreatic tumor, partial homology with calcitonin gene-related peptide (human), **140**, 827
- angiotensin-converting enzyme (bovine lung), **141**, 968
- apolipoprotein B-100 fragments resulting from cleavage with kallikrein (human plasma), **137**, 821
- aromatase (human placenta), **135**, 713
- atriopeptin cardiac and circulating forms (rabbit), **138**, 1263
- cytochrome P-450 (rat renal microsomes), **141**, 711
- cytochrome P-450 K from rabbit kidney and protein encoded by plasmid pP-450 PBc2, comparison, **141**, 728
- elongation factor 2 (rat liver), **134**, 1015
- galactosyltransferase (human milk), **138**, 224
- H⁺.K⁺-ATPase (ovine kidney), **138**, 185
- protein S100 subunits, homology with lymphocytic p36-associated 10-kDa polypeptide (swine), **134**, 248
- renal renin (human), **139**, 446
- spore coat proteins (*Dictyostelium discoideum*), **135**, 336
- angiotensin-converting enzyme inhibitor (porcine plasma), **139**, 52
- L-aspartate ammonia-lyase and fumarate hydratase, homology (*Bacillus subtilis*, *Escherichia coli*, *Pseudomonas fluorescens*), **138**, 568
- C1 inhibitor (human), **137**, 620
- Ca²⁺.Mg²⁺-ATPase nucleotide-binding domain (canine cardiac sarcoplasmic reticulum), **135**, 864
- carboxy-terminal
- atriopeptin cardiac and circulating forms (rabbit), **138**, 1263
- crystalline serine hydroxymethyltransferase (*Hyphomicrobium methylovorans*), **139**, 71
- coat proteins of DNA and RNA plant viruses, homology, **134**, 372
- cystatin C and protein variant in hereditary cerebral amyloid angiopathy, comparison (human), **136**, 548
- fructose-1,6-bisphosphatase, homology among various species, **135**, 374
- globular proteins, application in domain boundary prediction, **139**, 11
- glycogen synthase tryptic fragment containing protein C-terminus (rabbit skeletal muscle), **137**, 542
- group A streptococcal Pep M5 protein, homology with other coiled-coil proteins, **140**, 684
- human T-cell leukemia virus type I protease, comparison with other retroviruses (MT-2 cells), **139**, 129
- hypertrehalosaemic factor (cockroach), **140**, 674; **141**, 774
- inhibin subunits α , β_A , and β_B (human ovary), **135**, 957
- kinetensin, homology with serum albumin, neurotensin, and angiotensin (human), **136**, 983
- leucosulfakinin-II, homology with cholecystokinin and gastrin (*Leukophaea maderae*), **140**, 357
- methylamine dehydrogenase cofactor segments adjacent to linkage sites (*Methylobacterium* sp.), **141**, 562
- β_2 -microglobulin fragments in amyloid kidney stones from uremic patients, **136**, 665
- Novikoff hepatoma hexokinase C, comparison with yeast hexokinases (rat), **139**, 714
- peptide homologous to neuropeptide Y (coho salmon endocrine pancreas), **141**, 1084
- peptides from apolipoprotein B-100 surface-exposed segments on low-density lipoprotein particles (human), **140**, 250
- phospholamban structural unit (canine cardiac sarcoplasmic reticulum), **138**, 1044
- procarboxypeptidase A activation segment (porcine pancreas), **141**, 517
- proteins, comparison by principal coordinate analysis, **140**, 808
- segments surrounding tyrosine sulfation sites in proteins, **141**, 326
- 9-Aminoacridine, *see* Aminacrine
- δ -(L- α -Aminoadipyl)-L-cysteine
- cell-free synthesis (*Cephalosporium acremonium* extracts), **137**, 528
- 3-Aminobenzamide
- inhibition of phorbol ester-enhanced malignant transformation of embryonic fibroblasts (mouse), **134**, 1334
- γ -Aminobutyric acid
- stimulation of chloride uptake by membrane vesicles, effect of benzodiazepines and β -carbolines (rat cerebral cortex), **141**, 1
- γ -Aminobutyric acid receptors
- photoaffinity labeling with [³H]flunitrazepam and [³H]mus-

Subject Index

- cimol, comparison (bovine brain), **138**, 1308
- 2-Amino-3,8-dimethylimidazo[4,5-f]quinoxaline
detection in dialysate from patients with uremia, **138**, 1084
- N-(2-Aminoethyl)-5-isoquinolinesulfonamide
effect on neutrophil activation (human), **135**, 749
- α -Aminoisobutyric acid
transport by Na^+, K^+ -ATPase, basal and stimulated, oua-
bain effects (rat skeletal muscle), **134**, 1342
- Aminolevulinate aminotransferase
induction by phenylhydrazine, role of heme level (rat kid-
ney, liver), **138**, 742
inhibition by excess generation of endogenous heme (rat
liver), **138**, 751
- 5-Aminolevulinate synthase
mRNA levels, effects of testosterone and hemin (embryonic
chicken liver), **140**, 81
- δ -Aminolevulinate synthase, *see* 5-Aminolevulinate synthase
- 4'-Aminomethyl-4,5',8-trimethylpsoralen
insulin-conjugated derivative, insulin receptor-mediated
photocytotoxic effect on activated lymphocytes, **141**,
502
- 1-Amino-3,7,8-trichlorodibenzo-*p*-dioxin
antagonism of TCDD-induced myelotoxicity (murine bone
marrow), **139**, 747
- Ammonia
repression of nitrogenase (*Anabaena cylindrica*), **134**, 835
toxicity, effect of sodium benzoate (rat), **138**, 369
- Ammonium salts
quaternary, allergenicity (guinea pig), **135**, 221
- AMP (adenosine 5'-monophosphate)
fructose-1,6-bisphosphatase sensitivity, effect on enzyme
modification by chelators (fish, rat brain), **139**, 571
- Amphotericin B
restoration of stationary-phase hepatoma cell sensitivity to
dipyridamole (rat), **140**, 434
- Amsacrine
induced DNA cleavage, magnitude, correlation with sen-
sitivity to drug (human leukemia cells), **134**, 638
stimulation of topoisomerase II-induced cleavage sites in
c-myc proto-oncogene *in vivo* (N417 cells), **137**, 154
- α -Amylase
release from pancreatic acini, effect of chloroquine binding
to muscarinic receptors (rat), **137**, 664
signal peptides, secretion activities in *Escherichia coli* cells
(*Bacillus subtilis*), **134**, 624
- α -Amylase inhibitors
gene from *Streptomyces griseosporus*, cloning and expres-
sion in *S. lividans*, **141**, 1099
- Amyloid
fibril protein, from insulin-producing pancreatic tumor, par-
tial homology of N-terminal sequence with calcitonin
gene-related peptide (human), **140**, 827
fibrils, formation *in vitro* from synthetic peptides correspon-
ding to 17- and 28-residue sequences of Alzheimer's
disease β -protein (human), **141**, 782
P-component, fibronectin binding: localization of binding
site on C-terminal domain (human), **140**, 12
serum protein A, gene localization on chromosome 11 and
detection of DNA polymorphisms (human), **137**, 1196
- Amyloidotic polyneuropathy
familial, human transthyretin variant produced by trans-
genic mouse as disease model, **139**, 794
- Amylose-precipitating factor
regulation of starch-synthesizing enzymes (*Neurospora
crassa*), **135**, 701
- Androgen receptors
modification *in vitro* by cAMP (murine skeletal muscle),
135, 1069
- Androgens
-binding proteins, secretion by immature Sertoli cells, *in
vitro* effects of germ cells (rat), **141**, 861
- Angiogenesis
inhibition by bisantrene and mitoxantrone (rat avascular
cornea), **140**, 901
- Angiopathy
hereditary cerebral amyloid, cystatin C and protein variant,
amino acid sequences, comparison (human), **136**, 548
- Angiotensin I
amino acid sequence, homology with kinetensin (human),
136, 983
conversion to type II by peptidyl dipeptidase (bovine en-
dothelial cells), **134**, 770
- Angiotensin II
antiserum, purification and characterization by affinity
chromatography (rabbit), **136**, 685
contractile response of glomeruli, impairment in diabetes
mellitus (rat), **136**, 1185
effect on cytoplasmic free Ca^{2+} levels, comparison with
phenylephrine and vasopressin (rat hepatocytes), **140**,
107
generation by renal carboxypeptidase (human), **138**, 573
induced prostaglandin E_2 synthesis in mesangial cells, op-
posite effects of different concentrations of pertussis
toxin (rat), **141**, 39
stimulation of
aldosterone production, effect of osmolality (bovine ad-
renal glomerulosa cells), **139**, 46
protein phosphorylation, inhibition by atrial natriuretic
peptide (bovine adrenal glomerulosa cells), **140**, 814
tyrosine phosphorylation by casein kinase A (human eryth-
rocytes), **139**, 855
- Angiotensin-converting enzyme, *see* Dipeptidyl carboxypep-
tidase I
- Angiotensinogen
peptide analogs, effect of chain length on renin inhibition
(monkey), **139**, 982
- Anisylidithiolthione
inhibition of lipid peroxidation (rat hepatic microsomes),
135, 1015
- Ankyrin
tyrosine phosphorylation by casein kinase A (human eryth-
rocytes), **139**, 855
- Anthracene
phototoxicity, evaluation using *Paramecium aurelia*, **139**,
79
- Anthrapyrazoles
and aminoacridine derivative, effects on lipid peroxidation
(rabbit hepatic microsomes), **140**, 797
- Antibodies
actin, effect on IgM secretion by lymphoblastoid B-cell
line (human), **140**, 399
activation of B cells, associated intracellular Ca^{2+} release
and Ca^{2+} uptake, analysis with indo-1 (mouse), **137**,
500; *erratum*, **138**, 503

Subject Index

- auto-immune, recognition of initiator methionine tRNA from yeast (human), **139**, 275
- to *Bacillus cereus* phospholipase C, cross-reaction with phosphatidylcholine-preferring phospholipase C in human monocytic cells, **140**, 114
- binding to Na⁺/K⁺ pump-associated antigens, enhancement by intracellular K⁺ (ovine erythrocytes), **134**, 982
- elongation factor 2, preparation (rat liver), **134**, 1015
- enzyme-specific, in demonstration of guinea pig L-dopa decarboxylase and rat L-histidine decarboxylase immunohistochemical and immunochemical cross-reactivity, **141**, 306
- to hepatitis B virus surface antigens, production by administration of murabutide associated with synthetic pre-S viral peptide conjugated to toxoid carrier, **140**, 557
- HNK-1, reactive glucuronic acid, demonstration on brain glycoproteins (goldfish), **138**, 902
- to insulin-like growth factor I receptors, effect on insulin-like growth factor II stimulation of DNA synthesis (human fibroblasts), **139**, 501
- to insulin-like growth factor II receptors, generation (human pituitary adenoma), **138**, 304
- iron-sulfur protein, interaction with mitochondrial complex III (bovine heart), **136**, 376
- masked, naturally occurring in serum, recognition of mitotic spindle apparatus component (mouse), **136**, 145
- monoclonal, *see* Monoclonal antibodies
- peptide-specific, identification of protein-tyrosine kinase (human T cells), **140**, 392
- phospholipid, two-dimensional crystallization, induction by supported phospholipid bilayers, **134**, 819
- polyclonal
- anti-paratypic, cross-reactive properties, inhibition of morphine binding (rabbit), **136**, 1177
 - for comparison of regulatory and catalytic subunits of cAMP-dependent protein kinase (bovine heart, *Dicystostelium discoideum*), **136**, 651
 - in cytolocalization and subunit structure identification of 1,4-dihydropyridine component of neuronal Ca²⁺ channel (rabbit brain, skeletal muscle), **139**, 996
 - IgG, effect on autophosphorylation and tyrosine kinase activity of insulin-like growth factor receptors (human), **138**, 1023; *erratum*, **140**, 1152
 - triamcinolone acetone, and glucocorticoid receptors, steroid-binding domains, comparison (rat), **136**, 470
 - ubiquinone-binding protein, interaction with mitochondrial complex III (bovine heart), **136**, 376
- Antigens
- cytochrome *c*, properties and molecular size, comparison (bovine, mouse, rabbit, rat), **141**, 131
 - destabilization of target-sensitive immunoliposomes by binding: as rapid assay for viruses, **138**, 931
 - determinant groups, *see* Epitopes
 - GA733, gastric carcinoma-associated, isolation and characterization (human), **135**, 297
 - macrophage surface-directed monoclonal antibody, coordinate inhibition of NADPH oxidase activation and phosphorylation of membrane proteins (human neutrophils), **140**, 1121
 - middle T, polyoma virus
 - and mitogenic growth factors, cooperation in transformation of secondary cultured embryo cells (rat), **136**, 921
 - protein kinase C-promoted tyrosine residue phosphorylation (murine py6, rat pyF cells), **136**, 995
 - transforming ability, enhancement by protein kinase C stimulation (rat F111 cells), **140**, 1106
 - Na⁺/K⁺ pump-associated, binding to antibodies, enhancement by intracellular K⁺ (ovine erythrocytes), **134**, 982
 - PrP-related, localization in normal human brain (scrapie-infected hamster brain), **140**, 758
 - surface, hepatitis B virus
 - expression and secretion in *Escherichia coli*, **135**, 1042
 - expression by yeast actin gene promoter, **136**, 1063
 - modified P31 particles, synthesis in yeast after gene modification, **141**, 942
 - P31 gene, expression in *Saccharomyces cerevisiae*, **138**, 268; *erratum*, **140**, 774
 - recognition by antibodies produced by administration of murabutide associated with synthetic pre-S viral peptide conjugated to toxoid carrier, **140**, 557
 - T, simian virus 40, purification by DNA-Sepharose affinity chromatography, **134**, 915
 - Tac, on interleukin-2 receptor, epitope phosphorylation by protein kinase C (human T cells), **135**, 239
 - tubulin-like, superficial, recognition by *Trypanosoma cruzi* monoclonal antibody (mouse), **139**, 1176
- Antiserum
- angiotensin II, purification and characterization by affinity chromatography (rabbit), **136**, 685
 - pp60^{src}, detection of tyrosine kinase in plasma, serum, and erythrocyte extracts (human), **135**, 426
 - α_1 -Antitrypsin, *see* α_1 -Trypsin inhibitor
- Aorta
- atrial natriuretic polypeptide molecular size, comparison with kidney and adrenal gland by uv-photoaffinity labeling (rat), **136**, 200
- Apamin
- sensitive Ca²⁺-activated K⁺ channels, and Ca²⁺ channels in muscular dysgenic skeletal muscle cells, action potentials and contractile activity (mouse), **136**, 935
- Apocytochrome *c*
- heme-attaching enzyme, localization in yeast mitochondria, **141**, 1145
- Apolipoprotein A-IV
- affinity for lipoproteins during cholesterol esterification (human), **135**, 756
 - complexes with dimyristoyl phosphatidylcholine, binding to plasma membranes (rat liver), **139**, 122
- Apolipoprotein B
- cleavage by kallikrein: characterization of resulting fragments (human plasma), **137**, 821
 - identification as calcium-binding protein (Hep G2 cells), **137**, 493
 - low-density lipoprotein receptor binding domains, identification and proposed consensus sequence (human), **139**, 1078
 - surface-exposed segments in low-density lipoprotein particles, associated peptides, isolation and amino acid sequence (human), **140**, 250
- Apolipoprotein C-I
- gene sequences on chromosome 19, mapping (human), **136**, 876

Subject Index

Apolipoprotein C-II

N- α -palmitoyl (56-79) fragment, activation of lipoprotein lipase (human plasma), **137**, 1041

Apolipoprotein E

binding to highly reactive heparin: identification of heparin-binding domains (human), **134**, 783

deficiency, associated monocyte-macrophage mRNA levels (human), **134**, 937

gene sequences on chromosome 19, mapping (human), **136**, 876

induced lipid uptake by hepatocyte monolayers, role of glycosaminoglycan sulfates (rat), **141**, 158

mediation of hypertriglyceridemic very-low-density lipoprotein binding to low-density lipoprotein receptors, detection by ligand blotting technique (bovine adrenal gland), **139**, 333

radioimmunoassay with solid-phase staphylococcus protein A, use of pooled plasma as secondary standard (human), **138**, 455

Apolipoproteins

phospholipid-binding regions, proline-containing, magnitude of helical hydrophobic moment, **139**, 202

Apoproteins

heme covalent binding during destruction of microsomal cytochrome *P*-450 (rat liver), **138**, 193

and lipid constituents of high-density lipoproteins, role in high-density lipoprotein-induced cardiac prostacyclin synthesis (rabbit), **139**, 416

Arabinogalactan

oxidation by hydrogen peroxide, role in cell wall breakdown (larch), **141**, 238

Arabinose 1,5-bisphosphate

effect on 6-phosphofructo-1-kinase and fructose-1,6-bisphosphatase (rat liver), **138**, 159

Arachidonate

decreased content in plasma phospholipids from Usher's syndrome patients, **141**, 600

Arachidonic acid

activation of NADPH-oxidase, role of phospholipase *A*₂ in intact neutrophils and cell-free system, comparison (human), **138**, 1099

enhancement of prostaglandin *I*₃ production from eicosapentaenoic acid (human endothelial cells), **135**, 403

free, requirement for leukotriene *B*₄ synthesis by 12-hydroperoxyeicosatetraenoic acid-stimulated neutrophils (human), **138**, 589

inhibition of leukotriene *B*₄ production (human T cells), **135**, 670

lipoxygenase-mediated metabolism in macrophages, synergistic stimulation by A23187 and phorbol ester (mouse), **140**, 861

lipoxygenation, requirement for aggregated IgG-stimulated phagocytic degranulation by neutrophils (human), **136**, 310

oxygenation

by microsomal cytochrome *P*-450, hydroxyeicosatetraenoic acids formed during, absolute configuration (rat liver), **141**, 1007

products, effect on protein kinase C (human), **134**, 1215

and protein kinase C, coordinate actions in gonadotropin-releasing hormone-stimulated luteinizing hormone se-

cretion (rat pituitary cells), **134**, 134

release from

A23187-stimulated neutrophils, enhancement by phorbol ester (human), **141**, 399

macrophages, role of phospholipase C-diglyceride lipase system (mouse), **141**, 367

promyelocytic leukemic HL-60 cells, induction by dimethylsulfoxide (human), **140**, 515

thrombin-stimulated platelets, role of diacylglycerol lipase pathway (human), **134**, 1327

releasing systems in aortic endothelial cells, mechanisms (swine), **139**, 1098

transmembrane penetration and conversion to mono- and dihydroxy metabolites, enhancement by ethanol (human leukocytes), **140**, 782

Arginase

hepatic, cDNA

cloning and expression in *Escherichia coli* (rat), **136**, 955

isolation and demonstration of nonhomology between hepatic and renal genes (human), **141**, 53

Arginine

-containing peptides, vasodilating properties, dependence on endothelium (rat aorta), **141**, 446

effect on insulin release from cloned B-cell line HIT-T15 (hamster), **140**, 616

Arginine vasopressin

effect on atrial natriuretic factor release by atrial slices (rat), **135**, 1090

release by hypothalamo-neurohypophyseal complex in culture, effect of synthetic atrial natriuretic factor (rat), **134**, 652

stimulation of inositol trisphosphate accumulation in cultured aortic myocytes, modulation by protein kinase C (rat), **140**, 1093

Aromatase

amino acid composition and amino-terminal sequence (human placenta), **135**, 713

follicle-stimulating hormone-induced, in granulosa cells, modulation by inhibin and transforming growth factor, comparison (rat), **136**, 969

and NADPH-cytochrome *P*-450 reductase, reconstituted system, stereochemistry of estrogen biosynthesis (human placenta), **136**, 454

purification and properties (human placenta), **134**, 704

Aromatic-L-amino-acid decarboxylase

and rat L-histidine decarboxylase, immunohistochemical and immunochemical cross-reactivity (guinea pig), **141**, 306

Arsenite

induced changes in hepatic phosphate levels, ³¹P NMR *in vivo* study (rat), **139**, 228

Aryl hydrocarbon hydroxylase, *see* Unspecific monooxygenase

L-Ascorbate oxidase

reduction, effect on active site-associated copper ions (cucumber), **135**, 644

Ascorbic acid

-copper system, selective oxidation of imidazole ring in histidine residues, **138**, 659

plasma levels, effect of *Plasmodium vinckei* infection (mouse), **134**, 152

Subject Index

- Asialofetuin
cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
- Asialo-orosomucoid
dissociation after internalization by hepatocytes, mediation by distinct galactosyl receptor populations (rat), **140**, 43
- Asparaginase
complex, reversible self-phosphorylation, characterization of associated protein kinase and protein phosphatase activities (*Leptospira michotii*), **140**, 1135
- L-Aspartate
inhibition of chemoreceptor demethylation in absence of regulatory domain of demethylating enzyme (*Salmonella typhimurium*), **141**, 918
- Aspartate aminotransferase
in glutamine utilization pathway, intracellular distribution in lymphocytes (rat), **138**, 318
mRNA, mitochondrial, molecular cloning and expression regulation in regenerating liver (rat), **134**, 803
- Aspartate ammonia-lyase
and fumarate hydratase, sequence homology (*Bacillus subtilis*, *Escherichia coli*, *Pseudomonas fluorescens*), **138**, 568
- Aspartate carbamoyltransferase
binding to *N*-(phosphonacetyl)-L-aspartate, mechanism (*Escherichia coli*), **136**, 822
- D-Aspartic acid
content in myelin and myelin basic protein (bovine, human brain), **135**, 683
free, levels in various tissues, measurement (human, rat), **141**, 27
- Aspergillus niger*
detection of α -L-(1 \rightarrow 6)-fucosidase, **136**, 563
- Aspirin
effect on bleeding-induced changes in hemoglobin proportions (adult rat), **137**, 69
inhibition of phospholipase C, mechanism (human monocytes), **139**, 115
- Astroglial cells
chemotaxis, stimulation by bovine brain-derived growth factor (rat), **141**, 67
in culture, nerve growth factor synthesis and secretion (mouse), **136**, 57
- 2-5A synthetase, *see* 2',5'-Oligoadenylate synthetase
- ATP (adenosine 5'-triphosphate)
-ADP carrier, substrate and inhibitor binding, analysis with 6'-O-dansyl- γ -aminobutyl atracyloside (bovine cardiac mitochondria), **134**, 266
agonist evoked secretion, inhibition by antioxidants (human platelets), **139**, 509
cellular levels after photodynamic therapy, ^{31}P NMR spectroscopy (murine mammary tumor), **140**, 273
and DNA, binding to urea-transformed native estrogen receptor, evidence for separate binding sites (immature bovine uterus), **139**, 1250
effect on ion-channel activity of Na^+/K^+ -ATPase incorporated into solvent-free phospholipid bilayers (canine renal microsomes), **136**, 767
external, permeabilization of 3T6 cells, **134**, 453
-magnesium and uncomplexed ATP, observation in slow exchange by ^{31}P NMR, **134**, 1379
role in inositol (1,4,5)-trisphosphate interconversion to inositol (1,3,4,5)-tetrakisphosphate and (1,3,4)-trisphosphate (bovine adrenal glomerulosa cells), **139**, 259
stereochemistry in formyltetrahydrofolate synthetase reaction (*Clostridium cylindrosporum*), **139**, 967
stimulation of phosphatidylcholine synthesis in plasma membranes (rat adipocytes), **137**, 1
synthesis by temperature-sensitive mutant, characterization (*Escherichia coli*), **138**, 72
vasodilating effect, effect of bovine serum albumin (rat aortic rings), **138**, 1211
- ATP-ADP translocase
levels in cardiac mitochondria, determination with enzyme inhibitors (chicken, dog, rabbit, rat), **134**, 359
- ATP:AMP phosphotransferase, *see* Adenylate kinase
- ATPase
 Ca^{2+}
and calcium pump, activity in intestinal basolateral membranes, comparison (rat), **141**, 979
detergent-solubilized, from sarcoplasmic reticulum, nucleotide affinity for active site (rabbit muscle), **138**, 652
platelet membrane activity, effect of activating hormones (human), **138**, 215
solubilization and separation from Ca^{2+} -ryanodine receptor complex, **139**, 235
- Ca^{2+} , Mg^{2+} -
activation by calbindin $\text{D}_{28\text{K}}$ (human erythrocyte plasma membranes), **138**, 547
calmodulin-stimulated, direct regulatory effect of cholesterol (rabbit cardiac sarcolemma), **139**, 868
nucleotide-binding domain, amino acid sequence (canine cardiac sarcoplasmic reticulum), **135**, 864
- F_1
effect of anionic detergents (thermophilic bacterium PS3), **136**, 899
kinetic and structural changes with temperature (bovine cardiac mitochondria), **136**, 891
 γ subunit, detection of antibody-binding site (*Escherichia coli*), **137**, 468
- H^+ -
coupling factor I, associated high Mg^{2+} -ATPase activity, induction by octyl glucoside after preactivation with heat and dithiothreitol (spinach chloroplasts), **141**, 636
plasma membrane-associated, activity increase induced by cell proliferation (*Dictyostelium discoideum*), **141**, 1092
- H^+ , K^+ -, amino-terminal sequence and immunological cross-reactivity with Na^+ , K^+ -ATPase (porcine intestine), **138**, 185
- Mg^{2+} -, high activity associated with coupling factor I, induction by octyl glucoside after preactivation with heat and dithiothreitol (spinach chloroplasts), **141**, 636
mitochondrial, role in aerobic metabolism enhancement during perinatal cardiac development (rat), **141**, 319
- Na^+ , K^+ -
activity in renal basolateral membranes, effect of insulin (non-insulin-dependent diabetic rat), **139**, 1313
 α -aminoisobutyric acid transport, basal and stimulated, ouabain effects (rat skeletal muscle), **134**, 1342
immunological cross-reactivity with H^+ , K^+ -ATPase

Subject Index

- (ovine kidney), **138**, 185
- incorporated into solvent-free phospholipid bilayers, ion-channel activity, voltage and ATP effects (canine renal microsomes), **136**, 767
- inhibition by enterolactone (guinea pig, human heart), **134**, 1064
- membrane-bound, solubilization and purification (porcine kidney), **135**, 1008
- ouabain-like inhibitor, purification and characterization (bovine plasma), **135**, 1
- stimulation by insulin (frog skeletal muscle), **134**, 609
- ATPase inhibitors
 - antigenic determinant, localization (bovine heart), **137**, 364
 - purification by high-performance liquid chromatography (*Candida utilis*), **134**, 332
- ATP synthase
 - subunit 8, localization of hydrophilic C terminus (*Saccharomyces cerevisiae*), **138**, 78
- Atrial natriuretic factor
 - analogs with reduced ring size, synthesis and activity profiles, **138**, 880
 - cardiac and plasma levels
 - effects of chronically administered adrenergic and cholinergic agonists (rat), **136**, 510
 - measurement in spontaneously hypertensive and spontaneously hypertensive stroke-prone rats, **135**, 74
 - radioimmunoassay (hyper-, hypothyroid rats), **134**, 178
 - detection in lung (rat), **135**, 515
 - direct radioimmunoassay and clinical evaluation (human plasma), **139**, 1215
 - effect on prostaglandin and renin release by isolated glomeruli (rat kidney), **138**, 496
 - enhancement of cGMP levels in glomeruli and distal tubules (rat kidney), **136**, 947
 - hypothalamic, inhibitory role in vasopressin release after sodium loading (rat), **139**, 1285
 - hypoxia-induced release from isolated heart (rabbit, rat), **140**, 427
 - immunoreactive, low-molecular-weight forms, identification in anterior uvea and retina (rat eye), **134**, 1022
 - induced phosphorylation of placental membrane protein, comparison with guanosine 3':5'-cyclic monophosphate effect (human), **139**, 431
 - inhibition of
 - amiloride-sensitive sodium transport (LLC-PK₁ cells), **134**, 852
 - angiotensin II-stimulated protein phosphorylation (bovine adrenal glomerulosa cells), **140**, 814
 - like immunoreactivity, detection in parasympathetic ganglia (rat), **134**, 279
 - low-molecular-weight, characterization (rat brain), **135**, 728
 - mRNA, regulation by glucocorticoids *in vivo* (rat heart), **139**, 1047
 - phorbol ester-induced secretion, regulation by forskolin, Bay k 8644, and 8-bromo-cyclic GMP (rat heart), **139**, 266
 - plasma levels
 - increase during hypertension (salt-sensitive rat), **137**, 876
 - measurement in salt-sensitive Dahl rats, **136**, 411
 - precursor gene, isolation and nucleotide sequence (bovine), **136**, 396
 - radioimmunoassay
 - in plasma (human), **139**, 287
 - in plasma and urine (human), **137**, 231
 - regulation of steroidogenic responsiveness and cyclic nucleotide levels in Leydig cells *in vitro* (mouse), **138**, 399
 - release from
 - atrial slices, effects of agonists and steroids (rat), **135**, 1090
 - isolated perfused heart, effect of Bay k 8644 (rat), **138**, 1170
 - hypothalamus *in vitro*, radioimmunoassay (rat), **136**, 590
 - removal by perfused lungs *in situ*, measurement (rabbit), **140**, 868
 - segments 99-126 and 101-126, comparative biological activities (bovine, chicken, rabbit, rat), **135**, 987
 - suppression of pro-opiomelanocortin-derived peptide and growth hormone secretion from pituitary cells *in vitro* (rat), **135**, 1035
 - synthesis by cardiac atrioocytes, serum-mediated enhancement (rat), **140**, 151
 - synthetic, effect on arginine vasopressin release by hypothalamo-neurohypophyseal complex in culture (rat), **134**, 652
- α -Atrial natriuretic factor
 - antiparallel and parallel dimers, synthesis and biological properties (human), **141**, 665
 - binding to renal basolateral membranes, hypertension effects (human), **137**, 1079
 - effect on steroidogenesis in cultured bovine adrenocortical cells (human), **137**, 657
- Atrial natriuretic factor receptors
 - association with particulate guanylate cyclase (bovine lung), **140**, 101
 - identification and solubilization (human placenta), **135**, 480
 - negative coupling to adenylate cyclase in atrial and ventricular cardiocytes (rat), **138**, 427
 - regulation in cultured vascular smooth muscle cells (rat heart), **138**, 405
- α -Atrial natriuretic factor receptors
 - distribution in adrenal gland (bovine), **137**, 657
 - molecular size, comparison between kidney, aorta, and adrenal gland by uv-photoaffinity labeling (rat), **136**, 200
- Atriopeptin
 - cardiac and circulating forms, identification and characterization (rabbit), **138**, 1263
 - cardiac and plasma levels, effect of dietary salt and water intake manipulations (rat), **137**, 922
 - release from isolated heart, regulation by α_1 -adrenergic receptors (rat), **137**, 94
- Atriopeptin receptors
 - type-III, high-affinity, detection on pulmonary membranes (rabbit), **140**, 302
- Atropine
 - effect on carbachol-induced pepsinogen secretion from isolated gastric glands, comparison with pirenzepine (guinea pig), **136**, 193
- Auranofin
 - induction of stress proteins (human, murine, rat cells), **138**, 1074
- Aurintricarboxylic acid
 - inhibition of human T-cell lymphotropic virus type III/lymphadenopathy-associated virus cytotoxicity *in vitro*, **136**, 64

Subject Index

- Autolysin
 -defective *Streptococcus pneumoniae* mutant, gene molecular characterization, **137**, 614
- Autolysis
 activation of Ca^{2+} -activated neutral protease (rabbit muscle), **138**, 638
 EDTA-induced, effect of antibiotics (*Escherichia coli*), **134**, 338
- Autophosphorylation
 activity of N-ras proteins *in vitro* (NIH 3T3 cells), **139**, 771
 insulin-like growth factor I receptors, effect of human polyclonal antibody (human placenta, rat liver), **138**, 1023; *erratum*, **140**, 1152
 insulin receptors
 kinetic evidence for activating and nonactivating components, **134**, 1312
 in streptozotocin-induced diabetes, effect of short fast (rat), **140**, 850
 platelet-derived growth factor receptor, effect of protein kinase C activation (human skin, murine Swiss 3T3 fibroblasts), **137**, 343
- 5-Aza-2'-deoxycytidine
 growth inhibitory effects on HL-60 cells resistant to differentiation induction, **141**, 629
- N-4-Azido-2-nitrophenyl aminobutyl NADP⁺
 photolabeling of superoxide radical-generating protein in polymorphonuclear neutrophils (bovine blood), **139**, 85
- B**
- Bacillus subtilis*
 yeast centromeric plasmid application as shuttle vector, **135**, 915
- Bacteriophages
 lambda
 multiply branched DNA molecules, identification as intermediates of postreplicational repair pathway, **137**, 869
 resistant genes from *Brevibacterium albidum* and *Proteus vulgaris*, cloning into *Escherichia coli*, **140**, 1101
 -like particles, induction in pathogenic *Vibrio parahaemolyticus* strain by mitomycin C, **141**, 1249
- Bacteroides fragilis*
 outer membrane permeability, **141**, 292
- Bay k 8644
 effect on
 phorbol ester-induced atrial natriuretic peptide secretion (rat heart), **139**, 266
 secretion of atrial natriuretic polypeptide from isolated perfused heart (rat), **138**, 1170
- Benzodiazepines
 effect on GABA-stimulated chloride uptake by membrane vesicles (rat cerebral cortex), **141**, 1
- Benzo[a]pyrene
 metabolism and mutagenesis *in vitro*, inhibition by resorfin (rat hepatic microsomes), **136**, 555
- Benzoxazinones
 inhibition of serine proteases, effects of electron withdrawal and 5-substitution (human), **140**, 928
- (dl)-3-Benzyl-N-(methanesulfonyloxy) succinimide
 inactivation of human leukocyte elastase and α -chymotrypsin, **141**, 741
- N-Benzoyloxycarbonyl-glycine-proline-diazomethyl ketone
 inhibition of post-proline cleaving enzyme (bovine brain), **134**, 888
- Beryllium sulfate
 modulation of multisubstrate Ca^{2+} - and cAMP-independent kinase (bovine aortic smooth muscle), **136**, 789
- Bicarbonate
¹³C-labeled, exchange kinetics across erythrocyte membrane at equilibrium, NMR (human), **136**, 266
- Bicuculline
 induced status epilepticus, associated enhancement of inositol turnover (rat brain), **136**, 827
- Biopterin
 transient accumulation in human T cells and various cell lines, induction by phorbol ester, **141**, 494
- Biotin
 reactive derivatives, selective labeling of functional groups on erythrocyte membrane proteins and/or glycoproteins (human), **136**, 80
- Bisantrene
 inhibition of angiogenesis (rat avascular cornea), **140**, 901
- 1,3-Bis(2-chloroethyl)-1-nitrosourea, *see* Carmustine
- Bleomycin
 generation of DNA breaks, activation by KHSO_5 (bacteriophage λ), **136**, 1013
- Blood
 whole, leukotriene B₄ production induced by A23187 or serum-treated zymosan, comparison with production by isolated neutrophils (human), **137**, 334
- Blood groups
 P system, detection of galactosyltransferase in p-type lymphoblastoid cells (human), **137**, 1187
- Bombesin
 conformational flexibility and interaction with lipids, **141**, 99
 effect on phosphorylation of acid protein substrate specific for protein kinase C (Swiss 3T3 cells), **140**, 379
 related peptides, specific binding and mitogenic effects, inhibition by substance P antagonist (Swiss 3T3 cells), **137**, 135
- Bone marrow
 reticuloendotheliosis virus T-transformed cells, identification of v-rel oncogene-encoded protein (chicken), **134**, 716
 stem cells, expression of gene transferred by electroporation (mouse), **141**, 1018
- Bones
 fetal, resorption and prostaglandin E₂ production, stimulation by recombinant human interleukin 1 α (mouse), **138**, 618
- Bongkrekic acid
 binding to ADP/ATP carrier, analysis with 6'-O-dansyl- γ -aminobutyl acetylcholine (bovine cardiac mitochondria), **134**, 266
- Botulinum toxin
 type E derivative, and fragments, binding to gangliosides and free fatty acids (*Clostridium botulinum*), **140**, 1015
- Bradykinin
 and D-alanine²-D-leucine⁵ enkephalin, effects on inositol trisphosphate release from NCB-20 cells, **135**, 507

Subject Index

- effect on intracellular Ca^{2+} levels (rabbit papillary collecting tubules), **134**, 299
induction of changes in inositol trisphosphate levels (MDCK cells), **140**, 644
stimulation of inositol 1-phosphate accumulation in chromaffin cells (bovine adrenal gland), **135**, 566
- Brain**
amine levels, effect of sustained exercise (rat), **137**, 149
cortex, detection of cholecystokinin receptors (guinea pig), **137**, 1167
detection of endogenous tetrahydroisoquinoline and 1-methyl-tetrahydroisoquinoline (rat), **140**, 448
nerve growth factor mRNA, localization by *in situ* hybridization (mouse), **138**, 813
- 8-Bromo-cyclic GMP**
regulation of phorbol ester-induced atrial natriuretic peptide secretion (rat heart), **139**, 266
- α -Bungarotoxin receptors**
and acetylcholine-operated ion channel, localization on different molecules (human neuroblastoma cells), **137**, 1141
- Buserelin**
and estradiol and tamoxifen, combined effects on tumor cell growth *in vitro* (human breast cancer), **140**, 550
- 1,3-Butadiene**
epoxidation by chloroperoxidase (*Caldariomyces fumago*), **136**, 778
- Butylated hydroxytoluene**
inhibition of agonist evoked aggregation, ATP secretion, and cytosolic Ca^{2+} increase in aspirinated platelets (human), **139**, 509
- tert*-Butyl hydroperoxide**
-heme complex formed in Fe(III)-tetraphenylporphyrin system, characterization, **135**, 972
induction of
radical production in mitochondria (rat liver), **141**, 1123
vascular relaxation (rat aortic rings), **139**, 102
- Butyl nitrite**
inactivation of methyl cobalamine and 5-methyl tetrahydrofolate, NMR analysis, **135**, 201
- N*-*t*-Butyl- α -phenylnitron**
hydroxyl and superoxide radical adducts, [^{17}O]oxygen hyperfine structure, **141**, 622
- C**
- Cachectin**
-like activity of recombinant human tumor necrosis factor, analysis with monoclonal antibodies, **137**, 847
-tumor necrosis factor, stimulation of prostacyclin synthesis in cultured vascular endothelial cells (human), **141**, 482
- Calbindin $\text{D}_{28\text{K}}$**
stimulation of human erythrocyte plasma membrane Ca^{2+} , Mg^{2+} -ATPase *in vitro* (rat kidney), **138**, 547
- Calcification**
osteoblastic cells *in vitro*, inhibition by β_2 -microglobulin (mouse), **141**, 360
- Calcimedlin**
degradation, calcium effects (rat), **134**, 1269
- Calcineurin**
phosphotyrosyl protein phosphatase activity, regulation by
calmodulin and protease treatment (bovine brain), **140**, 320
- Calcinosis**
experimental arterial, protection by taurine (mouse), **140**, 679
- Calcitonin**
gene-related peptide
partial homology with N-terminal sequence of amyloid fibril protein from insulin-producing pancreatic tumor (human), **140**, 827
structure-activity relationship (human), **134**, 1306
- Calcium**
-activated calmodulin-dependent K^+ channel, identification in adipocyte plasma membranes (rat), **135**, 934
activation of protein kinase C in presence of ganglioside GM3 and 12-*O*-tetradecanoylphorbol-13-acetate, **138**, 865
addition to calmodulin, associated volume changes (bovine testes), **138**, 199
agonist evoked cytosolic increase, inhibition by antioxidants (human platelets), **139**, 509
ATP-dependent pump, and Ca^{2+} -ATPase, activity in intestinal basolateral membranes, comparison (rat), **141**, 979
binding to
apolipoprotein B, analysis (Hep G2 cells), **137**, 493
calmodulin-mastoparan complexes, detection of fluorometrically distinguishable intermediate, **135**, 1050
diphtheria toxin, ESR analysis, **139**, 991
-binding protein, identification in bovine heart, **139**, 596
contractile mechanism sensitivity in chemically skinned skeletal muscle fibers, enhancement by inositol 1,4,5-trisphosphate (rabbit), **135**, 662
cytosolic
homeostasis, role in macrophage secretion inhibition by tetanus toxin (human), **135**, 16
in platelets
effects of thrombin and ionomycin, analysis with fura-2 (human blood), **139**, 308
sodium fluoride-induced increase, inhibition by protein kinase C activation (human), **136**, 381
in smooth muscle cells
agonist-induced transients, measurement with fura-2 (hamster, rabbit, rat), **136**, 927
histamine effects (rat aorta), **135**, 172
norepinephrine effects (rat aorta), **135**, 9
prostaglandin effects (rat aorta), **136**, 247
-dependent inactivation of K^+ currents across photoreceptor membranes, protein kinase C effects (*Hermisenda crassicornis*), **134**, 1245; *erratum*, **140**, 774
-depleted neutrophils
NADPH oxidase activity and phosphoinositide turnover, effect of formyl-methionyl-leucyl-phenylalanine and concanavalin A (human), **140**, 1
phorbol ester-primed, phosphoinositide turnover and NADPH oxidase, independent activation by formyl-methionyl-leucyl-phenylalanine (human), **135**, 785
effect on degradation of calcium-binding proteins (rat), **134**, 1269
- extracellular**
effect on induction of heat shock and glucose-regulated

Subject Index

- proteins (hamster fibroblasts), **140**, 881
- enhancement of 1-oleoyl-2-acetyl-*sn*-glycerol-induced superoxide production (human neutrophils), **134**, 690
- role in
- retinoic acid effect on DNA synthesis in keratinocytes (murine epidermis), **138**, 483
 - sponge cell aggregation (*Microciconia prolifera*), **134**, 1319
- flux in fura-2-loaded platelets, induction by thrombin or ADP, kinetics (human), **136**, 1124
- free
- cytoplasmic levels
 - effects of phenylephrine, vasopressin, and angiotensin II, comparison (rat hepatocytes), **140**, 107
 - measurement in vascular smooth muscle cells by laser microfluorospectrometry (spontaneously hypertensive rat), **141**, 340
 - cytosolic levels, effect of 1,25-dihydroxyvitamin D₃ (avian cartilage cells), **141**, 1236
 - effect on parathyroid hormone secretion in electroporated parathyroid cells (bovine), **139**, 1233
 - intracellular levels in cultured chromaffin cells, carbamylcholine- and high K⁺-induced elevation, differential effects of protein kinase C activators (bovine adrenal), **139**, 903
- high-affinity binding to actin monomers, **135**, 607
- induction of
- alterations in proteolytic enzyme levels and distribution in erythrocytes (human), **138**, 87
 - phase separation of phosphatidylserine-containing liposomes, potentiation by micromolar levels of Zn²⁺ (bovine brain), **136**, 476
- inositol-trisphosphate, and cAMP signaling systems, dual activation by adrenocorticotropin (rat adrenal cells), **135**, 742
- intracellular
- bradykinin-induced increase (rabbit papillary collecting tubules), **134**, 299
 - effects of carbamylcholine, fluoride, prostaglandin F_{2α}, and TRH (canine thyroid cells), **141**, 569
 - heat shock-induced increase (CHO cells), **137**, 826
 - in isolated perfused heart, effects of ionophores and adrenergic agents (rabbit), **139**, 816
 - kyotorphin-induced increase (rat cerebral synaptosomes), **137**, 897
 - release and uptake by antibody-activated B cells, analysis with indo-1 (mouse), **137**, 500; *erratum*, **138**, 503
 - requirement for zymosan-stimulated phospholipase C activation (murine peritoneal macrophages), **134**, 367
 - role in phorbol ester-induced contraction of thoracic aorta (rabbit), **138**, 1362
- levels in
- B cells, effect of antigen-induced activation, **140**, 1079
 - monocytes, effect of κ elastin (human), **141**, 92
- and magnesium, sequential binding to bovine prothrombin fragment 1, associated spectral response, **141**, 1207
- mitochondrial transport, side effects of membrane potential probe safranin (rat liver), **135**, 189
- mobilization
- by 5,6-epoxyeicosatrienoic acid in anterior pituitary cells (rat), **139**, 1188
 - induction by formyl-methionyl-leucyl-phenylalanine, inhibition by phorbol ester (human neutrophils), **135**, 556
 - in neutrophils, formyl-methionyl-leucyl-phenylalanine-induced, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
 - phorbol ester-mediated suppression, reversal by protein kinase C inhibitor (thrombin-activated human platelets), **134**, 868
 - in platelets, cGMP effects (human), **135**, 1099
 - and phorbol ester, synergistic effect on K⁺-induced catecholamine release from chromaffin cells (bovine adrenal gland), **140**, 990
 - regulation of parathyroid hormone secretion, effect of pertussis toxin treatment (bovine parathyroid cells), **138**, 960
 - release from
 - endoplasmic reticulum, inositol trisphosphate-induced, inhibition by calmodulin (rat pancreatic islets), **141**, 418
 - erythrocyte membranes, induction by phosphorylation (human), **136**, 288
 - permeabilized mastocytoma cells, induction by inositol 1,4,5-trisphosphate (mouse), **135**, 46
 - role in
 - inositol (1,4,5)-trisphosphate interconversion to inositol (1,3,4,5)-tetrakisphosphate and (1,3,4)-trisphosphate (bovine adrenal glomerulosa cells), **139**, 259
 - pancreatic acinar cytotoxic injury induced by toxic agents (rat), **139**, 530
 - pepsinogen secretion from dispersed chief cells (guinea pig gastric chief cells), **137**, 281
 - ryanodine, receptor complex, solubilization and separation from Ca²⁺-ATPase, **139**, 235
 - spermatozoal levels before/after *in vitro* capacitation (rabbit), **134**, 664
 - transients in cultured smooth muscle cells, inhibition by pertussis toxin (rat aorta), **140**, 195
 - translocation in thyroid cells, effects of carbamylcholine, fluoride, prostaglandin F_{2α}, and TRH (dog), **141**, 569
 - transport by ionophorous peptides, detection in lymphocytes by quin-2 fluorescence (dog, human), **134**, 731
 - uptake by
 - chromaffin cells *in vitro*, inhibition by potassium (bovine adrenal gland), **134**, 1
 - intracellular compartments in permeabilized enterocytes, effect of inositol 1,4,5-trisphosphate (rat intestine), **139**, 612
 - neuronal cells, effect of dihydropyridine (rat brain), **139**, 581
 - thyroid cells, stimulation by thyrotropin (swine), **136**, 306
 - and zinc, micromolar levels, synergistic effects on membrane fusion (bovine brain), **137**, 101
- Calcium channels
- antagonist binding, inhibition by phorbol esters and protein kinase C (PC12 neural cells), **136**, 1049
 - and apamin-sensitive Ca²⁺-activated K⁺ channels in muscular dysgenic skeletal muscle cells, action potentials and contractile activity (mouse), **136**, 935
 - coupling to muscarinic receptors, induction by point-mutated p21^{ras} (murine fibroblasts), **141**, 591
 - neuronal

Subject Index

- 1,4-dihydropyridine component, cytolocalization and subunit structure identification with polyclonal antibodies (rabbit brain, skeletal muscle), **139**, 996
voltage-sensitive, effect of dihydropyridine agonists and antagonists (rat brain), **134**, 1038
protein kinase C regulatory role (PC-12 pheochromocytoma cells), **134**, 1298
- Calcium chloride
alteration of signal I_2 in oxygen-evolving photosystem II preparations (spinach chloroplasts), **134**, 272
- Calcium ionophores
activity of prostaglandin B_1 oligomeric derivatives, evaluation, **137**, 215
- Caldesmon
phosphorylation by protein kinase C (chicken smooth muscle), **132**, 56; *erratum*, **134**, 1029
regulation of smooth muscle actomyosin ATPase, **136**, 962
- Calmodulin
binding to
mastoparan, ^1H NMR analysis, **137**, 1069
retinal heat shock protein, analysis (rabbit), **139**, 974
-binding protein, 34-kDa, isolation and characterization (chicken gizzard smooth muscle), **141**, 20
 Ca^{2+} addition, associated volume changes (bovine testes), **138**, 199
degradation, calcium effects (rat), **134**, 1269
-dependent Ca^{2+} -activated K^+ channel, identification in adipocyte plasma membranes (rat), **135**, 934
effect on calcineurin phosphotyrosyl protein phosphatase activity (bovine brain), **140**, 320
effects of dequalinium, propylinium, and rhodamine (rat C6 astrocytoma cells), **135**, 574
gene, increased expression in dystrophic muscle (chicken), **137**, 507
inhibition by bisbenzylisoquinoline derivative, **140**, 461
inhibition of inositol trisphosphate-induced Ca^{2+} release from endoplasmic reticulum (rat pancreatic islets), **141**, 418
interaction with
bee venom melittin (bovine testis), **135**, 419
plasma membranes (rat liver), **135**, 323
lanthanide binding order, EPR study, **138**, 1243
-like activity, detection in purified retroviruses, **141**, 1077
-like protein, detection in *Bacillus subtilis*, **134**, 212
-mastoparan complexes, Ca^{2+} binding: detection of fluorometrically distinguishable intermediate, **135**, 1050
stimulation of cardiac sarcolemmal calcium pump, direct regulatory effect of cholesterol (rabbit), **139**, 868
thyroid cellular levels, thyrotropin-stimulated increases (swine), **136**, 306
- Calpain
-endogenous calpain inhibitor system, defective, in erythrocytes, characterization (Milan hypertensive rat), **139**, 341
types I and II
activation of tyrosine hydroxylase (bovine adrenal medulla), **134**, 749
isolation and analysis of phosphorylation *in vivo* (human K-562 cells), **136**, 1090
- Calpain inhibitor
activity decrease in red blood cells from Milan hypertensive rats, **138**, 1370
- Calspectin
-actin interaction, role of erythrocyte protein 4.1-related factor (bovine brain), **140**, 1051
- Calvarium
neonatal, resorption, stimulation by homogeneous interferon β -inducing 22K factor (mouse), **139**, 1150
- Captopril
binding to angiotensin-converting enzyme, inhibition by peptide analogs (rat lung), **135**, 770
- Carbachol
effect on
atrial natriuretic factor release by atrial slices (rat), **135**, 1090
cardiac and plasma atrial natriuretic factor levels after chronic administration (rat), **136**, 510
induction of
inositol 1-phosphate accumulation, effects of lithium and phorbol esters (NCB-20 cells), **136**, 622
pepsinogen secretion from isolated gastric glands, effects of pirenzepine and atropine, comparison (guinea pig), **136**, 193
stimulation of inositol 1-phosphate accumulation in chromaffin cells (bovine adrenal gland), **135**, 566
vasodilating effect, effect of bovine serum albumin (rat aortic rings), **138**, 1211
- Carbamylcholine
effect on Ca^{2+} intracellular levels and translocation (canine thyroid cells), **141**, 569
induced elevation of intracellular free calcium concentration in cultured chromaffin cells, differential effects of protein kinase C activators (bovine adrenal), **139**, 903
- Carbohydrates
content in A_1 adenosine receptors (hamster, rat brain), **140**, 406
- β -Carbolines
effect on GABA-stimulated chloride uptake by membrane vesicles (rat cerebral cortex), **141**, 1
- Carbon
methyl group, in thiamine pyrimidine moiety, origin during thiamine biosynthesis (*Salmonella typhimurium*), **134**, 1136
- Carbon monoxide
effect of low levels on EPR spectrum of periplasmic hydrogenase (*Desulfovibrio vulgaris*), **137**, 1086
- Carbon tetrachloride
intoxication, effect on ^{23}Na NMR spin-lattice relaxation times in various tissues (rat), **134**, 1079
- Carboxyatractyloside
binding to ADP/ATP carrier, analysis with 6'-*O*-dansyl- γ -aminobutyryl atractyloside (bovine cardiac mitochondria), **134**, 266
- 2-Carboxybenzaldehyde reductase
induction by phenobarbital (rat hepatocytes), **141**, 488
- 20-Carboxy-leukotriene B_4
and leukotriene B_4 , simultaneous quantitation by gas chromatography-mass spectroscopy (human neutrophils), **134**, 512
- Carboxylic acid
-retinal Schiff base hydrogen bonds, proton transfer thermodynamics, **138**, 819
- Carboxy-nifedipine
synthesis and application in affinity resin preparation for

Subject Index

- nitrendipine receptor, **134**, 922
- Carboxypeptidase A
precursor, primary structure of activation segment (porcine pancreas), **141**, 517
- Carboxypeptidases
renal, generation of angiotensin II (human), **138**, 573
- Carcinogenesis
human, analysis: coal tar metabolism in human skin grafted onto athymic nude mice as model system, **138**, 33
skin, evaluation of cholera toxin as promoter (mouse), **137**, 486
- Carcinogens
induced DNA strand breaks, detection by gravity-flow alkaline elution (MDBK cells), **137**, 1006
- Cardiomyopathy
effect on cardiac mitochondrial NADH-ubiquinone oxidoreductase subunit levels (human), **140**, 88
estrene treatment effects (genetically diabetic mouse), **139**, 56
- Carmustine
reaction with guanosine: evidence for mechanism of DNA modification, **139**, 31
- L-Carnitine
and halothane, antagonistic effects on mitochondrial energy-linked processes, role of long-chain acyl CoA (rat liver), **139**, 303
- Carnitine acetyltransferase
inhibition by hemiacetylcarnitinium (pigeon breast), **138**, 735
- Carnitine palmitoyltransferase
effect of starvation and hypothermia (newborn rat brown adipose tissue, liver), **135**, 589
- β -Carotene
association with regression of buccal pouch carcinoma and induction of macrophage tumor necrosis factor (hamster), **136**, 1130
- Cartilage
resorption, stimulation by homogeneous interferon β -inducing 22K factor (bovine nose), **139**, 1150
- Casein
gene expression, hydrocortisone transcriptional and post-transcriptional roles (rat mammary explants), **134**, 812
- β -Casein
phosphorylation sites, localization by fast atom bombardment mass spectrometry (buffalo), **140**, 28
- Casein kinase
erythrocyte type A, phosphorylation of protein tyrosine (human), **139**, 855
type I
and cAMP-dependent protein kinase, comparative effects on glycogen synthase behavior as substrate for hepatic synthase phosphatases, **139**, 1033
phosphorylation of isolated plasma membranes (AH-66 hepatoma ascites cells), **138**, 95
type II, -phosphorylated ribosomal proteins, identification (cryptobiotic *Artemia* sp. gastrulae), **135**, 347
- CAT₁₂, *see* 4-(Dodecyl dimethyl ammonium)-1-oxy-2,2,6,6-tetramethyl piperidine bromide
- Catalase
activity in muscle, liver, and lung tissues of inherited muscular dystrophy-affected chickens, **134**, 550
prevention of doxorubicin-induced killing of breast cancer cells (human), **135**, 330
- Cataracts
formation, associated covalent change of lens membrane major intrinsic polypeptide (human), **135**, 965
- Catecholamines
release from chromaffin cells, K⁺-induced, synergistic effect of Ca²⁺ and phorbol ester (bovine adrenal gland), **140**, 990
- Cathepsin L
protease specificity, comparison with major excreted protein of transformed cells (mouse), **139**, 156
- Cauliflower mosaic virus
DNA, detection of potential transcription control sequence by dinucleotide primed *in vitro* transcription, **138**, 17
- CB 954
formation of DNA interstrand crosslinks in Walker tumor cells (rat), **140**, 1073
- Cell cultures
long-term, myocytes, effect on myosin isozyme pattern (rat heart), **137**, 855
nonneural cells derived from fetal spinal cord, nitrendipine binding (mouse), **136**, 783
primary, periportal and perivenous hepatocytes, retention of zonal characteristics (rat), **139**, 1055
steady-state, temperature effects (*Escherichia coli*), **135**, 708
- Cell cycle
heat production as monitoring parameter (murine breast cancer cells), **140**, 414
- Cell fusion
role of electroporation (Chinese hamster ovary cells), **140**, 258
- Cell lines, *see also* Tumor cells
BHK, CCL 39, and Raji, growth process, role of 5'-deoxy-5'-methylthioadenosine and L-homocysteine metabolism to methionine, **135**, 131
- CHO (Chinese hamster ovary)
hydroxyurea uptake by diffusion mechanism, **134**, 1254
insulin and insulin-like growth factor I binding, effect of glucose deprivation, **140**, 821
- HOS (human osteosarcoma), transformation by Kirsten murine sarcoma virus, induction of ouabain sensitivity, **136**, 807
- hybrid murine RAW264-TG3-human leukocyte, chemotactic response to N-formyl-methionyl-leucyl-phenyl-alanine, **138**, 1001
- insulin-like growth factor II-producing, identification and characterization of membrane receptors for insulin-like growth factor II, **134**, 595
- Jurkat JMN, transient accumulation of pteridines, induction by phorbol ester, **141**, 494
- L cells (murine fibroblasts)
interferon- β binding, enhancement by lanthanide ions, **137**, 480
transformed with human interferon- γ cDNA gene, extracellular and intracellular expression of human INF- γ , **135**, 894
- NRK (normal rat kidney), K-ras-transformed, loss of epidermal growth factor binding, mechanism, **135**, 435
- Ob17, transformed by polyoma virus middle-T-only gene, extracellular intermediates, promotion of Ob17 preadipocyte proliferation and differentiation, **137**, 378

Subject Index

- Reh, transient accumulation of pteridines, induction by phorbol ester, **141**, 494
- TA1 (fibroblast/adipocyte), responsiveness to insulin and insulin-like growth factor receptors before and after differentiation, **137**, 552
- 293, transient accumulation of pteridines, induction by phorbol ester, **141**, 494
- Vero (African green monkey kidney)
cytotoxin purification and physicochemical properties (*Escherichia coli* 0157:H7), **139**, 424
membrane permeability changes induced by *Clostridium perfringens* type A enterotoxin in absence of extracellular Ca^{2+} , **141**, 704
- Cellulase
catalyzed hydrolysis, stereochemical course, ^1H NMR determination (*Cellulomonas fimi*), **139**, 487
extracellular, purification and partial characterization (*Cellulomonas fermentans*), **140**, 219
- Cellulose
oxidation by hydrogen peroxide, role in cell wall breakdown (orange), **141**, 238
- Ceramide
-oligosaccharide linkage in glycosphingolipids, identification and characterization of cleaving enzyme (leech), **141**, 346
- Ceramide glycanase
glycosphingolipid-cleaving, identification and characterization (leech), **141**, 346
- Cerebrospinal fluid
cyclo(His-Pro)-like immunoreactivity, distribution and characterization (human), **136**, 835
- Ceruloplasmin
copper from, binding and uptake by plasma membrane (rat tissues), **139**, 822
desialation by hepatic endothelium (rat), **140**, 94
- Cesium fluoride
high concentrations, differential effects on behavior of octadeoxynucleotides d(A-T)₄ and d(T-A)₄, **139**, 1158
- Chemoattractants
oxidation by responding cells, effect on efficacy (human neutrophils), **138**, 66
receptor signal transduction, mediation by pertussis/cholera toxin-sensitive N protein (human), **138**, 887
- Chemoreceptors
demethylation, inhibition by attractant stimuli in absence of regulatory domain of demethylating enzyme (*Salmonella typhimurium*), **141**, 918
- Chemotactic factors
neutrophil, generation by ethanol-metabolizing hepatocytes (human), **137**, 688
- Chemotaxis
fibroblasts and astroglial cells, stimulation by bovine brain-derived growth factor (bovine, human, rat), **141**, 67
human-mouse hybrid cells, to *N*-formyl-methionyl-leucyl-phenylalanine, **138**, 1001
neutrophils, formyl-peptide-induced, inhibition by myelin basic protein (human), **141**, 524
- Chief cells
intracellular calcium levels and pepsinogen secretion, effect of A23187 and ionomycin (guinea pig stomach), **137**, 281
- Chloramphenicol
effect on EDTA-induced autolysis (*Escherichia coli*), **134**, 338
- Chlorate
inhibition of protein sulfation (rat PC12 cells), **141**, 870
- Chloride
efflux from normal and cystic fibrosis fibroblasts, kinetic analysis (human), **134**, 233
GABA-stimulated uptake by membrane vesicles, effect of benzodiazepines and β -carbolines (rat cerebral cortex), **141**, 1
-sodium-potassium cotransport system, biochemical and pharmacological properties (chicken cardiac cells), **134**, 326
- Chloride peroxidase
epoxidation of alkenes (*Caldariomyces fumago*), **136**, 778
- 3-Chloroperoxybenzoic acid
induction of vascular relaxation (rat aortic rings), **139**, 102
- Chlorophyll *a*
photogeneration of superoxide ion *in vitro* in water/acetone, ESR study (spinach), **139**, 1124
- Chloroquine
binding to muscarinic receptors in pancreatic acini, effect on secretory responses (rat), **137**, 664
- N*-Chlorosulfonyl dicyclohexylamine
as spermidine synthase inhibitor, synthesis and effects on Molt4B cells, **141**, 718
- S*-(2-Chloro-1,1,2-trifluoroethyl)-L-cysteine
metabolism to hydrogen sulfide, role in induced mitochondrial toxicity (rat kidney), **138**, 707
- Chlorpromazine
effect on inositol incorporation into phosphoinositides and inositol phosphates (rat C6 glioma cells), **135**, 465
regulation of fluoropyrimidine metabolism (murine sarcoma cells), **138**, 1009
- Chlorpropamide
effect on insulin release from cloned B-cell line HIT-T15 (hamster), **140**, 616
- Chlorsulfuron
cytochrome *P*-450-dependent metabolism, induction by sulfonylurea (*Streptomyces griseolus*), **140**, 650
- Cholate
effect on F_1 -ATPase activity (thermophilic bacterium PS3), **136**, 899
selective inhibition of microsomal hepatic cholesterol epoxide hydrolase (mouse), **140**, 632
- Cholchicine
effect on insulin release from cloned B-cell line HIT-T15 (hamster), **140**, 616
- Cholecystokinin
amino acid sequence homology with leucosulfakinin II (*Leukophaea maderae*), **140**, 357
C-terminal octapeptide, *see* Sincalide
plasma levels, increase after intestinal administration of pancreatic enzyme secretion-stimulating peptide, radioimmunoassay (rat), **136**, 701
- Cholecystokinin receptors
detection in cerebral cortex (guinea pig), **137**, 1167
- Cholera toxin
adenylate cyclase sensitivity, increase by fasting, mechanism (rat fat cells), **138**, 1411
-catalyzed ADP-ribosylation
efficiency improvement by NADP (canine cardiac, rat

Subject Index

- hepatic membranes), **137**, 328
 substrate identification on cardiac cell membranes (rat), **138**, 1355
 GTP-activated adenylate cyclases as substrates, comparison, **134**, 928
 induction of epidermal hyperplasia: evaluation of tumor-promoting activity in skin carcinogenesis (mouse), **137**, 486
 sensitive N protein, mediation of chemoattractant receptor signal transduction (human), **138**, 887
- Cholesterol**
 direct regulatory effect on calmodulin-stimulated Ca^{2+} pump (rabbit cardiac sarcolemma), **139**, 868
 esterification, associated apolipoprotein A-IV affinity for lipoproteins (human), **135**, 756
 -lysophosphatidylcholine vesicles, structure, ^{31}P NMR spectroscopy, **139**, 25
- Cholesterol epoxide hydrolase**
 microsomal, selective inhibition by detergents and separation from xenobiotic epoxide hydrolase (murine liver), **140**, 632
- Cholesterol esterase**
 -catalyzed hydrolysis, mechanism and inhibition, analysis with phenyl-*n*-butylborinic acid, **134**, 386
- Cholesterol 7 α -hydroxylase**
 modulation by alkaline phosphatase and protein kinase (rat), **134**, 797
- Cholestyramine**
 stimulation of microsomal squalene synthetase, effect of diurnal cycle (rat liver), **138**, 335
- Cholic acid**
 conjugation with taurine and glycine by hepatic peroxisomes (rat), **138**, 167
- Chondrocytes**
 articular, stimulation by homogeneous interferon β -inducing 22K factor (human), **139**, 1150
 costal, synthesis of glycosaminoglycans, effect of human recombinant interleukin 1 (rat), **140**, 386
 embryonic sternum, identification of disulfide-bonded type X procollagen (chicken), **138**, 835
 expression of interleukin 1 genes: mechanism for autocrine control of cartilage matrix degradation (bovine, human), **141**, 904
 neonatal articular, proteoglycan synthesis *in vitro*, enhancement by interleukin 1-like factor (human), **134**, 539
- Chondroitin sulfate**
 E form, anticoagulant properties (human mast cells), **137**, 15
 oversulfated, synthesis in cultured peritoneal macrophages (human), **139**, 377
 sulfation, inhibition by protein kinase C activators (human promyelocytic leukemia cells), **135**, 909
- Chromaffin cells**
 calcium uptake and noradrenaline release *in vitro*, inhibition by potassium (bovine adrenal gland), **134**, 1
 granule membranes, $\text{p60}^{\text{c-src}}$ activity detection (bovine), **134**, 736
 inositol 1-phosphate accumulation, stimulation by bradykinin, carbachol, and histamine (bovine adrenal gland), **135**, 566
 intracellular free calcium levels, carbamylcholine- and high K^{+} -induced elevation, differential effects of protein kinase C activators (bovine adrenal), **139**, 903
 release of catecholamines, K^{+} -induced, synergistic effect of Ca^{2+} and phorbol ester (bovine adrenal gland), **140**, 990
- Chromate**
 reduction by aldehyde oxidase (rabbit liver), **137**, 8
- Chromatin**
 -histone integration, monitoring with double spin-labeled H1 and H5 (avian erythrocytes), **141**, 434
 nerve growth factor receptor identification with monoclonal antibody (human melanoma cells), **140**, 174
- Chromogranin A**
 cDNA, detection of 2.3kb mRNA in diverse neuroendocrine tissues (human), **137**, 418
- Chromoproteins**
 hemolymph, larval specific, purification and characterization (*Heliothis zea*), **134**, 580
- Chromosomes**
 assignment of genes for gastrin, thyrotropin β -subunit, and *c-erbB-2* by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
 11, localization of serum amyloid A gene (human), **137**, 1196
 localization of
 galactosyltransferase gene (human), **141**, 1185
 genes for acidic and basic fibroblast growth factors (human), **138**, 644
 metaphase, deoxyribonuclease I digestion, release of base pair DNA fragments 300 bp long (Chinese hamster V79 cells), **138**, 413
 pepsinogen A genes, close linkage (human), **138**, 289
 20, PrP gene on p arm, localization of homologous human gene (scrapie-infected hamster brain), **140**, 758
- Chymotrypsin**
 binding by α_1 -protease inhibitor, comparison with trypsin, **138**, 925
 effect on
 calcineurin phosphotyrosyl protein phosphatase activity (bovine brain), **140**, 320
 microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
 inhibition by benzoxazinones, effects of electron withdrawal and 5-substitution (human), **140**, 928
 type α , inactivation by (*dl*)-3-benzyl-*N*-(methanesulfonyloxy) succinimide, **141**, 741
- Cilia**
 olfactory, phosphoinositide turnover, stimulation by L-alanine or GTP (catfish), **137**, 36
- Cirsiliol**
 antiproliferative effects on leukemic cell lines (human), **140**, 832
- Cisplatin**
 high-melting adduct, synthesis, **138**, 758
 inhibition of DNA synthesis in leukemia cells, effect of serum (mouse), **138**, 631
- Citrate synthase**
 structural gene, molecular cloning (*Acinetobacter anitratum*), **141**, 797
- Clathrin**
 and hydrophobic domain of boundary membranes in endocytic vesicles, close association (bovine brain), **141**, 878

Subject Index

- Cloacin
cleavage of 16 S RNA 3'-terminal fragment (*Streptomyces aureofaciens*), **137**, 80
- Clomiphene
induction of DNA strand breaks (human leukemia cells), **138**, 1203
- Clonidine
-displacing substance, inhibition of vas deferens twitch response (rat brain), **134**, 184
enhancement of β -adrenergic ligand binding to pineal particulate fractions (rabbit), **138**, 1196; *erratum*, **139**, 1320
and growth hormone-releasing hormone, comparative effects on stimulation of growth hormone synthesis in neonatal pituitary (rat), **138**, 1223
 ^3H -labeled, specific binding in brown adipose tissue, α_2 -adrenergic effects (fetal rat), **138**, 1390
- Cloning
 α -amylase inhibitor gene from *Streptomyces griseosporus* in *S. lividans*, **141**, 1099
arginase, in *Escherichia coli* (rat liver), **136**, 955
cDNA encoding
catalytically active DNA polymerase α fragment (bovine thymus), **140**, 21
polypeptide similar to human insulin receptor precursor (*Drosophila* sp.), **141**, 474
reovirus serotype 1 Lang strain mRNA encoding major capsid surface polypeptide $\sigma 3$, **136**, 183
minor capsid polypeptides $\sigma 1a$ and nonstructural $\sigma 1bNS$, **140**, 508
 σNS polypeptides, **139**, 845
lipase gene from *Pseudomonas fragi*, **141**, 185
mitochondrial DNA fragment from *Lentinus edodes* in *Saccharomyces cerevisiae*: analysis of autonomous replication capability, **138**, 1110
mitochondrial glutamic-oxaloacetic transaminase mRNA (rat), **134**, 803
nerve growth factor genes: mRNA distribution in developing and adult tissues (chicken), **141**, 1116
phage λ -resistant genes from *Brevibacterium albidum* and *Proteus vulgaris* into *Escherichia coli*, **140**, 1101
proteinous α -amylase inhibitor gene from *Streptomyces griseosporus* in *S. lividans*, **141**, 1099
structural gene for citrate synthase (*Acinetobacter anitratum*), **141**, 797
- Coal tar
metabolism in human skin grafted onto athymic nude mice: as model system for analysis of carcinogenesis, **138**, 33
- Cobalt
-substituted basic blue protein plantacyanin, preparation and spectroscopic analysis (cucumber), **139**, 961
-substituted lectin, circular dichroism (lentil), **136**, 438
- Codeine
concentration estimation by comparative immunoremoval (bovine tissues), **137**, 316
- Coenzyme A
localization in purified peroxisomes: association with matrix protein (rat liver), **139**, 1195
- Collagen
interaction with proteodermatan sulfate, role of disulfide bridges in protein core (bovine skin), **138**, 1348
processing in *ras*-transfected mammary epithelial cells (mouse), **137**, 161
production and processing in fibroblasts, stimulation by transforming growth factor- β (human), **138**, 974
type III, gene expression, stimulation by *v-fos* (NIH 3T3 cells), **136**, 1042
type V, from FBJ virus-induced osteosarcoma, staining with concanavalin A (mouse), **136**, 857
type X
biosynthesis, spatiotemporal changes during development (embryonic chick sternum), **137**, 1125
translation *in vitro*, in system with pancreatic microsomes (chicken), **136**, 109
- Collagenase
cleavage site specificity, analysis, **139**, 1184
inhibition by peptide hydroxamic acids (human skin), **136**, 390
perfusion of hepatocytes, effect on membrane transferrin receptors (rat), **134**, 711
- Colloids
intraglandular, of intermediate lobe origin, detection of Met-enkephalin (bovine pituitary), **139**, 480
- Colony-stimulating factor
expression during monocytic differentiation of HL-60 cells, **141**, 924
macrophage-specific
purification from medium conditioned by pancreatic carcinoma cells (human), **138**, 238
stimulation of glucose uptake in bone marrow-derived cells (mouse), **138**, 445
production in MC3T3-E1 cells, stimulation by recombinant human interleukin-1 α and β , **141**, 285
synthesis by MC3T3-E1 osteoblasts (mouse), **134**, 400
- Complement
C3 and C4, processing and secretion, effects of acidotropic compounds (HepG2 cells), **136**, 603
C3, serine proteinase cleaving α and β chains, purification from erythrocyte membranes and partial characterization (human), **140**, 1113
- Complement inhibitors
C1, cDNA isolation and sequencing (human), **137**, 620
- Compound 48/80
induced histamine secretion in peritoneal mast cells, effect of phorbol ester pretreatment (rat), **137**, 1112
- Concanavalin A
and formyl-methionyl-leucyl-phenylalanine, effect on NADPH oxidase activity and phosphoinositide turnover in Ca^{2+} -depleted neutrophils (human), **140**, 1
induction of microvillous fusion and pinching-off (rat colon), **140**, 766
opsonization of phosphatidylinositol-containing liposomes, stimulation of phosphatidylinositol uptake by macrophages, **138**, 1090
-polysaccharide interaction, effect on K^+ currents through lipid bilayer membranes, **141**, 912
precipitation by high-mannose-type glycopeptide, stoichiometry (jack bean), **137**, 670
solutions, detection and quantitation of polysaccharides with electrical oscillator, **137**, 978
staining of type V collagen from FBJ virus-induced osteosarcoma (mouse), **136**, 857
stimulation of phosphatidylinositol turnover in thymocytes,

Subject Index

- effect on activation (mouse), **141**, 657
- Conductivity
in ultrathin films of serum albumin and dinitrophenylated derivatives, direct current measurements (bovine), **133**, 1152; *erratum*, **135**, 1150
- Conotoxin G1
surface-adsorbed, reaction products, mass spectrometry, **134**, 420
- Copper
-albumin complexes in bloodstream, functional role, **135**, 1064
-ascorbic acid system, selective oxidation of imidazole ring in histidine residues, **138**, 659
associated with L-ascorbate oxidase active site, effect of enzyme reduction (cucumber), **135**, 644
ceruloplasmin, binding and uptake by plasma membrane (rat tissues), **139**, 822
content in cytochrome-c oxidases, analysis by proton-induced X-ray emission (bovine, porcine, rat tissues), **140**, 1007
-cytokinin complexes, superoxide dismutase mimetic activity, **137**, 372
induction of luteinizing hormone release and desensitization of pituitary gonadotropes (rat), **136**, 417
-peroxydisulfate system, biomimetic oxidation of lignin model compounds, **139**, 1104
- Corpus luteum
regression, preservation of gonadotropin receptor subunit structure (rat), **137**, 546
- Corticosterone
plasma levels, effects of α -melanocyte-stimulating hormone *in vivo* (rat), **141**, 1279
uptake and metabolism by isolated hepatocytes (nonadrenalectomized rat), **135**, 160
- Corticotropin-releasing factor
binding to peripheral tissues, effect of adrenalectomy and glucocorticoid treatment (rat), **134**, 255
erythrocyte membrane binding sites, identification and modulation by chronic ethanol treatment (human, rat), **136**, 137
- Cortisol, *see* Hydrocortisone
- Cortisone
cytochrome P-450_{11 β} -dependent formation from cortisol (bovine), **136**, 369
- Coupling factor 1, *see* ATPase, H⁺-
- Creatine kinase
association with mitochondrial inner membrane surface in physiological salt solution, relative importance of salt composition and ionic strength (rat heart), **139**, 1262
B subunit, mRNA nucleotide sequence: coding and 3' non-coding region homology among species (canine brain), **138**, 392
levels in cardiac mitochondria, determination with enzyme inhibitors (chicken, dog, rabbit, rat), **134**, 359
mitochondrial, role in aerobic metabolism enhancement during perinatal cardiac development (rat), **141**, 319
M subunit, cDNA isolation and sequencing (human), **140**, 981
- Crystallization
two-dimensional, phospholipid antibodies, induction by supported phospholipid bilayers, **134**, 819
- Cucumber mosaic virus
satellite CARN 5, cDNA clones, enzymatic transcription to infectious RNAs, **135**, 290
- Cumene hydroperoxide
induction of vascular relaxation (rat aortic rings), **139**, 102
- Cyanate
effect on tropomyosin end-to-end polymerization (rabbit muscle), **134**, 64
- α -Cyanocinnamate
-sensitive pyruvate transporter, extraction from mitochondria and reconstitution (castor bean), **140**, 1036
- Cyanopindolol
¹²⁵I-labeled, binding to pineal particulate fractions, enhancement by melatonin and α -adrenergic compounds (rabbit), **138**, 1196; *erratum*, **139**, 1320
- Cyclic AMP
accumulation in pituitary tumor cells, induction by bacterial adenylate cyclase (rat), **136**, 463
and ACTH, comparative effects on epinephrine-induced melanosome aggregation (goldfish), **138**, 1081
activation of thyroglobulin gene in hormonally depleted FRTL-5 cells, **141**, 1261
cellular levels, regulation by atrial natriuretic factor *in vitro* (murine Leydig cells), **138**, 399
effect on
expression of *v-ras*^H oncogene linked to murine mammary tumor virus promoter, **134**, 436
formyl-methionyl-leucyl-phenylalanine-induced inositol phospholipid metabolism and Ca²⁺ mobilization (guinea pig neutrophils), **139**, 1272
parathyroid hormone secretion in electroporated parathyroid cells (bovine), **139**, 1233
and glucocorticoids, synergistic regulation of preproenkephalin mRNA levels in neuroblastoma-glioma hybrid cells, **139**, 1
and inositol-trisphosphate-Ca²⁺ signaling systems, dual activation by adrenocorticotropin (rat adrenal cells), **135**, 742
metabolism by thrombopathic platelets, effects of forskolin and 1-methyl-3-isobutyl-xanthine (dog), **140**, 595
modification of androgen receptors *in vitro* (murine skeletal muscle), **135**, 1069
monocyte levels
effect of κ elastin (human), **141**, 92
synergistic action of adenosine and formyl-methionyl-leucyl-phenylalanine (human), **138**, 1376
occurrence in archaeobacteria, **136**, 1078
production in brain, effect of phorbol esters (rat), **139**, 251
regulation of acetylcholine-activated channel life time in cultured myotubes (chicken embryo), **140**, 243
- Cyclic-AMP phosphodiesterase
in butyrate-sensitive and resistant mastocytoma cells, differential effects of sodium butyrate, **138**, 1030
- Cyclic GMP
cellular levels, regulation by atrial natriuretic factor *in vitro* (murine Leydig cells), **138**, 399
effects on serotonin secretion, inositol trisphosphate production, and Ca²⁺ mobilization in platelets (human), **135**, 1099
induced phosphorylation of placental membrane protein, comparison with atrial natriuretic factor effect (human), **139**, 431
inhibition of amiloride-sensitive sodium transport (LLC-PK₁ cells), **134**, 852

Subject Index

- monocyte levels, effect of κ elastin (human), **141**, 92
nephron segment levels, enhancement by atrial natriuretic peptide (rat kidney), **136**, 947
- Cycloadenosine
R- and *S*-8,5', yields in irradiated polyadenylic acid, analysis by HPLC, **134**, 883
- Cycloheximide
effect on
hexose transport reversal and insulin binding in glucose-starved fibroblasts after glucose refeeding (human), **140**, 844
phosphorylation of proteins crosslinked to poly(A)⁺ RNA from free polyribosomes (rat), **136**, 356
prevention of heat shock-induced destabilization of cytoplasmic murine mammary tumor virus RNA, **137**, 1028
- Cyclohexyladenosine receptors
phylogenetic distribution in nervous tissue, **137**, 182
- Cyclopentene
epoxidation by chloroperoxidase (*Caldariomyces fumago*), **136**, 778
- Cystatin C
and protein variant in hereditary cerebral amyloid angiopathy, amino acid sequences, comparison (human), **136**, 548
- Cysteamine
effect on pancreatic secretion of glucagon, insulin, and somatostatin (rat), **134**, 1291
- Cysteine
bound to labile adduct formed during inactivation of monoamine oxidase by 1-phenylcyclopropylamine, identification, **135**, 154
in NADPH-cytochrome *P*-450 reductase, localization and role in enzyme function, assessment by fluorescent titration (rat hepatic microsomes), **134**, 393
role in catalytic activity of β -hydroxy- β -methylglutaryl-CoA reductase (rat liver, yeast), **141**, 278
- Cysteine protease
role in dynorphin degradation by neuroblastoma cell membrane (mouse), **140**, 335
- Cystic fibrosis
effect on
fibroblast permeability to chloride, kinetic analysis (human), **134**, 233
sodium-potassium pump affinity for K⁺ in erythrocytes (human), **137**, 1021
- Cytidine
2',3'-*seco* nucleosides and nucleotides, behavior in various enzymatic systems, **137**, 453
- Cytidine diphosphate
incorporation into mixed DNA-RNA polymer (permeabilized hamster cells), **135**, 340
- Cytidylyl-3',5'-guanosine
cooperativity of ethidium ion in complex formation, **139**, 1164
- Cytochalasin B
effect on platelet serotonin release and clot retraction (human), **140**, 329
enhancement of 1-oleoyl-2-acetyl-*sn*-glycerol-induced superoxide production (human neutrophils), **134**, 690
- Cytochrome *aa*₃
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
- Cytochrome *b*
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
- Cytochrome *b*₅
binary complex formation with cytochrome *P*-450 LM₂ and NADPH-cytochrome *P*-450 reductase, mechanism (rabbit hepatic microsomes), **134**, 519
- Cytochrome *b*₅₅₅
microsomal, purification and partial characterization (*Catharanthus roseus*), **136**, 570
- Cytochrome *b*₅₆₆
role in mitochondrial superoxide formation (rat heart), **138**, 533
- Cytochrome *bc*₁
EPR spectroscopy (*Rhodospseudomonas sphaeroides*), **136**, 616
interaction with antibodies to ubiquinone-binding protein and iron-sulfur protein (bovine cardiac mitochondria), **136**, 376
ubiquinone-binding protein, cell-free synthesis (rat hepatic mitochondria), **138**, 1291
- Cytochrome *c*
somatic and testicular forms, antigenic properties and molecular size, comparison (bovine, mouse, rabbit, rat), **141**, 131
- Cytochrome *cc*₁
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
- Cytochrome-*c* oxidase
*aa*₃-type, in cyanobacterial membrane preparations, immunological identification (*Anacystis nidulans*), **136**, 235
copper, iron, and zinc contents, analysis by proton-induced X-ray emission (bovine, porcine, rat tissues), **140**, 1007
inhibition by psychosine (rat hepatic mitochondria), **137**, 323
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
pulsed, enhanced superoxide dismutase activity (bovine heart), **136**, 433
- Cytochrome-*c* peroxidase
electron transfer reaction at tin oxide electrodes, **139**, 1003
- Cytochrome oxidase, *see* Cytochrome-*c* oxidase
- Cytochrome *P*-450
aflatoxin B₁-specific, from 3-methylcholanthrene-treated liver, purification and characterization (hamster), **139**, 466
catalyzing lanosterol 14 α -demethylation, interaction with diniconazole enantiomers, stereoselectivity (yeast), **137**, 513
dependent microsomal drug metabolism, inhibition by recombinant tumor necrosis factor (mouse), **136**, 316
dependent sulfonylurea metabolism, induction by sulfonylurea (*Streptomyces griseolus*), **140**, 650
heme moiety, covalent binding to apoprotein during enzyme destruction (rat hepatic microsomes), **138**, 193
hepatic levels, reduction by polyribonoinosinic acid-polyribocytidylic acid treatment (mouse, rat), **134**, 632
inactivation by spironolactone (rat liver), **136**, 1162
induction by soybean flour (*Streptomyces griseus*), **141**, 405

Subject Index

- isozymes, interaction with NADPH-cytochrome *P*-450 reductase, spectrophotometric analysis (rabbit, rat), **137**, 437
- like pseudogene, isolation and characterization (rabbit liver), **134**, 499
- LM2b, purification and characterization (rabbit hepatic microsomes), **134**, 1101
- mediated enzyme, polyribonucleoside-polyribocytidylic acid-induced decreases, role of H-2 haplotypes (mouse), **138**, 246
- mRNA, synthesis induction by *N*-alkanes (*Candida maltosa*), **136**, 1148
- and NADPH-specific flavoprotein monomers in solution, reconstitution of hepatic monooxygenase system (rabbit), **139**, 883
- oxygenation of arachidonic acid, hydroxyeicosatetraenoic acids formed during, absolute configuration (rat hepatic microsomes), **141**, 1007
- phenobarbital-inducible
- binary complex formation with cytochrome *b*₅ and NADPH-cytochrome *P*-450 reductase, mechanism (rabbit hepatic microsomes), **134**, 519
 - microsome-bound, phosphorylation by cAMP-dependent protein kinase catalytic subunit (rabbit liver), **134**, 461
 - and protein encoded by plasmid pP-450 PBc2, N-terminal amino acid sequences, comparison (rabbit kidney), **141**, 728
- polycyclic hydrocarbon-inducible, induction by
- β -naphthoflavone in hepatocytes and hepatic hematopoietic cells (fetal rat), **141**, 299
 - phenobarbital (rat hepatoma cells), **137**, 120
- purification and characterization (rat renal microsomes), **141**, 711
- Cytochrome *P*-450₁₁₈
- catalysis of androgen conversion to estrogen (bovine adrenal), **140**, 530
 - dependence of 19-hydroxyandrostenedione-to-19-oxoandrostenedione and cortisol-to-cortisone conversion (bovine), **136**, 369
- Cytochrome-*P*-450 monooxygenase
- 14 α -demethylation of obtusifolioside (corn microsomes), **140**, 1064
- Cytokinin
- copper(II) complexes, superoxide dismutase mimetic activity, **137**, 372
- Cytomegalovirus
- human monoclonal antibodies to, generation of producing hybridomas, **137**, 273
- D**
- 6'-*O*-Dansyl- γ -aminobutyryl atractyloside
- fluorescent probe for analysis of substrate or inhibitor binding to ADP/ATP carrier (bovine cardiac mitochondria), **134**, 266
- 3-Deazaneplanocin
- inhibition of *S*-adenosylhomocysteine hydrolase: characterization and effects on promyelocytic leukemia cells (human), **135**, 688
- Debromoaplysiatoxin
- activation of protein kinase C. role of sulfatide (bovine brain), **138**, 153
- Decolorization
- hydrogen peroxide-dependent, poly R-481 by particulate fractions from *Phanerochaete chrysosporium*, **136**, 220
- Dec-2-ynoyl CoA
- effect on microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
- Deglycosylation
- γ -glutamyl transpeptidase, methodology (rat kidney), **135**, 304
- Dehydration
- metabolic, prostaglandin E₂, and cellular uptake, relationship to prostaglandin E₂-induced growth inhibition (murine leukemia cells), **139**, 808
- Delta toxin
- active fragment, purification and characterization (*Bacillus thuringiensis*), **141**, 106
- Demethylation
- chemoreceptors, inhibition by attractant stimuli in absence of regulatory domain of demethylating enzyme (*Salmonella typhimurium*), **141**, 918
- 14 α -Demethylation
- lanosterol, role of sterol 14-reductase (yeast), **134**, 659
 - obtusifolioside, catalysis by cytochrome *P*-450 monooxygenase (corn microsomes), **140**, 1064
- O*-Demethylation
- etoposide and teniposide, **135**, 215
- Deoxycholate
- effect on F₁-ATPase activity (thermophilic bacterium PS3), **136**, 899
- Deoxycorticosterone
- effect on atrial natriuretic factor release by atrial slices (rat), **135**, 1090
- 2-Deoxy-D-glucose
- inhibition of lymphadenopathy-associated virus/human T-lymphotropic virus type III glycoprotein expression, **141**, 33
 - insulin- and phorbol ester-stimulated uptake by fibroblasts, inhibition by sphingolipids, **138**, 463; *erratum*, **139**, 852
- 5'-Deoxy-5'-methylthioadenosine
- metabolism to methionine, role in cell growth (BHK, CCL 39, Raji cells), **135**, 131
- Deoxynucleoside triphosphates
- altered metabolism after DNA damage, role of poly(ADP-ribose) polymerase (L1210 cells), **137**, 1153
- 3-Deoxy-D-manno-2-octulosonic acid
- β -ketopyranosidic linkages, hydrolysis by bacteriophage-associated glycanase (*Escherichia coli*), **136**, 329
- Deoxyribonuclease I
- metaphase chromosome digestion, release of DNA fragment 300 bp long (Chinese hamster V79 cells), **138**, 413
- Dephosphorylation
- phosphorylation processes, role in phosphoenolpyruvate carboxylase diurnal regulatory changes (CAM plants), **136**, 151
- Depolarization
- potassium-induced, enhancement of muscarinic receptor-stimulated inositol tetrakisphosphate accumulation in

Subject Index

- cerebral cortical slices (rat), **141**, 1130
- smooth muscle cell monolayers containing fluorescent probes, effect on cytosolic Ca^{2+} levels (hamster, rabbit, rat), **136**, 927
- Depolymerization**
- actin filaments, in evaluation of filament length, **139**, 109
- Dequalinium**
- inhibition of calmodulin activity and cellular proliferation (rat C6 astrocytoma cells), **135**, 574
- Dermatan sulfate**
- L-iduronate ring conformation, solid-state ^{13}C NMR and X-ray diffraction analysis (porcine skin), **137**, 87
- Dermorphin**
- molecular mechanical simulations, **140**, 342
- Desialation**
- ceruloplasmin by hepatic endothelium (rat), **140**, 94
- Desipramine**
- effect on inositol incorporation into phosphoinositides and inositol phosphates (rat C6 glioma cells), **135**, 465
- Desmethylimipramine, *see* Desipramine
- Deuterium**
- based NMR spectroscopy in monitoring metabolic processes *in vivo*, evaluation (mouse), **137**, 579
- Dexamethasone**
- and cAMP, synergistic regulation of preproenkephalin mRNA levels in neuroblastoma-glioma hybrid cells, **139**, 1
- effect on
- atrial natriuretic factor release by atrial slices (rat), **135**, 1090
- 12-*O*-tetradecanoylphorbol 13-acetate-induced secretion of ACTH and prostaglandin E_2 by pituitary cells: role of Ca^{2+} (rat), **138**, 323
- induction of glucose-6-phosphatase in cultured hepatoma cells, **139**, 1130
- inhibition of activated macrophage antitumor potential by receptor-mediated mechanism (mouse), **136**, 130
- regulation of
- albumin mRNA levels in hepatocytes *in vitro* (*Xenopus laevis*), **136**, 8
- atrial natriuretic peptide mRNA *in vivo* (rat heart), **139**, 1047
- stimulation of protein kinase C (rat hepatocytes), **135**, 33
- uptake and metabolism by isolated hepatocytes (nonadrenalectomized rat), **135**, 160
- Dexamethasone phosphate**
- induction of tyrosine aminotransferase and ornithine decarboxylase *in vivo*, enhancement by diacylglycerol (adrenalectomized rat), **138**, 275
- Dextran**
- concanavalin A interaction, effect on K^+ currents through lipid bilayer membranes, **141**, 912
- in concanavalin A solutions, detection and quantitation with electrical oscillator, **137**, 978
- Diabetes**
- alloxan- or streptozotocin-induced, associated decreased expression of epidermal growth factor receptors (rat), **137**, 258
- associated mitochondrial protein phosphorylation and cardiomyopathy, effect of estrone treatment (mouse), **139**, 56
- effects on fructose-2,6-bisphosphatase, glucose-1,6-bisphosphatase, and 6-phosphofructo-2-kinase (rat liver), **136**, 498
- impairment of glomerular contractile response to angiotensin II (rat), **136**, 1185
- insulin-resistant type II, Na^+/K^+ -ATPase activity in renal basolateral membranes, effect of insulin (rat), **139**, 1313
- streptozotocin-induced
- associated insulin receptor autophosphorylation and kinase activity, effect of short fast (rat), **140**, 850
- and insulin treatment, effect on hexokinase II degradation in skeletal muscle (rat), **138**, 374
- Diacylglycerol**
- enhancement of tyrosine aminotransferase and ornithine decarboxylase *in vivo* induction by glucocorticoid (adrenalectomized rat), **138**, 275
- inhibition of chondroitin sulfate sulfation (human promyelocytic leukemia cells), **135**, 909
- stereospecificity, role in stimulus-response coupling in platelets (human), **140**, 1143
- Diacylglycerol lipase**, *see* Lipoprotein lipase
- Diadenosine tetraphosphate**
- activation of cytosol 5'-nucleotidase (*Artemia* embryo), **138**, 261
- cis*-Diamminedichloroplatinum(II), *see* Cisplatin
- trans*-Diamminedichloroplatinum(II)
- inhibition of DNA synthesis in leukemia cells, effect of serum (mouse), **138**, 631
- 1,4-Diazabicyclo[2.2.2]octane, *see* Triethylenediamine
- Diazepam**
- inhibition of K^+ -induced aldosterone secretion by glomerulosa cells (bovine adrenal), **135**, 994
- p*-Diazobenzoyl biocytin
- labeling of tyrosines and histidines in proteins, **138**, 872
- 1,2-Dibromoethane, *see* Ethylene dibromide
- Dibutyryl cAMP**
- inhibition of erythroleukemic cell differentiation (mouse), **134**, 845
- 3-(3',4'-Dichlorophenyl)-1,1-dimethylurea, *see* Diuron
- Dictyostelium discoideum**
- fluid-phase pinocytosis, kinetics, **138**, 1146
- proliferation, effect on plasma membrane ATPase activity, **141**, 1092
- 2',3'-Dideoxycytidine
- selective inhibition of human T-lymphotropic virus type III/lymphadenopathy-associated virus, **140**, 735
- 2',3'-Dideoxythymidine 5'-triphosphate
- and derivatives modified on 3' position, inhibition of murine leukemia virus reverse transcriptase activity, **140**, 498
- Diet**
- fasting, *see* Fasting
- induction of *c-myc* oncogene expression in liver (rat), **140**, 574
- lipid content, effect on nuclear membrane structure and function (murine liver), **135**, 655
- N,N*-Diethyl-2-[(4-phenylmethyl)phenoxy]-ethanamine HCl
- binding site, identification as growth-promoting histamine receptor mediating tamoxifen effects (rat liver), **134**, 601
- Diethylstilbestrol**
- effect on glutathione transferase expression in seminal ves-

Subject Index

- icles and pituitary gland (rat), **136**, 1057
- Differentiation-inducing factor
synthesis by MC3T3-E1 osteoblasts (mouse), **134**, 400
- Diffusion
adsorbed substrate on membrane surface, catalytic rate enhancement of alkaline phosphatase, **138**, 803
- α -Difluoromethylornithine
induction of embryonal carcinoma cell differentiation *in vitro* (human), **140**, 66
interaction with ornithine decarboxylase-antizyme complex, analysis (rat liver), **137**, 1101
pulmonary tumor sensitivity, relationship to ornithine decarboxylase mRNA levels (human), **134**, 572
- Diglyceride lipase, *see* Lipoprotein lipase
- Dihydroalprenolol
³H-labeled, binding to pineal particulate fractions, enhancement by melatonin and α -adrenergic compounds (rabbit), **138**, 1196; *erratum*, **139**, 1320
- Dihydroalprenolol receptors
detection in skeletal muscle (rat), **136**, 1071
- Dihydroanisoin
as lignin model compound, biomimetic oxidation by simple inorganic complexes, **139**, 1104
- trans*-1,2-Dihydrobenzene-1,2-diol dehydrogenase
and 3α -hydroxysteroid dehydrogenase: solubilization, separation, and partial purification (rat hepatic microsomes), **135**, 795
- Dihydrodiol dehydrogenase, *see trans*-1,2-Dihydrobenzene-1,2-diol dehydrogenase
- Dihydrodiols
-derivatized nitro-polycyclic aromatic hydrocarbons, binding to DNA, catalysis by xanthine oxidase, **141**, 245
- trans*-Dihydrodiols
identification as products formed during 5-methylchrysene oxidative metabolism at K-regions (rat hepatic microsomes), **141**, 734
- Dihydroflavin mononucleotide
cofactor activity for indoleamine 2,3-dioxygenase (murine epididymis), **137**, 1106
- Dihydrofolate reductase
antigenically-related high-molecular-weight nonfunctional protein, detection in brain (mouse), **141**, 258
- 9,10-Dihydro-9-(*S*-glutathionyl)-10-hydroxyphenanthrene diastereomers, stereoselective inhibition of glutathione *S*-transferase (rat liver), **141**, 892
- Dihydrolipoamide dehydrogenase
excess, role in reconstituted α -ketoglutarate dehydrogenase complex (*Escherichia coli*), **135**, 802
- Dihydropyridine
agonists and antagonists, effect on neuronal voltage-sensitive calcium channels (rat brain), **134**, 1038
effect on Ca^{2+} uptake by neuronal cells (rat brain), **139**, 581
- 1,4-Dihydropyridine receptors
neuronal Ca^{2+} channel, cytolocalization and subunit structure identification with polyclonal antibodies (rabbit brain, skeletal muscle), **139**, 996
- 3,4-Dihydroxy-3,4-dihydrobenzo[*a*]anthracene
hydroperoxide-dependent epoxidation by ovine seminal vesicle microsomes and hematin, **140**, 181
- 2-(1,2-Dihydroxyethyl)thieno[3,2-*g*]pterin
identification as dephosphorylated oxidation product of xanthine oxidase molybdenum cofactor, **135**, 256
- 7,12-Dihydroxymethylbenzo[*a*]anthracene
regioselective glutathione conjugation via reactive 7-hydroxymethyl sulfate ester (rat hepatic cytosol), **134**, 100
- 1,25-Dihydroxyvitamin D₃
effect on
cytosolic free Ca^{2+} levels (avian cartilage cells), **141**, 1236
tumor growth after inoculation with receptor-rich osteosarcoma cells (athymic nude mouse), **139**, 1292
inhibition of clonogenic growth of transformed cells, role of receptors, **139**, 136
- 1,25-Dihydroxyvitamin D₃ receptors
homologous up-regulation *in vivo* (rat), **137**, 742
- 4,4'-Diisothiocyano-2,2'-stilbene disulfonic acid
effect on sodium regulation of α_2 -adrenoreceptors (human platelets), **141**, 222
- Diltiazem
stimulation of low-density lipoprotein receptor synthesis (human skin fibroblasts), **136**, 845
- 7,12-Dimethylbenzo[*a*]anthracene, *see* 9,10-Dimethyl-1,2-benzanthracene
- 9,10-Dimethyl-1,2-benzanthracene
mammary tumor induction, anticarcinogenic effect of retinoids, relationship to cAMP-dependent protein kinase (rat), **135**, 116
- 1,1'-Dimethylferrocene
inhibition of agonist evoked aggregation, ATP secretion, and cytosolic Ca^{2+} increase in aspirinated platelets (human), **139**, 509
- Dimethylformamide
solubilization of iron-molybdenum cofactor of *Azotobacter vinelandii* nitrogenase, **139**, 740
- 17,21-Dimethyl-19-nor-4,9-pregnadiene-3,20-dione
effect on nuclear estrogen receptor turnover and synthesis, analysis by density-shift method (hamster uterine decidua cells), **135**, 98
and RU 38486, interaction with progesterone receptors, comparison (human breast cancer cells), **135**, 90
- 5,5-Dimethyl-1-pyrroline *N*-oxide
hydroxyl and superoxide radical adducts, [¹⁷O]oxygen hyperfine structure, **141**, 622
- 6,7-Dimethyl-8-ribityllumazine
formation by heavy riboflavin synthase (*Bacillus subtilis*), **139**, 1111
- Dimethyl sulfoxide
effect on P150 phosphorylation in adriamycin-resistant HL60 cells, **138**, 9
induction of
phospholipid methylation and arachidonic acid release by promyelocytic leukemic HL-60 cells (human), **140**, 515
soluble protein tyrosine phosphorylation (human peripheral blood T lymphocytes), **140**, 743
teratocarcinoma cell differentiation, specific expression of homoeobox-containing genes, **137**, 520
-water mixtures, hydration states of L-alanine and L-proline, ¹⁷O NMR, **135**, 363
- Dimyristoylphosphatidylcholine
complexes with apolipoprotein A-IV, binding to plasma membranes (rat liver), **139**, 122

Subject Index

- Dimyristoylphosphatidylserine
interaction with bombesin, effect on bombesin conformation, **141**, 99
- Diniconazole
enantiomers, stereoselective interaction with cytochrome P-450/14DM (yeast), **137**, 513
- 2,4-Dinitro-5-aziridinyl benzamide, *see* CB 1954
- Dinitrophenyl
induction of B-cell activation, role of phosphatidylinositol pathway, **140**, 1079
- S-Dinitrophenyl glutathione
and oxidized glutathione, transport mechanisms in erythrocytes, comparison (human), **139**, 538
- Diocanoyl glycerol
with diisopropyl fluorophospho-thrombin, stimulation of DNA synthesis (NIL cells), **141**, 650
enhancement of phosphoprotein B-50 phosphorylation in native synaptic plasma membranes (rat brain), **136**, 1007
- Dioleoylphosphatidylethanolamine
and glycophorin-containing liposomes, trypsin-induced destabilization, **141**, 973
- Dipeptidyl carboxypeptidase I
amino-terminal sequence analysis and preliminary characterization of hybridization-selected primary translation product (bovine lung), **141**, 968
binding to captopril, inhibition by peptide analogs (rat lung), **135**, 770
and glucocorticoid receptors, endothelial cell levels, reduction by fucosterol (bovine carotid artery), **139**, 348
inhibitory peptide, isolation and characterization (porcine plasma), **139**, 52
- 1,2-Diphenyl-1-(S-glutathionyl)-2-hydroxyethane
diastereomers, stereoselective inhibition of glutathione S-transferase (rat liver), **141**, 892
- Diphtheria toxin
binding of divalent cations, ESR analysis, **139**, 991
-synthetic pre-S viral peptide-murabutide conjugates, in production of antibodies recognizing hepatitis B virus surface antigens, **140**, 557
type B, peptide fragment, pH-dependent insertion in lipid membrane, conformational analysis, **136**, 160
- Dipyridamole
sensitivity of stationary-phase hepatoma cells, restoration by amphotericin B (rat), **140**, 434
- Disaccharides
penetration inhibition by *Pseudomonas aeruginosa* outer membranes, **134**, 106
- Distal tubules
cGMP levels, enhancement by atrial natriuretic peptide (rat liver), **136**, 947
- Distamycin, *see* Stallimycin
- Disuccinimidyl suberate
crosslinking of somatostatin receptors to radioligands (rat adrenal cortex), **139**, 757
- Disulfide bonds
extracellular, stabilization of acetylcholine receptor dimers (*Torpedo californica*), **139**, 830
in protein core, role in interaction of proteodermatan sulfate and collagen (bovine skin), **138**, 1348
- Dithiothreitol
role in induction of high Mg^{2+} -ATPase activity of coupling factor 1 by octyl glucoside (spinach chloroplasts), **141**, 636
- Diurnal cycle
effect on hepatic microsomal squalene synthetase stimulation by cholestyramine (rat), **138**, 335
- Diuron
binding sites in photosystem II, characterization (spinach), **141**, 682
- DNA
adenovirus type 2, chromatin-like complex with core protein VII, transcription *in vitro*, **136**, 86
and ATP, binding to urea-transformed native estrogen receptor, evidence for separate binding sites (immature bovine uterus), **139**, 1250
binding of
dihydrodiol-derivatized nitro-polycyclic aromatic hydrocarbons, catalysis by xanthine oxidase, **141**, 245
distamycin synthetic analog, characteristics, **140**, 626
toromycin, analysis, **136**, 885
binding sites on estrogen and glucocorticoid receptors, computer-based search, **139**, 281
cauliflower mosaic virus, potential transcription control sequence, detection by dinucleotide primed *in vitro* transcription, **138**, 17
cellular, 8-hydroxyguanine residue formation after 4-nitroquinoline 1-oxide exposure (Ehrlich ascites cells), **139**, 626
cleavage induction by
amsacrine, magnitude, correlation with sensitivity to drug (human leukemia cells), **134**, 638
bleomycin-iron III complex, activation by $KHSO_5$ (bacteriophage λ), **136**, 1013
metallodaunomycin, role of metal ions, **136**, 1
cleavage rates, calculation at different sites from partial digestion of end-labeled molecule, method reevaluation, **138**, 778
cloned, as probe for detection of *Plasmodium falciparum* malaria, **138**, 363
complementary
C1 inhibitor, isolation and sequencing (human), **137**, 620
chromogranin A, detection of 2.3kb mRNA in diverse neuroendocrine tissues (human), **137**, 418
cloned from cucumber mosaic virus satellite, enzymatic transcription to infectious RNAs, **135**, 290
creatine kinase M subunit, isolation and sequencing (human), **140**, 981
encoding catalytically active DNA polymerase α fragment, cloning and expression (bovine thymus), **140**, 21
encoding polypeptide similar to human insulin receptor precursor, cloning (*Drosophila* sp.), **141**, 474
galactosyltransferase, isolation and characterization (human milk), **139**, 163
to α_{2u} -globulin mRNA induced by phenobarbital, characterization (rat liver), **134**, 1182
glutamine synthetase, selection from renal library and restriction map construction (rat), **134**, 146
glutathione S-transferase H_a subunit 1, nucleotide sequence analysis (human liver), **141**, 229
heparin cofactor II, partial clone, isolation and characterization (human), **137**, 431
to hepatic arginase, isolation and demonstration of nonhomology between hepatic and renal genes

Subject Index

- (human), **141**, 53
- human interferon γ , transformed murine L cells, extracellular and intracellular expression of human INF- γ , **135**, 894
- human interleukin 2 receptor, expression in *Escherichia coli*, **141**, 804
- ligninase, identification of clones with synthetic oligonucleotide probes (*Phanerochaete chrysosporium*), **137**, 649
- neuron-specific and nonneuronal enolase, isolation and characterization (murine brain), **141**, 271
- 6-phosphofructokinase, isolation (human muscle), **135**, 615
- postmeiotically expressed clone encoding lactate dehydrogenase, nucleotide sequence (mouse), **136**, 760
- rat liver arginase, cloning and expression (*Escherichia coli*), **136**, 955
- to reovirus serotype-1 Lang strain mRNA encoding major capsid polypeptide $\sigma 3$, cloning and nucleotide sequence (*Escherichia coli*), **136**, 183
- minor capsid polypeptides $\sigma 1a$ and nonstructural $\sigma 1bNS$, cloning, **140**, 508
- σNS polypeptides, cloning and nucleotide sequence (human), **139**, 845
- sphingolipid activator protein 1, isolation and characterization (human), **134**, 989
- for substance P precursor, sequence analysis (rat), **139**, 1040
- damage
- associated alteration of deoxynucleoside triphosphate metabolism, role of poly(ADP-ribose) polymerase (L1210 cells), **137**, 1153
 - induction of alterations in glucose metabolism by poly(ADP-ribose) polymerase activation (murine L1210 cells), **134**, 227
 - induction by 1,2-dibromoethane at low doses (rat liver), **141**, 723
- double-stranded, binding to *recA* proteins, superhelicity effects (*Escherichia coli*), **136**, 404
- encoding serum amyloid A gene, detection of polymorphisms (human), **137**, 1196
- enhancer sequences, prediction of binding proteins by informational spectrum method, **141**, 831
- Epstein-Barr virus, fragment coding for nuclear antigen, transfer to normal human lymphocytes: gene expression without cell transformation, **134**, 1260
- fragments, 300 bp long, release from metaphase chromosomes after deoxyribonuclease I digestion (Chinese hamster V79 cells), **138**, 413
- genomic
- from rat islets, transfection of fibroblasts, induction of insulin release, **136**, 638; *erratum*, **139**, 852
 - sequences homologous to
 - human epidermal growth factor receptor cDNA, application in characterization of rodent EGF receptor transcripts (mouse), **141**, 1109
 - interleukin 2 cDNA, characterization (human placenta), **138**, 966
 - hepatitis B virus subtype *adw* pre-S2 region, nucleotide sequence, comparison with other subtypes, **135**, 382
 - hybrid segment containing human DNA polymerase α gene, identification (mouse), **139**, 637
 - 8-hydroxydeoxyguanosine accumulation after exposure to phorbol ester (human granulocytes), **137**, 841
 - interaction with
 - histone H4, effect of poly(ADP-ribose) (rat liver), **139**, 523
 - meso*-tetraaryl porphyrins bearing 1 to 4 positive charges, mechanism (bovine thymus), **141**, 643
 - interstrand crosslinking by CB 1954 in Walker tumor cells (rat), **140**, 1073
 - IVS 1 donor site consensus sequence, identification of mutation inducing β thalassemia (human), **139**, 709
 - length and sequence polymorphisms, detection in ribosomal gene spacer (*Tripneustes gratilla*), **137**, 834
 - linking number, modification by 5 S RNA gene-specific transcription factor, **134**, 1086
 - mediation of gene transfer into epidermal cells by electroporation (mouse), **137**, 244
 - methotrexate-induced lesions, inhibition by leucovorine (human colon adenocarcinoma cells), **138**, 54
 - mitochondrial, autonomous replicating sequences, cloning in *Saccharomyces cerevisiae* (*Lentinus edodes*), **138**, 1110
 - modification mechanism: guanosine reaction with 1,3-bis(2-chloroethyl)-1-nitrosourea, **139**, 31
 - multiply branched molecules, identification as intermediates of postreplicational repair pathway (bacteriophage λ), **137**, 869
 - mung bean, construction of genomic library in phage charon 4A: detection of repetitive sequences and β -tubulin gene, **137**, 788
 - 4-nitroquinoline-1-oxide adducts, lack of excision by cell extracts capable of excising pyrimidine dimers (bovine thymus), **140**, 775
 - plasmid
 - pBR322, restriction fragments, preferential nucleosome placement (*Escherichia coli*), **138**, 110
 - pPyLT1, maintenance of autonomous genetic elements after transfer in transgenic strains (mouse), **141**, 1162
 - strand scission by hemin-thiolate complexes, **136**, 645
 - recombinant junctions, screening with synthetic oligonucleotides, **138**, 945
 - reversible intercalator-induced frameshift mutations, effect of mutation in *uvrB* gene (*Salmonella typhimurium*), **138**, 505
 - RNA mixed polymer, incorporation of cytidine diphosphate (permeabilized hamster cells), **135**, 340
 - simian virus-40, replication inhibition by 1-*N*-acyl-3'-*N*-(trifluoroacetyl)kanamycin *in vitro* (CosI cells), **136**, 322
 - single-stranded
 - binding proteins, modulation of topoisomerase I relaxing activity (*Escherichia coli*), **137**, 795
 - pyrimidine dimer-dependent cleavage by T4 uv endonuclease, **141**, 204
 - specific sites, supercoil-dependent structure, recognition by chromosomal high-mobility-group protein (chicken), **138**, 910
 - strand breaks
 - carcinogen-induced, detection by gravity-flow alkaline elution (MDBK cells), **137**, 1006
 - induction by clomiphene (human leukemia cells), **138**, 1203

Subject Index

- T-, gene 7, expression in *Xenopus laevis* oocytes (*Agrobacterium tumefaciens*), **141**, 1242
transcriptionally active, affinity isolation (chicken erythrocytes), **137**, 474
uracil incorporation, methotrexate effect (HeLa cells, human lymphoblastoid cells), **135**, 886
yeast, localization of sequences affecting kanamycin resistance gene expression in *Escherichia coli*, **136**, 529
- DNA-directed DNA polymerase
catalytically active fragment, cloning and expression of encoding cDNA (bovine thymus), **140**, 21
gene, identification in hybrid DNA segment (human), **139**, 637
phosphatidylinositol-dependent activation (murine sarcoma cells), **135**, 880
sarcoma-associated, reactive monoclonal antibody, isolation and characterization (mouse), **141**, 60
and topoisomerase II levels in developing neuronal nuclei, independent regulation (rat brain), **138**, 1116
- DNA glycosylase
identification in *Drosophila melanogaster* embryos, **134**, 201
- DNA polymerase α , *see* DNA-directed DNA polymerase
- DNA polymerase β
inhibition by poly(ADP-ribosyl)ation *in vitro* (bovine thymus), **140**, 666
mRNA, nucleotide sequencing via cDNA cloning (human), **136**, 341
- DNA synthesis
hepatic, stimulation by prolactin administration (rat), **138**, 1138
inhibition by *cis*- and *trans*-diamminedichloroplatinum(II), effect of serum (murine leukemia cells), **138**, 631
in keratinocytes, effect of retinoic acid, role of extracellular Ca^{2+} (murine epidermis), **138**, 483
and poly(ADP-ribosyl)ation and mammary gland differentiation, interrelationships (mouse), **137**, 359
repair, in lymphocytes, inhibition by D-ribose (human), **138**, 673
stimulation by
diisopropyl fluorophospho-thrombin with dioctanoyl glycerol or phorbol myristate acetate (NIL cells), **141**, 650
insulin-like growth factor II, effect of anti-insulin-like growth factor I receptor antibody (human fibroblasts), **139**, 501
substance K-induced, correlation with *myc*-gene transcript levels (quail fibroblasts), **137**, 167
- Docosahexaenoate
decreased content in plasma phospholipids from Usher's syndrome patients, **141**, 600
- 4-(Dodecyltrimethyl ammonium)-1-oxyl-2,2,6,6-tetramethyl piperidine bromide
internalization by energized mitochondria (rat liver), **136**, 941
- L-Dopa decarboxylase, *see* Aromatic-L-amino-acid decarboxylase
- Dopamine
brain levels, effect of sustained exercise (rat), **137**, 149
- Dopamine β -monooxygenase
inhibition by aponeurocuprein (bovine brain), **138**, 1
- Dopamine receptors
D-1, extraction from striatal membranes (rat), **137**, 943
- Doxorubicin
effect on adenylate cyclase after chronic administration (murine heart), **136**, 745
induction of breast cancer cell killing, prevention by iron chelators and oxygen radical scavengers (human), **135**, 330
resistant HL60 cells, P150 phosphorylation, effects of dimethyl sulfoxide, retinoic acid, and 12-O-tetradecanoylphorbol-13-acetate, **138**, 9
- DPX-F6025
cytochrome P-450-dependent metabolism, induction by sulfonyleurea (*Streptomyces griseolus*), **140**, 650
- DT-diaphorase, *see* NAD(P)H dehydrogenase (quinone)
- Duroquinol oxidase
cyanide-resistant activity in submitochondrial particles, comparison with NADH and succinate oxidation (*Arum maculatum*), **135**, 677
- Dyes
synthetic, phototoxicity evaluation using *Paramecium aurelia*, **139**, 79
- Dynorphins
degradation by neuroblastoma cell membrane, role of cysteine protease (mouse), **140**, 335
- Dystrophy
retinal, associated reduction in rhodopsin phosphorylation (dog), **135**, 1022

E

- Ecdysterone
induction of 26-kDa and 27-kDa heat shock protein phosphorylation (*Drosophila melanogaster*), **141**, 426
- Edetic acid
induction of autolysis, effect of antibiotics (*Escherichia coli*), **134**, 338
modification of fructose-1,6-bisphosphatase activity, effect on 5'-AMP sensitivity (fish, rat brain), **139**, 571
- EDTA, *see* Edetic acid
- Eicosapentaenoic acid
effect on natural killer cell activity of lymphocytes (human), **138**, 1058
prostaglandin I_3 production from, enhancement by arachidonic acid (human endothelial cells), **135**, 403
- trans*-2-*cis*-8,11,14-Eicosatetraenoyl CoA
effect on microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
- Elastase
release from monocytes, effect of κ elastin (human), **141**, 92
- Elastin
lysine side chains, oxidation by canine myeloperoxidase and stimulated human neutrophils (chicken embryonic aorta), **135**, 451
and polypentapeptide, elastomer length, temperature dependence (bovine ligamentum nuchae), **141**, 749
quantitation by measurement of pentapeptide content (swine), **136**, 672
type κ , effects on monocytes (human), **141**, 92
- Electrical stimulation
effect on phosphoinositide metabolism (rat muscle), **136**, 863

Subject Index

- Electron paramagnetic resonance
periplasmic hydrogenase spectrum, effect of low carbon monoxide levels (*Desulfovibrio vulgaris*), **137**, 1086
- Electron transfer
cytochrome-c peroxidase at tin oxide electrodes, **139**, 1003
transplasmalemma, from hepatocytes, role in diferric transferrin reductase system (rat), **139**, 1117
- Electropermeabilization
role in cell fusion (Chinese hamster ovary cells), **140**, 258
- Electroporation
epidermal cells, for DNA-mediated gene introduction (mouse), **137**, 244
mediation of gene transfer into hematopoietic stem cells (mouse), **141**, 1018
- Ellagic acid
inhibition of glutathione S-transferase isoenzymes from human and rat tissues, comparison, **141**, 1170
- Elongation factor 2
purification, characterization, and antibody preparation (rat liver), **134**, 1015
specific cellular ADP-ribosyltransferase, characterization (rat liver), **139**, 1210
- Emphysema
treatment, peptide aldehydes and α -antitrypsin as elastase inhibitors, comparison (human), **139**, 896
- Emulgen 108
selective inhibition of microsomal hepatic cholesterol epoxide hydrolase (mouse), **140**, 632
- Encephalomyelitis
allergic, effect on phosphoinoside covalent linkage to myelin basic protein *in vitro* (rabbit, guinea pig), **136**, 426
- Encephalomyopathy
mitochondrial, effect on cardiac mitochondrial NADH-ubiquinone oxidoreductase subunit levels (human), **140**, 88
- Endo- β -N-acetylglucosaminidase F
deglycosylation of thyroid-stimulating hormone (bovine), **138**, 230
- Endo-1,4- β -glucanase, *see* Cellulase
- Endo-glycosidase F
and peptide N-glycosidase F, release of N-linked oligosaccharides from mammalian cells, applications, **140**, 967
- Endonucleases
T4 uv, pyrimidine dimer-dependent cleavage of single-stranded DNA, **141**, 204
- Endopeptidase
associated with neurohypophyseal secretory granules, cleavage of pro-oxytocin/neurophysin at paired basic residues (bovine), **134**, 1190
- Endoplasmic reticulum
calcium release, inositol trisphosphate-induced, inhibition by calmodulin (rat pancreatic islets), **141**, 418
- β -Endorphin
active and inactive forms, production in gastric antrum (rat), **134**, 14; *erratum*, **138**, 502
- Endothelial cells
arachidonic acid releasing systems, mechanisms (porcine aorta), **139**, 1098
cerebrovascular, glycogen metabolism, regulatory mechanism, comparison with smooth muscle (rat), **134**, 484
fibroblast growth factor-induced proliferation, inhibition by β -transforming growth factor (bovine aorta), **138**, 476
internalization and metabolism of endogenous heparin (bovine adrenal capillaries), **140**, 294
plasminogen activator inhibitor levels, enhancement by interleukin 1 (human), **139**, 720
proliferation
inhibition by transforming growth factor β *in vitro* (bovine aorta), **137**, 295
role of human platelet-derived acid-sensitive growth factor (bovine aorta), **138**, 795
prostacyclin synthesis *in vitro*, stimulation by cachectin/tumor necrosis factor (human umbilical vein), **141**, 482
stimulating macrophage factor, partial purification and properties (human), **134**, 344
vascular, dependence of relaxation by arginine-containing peptides (rat aorta), **141**, 446
- Endothelium-derived relaxing factor
stabilization and partial characterization (bovine aorta), **141**, 689
- δ -Endotoxin
cloned, biochemical and immunological characterization (*Bacillus thuringiensis* var. *israelensis*), **137**, 748
- Enkephalin receptors
and morphine receptors, identification in neuroblastoma cells (human), **137**, 813
- Enkephalins
cyclic and linear pseudopeptide analogs, synthesis and biological activity *in vitro*, **136**, 730
sulfation by sulfotransferase (*Eubacterium* A-44), **140**, 38
Tyr¹-substituted, fluorescent, selective binding to μ - and δ -opiate receptors (rat brain), **136**, 1170
- Enolase
neuron-specific and nonneuronal, cDNA isolation and characterization (murine brain), **141**, 271
- trans*-2-Enoyl-CoA reductase
in fatty acid chain elongation system, effects of competitive substrates, inhibitors, and proteases (rat hepatic microsomes), **140**, 74
- Enterocytes
permeabilized, Ca²⁺ uptake by intercellular compartments, effect of inositol 1,4,5-trisphosphate (rat intestine), **139**, 612
- Enterolactone
inhibition of Na⁺, K⁺-ATPase and displacement of ouabain from binding sites (guinea pig, human heart), **134**, 1064
- Enterotoxins
induction of membrane permeability changes in HeLa and Vero cells in absence of extracellular Ca²⁺ (*Clostridium perfringens* type A), **141**, 704
- Eosinophils
derived cytotoxic cationic protein, ribonuclease activity, **139**, 1239
- Epidermal cells
DNA-mediated gene introduction by electroporation (mouse), **137**, 244
- Epidermal growth factor
effect on hepatic lipogenic enzymes after chronic administration (hypophysectomized rat), **141**, 78
enhancement of c-myc mRNA accumulation in cultured thyroid cells (dog), **141**, 1066

Subject Index

- like polypeptide, synthesis and secretion by MCF-7 tumor cells (human), **136**, 300
- and polyoma virus middle T antigen, effect on secondary cultured embryo cell transformation (rat), **136**, 921
- potentiated phenotypic transformation of kidney cells, induction by platelet-derived growth factor-like polypeptide (rat), **141**, 1229
- Epidermal growth factor receptors
 - associated tyrosine kinase activity, phosphorylation of band 3 glycoprotein (human erythrocytes), **135**, 720
 - binding activity
 - decrease after chronic phenobarbital treatment (rat liver), **135**, 501
 - loss in K-ras-transformed renal cells, mechanism (rat), **135**, 435
 - decreased expression in alloxan- or streptozotocin-induced diabetes (rat liver), **137**, 258
 - relationship to terminal cellular differentiation (human epidermoid carcinoma A431), **140**, 837
 - transcripts in rodents, characterization with murine genomic probe, **141**, 1109
- Epinephrine
 - effect on
 - adenylate cyclase-phosphodiesterase system in thrombopathic and normal platelets (dog), **140**, 589
 - atrial natriuretic factor release by atrial slices (rat), **135**, 1090
 - cardiac and plasma atrial natriuretic factor levels after chronic administration (rat), **136**, 510
 - triglyceride synthesis and secretion in isolated hepatocytes (rat), **141**, 191
 - induction of melanosome aggregation, comparative effects of ACTH and cAMP (goldfish), **138**, 1081
- Epithelial cells
 - intestinal IEC-6, sodium-dependent concentrative nucleoside transport (rat), **140**, 1028
 - LLC-PK₁, amiloride-sensitive sodium transport, inhibition by atrial natriuretic factor and cGMP, **134**, 852
 - mammary, ras-transfected, collagen processing (mouse), **137**, 161
 - MDCK, Forssman glycolipid flow between: model of tight junction structure, **138**, 1163
 - small intestinal, protein kinase C identification and characterization (rabbit, rat), **139**, 875
 - stimulation by
 - human fibroblast growth factors, heparin effects (bovine lens), **137**, 1205
 - synthetic heparin pentasaccharide, enhancement by human acidic fibroblast growth factors (bovine lens), **139**, 728
- Epitopes
 - ATPase inhibitor, localization (bovine heart), **137**, 364
 - transcortin, phylogenetic analysis with monoclonal antibodies (human), **138**, 1037
- Epoxidation
 - aldrin, lipid peroxidation-dependent, analysis in hepatic microsomes, hepatocytes, and granuloma cells (rat), **138**, 24
 - hydroperoxide-dependent, 3,4-dihydroxy-3,4-dihydrobenzo[*a*]anthracene by ovine seminal vesicle microsomes and hematin, **140**, 181
- Epoxide hydrolase
 - microsomal, expression by human-murine hybrid cells, quantitation, **137**, 775
 - xenobiotic, associated activity of hepatic microsomal cholesterol epoxide hydrolase, enhancement by detergents (mouse), **140**, 632
- 5,6-Epoxyeicosatrienoic acid
 - mobilization of Ca²⁺ in anterior pituitary cells (rat), **139**, 1188
- 9,10-Epoxy-12-octadecenoate
 - biosynthesis by leukocytes in lung lavages after hyperoxic exposure (rat), **134**, 1071; *erratum*, **135**, 1150
 - synthesis by neutrophils (dog, guinea pig, human), **137**, 424
- Epsstein-Barr virus
 - DNA fragment coding for nuclear antigen, transfer to normal human lymphocytes: gene expression without cell transformation, **134**, 1260
 - transformed B cells, detection of interleukin-1 α mRNA (human), **139**, 353
- Erythrocytes
 - calpain inhibitor activity (Milan hypertensive rat), **138**, 1370
 - defective calpain-endogenous calpain inhibitor system, characterization (Milan hypertensive rat), **139**, 341
 - inside-out vesicles, transglutaminase substrates, identification by immunoblotting (human), **134**, 685
 - membranes
 - bicarbonate exchange kinetics at equilibrium, ¹³C NMR (human), **136**, 266
 - corticotropin-releasing factor binding sites, identification and modulation by chronic ethanol treatment (human, rat), **136**, 137
 - ozone-induced oxidation, uric acid protective effects (bovine), **137**, 536
 - phosphatidic acid metabolism, activation by perfringolysin O (human), **137**, 23
 - phosphorylation-induced Ca²⁺ release (human), **136**, 288
 - proteins and/or glycoproteins, functional group labeling with biotin derivatives and ¹²⁵I-labeled streptavidin (human), **136**, 80
 - occurrence of (ADP-ribose)_n glycohydrolase (human), **136**, 1110
 - outer membrane bilayer, phosphatidylserine localization (newborn human), **136**, 914
 - polyamines, intracellular location (human), **134**, 1365
 - protein tyrosine phosphorylation by casein kinase A (human), **139**, 855
 - proteolytic enzyme levels and subcellular distribution, Ca²⁺-induced alterations (human), **138**, 87
 - sickle cell, hematoporphyrin-sensitized photohemolysis, inhibition by lawsone (human), **139**, 780
 - sodium-potassium pump, decreased affinity for K⁺ in cystic fibrosis patients and heterozygotes, **137**, 1021
 - transport systems for S-dinitrophenyl glutathione and oxidized glutathione, comparison (human), **139**, 538
- Erythro-9-[3-(2-hydroxynonyl)]adenine
 - effect on actin polymerization and nucleotide exchange, **137**, 1181
- Escherichia coli*
 - adenylate kinase localization, **134**, 527
 - cDNA cloning and expression of rat liver arginase, **136**, 955

Subject Index

- cloning of phage λ -resistant genes from *Brevibacterium albidum* and *Proteus vulgaris*, **140**, 1101
- expression of
human interleukin-2 receptor cDNA, **141**, 804
kanamycin resistance gene, effect of specific yeast sequences, **136**, 529
and secretion of hepatitis B viral surface antigen, **135**, 1042
produced ornithine carbamyltransferase precursor, uptake and processing *in vitro* by rat cardiac and hepatic mitochondria, **134**, 21
related promoter genes, nalidixic acid effects on expression, **135**, 411
steady-state cultures, temperature effects, **135**, 708
yeast centromere plasmid application as shuttle vector, **135**, 915
- Esterase
Ca²⁺-dependent, identification and isolation (human heart), **136**, 773
- Esterification
cholesterol, associated apolipoprotein A-IV affinity for lipoproteins (human), **135**, 756
- Estradiol
and buserelin and tamoxifen, combined effects on tumor cell growth *in vitro* (human breast cancer), **140**, 550
secretion by immature Sertoli cells, *in vitro* effects of germ cells (rat), **141**, 861
- Estrogen
biosynthesis by reconstituted aromatase-cytochrome P-450 system, stereochemistry (human placenta), **136**, 454
induced glycoprotein secreted by breast cancer cells, identification as lysosomal acidic protease (human), **138**, 102
-like activity of phenol red, analysis in cultured anterior pituitary cells (rat), **141**, 885
regulation of albumin mRNA levels in hepatocytes *in vitro* (*Xenopus laevis*), **136**, 8
- Estrogen receptors
breast tumor cell, high-affinity hydroxytriphenylethylenes, inhibition of cell proliferation, mechanism (human), **140**, 523
native, urea-induced transformation: evidence for separate DNA- and ATP-binding sites (immature bovine uterus), **139**, 1250
nuclear, turnover and synthesis, R 5020 effects, analysis by density-shift method (hamster uterine decidual cells), **135**, 98
protein and phospholipid kinase activities (MCF-7 cells), **135**, 597
specific binding to chicken ovalbumin gene (bovine uterus), **139**, 396
steroid and DNA binding sites, computer-based search, **139**, 281
- Estrone
cytochrome P-450_{11 β} -catalyzed formation from 19-oxoandrostenedione, **140**, 530
effect on mitochondrial protein phosphorylation and cardiomyopathy (genetically diabetic mouse), **139**, 56
- Ethacrynic acid
selective inhibition of leukotriene C₄ synthesis (human neutrophils), **137**, 195
- Ethanol
effect on
corticotropin-releasing factor binding sites on erythrocyte membranes (human, rat), **136**, 137
prolactin binding sites in plasma membrane and Golgi fractions (rat liver, prostate), **134**, 1122
enhancement of arachidonic acid transmembrane penetration and activation of 5-lipoxygenase pathway (human leukocytes), **140**, 782
-metabolizing hepatocytes, generation of neutrophil chemotactic factor (human), **137**, 688
- Ethidium ion
cooperativity in formation of complexes with cytidylyl-3',5'-guanosine, **139**, 1164
- Ethoxycoumarin deethylase
inhibition by recombinant tumor necrosis factor (mouse), **136**, 316
- O-(4-Ethoxylbutyl)-berbamine
inhibition of calmodulin, **140**, 461
- Ethylene dibromide
induction of DNA damage at low doses (rat liver), **141**, 723
- Ethylketocyclazocine receptors
effects of swainsonine and tunicamycin, comparison (NCB-20 cells), **134**, 1387
- N-Ethylmaleimide
desensitization of K⁺/H⁺ antiporter pH dependence (*Vibrio alginolyticus*), **136**, 1030
- Etioplasts
detection of efficient and prolonged *in vitro* translation system (cucumber), **135**, 248
- Etoposide
peroxidative free radical formation and O-demethylation, **135**, 215
- Evans blue
inhibition of human T-cell lymphotropic virus type III/lymphadenopathy-associated virus cytotoxicity *in vitro*, **136**, 64
- Exercise
effect on hepatic glutathione levels (rat), **139**, 926
stimulation of α -aminoisobutyric acid transport by Na⁺,K⁺-ATPase, ouabain effects (rat skeletal muscle), **134**, 1342
sustained, effect on plasma aromatic and branched-chain amino acid levels and brain amine content (rat), **137**, 149
- Exo-1,4- β -glucanase
catalyzed hydrolysis, stereochemical course, ¹H NMR determination (*Cellulomonas fimi*), **139**, 487
- Extracellular matrix
defined components, melanoma cell adhesion (mouse), **139**, 515
- Eye-derived growth factor I
comparison with fibroblast growth factor purified from adult retina (bovine), **136**, 102

F

- Factor XIII
separation from fibrinogen by fibrinogen polymer formation (human plasma), **134**, 678

Subject Index

- Fast atom bombardment mass spectrometry
 high-sensitivity peptide assay: direct analysis of aqueous solutions, **141**, 1058
 in localization of phosphorylation sites on β -casein (butfalo), **140**, 28
- Fasting
 effect on
 adenylate cyclase sensitivity to stimulatory agonists, mechanism (rat), **138**, 1411
 carnitine palmitoyltransferase (newborn rat brown adipose tissue, liver), **135**, 589
 short-term, effect on insulin receptor autophosphorylation and kinase activity in streptozotocin-induced diabetes (rat), **140**, 850
- Fatty acid-CoA synthetase
 identification in bovine rod outer segments, **138**, 988
- Fatty acids
 -binding proteins from heart, liver, and intestine, purification and tissue expression (rat), **137**, 929
 blood levels, effects of recombinant DNA-derived methionyl human growth hormone 20-kDa variant (hypophysectomized rat), **138**, 342
 chain elongation system, *trans*-2-enoil CoA reductase component, effects of competitive substrates, inhibitors, and proteases (rat hepatic microsomes), **140**, 74
 composition in nuclear membranes, effect of dietary lipid content (murine liver), **135**, 655
 cotranslational attachment to nascent peptides in gastric mucus glycoprotein (rat), **141**, 387
 covalent attachment to epidermal macromolecules, characterization (swine), **137**, 992
 free, binding to *Clostridium botulinum* type E derivative toxin and fragments, **140**, 1015
 synthesis by
 isolated hepatocytes, effects of ionomycin and phorbol ester (rat), **139**, 403
 periportal and perivenous hepatocytes, zonal characteristics in primary culture, analysis (rat), **139**, 1055
- Fc receptors
 bacterial, binding of IgG, effect of various dipeptides, **135**, 1105
 -Fc receptor interactions, detection with spin-labeled IgG heterosaccharides in model immune complexes (mouse), **137**, 237
- Ferredoxin-NADP⁺ reductase
 inhibition by hexacyanochromate(III) ion (spinach), **141**, 578
- Ferricyanide
 stimulation of NADH oxidation (HeLa cells), **135**, 110
- Ferritin
 synthesis, regulatory mechanisms in malignant and non-malignant lymphoid cells (human), **139**, 652
- Fertilization
 src-related tyrosine-protein kinase activity, quantitative analysis (sea urchin egg), **138**, 349
- α -Fetoprotein
 synthesis in transformed fetal hepatocytes (rat), **135**, 844
- Fibrinogen
 binding to platelets, inhibition by ajoene (human), **141**, 145
 gene expression in megakaryocytes (rat), **140**, 543
 polymer formation, for factor XIII separation (human plasma), **134**, 678
- Fibrin-stabilizing factor, *see* Factor XIII
- Fibroblast growth factor
 acidic
 amino acid sequence analysis
 bovine brain, **133**, 554; *erratum*, **134**, 1419
 human brain, **138**, 611; **140**, 874
 amino-terminal sequence and mitogenic activity (human brain), **135**, 541
 chromosomal localization (human), **138**, 644
 enhancement of synthetic heparin pentasaccharide stimulation of nonvascular cells (human), **139**, 728
 induction of endothelial cell proliferation, inhibition by β -transforming growth factor (bovine aorta), **138**, 476
 related heparin-binding growth factors, isolation from established human tumor line lysates (bovine brain), **139**, 861
 stimulation of nonvascular cells, heparin effects (human), **137**, 1205
- basic
 amino-terminal extended form, isolation and characterization (bovine pituitary), **138**, 580
 amino-terminal sequence and mitogenic activity (human brain), **135**, 541
 chromosomal localization (human), **138**, 644
 induction of endothelial cell proliferation, inhibition by β -transforming growth factor (bovine aorta), **138**, 476
 stimulation of nonvascular cells, heparin effects (human), **137**, 1205
 -like substance, purification from adult retina, comparison with eye-derived growth factor I (bovine), **136**, 102
- Fibroblasts
 buoyant lysosomal fractions, *N*-acetylneuraminic acid accumulation in patients with generalized *N*-acetylneuraminic acid storage disease, **139**, 838
 carcinogen-exposed, poly(ADP-ribose) biosynthesis and suicidal NAD⁺ depletion, relationship (mouse), **138**, 1051
 chemotaxis, stimulation by bovine brain-derived growth factor (bovine, human), **141**, 67
c-myc oncogene expression, senescence-associated changes (human), **135**, 105
 collagen production and processing, stimulation by transforming growth factor- β (human), **138**, 974
c-ras oncogene expression, decrease during *in vitro* senescence (human), **137**, 917
 cultured, uptake of γ -irradiated low-density lipoprotein (Chinese hamster), **138**, 938
 cystic fibrosis and normal, chloride efflux, kinetic analysis (human), **134**, 233
 embryonic
 in culture, secretion of low-molecular-weight growth inhibitor (chicken), **139**, 1243
 phorbol ester-enhanced malignant transformation, inhibition by 3-aminobenzamide (mouse), **134**, 1334
 transformation induction by mitogenic growth factors with polyoma virus middle T antigen (rat), **136**, 921
 glucose-starved, hexose transport reversal by glucose re-feeding, effects of cycloheximide and tunicamycin (human), **140**, 844
 insulin release, induction by transfection with genomic rat islet DNA (mouse), **136**, 638; *erratum*, **139**, 852
 intercellular junctional communication, reversible inhibi-

Subject Index

- tion by glycyrrhetic acid (human), **134**, 29
- Kirsten and Rous sarcoma virus-transformed, ^{31}P NMR spectra (chicken, rat), **138**, 699
- low-density lipoprotein metabolism *in vitro*, effect of phospholipid polar head group manipulation (human), **137**, 43
- low-density lipoprotein receptors, synthesis, stimulation by diltiazem and verapamil (human skin), **136**, 845
- α_2 -macroglobulin synthesis and secretion (human lung), **134**, 57
- NIH 3T3, Kirsten virus-transformed, major excreted protein, protease specificity, comparison with cathepsin L (mouse), **139**, 156
- in primary cultures, heat shock response, comparison with established cell lines (human), **141**, 46
- ras* oncogene-transformed, extraction of hemolytic factor (human), **140**, 365
- serial divisions, effect on actin, globin, and interferon gene expression (human), **141**, 112
- stimulation by
- human fibroblast growth factors, heparin effects (hamster), **137**, 1205
 - synthetic heparin pentasaccharide, enhancement by human acidic fibroblast growth factors (hamster), **139**, 728
- 3T3, growth-stimulating activity of peptide purified from rat pancreatic juice, **139**, 545
- 3T6
- clonogenic growth, inhibition by 1,25-dihydroxyvitamin D_3 , role of receptors (mouse), **139**, 136
 - permeabilization by external ATP, **134**, 453
- transfected with plasmid containing cell cycle regulatory sequence, methylation pattern analysis (hamster), **135**, 942
- Fibroin
- silk, entrapment of phenylalanine ammonia-lyase, for protection from proteolysis (*Rhodotorula glutinis*), **141**, 165
- Fibronectin
- binding to amyloid P component: binding site localization on 31-kDa C-terminal domain (human), **140**, 12
 - isotypes expressed in adult and fetal pulmonary tissues, comparison (human), **141**, 1012
- Filaments
- Alzheimer's disease-associated, self-assembly of microtubule-associated protein τ from porcine brain into similar filaments, **141**, 790
- Films
- protein, ultrathin, direct current conductivity measurements (bovine), **133**, 1152; *erratum*, **135**, 1150
- Flavin
- oxidized, detection in reduced EH_2 intermediate of glutathione reductase, **136**, 22
- Flavin-adenine dinucleotide
- effect on 5-lipoxygenase activity (rat neutrophil extracts), **141**, 534
- Flavine mononucleotide
- effect on 5-lipoxygenase activity (rat neutrophil extracts), **141**, 534
- Flavobacterium* sp.
- pentachlorophenol catabolism, analysis, **141**, 825
- Flavoprotein
- NADPH-specific monomers, and cytochrome *P*-450 in solution, reconstitution of hepatic monooxygenase system (rabbit), **139**, 883
- Flow cytofluorimetry
- fluorescein fluorescence polarization, in lymphocyte activation assay (BALB/c mouse), **136**, 1021
- Flunitrazepam
- ^3H -labeled, photoaffinity labeling of γ -aminobutyric acid receptor binding site, comparison with [^3H]muscimol binding (bovine brain), **138**, 1308
- N*-2-Fluorenylacetylacetamide
- inducible cytochrome *P*-450, induction by phenobarbital (rat hepatoma cells), **137**, 120
- Fluorescein
- fluorescence polarization with flow cytofluorimetry, in lymphocyte activation assay (BALB/c mouse), **136**, 1021
- Fluorescence
- fluorescein, polarization measurement by flow cytofluorimetry, in lymphocyte activation assay (BALB/c mouse), **136**, 1021
 - steady-state emission from 5-iodoacetamidofluorescein bound to hemoglobin, **138**, 489
- Fluoride
- effect on Ca^{2+} intracellular levels and translocation (canine thyroid cells), **141**, 569
- Fluoropyrimidine
- metabolism, regulation by chlorpromazine (murine sarcoma cells), **138**, 1009
- Fodrin
- binding proteins, role in structure of neuronal postsynaptic density, regulation by phosphorylation (rat brain), **138**, 59
- Folinic acid-SF
- effect on methotrexate induction of DNA lesions (human colon adenocarcinoma cells), **138**, 54
- Follicle-stimulating hormone
- induced aromatase activity in granulosa cells, modulation by transforming growth factor and inhibin, comparison (rat), **136**, 969
 - secretion from pituitary
 - in monolayer system, stimulation by type β transforming growth factor (rat), **135**, 950
 - stimulation by homodimer of inhibin A β -subunits (pig), **138**, 1129 - α -subunit secreted by tumor cell lines, phosphorylation (human), **140**, 143
 - suppressible inhibiting factor of testicular steroidogenic activity, detection in extracts from Sertoli's cells (porcine testis), **139**, 169
- Follicular fluids
- demonstration of inhibin high-molecular-weight forms with monoclonal antibodies (bovine), **136**, 1103
 - identification of insulin-like growth factor II (human), **140**, 536
- Formate-tetrahydrofolate ligase
- catalyzed reaction, nucleotide stereochemistry, analysis (*Clostridium cylindrosporium*), **139**, 967
- Formyl-methionyl-leucyl-phenylalanine
- and adenosine, synergistic induction of cAMP levels in monocytes (human), **138**, 1376
 - chemotactic, stimulation of neutrophils, effect on translocation of protein kinase C (guinea pig), **139**, 1169
 - and concanavalin A, effect on NADPH oxidase activity

Subject Index

- and phosphoinositide turnover in Ca^{2+} -depleted neutrophils (human), **140**, 1
- identification as chemoattractant for human-mouse hybrid cells, **138**, 1001
- independent activation of phosphoinositide turnover and NADPH oxidase in Ca^{2+} -depleted phorbol ester-primed neutrophils (human), **135**, 785
- induction of
- neutrophil chemotaxis, inhibition by myelin basic protein (human), **141**, 524
 - respiratory burst, phosphoinositide hydrolysis, and Ca^{2+} mobilization, phorbol ester effects (human neutrophils), **135**, 556
 - oxidation by responding cells, effect on efficacy (human neutrophils), **138**, 66
- Formyltetrahydrofolate synthetase, *see* Formate-tetrahydrofolate ligase
- Forskolin
- adenylate cyclase sensitivity, increase by fasting, mechanism (rat fat cells), **138**, 1411
 - effect on
 - adenylate cyclase-phosphodiesterase system in thrombopathic and normal platelets (dog), **140**, 589
 - cAMP metabolism by thrombopathic platelets (dog), **140**, 595
 - enhancement of c-myc mRNA accumulation in cultured thyroid cells (dog), **141**, 1066
 - regulation of phorbol ester-induced atrial natriuretic peptide secretion (rat heart), **139**, 266
 - stabilizing effect on interaction between guanine nucleotide-binding stimulatory protein and catalytic unit of adenylate cyclase system (rat erythrocytes), **137**, 190
- Free radicals
- in lipoxygenase/linoleic acid system, characterization, **141**, 614
 - production in mitochondria, induction by hydroperoxide (rat liver), **141**, 1123
- β -Fructofuranosidase
- cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
- Fructose-bisphosphatase
- activity in brown and white adipose tissue (rat), **138**, 687
 - amino acid sequence homology among various species, **135**, 374
 - effects of
 - arabinose- and ribose-1,5-bisphosphate (rat liver), **138**, 159
 - zinc-deficient diet (rat liver), **134**, 944 - modification by chelators, effect on 5'-AMP sensitivity (fish, rat brain), **139**, 571
 - and phosphofructokinase, interaction at physiological concentrations, effect on kinetic behavior (rabbit muscle), **135**, 852
- Fructose-2,6-bisphosphatase
- effects of
 - diabetes (rat liver), **136**, 498
 - tolbutamide (rat liver), **139**, 687
- Fructose 2,6-bisphosphate
- activation of pyrophosphate-dependent sucrose metabolism in sucrose-importing plants, **141**, 440
 - role in regulation of basal- and thrombin-stimulated glycolysis (human platelets), **138**, 666
- Fructose-bisphosphate aldolase
- type A, mRNA expression in adult and fetal tissues and hepatoma (human), **134**, 1093
- Fructose 6-phosphate
- shunt activity in intact lens, dynamic assessment by proton NMR spectroscopy (rabbit), **138**, 1068
- Fructose-6-phosphate, 2-kinase, *see* 6-Phosphofructo-2-kinase
- FSH, *see* Follicle-stimulating hormone
- α -L-Fucosidase
- purification and acrosome reaction promoting property (bovine seminal plasma), **137**, 1061
- α -L-(1 \rightarrow 6)-Fucosidase
- detection and purification (*Aspergillus niger*), **136**, 563
- Fucosterol
- effect on angiotensin-converting enzyme levels and glucocorticoid receptors in endothelial cells (bovine carotid artery), **139**, 348
- Fumarate hydratase
- and L-aspartate ammonia-lyase, sequence homology (*Bacillus subtilis*, *Escherichia coli*, *Pseudomonas fluorescens*), **138**, 568
- Fumarate reductase, *see* Succinate dehydrogenase
- Fura-2
- in analysis of thrombin and ionomycin effects on platelet cytosolic Ca^{2+} levels (human blood), **139**, 308
 - loaded platelets, thrombin- and ADP-induced Ca^{2+} flux, kinetics (human), **136**, 1124
 - in smooth muscle cell monolayers, measurement of agonist-induced transients in cytosolic Ca^{2+} levels (hamster, rabbit, rat), **136**, 927
- Fura-2-am
- interaction with sarcoplasmic reticulum (rabbit skeletal muscle), **138**, 1153

G

GABA, *see* γ -Aminobutyric acid

D-Galactose

- specific lectin, precipitation by triantennary complex-type oligosaccharide (*Erythrina indica*), **141**, 963

Galactosides

- binding to lactose transporter from *Escherichia coli*, stoichiometry, confirmation by quantitative immunoprecipitation, **134**, 975

β -Galactosides

- binding lectin, 14-kDa, mRNA nucleotide sequence (chicken embryo), **134**, 51

Galactosyl receptors

- mediation of asialo-orosomucoid dissociation after internalization by hepatocytes: detection of distinct pathways (rat), **140**, 43

Galactosyltransferase, *see* Lactose synthase

Galanin

- inhibition of glucose-stimulated insulin release from pancreatic islets *in vitro*, mechanism (mouse), **140**, 1059

Gangliosides

- binding to *Clostridium botulinum* type E derivative toxin and fragments, **140**, 1015

- GM3, activation of protein kinase C in presence of calcium

Subject Index

- and 12-*O*-tetradecanoylphorbol-13-acetate, **138**, 865
- Gas chromatography-mass spectrometry
- 4-hydroxyalkenals in various tissues (rat), **139**, 144
- simultaneous quantitation of leukotriene B₄ and ω -oxidation products (human neutrophils), **134**, 512
- Gastric glands
- pentagastrin binding, characterization (guinea pig), **134**, 895
- Gastrin
- amino acid sequence homology with leucosulfakinin-II (*Leucophaea maderae*), **140**, 357
- gene, chromosomal assignment by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
- GDP (guanosine 5'-diphosphate)
- complex with Ha-*ras* oncogene-encoded p21, ³¹P NMR, **135**, 549
- inhibition of microtubule elongation, **137**, 351
- stimulation of [³H]phosphatidylinositol 4,5-bisphosphate hydrolysis by phospholipase C (human platelets), **140**, 728
- Gelsolin
- plasma, actin monomer sequential binding, inhibition by vitamin D-binding protein, **136**, 72
- Genes
- acid phosphatase, effect of vector type, host strains, and transcription terminator on expression (yeast), **140**, 602
- actin
- expression in fibroblasts, effect of serial divisions (human), **141**, 112
- promoter sequences, application in expression of hepatitis B surface antigen (*Saccharomyces cerevisiae*), **136**, 1063
- acute phase protein, association with hepatic nuclear matrix during inflammation (rat), **137**, 633
- α -amylase inhibitor, from *Streptomyces griseosporus*, cloning and expression in *S. lividans*, **141**, 1099
- androgen-dependent, relationship to nuclear matrix, basis for prediction of transcriptional activity (rat prostate), **138**, 1334
- apolipoproteins CI and E, on chromosome 19, mapping (human), **136**, 876
- arginase, from kidney and liver, homology (human), **141**, 53
- atrial natriuretic peptide precursor, isolation and nucleotide sequence (bovine), **136**, 396
- from autolysin-defective *Streptococcus pneumoniae* mutants, molecular characterization, **137**, 614
- calcitonin, related peptide
- partial homology with N-terminal sequence of amyloid fibril protein from insulin-producing pancreatic tumor (human), **140**, 827
- structure-activity relationship (human), **134**, 1306
- calmodulin, increased expression in dystrophic muscle (chicken), **137**, 507
- casein, expression, hydrocortisone transcriptional and post-transcriptional roles (rat mammary explants), **134**, 812
- c-erbB-2*, chromosomal assignment by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
- c-fos*, nucleosome relaxation during differentiation (HL-60 cells), **141**, 213
- c-Ha-ras*, expression in
- aflatoxin B₁-induced hepatocellular carcinomas (rat), **138**, 858
- early second trimester fetal tissues (human), **141**, 510
- citrate synthase, molecular cloning (*Acinetobacter anitratum*), **141**, 797
- c-Ki-ras*
- activated, point mutation-induced amplification (human pancreatic adenocarcinoma cells), **140**, 167
- codon 12 GGT-CGT point mutation, occurrence in lung cancer (human), **140**, 699
- c-mos* proto-oncogenic, expression in undifferentiated teratocarcinoma cells, **140**, 477
- c-myc*
- corresponding mRNA, accumulation in cultured thyroid cells, enhancement by epidermal growth factor, forskolin, phorbol ester, and thyrotropin (dog), **141**, 1066
- expression in
- aflatoxin B₁-induced hepatocellular carcinomas (rat), **138**, 858
- fibroblasts, senescence-associated changes (human), **135**, 105
- liver, induction by dietary manipulation (rat), **140**, 574
- first exon, role in protein synthesis regulation (human), **140**, 313
- methylation, changes during aging process (mouse), **139**, 1299
- nucleosome relaxation during differentiation (HL-60 cells), **141**, 213
- point mutation-induced amplification (human pancreatic adenocarcinoma cells), **140**, 167
- serial expression during hematopoiesis *in vitro* (human), **135**, 521
- topoisomerase II-induced cleavage sites, *in vivo* stimulation by antitumor drugs (N417 cells), **137**, 154
- transcript levels, correlation with substance K-induced DNA synthesis (quail fibroblasts), **137**, 167
- codon bias, protection of protein spatial structures from destabilization by random single-point mutations, **139**, 1094
- colony-stimulating factor, expression during monocytic differentiation of HL-60 cells, **141**, 924
- c-ras*, expression decrease during *in vitro* senescence (human fibroblasts), **137**, 917
- c-src*-related tyrosine-protein kinase, quantitation (sea urchin egg), **138**, 349
- DNA-mediated transfer into epidermal cells by electroporation (mouse), **137**, 244
- DNA polymerase α , identification in hybrid DNA segment (human), **139**, 637
- EJ-*ras*^H, transfected mammary epithelial cells, collagen processing (mouse), **137**, 161
- encoding hydrogen-oxidizing enzymes
- conjugal transfer from *Alcaligenes hydrogenophilus* to *Pseudomonas oxalaticus*, **137**, 108
- isolation from *Alcaligenes hydrogenophilus* and expression in *Pseudomonas oxalaticus*, **137**, 114
- fibrinogen, expression in megakaryocytes (rat), **140**, 543
- fibroblast growth factor, chromosomal localization (human), **138**, 644
- galactosyltransferase, chromosomal localization (human), **141**, 1185

Subject Index

- gastrin, chromosomal assignment by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
- globin
- expression in fibroblasts, effect of serial divisions (human), **141**, 112
 - type β , supercoil-dependent structure, recognition by chromosomal high-mobility-group protein (chicken), **138**, 910
 - types β and ϵ , 5'-flanking sequences, role in expression specificity (human, murine erythroleukemia cells; nonerythroid cells), **134**, 128
- heterologous, cloned in *Streptomyces* plasmid pIJ702, structural stability, **140**, 372
- homoeobox-containing, specific expression during drug-induced teratocarcinoma cell differentiation, **137**, 520
- H-ras, expression in transformed NIH 3T3 cells, effect on cAMP-dependent protein kinase activity, **139**, 787
- 21-hydroxylase, restriction mapping and restriction fragment length polymorphisms (human), **136**, 722; *erratum*, **138**, 502
- IncFII plasmid, *polA* mutation, enhancement of instability of basic replicon, **139**, 918
- inhibin A subunit, expression in immature ovaries, stimulation by pregnant mare serum gonadotropin (rat), **138**, 1191
- interferon α , sequence and expression: homology with enhancer elements in regulatory region (mouse), **138**, 826
- interferon β , expression in fibroblasts, effect of serial divisions (human), **141**, 112
- interleukin 1, expression in chondrocytes: mechanism for autocrine control of cartilage matrix degradation (bovine, human), **141**, 904
- kanamycin resistance, expression in *Escherichia coli*, effect of specific yeast sequences, **136**, 529
- lipase, molecular cloning and nucleotide sequence analysis (*Pseudomonas fragi*), **141**, 185
- mdr1*, associated with multidrug resistance, encoding of P-glycoprotein (KB cells), **141**, 956
- middle-T-only, polyoma virus, transformed Ob17 cells, extracellular intermediates, promotion of Ob17 preadipocyte proliferation and differentiation, **137**, 378
- nah* operon in plasmid NAH7, regulation by *nahR* region, **141**, 986
- nerve growth factor, cloning: mRNA distribution in developing and adult tissues (chicken), **141**, 1116
- N-ras, expression in
- aflatoxin B₁-induced hepatocellular carcinomas (rat), **138**, 858
 - early second trimester fetal tissues (human), **141**, 510
- oncogenic
- expression in PLC/PRF/5 hepatoma cells, Northern blot analyses (human), **135**, 262
 - tyrosine kinase products, relationship to heat shock protein hsp 90, **138**, 1298
- ovalbumin, specific binding of estrogen receptor (chicken), **139**, 396
- pepsinogen A, chromosomal, close linkage (human), **138**, 289
- P31, hepatitis B virus
- expression in *Saccharomyces cerevisiae*, **138**, 268; *erratum*, **140**, 774
 - modification, subsequent synthesis of modified particles in yeast, **141**, 942
- phage λ -resistant, from *Brevibacterium albidum* and *Proteus vulgaris*, cloning into *Escherichia coli*, **140**, 1101
- 6-phosphofructokinase, genomic clones, isolation and characterization (human liver), **141**, 374
- phosphoenolpyruvate carboxykinase, expression regulation by insulin during preadipose cell differentiation, **138**, 468
- pol*, endogenous A-type retrovirus, relationship to mRNA from colon tumor and surrounding mucosa (human), **139**, 1071
- promoter, nalidixic acid effects on expression of related types (*Escherichia coli*), **135**, 411
- protease, human T-cell leukemia virus type I, identification, **139**, 129
- proto-oncogenic, sequential expression during differentiation of erythroleukemia cells (mouse), **135**, 1112
- PrP on p arm of chromosome 20, localization of homologous human gene (scrapie-infected hamster brain), **140**, 758
- pseudo-, cytochrome P-450-like, isolation and characterization (rabbit liver), **134**, 499
- ras*, encoded proteins, effect on phosphorylation of mitochondrial membrane proteins (rat liver), **135**, 316
- rRNA, number and organization (*Mycobacteria* sp.), **136**, 1136
- sal* operon in plasmid NAH7, regulation by *nahR* region, **141**, 986
- for serine-rich proteins, transcriptional expression, modulation by glucose (*Saccharomyces cerevisiae*), **138**, 297
- serum amyloid A, localization on chromosome 11 and detection of DNA polymorphisms (human), **137**, 1196
- 7, in *Agrobacterium tumefaciens* T-DNA, expression in *Xenopus laevis* oocytes, **141**, 1242
- spacer, detection of length and sequence polymorphisms (*Triptoneustes gratilla* ribosomes), **137**, 834
- 5 S RNA, specific transcription factor, modification of DNA linking number, **134**, 1086
- synthetic, assembly *in vitro*, **134**, 407
- thyroglobulin
- activation by thyrotropin and cAMP in hormonally depleted FRTL-5 cells, **141**, 1261
 - expression, relationship to gene methylation pattern (human), **134**, 1109
- thyrotropin β -subunit, chromosomal assignment by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
- transcriptionally active, affinity isolation (chicken erythrocytes), **137**, 474
- transferred into hematopoietic stem cells by electroporation, expression (mouse), **141**, 1018
- β -tubulin, detection in recombinant phages constructed from mung bean genomic DNA, **137**, 788
- U3 RNA
- sequence homology with vertebrate U3 RNA (*Drosophila melanogaster*), **138**, 512
 - small nucleolar, isolation and characterization (human), **137**, 1133
- uvrB*, mutation, effect on reversible DNA intercalator-induced frameshift mutations (*Salmonella typhimurium*), **138**, 505

Subject Index

- v-fos*, stimulation of α_1 (III) collagen gene expression (NIH 3T3 cells), **136**, 1042
v-ras^H, linked to murine mammary tumor virus promoter, expression, cAMP effect, **134**, 436
v-rel, protein product, identification in reticuloendotheliosis virus T-transformed bone marrow cells, expression in Cos1 cells (chicken), **134**, 716
v-src, encoded protein, related tyrosine kinase phosphatidylinositol kinase activity, isolation (*Narke japonica*), **139**, 473
- Genistein
 induction of cytochrome P-450 (*Streptomyces griseus*), **141**, 405
- Germ cells
 effects on *in vitro* secretion of androgen-binding protein and estradiol-17 β by immature Sertoli's cells (rat), **141**, 861
- Glibenclamid, *see* Glyburide
- α -Globin
 gene expression in fibroblasts, effect of serial divisions (human), **141**, 112
 transcriptionally active gene, affinity isolation (chicken erythrocytes), **137**, 474
- β -Globin
 gene
 5'-flanking sequences, role in expression specificity (human, murine erythroleukemia cells; nonerythroid cells), **134**, 128
 supercoil-dependent structure, recognition by chromosomal high-mobility-group protein (chicken), **138**, 910
- ϵ -Globin
 gene, 5'-flanking sequences, role in expression specificity (human, murine erythroleukemia cells; nonerythroid cells), **134**, 128
- Globins
 mRNA storage during protein synthesis inhibition by heme deprivation (rabbit), **134**, 1048
- α_{2u} -Globulin
 mRNA, phenobarbital-induced: characterization of cDNA clone (rat liver), **134**, 1182
- Glomeruli
 cGMP levels, enhancement by atrial natriuretic peptide (rat kidney), **136**, 947
 diabetic, impaired contractile responsiveness to angiotensin II (rat), **136**, 1185
- Glomerulosa cells
 aldosterone secretion, K⁺-induced, inhibition by diazepam (bovine adrenal), **135**, 994
 formation of inositol phosphates, role of guanyl nucleotide-binding proteins (rat adrenal gland), **140**, 941
- Glucagon
 binding to isolated epididymal adipocytes, insulin effects (rat), **141**, 812
 pancreatic secretion, cysteamine effect (rat), **134**, 1291
- Glucan 1,4- α -glucosidase
 mRNA, translation *in vitro* (rat intestine), **134**, 37
- β -1,3-Glucan receptors
 and peptidoglycan receptors, identification in prophenol-oxidase-activating system (silkworm), **141**, 1177
- Glucoamylase, *see* Glucan 1,4- α -glucosidase
- Glucocerebrosidase
 routing to lysosomes, role of oligosaccharide chains (U937 cells), **141**, 452
- Glucocorticoid receptors
 and angiotensin-converting enzyme, endothelial cell levels, reduction by fucosterol (bovine carotid artery), **139**, 348
 binding of RU 38486 (rat liver, thymocytes), **134**, 44
 complexes, binding to hepatic nuclear matrix, evaluation (rat), **137**, 640
 steroid and DNA binding sites, computer-based search, **139**, 281
 and triamcinolone acetonide antibodies, steroid-binding domains, comparison (rat), **136**, 470
 unactivated complex, association with RNA (human lymphoid cells), **140**, 204
- Glucocorticoids
 effect on corticotropin-releasing factor binding to peripheral tissues (rat), **134**, 255
- Glucokinase
 anomeric specificity, analysis, **141**, 931
 hepatic activity, regulatory factors (chicken), **139**, 1086
- Gluconeogenesis
 effect of RU 38486 (rat liver), **134**, 44
 by periportal and perivenous hepatocytes, retention of zonal characteristics in primary culture (rat), **139**, 1055
- Glucose
 activation of glycogen synthase, role of glucose 6-phosphate (rat hepatocytes), **141**, 1195
 anomeric forms produced by glucose-6-phosphatase and used by glucokinase, analysis, **141**, 931
 blood levels, effects of recombinant DNA-derived methionyl human growth hormone 20-kDa variant (hypophysectomized rat), **138**, 342
 deprivation, effect on insulin and insulin-like growth factor I binding to CHO cells, **140**, 821
 effect on
 insulin release
 after α_2 -adrenergic activation (murine pancreas), **139**, 557
 from cloned B-cell line HIT-T15 (hamster), **140**, 616
 pancreatic preproinsulin mRNA levels after oral administration (rat), **141**, 1201
 self-transport in Ehrlich ascites tumor cells (mouse), **134**, 1231
 induction of hexose transport reversal in glucose-starved fibroblasts, effects of cycloheximide and tunicamycin (human), **140**, 844
 metabolism, alterations induced by DNA damage-mediated poly(ADP-ribose) polymerase activation (murine L1210 cells), **134**, 227
 -regulated proteins, induction in fibroblasts, effect of extracellular Ca²⁺ and temperature (hamster), **140**, 881
 regulation of serine-rich protein gene expression (*Saccharomyces cerevisiae*), **138**, 297
 repression and derepression, effect on intracellular distribution of enzymes (*Saccharomyces cerevisiae*), **134**, 285; *erratum*, **136**, 843
 stimulation of insulin release from pancreatic islets *in vitro*, inhibition by galanin, mechanism (mouse), **140**, 1059
 uptake in bone marrow-derived macrophages, stimulation by colony-stimulating factor 1 (mouse), **138**, 445
- Glucose-1,6-bisphosphatase
 effect of diabetes (rat liver), **136**, 498

Subject Index

- D-Glucose dehydrogenase (pyrroloquinoline-quinone)
 holoenzyme formation, essential role of 9-carboxyl group of pyrroloquinoline quinone as prosthetic group, **139**, 1279
- Glucose-6-phosphatase
 activity in brown and white adipose tissue (rat), **138**, 687
 anomeric specificity, analysis, **141**, 931
 induction by dexamethasone in cultured hepatoma cells, **139**, 1130
- Glucose 6-phosphate
 role in glycogen synthase activation by glucose (rat hepatocytes), **141**, 1195
- Glucose-6-phosphate dehydrogenase
 effects of epidermal growth factor, growth hormone, and thyroxine after chronic administration (hypophysectomized rat), **141**, 78
- β -Glucosidase
 -catalyzed hydrolysis, stereochemical course, ^1H NMR determination (almond), **139**, 487
- Glucuronic acid
 in brain glycoproteins, HNK-1 antibody-reactive, demonstration (goldfish), **138**, 902
- β -Glucuronidase
 cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
 release from monocytes, effect of κ elastin (human), **141**, 92
- Glutamate-ammonia ligase
 adenylylated, conversion to phosphotyrosyl form by micrococcal nuclease (*Mycobacterium smegmatis*), **137**, 716
 cDNA, selection from renal library and restriction map construction (rat), **134**, 146
- Glutamic-oxaloacetic transaminase, *see* Aspartate aminotransferase
- Glutamine
 utilization pathway in lymphocytes, intracellular distribution of enzymes (rat), **138**, 318
- Glutamine synthetase, *see* Glutamate-ammonia ligase
- γ -Glutamyltransferase
 deglycosylated and native forms, biochemical and immunologic analysis (rat kidney), **135**, 304
 subunits, electrophoretic behavior and molecular weights, effect of enzyme glycosylation (bovine, human, rat kidney), **141**, 1189
- γ -Glutamyl transpeptidase, *see* γ -Glutamyltransferase
- Glutathione
 abnormal metabolism in platelets, role in deficient leukotriene C_4 synthesis (human), **140**, 857
 hepatic, severe depletion during physical exercise (rat), **139**, 926
 inhibition of 4-nitrobenzyl mercaptan S-sulfate covalent binding to cytosolic proteins (rat liver), **137**, 1055
 oxidized, and S-dinitrophenyl glutathione, transport mechanism in erythrocytes, comparison (human), **139**, 538
 regioselective conjugation of 7,12-dihydroxymethylbenz[a]anthracene via reactive 7-hydroxymethyl sulfate ester (rat hepatic cytosol), **134**, 100
 role in
 stress protein induction (fetal rat fibroblasts), **134**, 492
 sulfite detoxication (bovine kidney), **136**, 1036
- Glutathione peroxidase
 activity in muscle, liver, and lung tissues of inherited muscular dystrophy-affected chickens, **134**, 550
- Glutathione reductase
 activity in muscle, liver, and lung tissues of inherited muscular dystrophy-affected chickens, **134**, 550
 reduced intermediates, structure, resonance Raman spectroscopy, **136**, 22
- Glutathione transferase
 activity in muscle, liver, and lung tissues of inherited muscular dystrophy-affected chickens, **134**, 550
 cationic, characterization (human erythrocytes), **137**, 1174
 expression in seminal vesicles and pituitary gland, effect of xenobiotics (rat), **136**, 1057
 H_a subunit 1 cDNA, nucleotide sequence analysis (human liver), **141**, 229
 isoenzymes from human and rat tissues, differential inhibition by plant phenols, **141**, 1170
 major hepatic form, developmental and hormonal regulation (male mouse), **140**, 581
 stereoselective inhibition by reaction products (rat liver), **141**, 892
- Glyburide
 effect on insulin release from cloned B-cell line HIT-T15 (hamster), **140**, 616
- Glycanase
 bacteriophage-associated, hydrolysis of 3-deoxy-D-manno-2-octulosonic acid β -ketopyranosidic linkages (*Escherichia coli*), **136**, 329
- Glyceraldehyde-3-phosphate dehydrogenase
 domain boundaries, prediction from amino acid sequence (pig), **139**, 11
- Glycine
 conjugation with cholic acid by hepatic peroxisomes (rat), **138**, 167
 glycol derivative, effect on IgG binding to bacterial Fc receptors, **135**, 1105
- Glycogen
 -concanavalin A interaction, effect on K^+ currents through lipid bilayer membranes, **141**, 912
 deposition in cultured hepatocytes, comparative effects of insulin and insulin-like growth factors (chick embryo), **134**, 427
 metabolism in cultured cerebromicrovascular smooth muscle and endothelium, regulatory mechanism, comparison (rat), **134**, 484
- Glycogen phosphorylase
 calcium-dependent activation by nerve growth factor (rat pheochromocytoma PC12 cells), **138**, 917
 regulation by amylose-precipitating factor (rabbit muscle), **135**, 701
- Glycogen(starch) synthase
 C-terminus, identification (rabbit skeletal muscle), **137**, 542
 kinetic properties after phosphorylation by glycogen synthase kinase 4 (rabbit, rat skeletal muscle), **134**, 1129
- Glycogen synthase, *see* Starch(bacterial glycogen) synthase
- Glycogen synthase I, *see* Glycogen(starch) synthase
- Glycogen synthase kinase 4, *see* Protein kinase
- Glycolipids
 Forssman, flow between epithelial cells: model of tight junction structure (MDCK cells), **138**, 1163
- Glycolysis
 basal- and thrombin-stimulated, regulatory role of fructose 2,6-bisphosphate (human platelets), **138**, 666

Subject Index

Glycopeptides

high-mannose-type, precipitation of concanavalin A, stoichiometry, **137**, 670

-hydrolyzing α -L-(1 \rightarrow 6)-fucosidase, detection and purification (*Aspergillus niger*), **136**, 563

Glycophorin

and dioleoylphosphatidylethanolamine-containing liposomes, trypsin-induced destabilization, **141**, 973

Glycoproteins

amyloid P component, isolation from Engelbreth-Holm-Swarm tumor (mouse), **134**, 995

band 3, phosphorylation by tyrosine kinase activity associated with EGF receptors and *src* proteins (human erythrocytes), **135**, 720

detergent-insoluble, 120- and 80-kDa, as markers of differentiation and adhesion in B16 melanoma cells, **140**, 638

and/or erythrocyte membrane proteins, functional group labeling with biotin derivatives and 125 I-labeled streptavidin (human), **136**, 80

estrogen-induced, secreted by breast cancer cells, identification as lysosomal acidic protease (human), **138**, 102

HNK-1 antibody-reactive, glucuronic acid, demonstration (goldfish brain), **138**, 902

lymphadenopathy-associated virus/human T-lymphotropic virus type III, expression, suppression by glycosylation inhibitors, **141**, 33

nascent peptides, cotranslational attachment of fatty acids (rat gastric mucus), **141**, 387

P170, encoding by *mdr1* gene associated with multidrug resistance (KB cells), **141**, 956

polygalactosamine-containing, occurrence of Gal β (1 \rightarrow 3) GalNAc-Ser/Thr in linkage region (*Cordyceps ophioglossoides*), **140**, 350

surface, variant incorporated into liposomes, interactions with lectins (*Trypanosoma brucei brucei*), **141**, 1274

Glycosaminoglycans

synthesis in costal chondrocytes, effect of human recombinant interleukin 1 (rat), **140**, 386

Glycosaminoglycan sulfates

role in apolipoprotein E-induced lipid uptake by hepatocyte monolayers (rat), **141**, 158

Glucose phosphotransferase

-phosphoenolpyruvate system, III^{gic} protein, role in adenylate cyclase inhibition by inorganic orthophosphate (*Escherichia coli*), **141**, 1138

Glycosphingolipids

-cleaving enzyme, identification and characterization (leech), **141**, 346

Glycosylation

γ -glutamyl transpeptidase, effect on electrophoretic behavior and molecular weights of subunits (bovine, human, rat kidney), **141**, 1189

prolactin, inhibition of insulin-like activity in adipose tissue *in vitro* (rat), **134**, 764

Glycyrrhetic acid

reversible inhibition of intercellular junctional communication (human fibroblasts), **134**, 29

Glyoxalase I. *see* Lactoylglutathione lyase

Golgi apparatus

membrane fraction, detection of putative opioid-peptide processing activity (rat), **138**, 356

prolactin binding sites, ethanol effects (rat liver, prostate), **134**, 1122

Gonadotropin

human chorionic, *see* Human chorionic gonadotropin

-inducible high-density lipoprotein receptors, identification in solubilized ovarian membranes (rat), **134**, 1006

pregnant mare serum, stimulation of inhibin A-subunit gene expression in immature ovaries (rat), **138**, 1191

Gonadotropin receptors

subunit structure, preservation during corpus luteum regression (rat), **137**, 546

Gonadotropin-releasing hormone

and analogs, *in vitro* and *in vivo* effects, comparison (chicken, mammal, salmon), **138**, 419

enhancement of polyphosphoinositide hydrolysis (rat pituitary cells), **134**, 861

stimulation of luteinizing hormone secretion

arachidonic acid and protein kinase C coordinate actions (rat pituitary), **134**, 134

effect of wheat germ agglutinin (rat pituitary), **140**, 750

Gramicidin

alteration of accessibility of surface thiols (human lymphocytes), **140**, 999

Granulosa cells

accumulation and catabolism of progesterone *in vitro*, effects of luteinizing hormone and human chorionic gonadotropin (rat), **141**, 1025

progesterone metabolism, effects of prostaglandins E₂ and F_{2 α} (rat), **135**, 764

Growth factors, *see also specific growth factors*

brain-derived

chemotactic activity for fibroblasts and astroglial cells (bovine, human, rat), **141**, 67

transforming growth factor activity (NRK cells), **139**, 619

effect on protein kinase C translocation from soluble to particulate cell fraction (rat hepatocytes), **138**, 1328

heparin-binding, related to bovine acidic brain fibroblast growth factor, isolation from established tumor cell lysates (human), **139**, 861

macrophage-derived, endothelial cell-stimulating, partial purification and properties (human), **134**, 344

Growth hormone, *see* Somatotropin

Growth hormone-releasing factor

amphiphilic analogs: peptide design and biological activity *in vivo* (human), **139**, 763

and clonidine, comparative effects on stimulation of growth hormone biosynthesis in neonatal pituitary (rat), **138**, 1223

immunoreactive, detection in milk (human), **135**, 1084

Growth inhibitor

low-molecular-weight, secretion in cultures of embryo fibroblasts (chicken), **139**, 1243

GTP (guanosine 5'-triphosphate)

-activated adenylate cyclases, comparison as cholera toxin substrates, **134**, 928

-binding domain, exchangeable, localization in β -tubulin (bovine brain), **135**, 1135

complex with Ha-*ras* oncogene-encoded p21, 31 P NMR, **135**, 549

effect on

prostaglandin E₂ receptor binding affinity (porcine brain), **135**, 282

Subject Index

- vincristine-tubulin complex formation and stability in tumor cytosols (human), **135**, 695
- inhibition of platelet-activating factor binding to neutrophil membranes (human), **141**, 353
- stimulation of
- [³H]phosphatidylinositol 4,5-bisphosphate hydrolysis by phospholipase C (human platelets), **140**, 728
 - phosphoinositide hydrolysis in isolated platelet membranes (human), **137**, 801
 - phosphoinositide turnover in olfactory cilia (catfish), **137**, 36
- GTPase
- activity of N-ras proteins *in vitro* (NIH 3T3 cells), **139**, 771
 - thyrotropin-releasing hormone-induced activity, mediation by guanine nucleotide-binding protein (GH₃ cells), **138**, 1383
- Guanine nucleotide exchange factor
- purification, properties, and role in protein synthesis initiation (HeLa cells), **134**, 1160
- Guanine nucleotides
- binding proteins
 - ADP-ribosylated by pertussis toxin, detection in spermatozoa (invertebrates, vertebrates), **138**, 728; *addendum*, **140**, 773
 - mediation of thyrotropin-releasing hormone-induced GTPase activity and phosphoinositide metabolism (GH₃ cells), **138**, 1383
 - role in formation of inositol phosphates in adrenal glomerulosa cells (rat), **140**, 941
 - stimulatory, functional coupling with adenylate cyclase catalytic unit, relationship to enzyme inhibition by pentobarbital (rat brain), **140**, 237
 - dependent regulatory protein, stimulation of phospholipase C binding to substance P receptors (rat parotid gland), **136**, 362
 - effect on quinuclidinyl benzilate binding to muscarinic receptors (rat heart), **134**, 172
- Guanosine
- reaction with 1,3-bis(2-chloroethyl)-1-nitrosourea: evidence for mechanism of DNA modification, **139**, 31
- Guanosine 3':5'-cyclic monophosphate, *see* Cyclic GMP
- Guanylate cyclase
- particulate, association with atrial natriuretic peptide receptors (bovine lung), **140**, 101
 - phosphate loss during fertilization, stoichiometry (*Arbacia punctulata* sperm), **137**, 1148
- ## H
- Halobacterium*
- H. halobium*, flash spectrophotometric identification of rhodopsin-like pigment, **139**, 389
 - H. volcanii*, occurrence of cAMP, **136**, 1078
- Haloethane
- and L-carnitine, antagonistic effects on mitochondrial energy-linked processes, role of long-chain acyl CoA (rat liver), **139**, 303
- Haplotypes
- H-2, role in polyriboinosinic-polyribocytidylic acid effect on xanthine oxidase and cytochrome P-450-mediated enzymes (mouse), **138**, 246
- Heart
- atrial natriuretic factor levels
 - decrease in spontaneously hypertensive and spontaneously hypertensive stroke-prone rats, **135**, 74
 - effect of chronically administered adrenergic and cholinergic agonists (rat), **136**, 510
 - radioimmunoassay (hyper-, hypothyroid rats), **134**, 178
 - atriopeptin levels, effect of dietary salt and water intake manipulations (rat), **137**, 922
 - detection of Ca²⁺-dependent hydrolases (human), **136**, 773
 - hypertrophy, stress-induced, role of initial protein synthesis suppression (rat), **135**, 857
 - myocardium
 - hypoxia-, hyperkalemia-, and acidosis-induced changes, procaine effects (dog), **134**, 900
 - phosphate metabolite levels, rhythmic changes during cardiac cycle, depth-selected EKG-gated ³¹P NMR analysis *in vivo* (rat), **135**, 808
 - reperfusion injury, allopurinol beneficial action, mechanism, **137**, 76
 - release of
 - atriopeptin *in vitro*, regulation by α₁-adrenergic receptor (rat), **137**, 94
 - prostaglandins *in vitro*, enhancement by cytochrome P-448 inducers (chick embryo), **136**, 582
 - transcriptional and translational processes, heat shock-induced changes (dog), **134**, 166
- Heat shock
- associated inhibition of protein synthesis, role of eIF-2α phosphorylation (HeLa cells), **138**, 519
 - associated proteins
 - induction in fibroblasts, effect of extracellular Ca²⁺ and temperature (hamster), **140**, 881
 - retinal, association with cytoskeletal elements and binding to calmodulin, analysis (rabbit), **139**, 974
 - destabilization of cytoplasmic murine mammary tumor virus RNA, prevention by cycloheximide, **137**, 1028
 - effect on primary cell cultures and established cell lines, comparison (human), **141**, 46
 - induction of
 - intracellular Ca²⁺ and inositol trisphosphate increases (CHO cells), **137**, 826
 - phosphorylation of 26- and 27-kDa heat shock proteins (*Drosophila melanogaster*), **141**, 426
 - specific protein synthesis by HeLa cells exposed to 42°C, **137**, 957
 - transcriptional and translational changes (canine heart), **134**, 166
- Hematin
- hydroperoxide-dependent epoxidation of 3,4-dihydroxy-3,4-dihydrobenzo[*a*]anthracene, **140**, 181
 - induced hepatic heme oxygenase isoforms, comparison with splenic isoforms (rat), **141**, 528
- Hematopoiesis
- associated serial expression of *c-myc* oncogene (human), **135**, 521
- Hematopoietic cells
- cytochrome P-450c induction by β-naphthoflavone, analysis (fetal rat liver), **141**, 299
- Hematoporphyrin
- derivatives, phototoxicity enhancement by selenium (murine bladder tumor cells), **140**, 1086

Subject Index

- phototoxicity evaluation using *Paramecium aurelia*, **139**, 79
- Heme**
- attachment to apocytochrome *c*, catalyzing enzyme, localization in yeast mitochondria, **141**, 1145
 - deprivation, induction of protein synthesis inhibition, promotion of globin mRNA storage (rabbit), **134**, 1048
 - endogenous, excess levels, inhibition of mitochondrial L-alanine:4,5-dioxovalerate transaminase (rat liver), **138**, 751
 - hepatic and renal levels, effect on L-alanine:4,5-dioxovalerate transaminase induction by phenylhydrazine (rat), **138**, 742
 - in microsomal cytochrome *P*-450, covalent binding to apo-protein during enzyme destruction (rat liver), **138**, 193
 - peroxide complex formed in Fe(III)-tetraphenylporphyrin system, characterization, **135**, 972
- Heme oxygenase**
- splenic isoforms, characterization and comparison with hematin-induced and constitutive hepatic isoforms (rat), **141**, 528
- Hemiacetylcarnitinium**
- inhibition of carnitine acetyltransferase (pigeon breast), **138**, 735
- Hemin**
- effect on δ -aminolevulinate synthase mRNA levels (embryonic chicken liver), **140**, 81
 - polyethylene glycol-modified, peroxidase activity in organic solvents, **138**, 283
 - thiolate complexes, scission of plasmid DNA strands, **136**, 645
- Hemocytes**
- coagulating lectin, purification and characterization (*Manduca sexta*), **137**, 729
- Hemoglobin**
- bound 5-iodoacetamidofluorescein, steady-state fluorescence emission (human), **138**, 489
 - domain boundaries, prediction from amino acid sequence (human), **139**, 11
 - hematic proportions, bleeding-induced changes, effect of acetylsalicylic acid (adult rat), **137**, 69
 - unstable, detection (mouse), **139**, 551
- Hemolytic factor**
- purification from *ras* oncogene-transformed fibroblasts (human), **140**, 365
- Heparin**
- binding growth factors related to bovine acidic brain fibroblast growth factor, detection in tumor cell lysates (human), **139**, 861
 - cofactor II, partial cDNA clone, isolation and characterization (human), **137**, 431
 - effect on nonvascular cell stimulation by human acidic and basic fibroblast growth factors, **137**, 1205
 - endogenous, internalization and metabolism (bovine adrenal capillary cells), **140**, 294
 - fragments, partial structure, mass spectrometric method for rapid determination (porcine mucosa), **139**, 18
 - highly reactive, binding to apolipoprotein E: identification of heparin-binding domains (human), **134**, 783
 - releasable lipoprotein lipase, activation by cAMP-dependent protein kinase (rat heart), **135**, 196
 - synthetic pentasaccharide, stimulation of nonvascular cells, enhancement by human acidic fibroblast growth factors (bovine lens epithelial cells, hamster fibroblasts), **139**, 728
- Hepatitis B virus**
- subtype *adw*, pre-S2 region nucleotide sequence, comparison with other subtypes, **135**, 382
 - surface antigens
 - expression and secretion in *Escherichia coli*, **135**, 1042
 - expression by yeast actin gene promoter, **136**, 1063
 - P31 gene, expression in *Saccharomyces cerevisiae*, **138**, 268; *erratum*, **140**, 774
 - recognition by antibodies produced by administration of murabutide associated with synthetic pre-S viral peptide conjugated to toxoid carrier, **140**, 557
 - synthesis of modified P31 particles in yeast after gene modification, **141**, 942
 - transcripts produced in HEP 3B cells, analysis (human), **138**, 131
- Hepatobiliary disease**
- diagnosis: implications of C-terminal parathyroid hormone peptide detection after biliary ligation (rat), **139**, 633
- Hepatocyte growth factor**
- high-molecular-weight, detection in human serum before and after hepatic resection, **139**, 361
- Hepatocytes, see also Liver**
- albumin mRNA levels, regulation by dexamethasone and estrogen *in vitro* (*Xenopus laevis*), **136**, 8
 - ATP depletion, induction by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and 1-methyl-4-phenylpyridine (rat), **137**, 310
 - carcinogen-exposed, poly(ADP-ribose) biosynthesis and suicidal NAD⁺ depletion, relationship (rat), **138**, 1051
 - cytochrome *P*-450c induction by β -naphthoflavone, analysis (fetal rat), **141**, 299
 - cytotoxic effects of 1-methyl-4-phenylpyridine and paraquat, comparison (rat), **137**, 303
 - detection of cytoplasmic RNA complementary to pre-albumin intron V (rat), **136**, 989
 - ethanol-metabolizing, generation of neutrophil chemotactic factor (human), **137**, 688
 - intracellular localization of insulin degradation products (rat), **137**, 1034
 - lipid uptake, apolipoprotein E-induced, role of glycosaminoglycan sulfates (rat), **141**, 158
 - membranes, transferrin receptor recovery after perfusion with collagenase (rat), **134**, 711
 - periportal and perivenous, retention of zonal characteristics in primary culture (rat), **139**, 1055
 - phorbol ester-induced translocation of protein kinase C from soluble to particulate cell fraction (rat), **138**, 1328
 - plasma protein synthesis and glycogen deposition, comparative effects of insulin and insulin-like growth factors (chick embryo), **134**, 427
 - in primary culture, platelet-derived growth inhibitors, detection and properties (rat), **134**, 755
 - sinusoidal, receptor-mediated endocytosis of malondialdehyde-modified high-density lipoprotein (rat), **137**, 29
 - substances incorporated by endocytosis, localization by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
 - transformed, α -fetoprotein synthesis (fetal rat), **135**, 844

Subject Index

- transplasmalemma electron transport, role in diferric transferrin reductase system (rat), **139**, 1117
- triglyceride synthesis and secretion *in vitro*, epinephrine effects (rat), **141**, 191
- uptake and metabolism of glucocorticoids (nonadrenalectomized rat), **135**, 160
- Herpes simplex virus
- monoclonal antibody-producing human hybridomas, generation after antigen stimulation *in vitro*, **135**, 495
 - types 1 and 2, infection of BHK cells, neomycin effects, **141**, 198
- Heterosaccharides
- spin-labeled, in IgG Fc region, dynamic properties, analysis in model immune complexes (mouse), **137**, 237
- 3,4,5,3',4',5'-Hexachlorobiphenyl
- enhancement of cardiac prostaglandin release *in vitro* (chick embryo), **136**, 582
- trans*-2-Hexadecanoyl CoA
- effect on microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
- Hexapeptides
- trichotoxin C-terminal, crystal structure, **139**, 590
- Hexokinase
- activity in brown and white adipose tissue (rat), **138**, 687
 - assay in brown adipose tissue during aging (rat), **140**, 419
 - intracellular distribution during glucose repression and derepression (*Saccharomyces cerevisiae*), **134**, 285; *erratum*, **136**, 843
 - isozyme C from Novikoff hepatoma, amino acid sequence, comparison with yeast hexokinases (rat), **139**, 714
 - type II, degradation in skeletal muscle, effects of streptozotocin-induced diabetes and insulin treatment (rat), **138**, 374
- Hexose monophosphate, *see* Fructose 6-phosphate
- Hexose transport
- in plasma membrane vesicles, characterization (rat L6 myoblasts), **141**, 124
- High-density lipoproteins
- apoprotein and lipid constituents, role in high-density lipoprotein-induced cardiac prostacyclin synthesis (rabbit), **139**, 416
 - malondialdehyde-modified, receptor-mediated endocytosis by sinusoidal hepatocytes (rat), **137**, 29
- High-performance liquid chromatography
- in preparation of [¹²⁵I]growth hormone for receptor studies (human), **134**, 671
 - purification of protein kinase C (rat brain), **135**, 636; *erratum*, **136**, 844
- Histamine
- compound 48/80-induced secretion by peritoneal mast cells, effect of phorbol ester pretreatment (rat), **137**, 1112
 - effect on cytosolic free Ca²⁺ levels in cultured smooth muscle cells (rat aorta), **135**, 172
 - stimulation of inositol 1-phosphate accumulation in chromaffin cells (bovine adrenal gland), **135**, 566
- Histamine receptors
- growth-promoting, mediating tamoxifen effects, identification as antiestrogen binding site (rat liver), **134**, 601
- H₁
- binding of mepyramine, characterization (bovine retina), **135**, 445
 - hepatic, molecular size determination by radiation inactivation (vertebrates), **137**, 593
- Histidine
- glycyl derivative, effect on IgG binding to bacterial Fc receptors, **135**, 1105
 - imidazole ring, selective oxidation by ascorbic acid-copper ion system, **138**, 659
 - modification of fructose-1,6-bisphosphatase activity, effect on 5'-AMP sensitivity (fish, rat brain), **139**, 571
 - at neutral endopeptidase active site, identification (rabbit kidney), **139**, 733
 - in proteins, labeling with *p*-diazobenzoyl biocytin, **138**, 872
 - role in catalytic activity of β -hydroxy- β -methylglutaryl-CoA reductase (rat liver, yeast), **141**, 278
- L-Histidine decarboxylase
- and guinea pig L-dopa decarboxylase, immunohistochemical and immunochemical cross-reactivity (rat), **141**, 306
- Histidyl-proline diketopiperazine
- like immunoreactivity in cerebrospinal fluid, distribution and characterization (human), **136**, 835
 - specific binding to hepatic membranes (rat), **134**, 443
- Histones
- chromatin integration, monitoring with double spin-labeled H1 and H5 (avian erythrocytes), **141**, 434
 - H4, interaction with DNA, effect of poly(ADP-ribose) (rat liver), **139**, 523
 - mRNAs, isolation and characterization (murine testis), **136**, 1116
- L-Homocysteine
- metabolism to methionine, role in cell growth (BHK, CCL 39, Raji cells), **135**, 131
- HTLV-III, *see* Human T-lymphotropic virus type III
- Human chorionic gonadotropin
- effect on progesterone accumulation and catabolism in cultured granulosa cells (rat), **141**, 1025
 - intracellular forms, analysis in normal noncultured first-trimester placenta, **137**, 443
 - α -subunit secreted by tumor cell lines, phosphorylation (human), **140**, 143
 - subunits, divergent metabolism in ovarian lysosomes (rat), **134**, 1197
- Human T-cell leukemia virus type I
- protease gene, identification and structural comparison with other retroviral proteases (MT-2 cells), **139**, 129
 - transformed T cells, detection of interleukin-1 α mRNA (human), **139**, 353
- Human T-lymphotropic virus type III
- envelope proteins, detection of hexapeptide homologous to interleukin-2 receptor binding region (murine spleen), **139**, 367
 - lymphadenopathy-associated virus
 - cytotoxicity, inhibition by aurintricarboxylic acid *in vitro*, **136**, 64
 - glycoprotein expression, suppression by glycosylation inhibitors, **141**, 33
 - selective inhibition by 2',3'-dideoxycytidine, **140**, 735
- Hyaluronic acid
- induction of interleukin 1 production by monocytes and macrophages (human, rabbit), **140**, 715
- Hybrid cells
- human-murine, expression of microsomal epoxide hydro-

Subject Index

- lase, quantitation, **137**, 775
- Hybridization
in nerve growth factor mRNA *in situ* localization (murine brain), **138**, 813
- Hybridomas
producing monoclonal antibodies against
cytomegalovirus, generation (human), **137**, 273
herpes simplex virus, generation after stimulation *in vitro* (human), **135**, 495
- Hydration
L-alanine and L-proline, in water/dimethyl sulfoxide mixtures, ^{17}O NMR, **135**, 363
sodium hyaluronate, NMR analysis of oriented sample, **141**, 73
- Hydrocarbons
polycyclic aromatic, *see* Polycyclic aromatic hydrocarbons
- Hydrocortisone
cytochrome P-450_{11 β} -dependent conversion to cortisone (bovine), **136**, 369
transcriptional and post-transcriptional roles in casein gene expression (rat mammary explants), **134**, 812
- Hydrogen
atomic, formation during lipid peroxidation, ESR spectroscopy, **141**, 381
-oxidizing enzymes, encoding genes
conjugal transfer from *Alcaligenes hydrogenophilus* to *Pseudomonas oxalaticus*, **137**, 108
isolation from *Alcaligenes hydrogenophilus* and expression in *Pseudomonas oxalaticus*, **137**, 114
-potassium antiporter, pH dependence, desensitization by N-ethylmaleimide (*Vibrio alginolyticus*), **136**, 1030
-sodium exchange
in basolateral membranes, detection by fluorescence quenching (rabbit small intestine), **134**, 827
osmotic activation, protein kinase C role (rat thymic lymphocytes), **134**, 8
in platelets, induction by 1-oleoyl-2-acetyl glycerol and phorbol ester (human), **141**, 13
- Hydrogenase
hydrogen-oxidizing, encoding genes
conjugal transfer from *Alcaligenes hydrogenophilus* to *Pseudomonas oxalaticus*, **137**, 108
isolation from *Alcaligenes hydrogenophilus* and expression in *Pseudomonas oxalaticus*, **137**, 114
kinetic determination in H₂ evolution direction by rapid electrocatalytic procedure, **135**, 928
multiple intrinsic membrane nickel-containing, isolation and characterization (*Desulfovibrio vulgaris*), **139**, 701
periplasmic, EPR spectrum, effect of low carbon monoxide levels (*Desulfovibrio vulgaris*), **137**, 1086
soluble, isolation and characterization (*Desulfovibrio africanus*), **139**, 658
- Hydrogen bonds
carboxylic acid-retinal Schiff base, proton transfer thermodynamics, **138**, 819
formation between triacetyl glycerol and water, detection by high-pressure infrared spectrometry, **134**, 140
- Hydrogen peroxide
dependence of
3,4-dihydroxy-3,4-dihydrobenzo[*a*]anthracene epoxidation by ovine seminal vesicle microsomes and hematin, **140**, 181
poly R-481 decolorization by particulate fractions from *Phanerochaete chrysosporium*, **136**, 220
-generating system, NADPH-dependent, from thyroid, solubilization and characterization, **141**, 839
-hypochlorous acid reaction, singlet oxygen production, enhancement by 1,4-diazabicyclo[2.2.2]octane, **134**, 777
induction of vascular relaxation (rat aortic rings), **139**, 102
lactoperoxidase-catalyzed metabolism, role of superoxide radicals, **139**, 327
oxidation of cell wall polysaccharides, role in cell wall breakdown (larch, orange), **141**, 238
peroxidase-dependent oxygen transfer to thioanisoles, **135**, 166
production
during macrophage respiratory burst, enhancement by interferon γ , antagonistic effects of interferons α and β (human), **140**, 948
in monocytes, effect of κ elastin (human), **141**, 92
- Hydrolysis
glycosidase-catalyzed, stereochemical course, ^1H NMR determination, **139**, 487
phosphoinositides
 α_1 -adrenergic receptor-stimulated, and aortic contraction, inhibition by phorbol esters (rat), **134**, 970
induced by formyl-methionyl-leucyl-phenylalanine, inhibition by phorbol ester (human neutrophils), **135**, 556
in isolated platelet membranes, stimulation by GTP and cytosol (human), **137**, 801
by platelet-derived growth factor receptor, effect of protein kinase C activation (human skin, murine Swiss 3T3 fibroblasts), **137**, 343
phosphorylcholine, quantitation in homogenates of temporal regions of Alzheimer's brain (human), **139**, 315
polyphosphoinositides
coupling to muscarinic receptors, induction by point-mutated p21^{ras} (murine fibroblasts), **141**, 591
enhancement by gonadotropin-releasing hormone (rat pituitary cells), **134**, 861
protein-bound ADP-ribose in skeletal muscle, catalyzing enzyme, identification (rat), **139**, 932
- Hydroperoxide, *see* Hydrogen peroxide
- 5-Hydroperoxyeicosatetraenoic acid
effect on 5-lipoxygenase activity (rat neutrophil extracts), **141**, 534
- 12-Hydroperoxyeicosatetraenoic acid
-stimulated neutrophils, leukotriene B₄ synthesis, requirement of free arachidonic acid (human), **138**, 589
- 15-Hydroperoxyeicosatetraenoic acid
induction of vascular relaxation (rat aortic rings), **139**, 102
- Hydroxamic acids
peptide, inhibition of skin collagenase (human), **136**, 390
- N-Hydroxy-2-acetylaminofluorene
inhibition of aldehyde oxidase (rabbit liver), **138**, 312
- ω -Hydroxyacids
covalent attachment to epidermal macromolecules, characterization (swine), **137**, 992
- ω -Hydroxyacylsphingosine
covalent attachment to epidermal macromolecules, characterization (swine), **137**, 992
- 4-Hydroxyalkenals
tissue levels, determination as pentafluorobenzyl oxime de-

Subject Index

- rivatives by gas chromatography-mass spectrometry (rat), **139**, 144
- 19-Hydroxyandrostendione
cytochrome *P*-450_{11β}-dependent conversion to 19-oxoandrostenedione (bovine), **136**, 369
- 8-Hydroxydeoxyguanosine
accumulation in DNA of phorbol ester-exposed granulocytes (human), **137**, 841
- 17β-Hydroxy-11β-(4-dimethylaminophenyl)-17α-(1-propynyl)-estra-4,9-diene-3-one
physiological action and receptor binding (rat liver, thymocytes), **134**, 44
and R 5020, interaction with progesterone receptors, comparison (human breast cancer cells), **135**, 90
- 20-Hydroxyecdysone
induction of β-tubulin subunit synthesis (*Drosophila melanogaster*), **134**, 191
- 5-Hydroxyeicosatetraenoic acid
synthesis in A23187-stimulated neutrophils, enhancement by phorbol ester (human), **141**, 399
- 12-Hydroxy-5,8,10,14-eicosatetraenoic acid
in platelets and psoriatic lesions, stereochemical differences (human), **136**, 169
- Hydroxyeicosatetraenoic acids
formed during arachidonic acid oxygenation by microsomal cytochrome *P*-450, absolute configuration (rat liver), **141**, 1007
- 9-Hydroxyellipticine
stimulation of topoisomerase II-induced cleavage sites in *c-myc* protooncogene *in vivo* (N417 cells), **137**, 154
- 8-Hydroxyguanine
formation in cellular DNA after 4-nitroquinoline 1-oxide exposure (Ehrlich ascites cells), **139**, 626
- 5-Hydroxyindoleacetic acid
brain levels, effect of sustained exercise (rat), **137**, 149
- 3-Hydroxykynurenine
role as oxygen radical scavenger in *Aldrichina grahami*, **139**, 666
- 11β-Hydroxylase
adrenal activity, effects of α-melanocyte-stimulating hormone *in vivo* (rat), **141**, 1279
- 18-Hydroxylase
adrenal activity, effects of α-melanocyte-stimulating hormone *in vivo* (rat), **141**, 1279
- 21-Hydroxylase
genes, restriction mapping and restriction fragment length polymorphisms (human), **136**, 722; *erratum*, **138**, 502
- 20-Hydroxy-leukotriene B₄
and leukotriene B₄, simultaneous quantitation by gas chromatography-mass spectroscopy (human neutrophils), **134**, 512
- 3'-Hydroxyl group
in penultimate position of 2',5'-phosphodiesterase, role in oligonucleotide degradation, **141**, 555
- Hydroxyl radicals
formation
during lipid peroxidation, ESR spectroscopy, **141**, 381
from 1-methyl-4-phenylpyridinium ion in presence of NADPH-cytochrome *P*-450 reductase and oxygen, **135**, 583
generation by *Trypanosoma cruzi* extracts, mediation by primaquine, **135**, 1029
spin trap adducts, [¹⁷O]oxygen hyperfine structure, **141**, 622
- Hydroxymethylglutaryl-CoA reductase
catalytic activity, role of cysteine and histidine (rat liver, yeast), **141**, 278
regulation by insulin (rat), **137**, 201
- 7-Hydroxymethyl sulfate ester
role in regioselective glutathione conjugation of 7,12-dihydroxymethylbenz[*a*]anthracene (rat hepatic cytosol), **134**, 100
- β-Hydroxynorvaline
inhibition of lymphadenopathy-associated virus/human T-lymphotropic virus type III glycoprotein expression, **141**, 33
- N*-(4-Hydroxyphenyl) retinamide
anticarcinogenic effect on mammary tumor induction by 7,12-dimethylbenz[*a*]anthracene, relationship to cAMP-dependent protein kinase (rat), **135**, 116
- Hydroxyproline
X-ray crystal structure data, analysis, **140**, 895
- 15-Hydroxyprostaglandin dehydrogenase (NAD⁺)
isolation and structural analysis (human placenta), **135**, 1058
- 3α-Hydroxysteroid dehydrogenase
and dihydrodiol dehydrogenase, solubilization, separation, and partial purification (rat hepatic microsomes), **135**, 795
- Hydroxytriphenylethylenes
with high affinity for estrogen receptor, inhibition of tumor cell proliferation, mechanism (human breast cancer), **140**, 523
- 5-Hydroxytryptamine, *see* Serotonin
- Hydroxyurea
uptake by diffusion mechanism (Chinese hamster ovary cells), **134**, 1254
- 25-Hydroxyvitamin D₃
metabolism in kidney, effect of tumor-promoting phorbol ester (chicken), **139**, 495
- Hyperglycemia
induced sorbitol accumulation and *myo*-inositol depletion in cultured conceptus, effects of aldose reductase inhibitors (rat), **140**, 974
- Hyperkalemia
induced changes in myocardial tissues, procaine effects (dog), **134**, 900
- Hyperoxia
associated leukotoxin biosynthesis by leukocytes in lung lavages (rat), **134**, 1071; *erratum*, **135**, 1150
- Hyperplasia
epidermal, induction by cholera toxin: evaluation of tumor-promoting activity in skin carcinogenesis (mouse), **137**, 486
- Hypertension
associated increase in plasma atrial natriuretic factor levels (salt-sensitive rat), **137**, 876
effect on human α-atrial natriuretic factor binding to renal basolateral membranes (rat), **137**, 1079
erythrocytes, activity decrease by calpain inhibitor (Milan hypertensive rat), **138**, 1370
- spontaneous
effect on protein kinase C activity in platelets, **141**, 769
and spontaneous stroke-prone, associated atrial natriuretic factor levels in left atrium and plasma (rat), **135**, 74

Subject Index

- Hypertrehalosaemic factor**
 amino acid sequence (cockroach corpus cardiacum), **141**, 774
 isolation and amino acid sequence (*Blaberus discoidalis*), **140**, 674
- Hypertrophy**
 cardiac, stress-induced, role of initial protein synthesis suppression (rat), **135**, 857
- Hypochlorous acid**
 -hydrogen peroxide reaction, singlet oxygen production, enhancement by 1,4-diazabicyclo[2.2.2]octane, **134**, 777
- Hypothalamus**
 -neurohypophysis complex in culture, arginine vasopressin release, effect of synthetic atrial natriuretic factor (rat), **134**, 652
 release of atrial natriuretic factor *in vitro*, radioimmunoassay (rat), **136**, 590
- Hypothermia**
 effect on carnitine palmitoyltransferase (newborn rat brown adipose tissue, liver), **135**, 589
- Hypoxia**
 induction of
 atrial natriuretic factor release from isolated heart (rabbit, rat), **140**, 427
 changes in myocardial tissues, procaine effects (dog), **134**, 900
- I**
- L-Iduronate**
 ring conformation in dermatan sulfate, solid-state ^{13}C NMR and X-ray diffraction analysis (porcine skin), **137**, 87
- Imidazole**
 modification of fructose-1,6-bisphosphatase activity, effect on 5'-AMP sensitivity (fish, rat brain), **139**, 571
- Immune complexes**
 model, in analysis of dynamic properties of spin-labeled heterosaccharides in IgG Fc region (mouse), **137**, 237
- Immunoassay**
 lectin-binding, detection of glycosylated growth hormone in pituitary gland (human), **140**, 491
- Immunoglobulins**
 IgG
 aggregated, stimulation of phagocytic degranulation by neutrophils, requirement for arachidonic acid lipooxygenation (human), **136**, 310
 binding to bacterial Fc receptors, effect of various dipeptides, **135**, 1105
 Fc region, spin-labeled heterosaccharides, analysis of dynamic properties in model immune complexes (mouse), **137**, 237
 -ricin A conjugate, mannose receptor-mediated uptake by hepatic nonparenchymal cells, measurement (rat), **137**, 892
 IgM, secretion by lymphoblastoid B-cell line, effect of actin antibodies (human), **140**, 399
- Indo-1**
 in analysis of intracellular Ca^{2+} release and Ca^{2+} uptake in antibody-activated B cells (mouse), **137**, 500; *erratum*, **138**, 503
- Indoleamine 2,3-dioxygenase**
 cofactor activities of dihydroflavin mononucleotide and tetrahydrobiopterin (murine epididymis), **137**, 1106
- Inflammation**
 experimental, association of acute phase protein genes with nuclear matrix (rat liver), **137**, 633
- Inhibin**
 detection in placenta (human), **140**, 485
 high-molecular-weight forms, demonstration with monoclonal antibodies (bovine follicular fluid), **136**, 1103
 modulation of FSH-induced aromatase in granulosa cells, comparison with transforming growth factor activity (rat), **136**, 969
 subunits α , β_A , and β_B , amino acid sequence (human ovary), **135**, 957
 type A
 subunit gene expression in immature ovaries, stimulation by pregnant mare serum gonadotropin (rat), **138**, 1191
 β -subunit homodimer, stimulation of pituitary secretion of follicle-stimulating hormone (pig), **138**, 1129
- Initiation factors**
 eIF-2 α , subunit phosphorylation, role in inhibition of protein synthesis under heat shock conditions (HeLa cells), **138**, 519
- Inosine**
 carbocyclic, metabolism and cytotoxicity in promastigotes of *Leishmania donovani* and *L. tropica*, **134**, 1114
- Inositides**
 metabolism in photoreceptor membranes, effect of cationic amphiphilic drugs (bovine), **134**, 378
- myo-Inositol**
 depletion in cultured conceptus, hyperglycemia-induced, effects of aldose reductase inhibitors (rat), **140**, 974
 incorporation into
 opsin *in vivo* (frog), **136**, 815
 phosphoinositides and inositol phosphates, effects of cationic amphiphilic drugs (rat C6 glioma cells), **135**, 465
- Inositol 1-phosphate**
 accumulation in chromaffin cells, stimulation by bradykinin, carbachol, and histamine (bovine adrenal gland), **135**, 566
 carbachol-induced accumulation, effects of lithium and phorbol esters (NCB-20 cells), **136**, 622
- Inositol phosphates**
 formation in adrenal glomerulosa cells, role of guanyl nucleotide-binding proteins (rat), **140**, 941
 incorporation of inositol, effects of cationic amphiphilic drugs (rat C6 glioma cells), **135**, 465
 metabolism in brain, enhancement during bicuculline-induced status epilepticus (rat), **136**, 827
- Inositol (1,3,4,5)-tetrakisphosphate**
 formation from inositol (1,4,5)-trisphosphate, calcium sensitivity and ATP dependence (bovine adrenal glomerulosa cells), **139**, 259
 muscarinic receptor-stimulated accumulation in cerebral cortical slices, enhancement by K^+ -induced depolarization (rat), **141**, 1130
- Inositol 1,4,5-trisphosphate**
 accumulation in cultured aortic myocytes, arginine vasopressin-induced, modulation by protein kinase C (rat), **140**, 1093

Subject Index

- calcium, and cAMP signaling systems, dual activation by adrenocorticotropin (rat adrenal cells), **135**, 742
- cellular levels, bradykinin-induced changes (MDCK cells), **140**, 644
- effect on Ca^{2+} uptake by intercellular compartments in permeabilized enterocytes (rat intestine), **139**, 612
- enhancement of Ca^{2+} sensitivity of contractile mechanism in chemically skinned skeletal muscle fibers (rabbit), **135**, 662
- heat shock-induced increase (CHO cells), **137**, 826
- induction of Ca^{2+} release from
 - endoplasmic reticulum, inhibition by calmodulin (rat pancreatic islets), **141**, 418
 - permeabilized mastocytoma cells (mouse), **135**, 46
- interaction with spermine, **135**, 58
- interconversion to inositol (1,4,5)-tetrakisphosphate and (1,4,5)-trisphosphate, calcium sensitivity and ATP dependence (bovine adrenal glomerulosa cells), **139**, 259
- production in platelets, inhibition by cGMP (human), **135**, 1099
- release from NCB-20 cells, effects of D-Ala²-D-Leu⁵-enkephalin and bradykinin, **135**, 507
- D-myo-Inositol-1,4,5-trisphosphate phosphatase
 - characterization (rat brain), **134**, 351
- Insulin
 - action mechanism, role of phosphorylase phosphatase catalytic subunit inhibition by oxalyl thioesters, **134**, 93
 - antilipolytic effect, requirement of phosphodiesterase activation (human adipocytes), **141**, 1157
 - binding
 - analysis in skeletal muscle plasma membranes (normal, obese-diabetic mice), **137**, 286
 - to CHO cells, effect of glucose deprivation, **140**, 821
 - in glucose-starved fibroblasts after glucose refeeding, effects of cycloheximide and tunicamycin (human), **140**, 844
 - brain levels, effect on downregulation of fetal insulin receptors (rabbit), **136**, 208
 - concomitant binding, degradation, and pyruvate dehydrogenase activation in phytohemagglutinin-stimulated T cells (human), **134**, 412
 - conjugated psoralen derivative, insulin receptor-mediated photocytotoxic effect on activated lymphocytes, **141**, 502
 - degradation products, intracellular localization (rat hepatocytes), **137**, 1034
 - effect on
 - glucagon binding to isolated epididymal adipocytes (rat), **141**, 812
 - Na^+/K^+ -ATPase activity in renal basolateral membranes (non-insulin-dependent diabetic rat), **139**, 1313
 - protein kinase C translocation from soluble to particulate cell fraction (rat hepatocytes), **138**, 1328
 - induction of ribosomal protein S6 kinase (3T3-L1 cells), **137**, 702
 - and insulin-like growth factors, comparative effects on plasma protein synthesis and glycogen deposition in cultured hepatocytes (chick embryo), **134**, 427
 - interaction with somatomedin receptors in fetal cortical neurons (mouse), **135**, 228
 - [Leu^{A3}]-, receptor binding and negative cooperative effects, **137**, 250
 - like effects
 - growth hormone 20-kDa variant, effect of receptor binding properties (human), **134**, 159
 - prolactin in adipose tissue *in vitro*, inhibition by glycosylation (rat), **134**, 764
 - recombinant DNA-derived methionyl human growth hormone 20-kDa variant, characterization (hypophysectomized rat), **138**, 342
 - mRNA nucleotide sequences, occurrence of short complementary sequences, **139**, 94
 - and phorbol ester, effects on cytosolic protein kinase C, comparison (rat adipocytes), **135**, 1119
 - regulation of
 - 3-hydroxy-3-methylglutaryl-CoA reductase (rat), **137**, 201
 - phosphoenolpyruvate carboxykinase gene during preadipose cell differentiation, **138**, 468
 - related factor secreted by 1246-3A teratoma cells, characterization (mouse), **135**, 533
 - release from
 - cloned B-cell line HIT-T15, effects of glucose, amino acids, sulfonylureas, and colchicine (hamster), **140**, 616
 - fibroblasts, induction by transfection with genomic rat islet DNA (mouse), **136**, 638; *erratum*, **139**, 852
 - intact and permeabilized islets of Langerhans, effect of polymyxin B (rat), **136**, 1001
 - pancreatic islets
 - after α_2 -adrenergic activation, effect of glucose (mouse), **139**, 557
 - glucose-stimulated, inhibition by galanin *in vitro*, mechanism (mouse), **140**, 1059
 - stimulation by lysophosphatidylinositol (rat), **138**, 720
 - perfused pancreas, cysteamine effect (rat), **134**, 1291
 - stimulatable serine/threonine kinase: identification of muscle protein substrates related to microtubule-associated protein 2 (3T3-L1 adipocytes), **134**, 565
 - stimulated serine kinase, detection in extracts of IM-9 lymphocytes, **139**, 179
 - stimulation of
 - α -aminoisobutyric acid transport by Na^+/K^+ -ATPase, ouabain effects (rat skeletal muscle), **134**, 1342
 - 2-deoxyglucose uptake by fibroblasts, inhibition by sphingolipids (mouse), **138**, 463; *erratum*, **139**, 852
 - Na^+/K^+ -ATPase (frog skeletal muscle), **134**, 609
 - phosphatidylcholine synthesis in plasma membranes (rat adipocytes), **137**, 1
 - pyruvate kinase and pyruvate kinase phosphatase (rat hepatocytes), **134**, 292
 - sulfation by sulfotransferase (*Eubacterium* A-44), **140**, 38
 - surface-adsorbed, reaction products, mass spectrometry (bovine), **134**, 420
 - treatment of streptozotocin-induced diabetes, effect on hexokinase II degradation in skeletal muscle (rat), **138**, 374
- Insulin-like growth factor I
 - binding to CHO cells, effect of glucose deprivation, **140**, 821
 - covalent crosslinking to specific serum-derived inhibitor (human), **137**, 411
 - growth hormone-dependent, binding protein from plasma, comparison to amniotic fluid IGF-binding protein

Subject Index

- (human), **139**, 1256
 and insulin, comparative effects on plasma protein synthesis and glycogen deposition in cultured hepatocytes (chick embryo), **134**, 427
 interaction with somatomedin receptors in fetal cortical neurons (mouse), **135**, 228
 purification by affinity chromatography with amniotic fluid somatomedin-binding protein (human), **136**, 253
 stimulation of Leydig's cell steroidogenesis *in vitro* (rat), **137**, 950
- Insulin-like growth factor II**
 binding to adipocytes, stimulation by vanadate (rat), **138**, 174
 growth hormone-dependent, binding protein from plasma, comparison to amniotic fluid IGF-binding protein (human), **139**, 1256
 identification in seminal and follicular fluids (human), **140**, 536
 and insulin, comparative effects on plasma protein synthesis and glycogen deposition in cultured hepatocytes (chick embryo), **134**, 427
 interaction with somatomedin receptors in fetal cortical neurons (mouse), **135**, 228
 stimulation of DNA synthesis, effect of anti-insulin-like growth factor I receptor antibody (human fibroblasts), **139**, 501
- Insulin-like growth factor I receptors**
 affinity-purified, characterization (human placenta), **138**, 1257
 antibodies, effect on insulin-like growth factor II stimulation of DNA synthesis (human fibroblasts), **139**, 501
 autophosphorylation and tyrosine kinase activity, effect of human polyclonal antibody (human placenta, rat liver), **138**, 1023; *erratum*, **140**, 1152
 characterization and fibroblast/adipocyte responsiveness before and after differentiation, **137**, 552
 detection in Madin-Darby canine kidney cells, **134**, 120
 ligand-binding subunit, properties, **136**, 45
 purification and characterization (human placenta), **137**, 695
 species distinction with monoclonal antibody: antibody identification, **138**, 1341
 tyrosine kinase-associated activity (rat brain), **134**, 532
- Insulin-like growth factor II receptors**
 characterization and fibroblast/adipocyte responsiveness before and after differentiation, **137**, 552
 detection in Madin-Darby canine kidney cells, **134**, 120
 identification in insulin-like growth factor II-producing cell lines and characterization, **134**, 595
 purification and antibody generation (human pituitary adenoma), **138**, 304
- Insulin receptors**
 autophosphorylation
 and kinase activity in streptozotocin-induced diabetes, effect of short fast (rat), **140**, 850
 kinetic evidence for activating and nonactivating components, **134**, 1312
 binding activity, decrease after chronic phenobarbital treatment (rat liver), **135**, 501
 characterization and fibroblast/adipocyte responsiveness before and after differentiation, **137**, 552
 dimer and tetramer binding characteristics (rat liver), **135**, 458
 -mediated photocytotoxicity: killing of activated lymphocytes by insulin-conjugated psoralen derivative, **141**, 502
 mRNA nucleotide sequences, occurrence of short complementary sequences, **139**, 94
 phosphorylation, analysis in skeletal muscle plasma membranes (normal, obese-diabetic mice), **137**, 286
 precursor-like polypeptide in *Drosophila*, cloning of encoding cDNA (human), **141**, 474
 regulation in fetal brain, effect of insulin levels (rabbit), **136**, 208
 tyrosine kinase-associated activity (rat brain), **134**, 532
- Interferon α**
 antagonistic effect on interferon γ -induced enhancement of H_2O_2 production during macrophage respiratory burst (human), **140**, 948
 gene sequence and expression: homology with enhancer elements in gene regulatory region (mouse), **138**, 826
 multiple molecular forms, induction of antiviral and 2',5'-oligoadenylate synthetase activities, comparison (MDBK cells), **141**, 1042
- Interferon α_2**
 Ala^{30,32,33} analog, biological and receptor-binding properties (human), **134**, 1412
 functionally important conserved amino acids, identification with analogs produced from synthetic genes (human), **134**, 1404
- Interferon α/β**
 induction by transformed cells, inhibition by retinoic acid (murine L-929, human WISH cells), **138**, 47
 and interferon γ , antagonistic effects on mannosyl-fucosyl receptors (murine peritoneal macrophages), **136**, 737
- Interferon β**
 antagonistic effect on interferon γ -induced enhancement of H_2O_2 production during macrophage respiratory burst (human), **140**, 948
 binding to human A549 and murine L cells, enhancement by lanthanide ions, **137**, 480
 gene expression in fibroblasts, effect of serial divisions (human), **141**, 112
 -inducing 22K factor, homogeneous, connective tissue cell-stimulating activities (bovine, human, neonatal mouse), **139**, 1150
- Interferon γ**
 enhancement of
 H_2O_2 production during macrophage respiratory burst, antagonistic effects of interferons α and β (human), **140**, 948
 neutrophil oxidative metabolism (human), **138**, 1276
 extracellular and intracellular expression by murine L cells transformed with human IFN- γ cDNA gene, **135**, 894
 induced differentiation of HL-60 cells, associated appearance of membrane-bound tyrosine kinase, **140**, 908
 and interferon α/β , antagonistic effects on mannosyl-fucosyl receptors (murine peritoneal macrophages), **136**, 737
 recombinant, phosphorylation sites, identification (human), **136**, 679
 transport into cell nucleus, role of nuclear membrane receptors (mouse), **138**, 254
- Interferons**
 induction of 2-5A synthetase and 2-5A binding protein,

Subject Index

- age-related differences (guinea pig macrophages), **140**, 455
- Interleukin 1**
- enhancement of plasminogen activator inhibitor levels in cultured endothelial cells (human), **139**, 720
 - genes, expression in chondrocytes: mechanism for autocrine control of cartilage matrix degradation (bovine, human), **141**, 904
 - human recombinant, effect on glycosaminoglycan synthesis (rat costal chondrocytes), **140**, 386
 - like factor, effect on proteoglycan synthesis in cultured articular chondrocytes (human), **134**, 539
 - production by
 - monocytes and macrophages, endogenous induction by hyaluronic acid (human, rabbit), **140**, 715
 - peritoneal macrophages, protein kinase C role (mouse), **135**, 355
 - resting macrophages, stimulation by tumor necrosis factor (mouse), **136**, 94
 - recombinant human, type α and β , stimulation of macrophage colony-stimulating activity and prostaglandin E_2 production (MC3T3-E1 cells), **141**, 285
- Interleukin 1 α**
- mRNA, detection in virus-transformed T and B cells (human), **139**, 353
 - recombinant, stimulation of prostaglandin E_2 production and bone resorption in fetal mouse (human), **138**, 618
- Interleukin 2**
- cDNA, homologous genomic DNA sequences, characterization (human placenta), **138**, 966
 - receptor binding region, detection of homologous hexapeptide in human T-lymphotropic virus type III envelope protein (murine spleen), **139**, 367
 - recombinant human, and methionyl IL-2, produced in *Escherichia coli*, separation, **135**, 837
 - stimulation of pp67 phosphorylation (murine NK-7 cells), **135**, 487
- Interleukin-2 receptors**
- human, cDNA expression in *Escherichia coli*, **141**, 804
 - Tac antigen epitope, phosphorylation by protein kinase C (human T cells), **135**, 239
- Interleukin 3**
- like activity, detection in supernatants of lipopolysaccharide-stimulated splenocytes (mouse), **138**, 1322
- Interleukin-3 receptors**
- surface, affinity isolation (murine FDC-P1, NSF-60 cells), **135**, 870
- Intermediate filaments**
- and τ proteins, association with retinal heat shock protein, analysis (rabbit), **139**, 974
- Intestine**
- jejunum, transport of intact nonapeptide (rabbit), **137**, 682
- Invertase**, *see* β -Fructofuranosidase
- Involucrin**
- action as transglutaminase substrate at multiple sites (human keratinocytes), **136**, 51
- Iodide**
- fluxes in FRTL-5 cells, activation mediated by α_1 -adrenergic and thyrotropin receptors, role of pertussis toxin-sensitive G proteins, analysis, **141**, 1000
- Iodine**
- radioactive, labeling of transforming growth factor β (pig), **138**, 714
- 5-Iodoacetamidofluorescein**
- hemoglobin-bound, steady-state fluorescence emission, **138**, 489
- Iodothyronine 5'-deiodinase**
- basal and noradrenaline-induced activity in brown adipose tissue, reduction during pregnancy and lactation (rat), **138**, 1315
- Ionomycin**
- alteration of accessibility of surface thiols (human lymphocytes), **140**, 999
 - effect on
 - intracellular calcium levels and pepsinogen secretion (guinea pig gastric chief cells), **137**, 281
 - platelet cytosolic Ca^{2+} levels, analysis with fura-2 (human blood), **139**, 308
 - induction of sponge cell aggregation without extracellular calcium (*Microciconia prolifera*), **134**, 1319
 - and phorbol ester, effects on fatty acid synthesis by isolated hepatocytes (rat), **139**, 403
- Ionophore A23187**
- alteration of accessibility of surface thiols (human lymphocytes), **140**, 999
 - effect on
 - intracellular calcium levels and pepsinogen secretion (guinea pig gastric chief cells), **137**, 281
 - phorbol dibutyrate receptor affinity states, role in phorbol-induced B-cell activation (human), **138**, 554
 - protein synthesis inhibition by bovine cell-surface sialoglycopeptide (murine 3T3 cells), **136**, 976
 - induction of
 - damage in pancreatic acinar membranes, role of calcium (rat), **139**, 530
 - leukotriene B_4 production in whole blood, comparison with production by isolated neutrophils (human), **137**, 334
 - prostaglandin E_2 synthesis in mesangial cells, opposite effects of different concentrations of pertussis toxin (rat), **141**, 39
 - sponge cell aggregation without extracellular calcium (*Microciconia prolifera*), **134**, 1319
 - and phorbol ester, synergistic stimulation of lipoxygenase-mediated arachidonic acid metabolism (murine macrophages), **140**, 861
 - stimulated neutrophils, arachidonic acid release and leukotriene synthesis, enhancement by phorbol ester (human), **141**, 399
 - stimulation of phosphate accumulation in cultured renal tubular cells, synergistic effects of phorbol esters (mouse), **136**, 177
- Iron**
- chromophore, in purple acid phosphatase, oxidation during enzyme interaction with phosphate (bovine spleen), **136**, 490
 - content in cytochrome-c oxidases, analysis by proton-induced X-ray emission (bovine, porcine, rat tissues), **140**, 1007
 - effect on plasma protein kinase activity (human), **138**, 693
 - generation of DNA breaks, activation by $KHSO_5$ (bacteriophage λ), **136**, 1013
 - molybdenum cofactor of *Azotobacter vinelandii* nitrogenase, solubilization in dimethylformamide and

Subject Index

- acetonitrile, **139**, 740
 -tetraphenylporphyrin system, generated hemoprotein-peroxide complex, characterization, **135**, 972
 Islets of Langerhans, *see* Pancreas
N-6-Isopentenyl adenine
 role in tumor cell growth (mouse), **137**, 208
 Isopropanol, *see* Isopropyl alcohol
 Isopropyl alcohol
 induced denaturation, in detection of unstable hemoglobin (mouse), **139**, 551
 Isopropyl-oxazolopyrido-carbazole
 induced frameshift mutations, effect of mutation in *uvrB* gene (*Salmonella typhimurium*), **138**, 505
 Isoproterenol
 effect on
 adenylate cyclase activity and [³H]clonidine specific binding in brown adipose tissue (fetal rat), **138**, 1390
 atrial natriuretic factor release by atrial slices (rat), **135**, 1090
 cardiac and plasma atrial natriuretic factor levels after chronic administration (rat), **136**, 510
 1-(5-Isoquinolinylsulfonyl)-2-methylpiperazine
 effect on neutrophil activation (human), **135**, 749
 reversal of phorbol ester-mediated suppression of neutrophil responses to chemoattractants (rabbit), **137**, 50
 polyphosphoinositide metabolism and cytosolic Ca²⁺ mobilization (thrombin-activated human platelets), **134**, 868
- K**
- Kallikrein
 cleavage of apolipoprotein B-100: characterization of resulting fragments (human plasma), **137**, 821
 Kanamycin
 resistance gene, expression in *Escherichia coli*, effect of specific yeast sequences, **136**, 529
 Ketoconazole
 effects on sterol biosynthesis (*Trypanosoma cruzi* epimastigotes), **136**, 851
 α-Ketoglutarate dehydrogenase
 reconstituted complex, role of excess lipoyl dehydrogenase (*Escherichia coli*), **135**, 802
 β-Ketopyranosidic linkages
 in 3-deoxy-D-manno-2-octulosonic acid, hydrolysis by bacteriophage-associated glycanase (*Escherichia coli*), **136**, 329
 Ketosugars
 -specific agglutinin, isolation (rat uterus), **136**, 116
 Kidney
 androgen response, role of putrescine (mouse), **138**, 771
 atrial natriuretic polypeptide molecular size, comparison with aorta and adrenal gland by uv-photoaffinity labeling (rat), **136**, 200
 dysfunction, effect on mitochondrial cytochrome and polypeptide composition (human), **137**, 911
 Kidney stones
 amyloid, from uremic patients, detection of β₂-microglobulin fragments, **136**, 665
 Kinetensin
 amino acid sequence, homology with serum albumin, neurotensin, and angiotensin (human), **136**, 983
 Kirsten murine sarcoma virus
 transformation of
 human HOS cells, induction of ouabain sensitivity, **136**, 807
 rat renal cells, associated loss of epidermal growth factor binding, mechanism, **135**, 435
 -transformed fibroblasts, ³¹P NMR spectra (chicken, rat), **138**, 699
 Kyotorphin
 induction of increase in intracellular Ca²⁺ levels (rat cerebral synaptosomes), **137**, 897
 sulfation by sulfotransferase (*Eubacterium A-44*), **140**, 38
- L**
- α-Lactalbumin
 labeling with 3-(trifluoromethyl)-3-(*m*-[¹²⁵I]iodophenyl) diazirine, effects of protein and NaCl levels (bovine), **138**, 604
 L-Lactate dehydrogenase
 isozyme X, postmeiotically expressed encoding cDNA clone, nucleotide sequence (mouse), **136**, 760
 Lactation
 effect on basal and noradrenaline-induced iodothyronine 5'-deiodinase activity in brown adipose tissue (rat), **138**, 1315
 Lactoferrin
 cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
 Lactoperoxidase, *see* Peroxidase
 Lactose synthase
 detection in p-type lymphoblastoid cells (human), **137**, 1187
 encoding cDNA, isolation and characterization (human), **139**, 163
 gene, chromosomal localization (human), **141**, 1185
 isolation and amino-terminal sequence determination (human milk), **138**, 224
 Lactose transporter
 quantitative immunoprecipitation: confirmation of galactoside binding stoichiometry (*Escherichia coli*), **134**, 975
 Lactoylglutathione lyase
 purification and characterization (*Pseudomonas putida*), **141**, 993
 Lanosterol
 14α-demethylation, role of sterol 14-reductase (yeast), **134**, 659
 Lanosterol cyclases
 inhibition by *N*-[(1,5,9)-trimethyl-decyl]-4α,10-dimethyl-8-aza-*trans*-decal-3β-ol, **138**, 764
 Lanthanides
 order of binding to calmodulin, EPR study, **138**, 1243
 Lanthanum chloride
 enhancement of interferon β binding to human A549 and murine L cells, **137**, 480
 Lapachol
 inhibition of NAD(P)H:quinone dehydrogenase (rat liver), **137**, 781
 Laser irradiation
 helium-neon, in treatment of cutaneous wounds, effect on

Subject Index

- type I and III procollagen mRNA levels (pig), **138**, 1123
- α -Latrotoxin
effect on membrane permeability to cations (rat PC12 neurosecretory cells), **134**, 320
- LAV, *see* Lymphadenopathy-associated virus
- Lawsone
inhibition of hematoporphyrin-sensitized photohemolysis (human sickle cell erythrocytes), **139**, 780
- Lead
-containing thionein, related protein, immunological analysis (rat liver), **136**, 535
- Lectins
-binding immunoassay, detection of glycosylated growth hormone in pituitary gland (human), **140**, 491
cobalt-substituted, circular dichroism (lentil), **136**, 438
D-galactose-specific, precipitation by triantennary complex-type oligosaccharide (*Erythrina indica*), **141**, 963
 β -galactoside-binding, 14-kDa, mRNA nucleotide sequence (chicken embryo), **134**, 51
interactions with variant surface glycoproteins incorporated into liposomes (*Trypanosoma brucei brucei*), **141**, 1274
M13, hemocyte-coagulating, purification and characterization (*Manduca sexta*), **137**, 729
release from fatty acid auxotroph grown in presence of oleic acid (*Saccharomyces cerevisiae*), **136**, 596
- Leishmania*
promastigotes, carbocyclic inosine metabolism and cytotoxicity (*L. donovani*, *L. tropica*), **134**, 1114
- Lens
hexose monophosphate shunt activity, dynamic assessment by proton NMR (rabbit), **138**, 1068
- Leucine
effect on insulin release from cloned B-cell line HIT-T15 (hamster), **140**, 616
glycyl derivative, effect on IgG binding to bacterial Fc receptors, **135**, 1105
- Leucopyrokinin
active fragments and analogs, structure-function analysis (*Leucophaea maderae*), **137**, 936
- Leucosulfakinin II
amino acid sequence, homology with cholecystokinin and gastrin (*Leucophaea maderae*), **140**, 357
- Leucovorine, *see* Folinic acid-SF
- Leukocyte elastase
inhibition by
benzoxazinones, effects of electron withdrawal and 5-substitution (human), **140**, 928
(*dl*)-3-benzyl-N-(methanesulfonyloxy) succinimide (human), **141**, 741
peptide aldehydes and α_1 -antitrypsin for treatment of emphysema, comparison (human), **139**, 896
- Leukocytes
circulating, effects of human recombinant tumor necrosis factor (mouse), **141**, 818
5-lipoxygenase pathway, activation after ethanol enhancement of arachidonic acid transmembrane penetration (human), **140**, 782
in lung lavages, leukotoxin biosynthesis (hyperoxia-exposed rat), **134**, 1071; *erratum*, **135**, 1150
polymorphonuclear, *see* Neutrophils
- Leukotriene A₄ hydrolase
tissue distribution (guinea pig), **135**, 139
- Leukotriene B₄
antagonistic effects of SM-9064 (mouse, rat), **138**, 540
and ω -oxidation products, simultaneous quantitation by gas chromatography-mass spectroscopy (human neutrophils), **134**, 512
synthesis in
A23187-stimulated neutrophils, enhancement by phorbol ester (human), **141**, 399
12-hydroperoxyeicosatetraenoic acid-stimulated neutrophils, requirement of free arachidonic acid (human), **138**, 589
stimulated whole blood and isolated neutrophils, comparison (human), **137**, 334
T cells, inhibition by arachidonic acid (human), **135**, 670
- 6-*trans*-Leukotriene B₄
metabolism by polymorphonuclear leukocytes, pathways (human), **136**, 707
- Leukotriene C₄
deficient synthesis in platelets, role of abnormal glutathione metabolism (human), **140**, 857
synthesis in neutrophils, selective inhibition by ethacrynic acid (human), **137**, 195
- Leukotriene C₄ synthase
assay in cell-free extracts (guinea pig, human, rat), **134**, 85
- Levorphanol
conformation at monoclonal antibody binding sites, ¹H transferred nuclear Overhauser effect measurements, **141**, 1267
- Leydig's cells
coculture with Sertoli's cells, effect on steroidogenesis (swine), **134**, 957
steroidogenesis
and cyclic nucleotide levels, regulation by atrial natriuretic factor *in vitro* (mouse), **138**, 399
stimulation by insulin-like growth factor I *in vitro* (rat), **137**, 950
- Ligninase
cDNA clones, identification with synthetic oligonucleotide probes (*Phanerochaete chrysosporium*), **137**, 649
- Linoleic acid
-lipoxygenase system, characterization of free radicals, **141**, 614
- Linolenic acid
ozone-induced oxidation, uric acid protective effects, **137**, 536
- Lipase, *see* Triacylglycerol lipase
- Lipids
and apoprotein constituents of high-density lipoproteins, role in high-density lipoprotein-induced cardiac prostacyclin synthesis (rabbit), **139**, 416
dietary content, effect on nuclear membrane structure and function (murine liver), **135**, 655
linked oligosaccharide, acetylated and pyruvylated, synthesis *in vitro* (*Rhizobium trifolii*), **134**, 205
peroxidation
effects of anthrapyrazoles and aminoacridine derivative (rabbit hepatic microsomes), **140**, 797
enhancement by polyribonucleosinic acid-polyribocytidylic acid treatment (mouse, rat), **134**, 632
formation of hydrogen atoms and hydroxyl radicals, ESR spectroscopy, **141**, 381
in hepatic microsomes, hepatocytes, and granulosa cells,

Subject Index

- relationship to aldrin epoxidation (rat), **138**, 24
- inhibition by anisylthiolthione (rat hepatic microsomes), **135**, 1015
- transfer protein, isolation from hemolymph (*Manduca sexta*), **136**, 260
- uptake by hepatocyte monolayers, apolipoprotein E-induced, role of glycosaminoglycan sulfates (rat), **141**, 158
- Lipocortins
 - phosphorylation *in vitro* by protein kinase C (bovine), **141**, 547
- Lipolysis
 - inhibition by insulin, requirement of phosphodiesterase activation (human adipocytes), **141**, 1157
- Lipopolysaccharides
 - bacterial, activation of macrophages, role of lipoxygenases (mouse), **138**, 842
- Lipoprotein lipase
 - activation by *N*- α -palmitoyl (56–79) fragment of apolipoprotein C-II (human plasma), **137**, 1041
 - catalyzed hydrolysis, mechanism and inhibition, analysis with phenyl-*n*-butylborinic acid, **134**, 386
 - and growth arrest, coupling during adipose conversion of preadipocytes, **137**, 903
 - heparin-releasable, activation by cAMP-dependent protein kinase (rat heart), **135**, 196
 - pathway, role in arachidonic acid release from thrombin-stimulated platelets (human), **134**, 1327
 - phospholipase C system, role in arachidonic acid release in macrophages (mouse), **141**, 367
- Lipoprotein receptors
 - high-density, gonadotropin-inducible, identification in solubilized ovarian membranes (rat), **134**, 1006
 - low-density, synthesis, stimulation by diltiazem and verapamil (human skin fibroblasts), **136**, 845
 - mediation of malondialdehyde-modified high-density lipoprotein endocytosis (rat sinusoidal hepatocytes), **137**, 29
- Lipoproteins
 - affinity for apolipoprotein A-IV during cholesterol esterification (human), **135**, 756
 - hemolymph, larval specific, purification and characterization (*Heliothis zea*), **134**, 580
 - high-density, *see* High-density lipoproteins
 - low-density, *see* Low-density lipoproteins
 - very-low-density, β -migrating, metabolism in peritoneal macrophages, inhibition by prostaglandin E₂ (rat), **137**, 461
- Liposomes
 - composed of dioleoylphosphatidylethanolamine and glycophorin, trypsin-induced destabilization, **141**, 973
 - incorporated variant surface glycoproteins, interactions with lectins (*Trypanosoma brucei brucei*), **141**, 1274
 - phosphatidylethanolamine, stabilized with acylated antibodies, destabilization by antigen binding: as rapid assay for viruses, **138**, 931
 - phosphatidylinositol-containing, opsonized by concanavalin A, stimulation of phosphatidylinositol uptake by macrophages, **138**, 1090
 - phosphatidylserine-containing, Ca²⁺-induced phase separation, potentiation by micromolar levels of Zn²⁺ (bovine brain), **136**, 476
 - pH-sensitive, 1-oleoyl-2-acetyl glycerol incorporation, stimulation of superoxide release in neutrophils (guinea pig), **136**, 242
 - polyphosphoinositide-containing, interaction with spermine, **135**, 58
- Lipotropic hormone
 - isolation and primary structure (*Heliothis zea* corpora cardiaca), **135**, 622
 - type II, sequence analysis by fast atom bombardment mass spectrometry (locust corpus cardiacum), **134**, 723
- Lipoxin A
 - activation of protein kinase C (human), **134**, 1215
- Lipoxygenase
 - effects of FAD, FMN, and 5-hydroperoxy eicosatetraenoic acid (rat neutrophil extracts), **141**, 534
 - linoleic acid system, characterization of free radicals, **141**, 614
 - mediation of arachidonic acid metabolism in macrophages, synergistic stimulation by A23187 and phorbol ester (mouse), **140**, 861
 - pathway in leukocytes, activation after ethanol enhancement of arachidonic acid transmembrane penetration (human), **140**, 782
 - role in macrophage activation by endotoxin (mouse), **138**, 842
- Lipoyl dehydrogenase, *see* Dihydrolipoamide dehydrogenase
- Lithium
 - effect on carbachol-induced inositol 1-phosphate accumulation (NCB-20 cells), **136**, 622
- Liver, *see also* Hepatocytes
 - isolated perfused, clearance of acute-phase plasma proteins without or with high-mannose, hybrid, or complex-type oligosaccharide side chains (rat), **141**, 949
 - metabolic activity of key enzymes of carbohydrate ketone body, adenosine, and glutamine, comparison with adipose tissues (rat), **138**, 687
 - nonparenchymal cells, mannose receptor-mediated uptake of ricin A-IgG conjugate, measurement (rat), **137**, 892
- Low-density lipoprotein receptors
 - apolipoprotein E-mediated binding of hypertriglyceridemic very-low-density lipoproteins, detection by ligand blotting technique (bovine adrenal gland), **139**, 333
 - binding domains, in apolipoprotein B-100, identification and proposed consensus sequence (human), **139**, 1078
- Low-density lipoproteins
 - binding to hepatic membrane receptors, inhibition by serum phosphorylcholine-binding protein (rat), **141**, 151
 - hypertriglyceridemic, apolipoprotein E-mediated binding to low-density lipoprotein receptors, detection by ligand blotting technique (bovine adrenal gland), **139**, 333
 - γ -irradiated, uptake by cultured fibroblasts and peritoneal macrophages (human), **138**, 938
 - metabolism in cultured fibroblasts, effect of phospholipid polar head group manipulation (human), **137**, 43
 - particles, apolipoprotein B-100 surface-exposed segments, associated peptides, isolation and amino acid sequence (human), **140**, 250
- Lubrol PX, *see* Polidocanol
- Luciferase
 - firefly, *see* *Photinus*-luciferin 4-monooxygenase (ATP-hydrolyzing)

Subject Index

- Lucigenin
in detection of active oxygen (rat hepatocytes), **140**, 468
- Lung
atrial natriuretic factor detection (rat), **135**, 515
perfused, removal of atrial natriuretic peptide *in situ*, measurement (rabbit), **140**, 868
- Luteinizing hormone
effect on progesterone accumulation and catabolism in cultured granulosa cells (rat), **141**, 1025
gonadotropin-releasing hormone-stimulated secretion, arachidonic acid and protein kinase C coordinate actions (rat pituitary cells), **134**, 134
regulation of prolactin receptors in mammary gland (rat), **135**, 1076
release by pituitary, stimulation by
copper (rat), **136**, 417
gonadotropin-releasing hormone, effect of wheat germ agglutinin (rat), **140**, 750
 α -subunit secreted by tumor cell lines, phosphorylation (human), **140**, 143
- Luteinizing hormone receptors
expression in ovarian granulosa cells, effect of transforming growth factor type β (rat), **139**, 800
- Luteinizing hormone-releasing hormone
antagonists containing D-3-pyridinyl-alanine⁶, increased potency, **137**, 709
- Lymphadenopathy-associated virus
-human T-cell lymphotropic virus type III
cytotoxicity, inhibition by aurointricarboxylic acid *in vitro*, **136**, 64
glycoprotein expression, suppression by glycosylation inhibitors, **141**, 33
selective inhibition by 2',3'-dideoxycytidine, **140**, 735
- Lymphocytes
activated, insulin receptor-mediated photocytotoxic effect of insulin-conjugated psoralen derivative, **141**, 502
activation assay with flow cytometry of fluorescein fluorescence polarization (BALB/c mouse), **136**, 1021
- B-cell
antibody-activated, intracellular Ca^{2+} release and Ca^{2+} uptake, analysis with indo-1 (mouse), **137**, 500; *erratum*, **138**, 503
antigen-induced activation, role of phosphatidylinositol pathway, **140**, 1079
HIT-T15, cloned, insulin release, effects of glucose, amino acids, sulfonylureas, and colchicine (hamster), **140**, 616
from patients with adenine phosphoribosyltransferase deficiencies, establishment and characterization, **137**, 998
protein kinase C redistribution and proteolytic cleavage during mitogenesis (human), **135**, 146
synergistic activation by phorbol dibutyrate and ionophore A23187 (human), **138**, 554
virus-transformed, detection of interleukin-1 α mRNA, **139**, 353
calcium transport by ionophorous peptides, detection by quin-2 fluorescence (dog, human), **134**, 731
DNA repair synthesis, inhibition by D-ribose (human), **138**, 673
glutamine utilization pathway, intracellular distribution of enzymes (rat), **138**, 318
interaction with *Salmonella minnesota*, energy dependence and role of thiol groups, **137**, 222
natural killer cell activity, inhibition by eicosapentaenoic acid (human), **138**, 1058
potassium channels, inhibition by retinoic acid (human), **138**, 560
proliferating, 48-kDa protein, isolation and characterization (human, mouse), **134**, 1238
protein-tyrosine kinase detection by renaturation in polyacrylamide gels, **134**, 963
p-type, detection of galactosyltransferase (human), **137**, 1187
surface thiols, accessibility, alteration by ionophores or OKT-3 antibodies (human), **140**, 999
- T-cell
activation-induced synthesis of sialoglycoconjugates, selective covalent modification (human), **135**, 473
cytosolic and particulate protein kinase C, stimulation by phytohemagglutinin (human), **137**, 128
leukotriene B₄ production, inhibition by arachidonic acid (human), **135**, 670
phytohemagglutinin-stimulated, concomitant binding and degradation of and pyruvate dehydrogenase activation by insulin (human), **134**, 412
protein synthesis, polymyxin B effect (human), **135**, 649
transient accumulation of pteridines, induction by phorbol ester (human), **141**, 494
virus-transformed, detection of interleukin-1 α mRNA, **139**, 353
- thymic
phosphatidylinositol turnover, stimulation by concanavalin A, effect on activation (mouse), **141**, 657
sodium-hydrogen exchange, osmotic activation, protein kinase C role (rat), **134**, 8
transfer of Epstein-Barr virus DNA fragment coding for nuclear antigen: gene expression without cell transformation (human), **134**, 1260
- Lysine
side chains in elastin, oxidation by canine myeloperoxidase and stimulated human neutrophils (chicken embryonic aorta), **135**, 451
- Lysolecithin
interaction with bombesin, effect on bombesin conformation, **141**, 99
- Lysophosphatidic acid
and lysophosphatidylinositol, effect on insulin release from pancreatic islets, comparison (rat), **138**, 720
- Lysophosphatidylcholine
-cholesterol vesicles, structure, ³¹P NMR spectroscopy, **139**, 25
micelles, structural transition into multilaminar vesicles, induction by myelin basic protein (bovine brain), **134**, 313
- Lysophosphatidylinositol
stimulation of insulin release from pancreatic islets (rat), **138**, 720
- Lysosomes
fibroblast buoyant fractions, N-acetylneuraminic acid accumulation in patients with generalized N-acetylneuraminic acid storage disease, **139**, 838
glucocerebrosidase routing to, role of oligosaccharide chains (U937 cells), **141**, 452

Subject Index

- ovarian, divergent metabolism of human chorionic gonadotropin subunits (rat), **134**, 1197
- sialic acid metabolism, defect in Salla disease (rat liver), **137**, 175
- Lysozyme**
- secretion by macrophages, inhibition by tetanus toxin, role of cytosolic Ca^{2+} homeostasis (human), **135**, 16
- M**
- α_2 -Macroglobulin
- without or with high-mannose, hybrid, or complex-type oligosaccharide side chains, clearance by isolated perfused liver (rat), **141**, 949
- synthesis and secretion by fibroblasts (human lung), **134**, 57
- Macromolecules**
- epidermal, covalent attachment to ω -hydroxyacid derivatives (pig), **137**, 992
- Macrophages**
- activated, antitumor potential, receptor-mediated inhibition by dexamethasone (mouse), **136**, 130
- activation by endotoxin, role of lipoxygenases (mouse), **138**, 842
- bone marrow-derived, glucose uptake, stimulation by colony-stimulating factor 1 (mouse), **138**, 445
- colony-stimulating factor purification from medium conditioned by pancreatic carcinoma cells (human), **138**, 238
- derived factor, endothelial cell-stimulating, partial purification and properties (human), **134**, 344
- lysozyme secretion, inhibition by tetanus toxin, role of cytosolic Ca^{2+} homeostasis (human), **135**, 16
- peritoneal
- cultured, synthesis of oversulfated chondroitin sulfate proteoglycan (human), **139**, 377
- interleukin 1 and prostaglandin E_2 production, stimulation by tumor necrosis factor (mouse), **136**, 94
- lipoxygenase-mediated arachidonic acid metabolism, synergistic stimulation by A23187 and phorbol ester (mouse), **140**, 861
- mannosyl-fucosyl receptors, antagonistic effects of interferons α/β and γ (mouse), **136**, 737
- metabolism of β -migrating very-low-density lipoprotein, inhibition by prostaglandin E_2 (rat), **137**, 461
- production of interleukin 1, protein kinase C role (mouse), **135**, 355
- prostaglandin synthesis, regulatory role of protein kinase C (mouse), **139**, 604
- uptake of γ -irradiated low-density lipoprotein (mouse), **138**, 938
- production of tumor necrosis factor, regulation by prostaglandin E_2 (mouse), **137**, 404
- release of arachidonic acid, role of phospholipase C-diglyceride lipase system (mouse), **141**, 367
- respiratory burst, associated H_2O_2 production, enhancement by interferon γ , antagonistic effects of interferons α and β (human), **140**, 948
- tumor necrosis factor induction, association with β -carotene (hamster), **136**, 1130
- uptake of phosphatidylinositol-containing liposomes, stimulation by concanavalin A opsonization, **138**, 1090
- Magnesium**
- ATP and uncomplexed ATP, observation in slow exchange by ^{31}P NMR, **134**, 1379
- and calcium, sequential binding to bovine prothrombin fragment 1, associated spectral response, **141**, 1207
- fluxes in mitochondria, quinine effect (rat liver), **135**, 830
- high-affinity binding to actin monomers, **135**, 607
- role in ATP/Mg-dependent phosphorylase phosphatase activation by kinase F_A , **136**, 16
- uptake and release from cerebral cortical synaptosomes characterization and properties (guinea pig), **136**, 906
- Maize streak virus**
- and satellite tobacco necrosis virus, coat protein sequence homology, **134**, 372
- Major intrinsic polypeptide**
- lens membrane, covalent change during senile cataractogenesis (human), **135**, 965
- Malaria**
- Plasmodium falciparum*, detection with cloned DNA probe, **138**, 363
- Malate dehydrogenase**
- mitochondrial, and precursor, conformation, comparison (CHO cells), **141**, 313
- NADP $^{+}$ -linked in glutamine utilization pathway, intracellular distribution in lymphocytes (rat), **138**, 318
- Malate dehydrogenase (oxaloacetate-decarboxylating) (NADP $^{+}$)**
- effects of epidermal growth factor, growth hormone, and thyroxine after chronic administration (hypophysectomized rat), **141**, 78
- Malic enzyme, see Malate dehydrogenase (oxaloacetate-decarboxylating) (NADP $^{+}$)**
- Malondialdehyde**
- modified high-density lipoprotein, receptor-mediated endocytosis by sinusoidal hepatocytes (rat), **137**, 29
- Mammary gland**
- differentiation, and DNA synthesis and poly(ADP-ribosylation), interrelationships (mouse), **137**, 359
- normal, preneoplastic, or neoplastic cells, metabolic cooperation with drug-resistant tumor cells *in vitro*, ouabain effects (mouse), **140**, 955
- Manganese**
- adenylate cyclase sensitivity, increase by fasting, mechanism (rat fat cells), **138**, 1411
- binding to diphtheria toxin, ESR analysis, **139**, 991
- Mannan**
- in concanavalin A solutions, detection and quantitation with electrical oscillator, **137**, 978
- Mannose**
- containing glycopeptide D3, precipitation of concanavalin A, stoichiometry, **137**, 670
- toxicity in honeybees, mechanism, **140**, 51
- Mannose receptors**
- mediation of ricin A–IgG conjugate uptake by hepatic non-parenchymal cells, measurement (rat), **137**, 892
- Mannosidase II**
- detection in mung bean seedling extracts, **134**, 1395
- Mannosyl-fucosyl receptors**
- antagonistic effects of interferons α/β and γ (murine peritoneal macrophages), **136**, 737
- Mass spectrometry**
- in determination of position of *O*-acylated residues in

Subject Index

- oligosaccharide sequences, **135**, 1126
- fast atom bombardment, *see* Fast atom bombardment mass spectrometry
- heparin fragments, in determination of partial structure (porcine mucosa), **139**, 18
- reaction products of surface-adsorbed peptides and proteins, **134**, 420
- Mast cells
 - peritoneal, compound 48/80-induced histamine secretion, effect of phorbol ester pretreatment (rat), **137**, 1112
- Mastoparan
 - binding by calmodulin, ^1H NMR analysis, **137**, 1069
 - calmodulin complexes, Ca^{2+} binding: detection of fluorometrically distinguishable intermediate, **135**, 1050
- Megakaryocytes
 - fibrinogen gene expression (rat), **140**, 543
- α -Melanocyte-stimulating hormone, *see* α -Melanotropin
- Melanoma growth factor
 - acid-sensitive, role in human melanoma and bovine endothelial cell proliferation (human platelets), **138**, 795
- Melanosomes
 - aggregation, epinephrine-induced, comparative effects of ACTH and cAMP (goldfish), **138**, 1081
- α -Melanotropin
 - effects on zona glomerulosa *in vivo* (rat), **141**, 1279
 - fragment analogs, synthesis and properties, **137**, 722
- Melatonin
 - enhancement of β -adrenergic ligand binding to pineal particulate fractions (rabbit), **138**, 1196; *erratum*, **139**, 1320
- Melittin
 - cytolysis of spleen, bone marrow, and leukemia cells, relationship to membrane structure (mouse), **139**, 222
 - induced damage in pancreatic acinar membranes, role of Ca^{2+} (rat), **139**, 530
 - interaction with bovine calmodulin (bee venom), **135**, 419
- Mellitin, *see* Melittin
- Membrane metallo-endopeptidase
 - active site, detection of histidine residue (rabbit kidney), **139**, 733
- Membranes
 - alkaline phosphatase catalytic rate enhancement by diffusion of adsorbed substrate on membrane surface, **138**, 803
 - basal, identification of 76-kDa β -adrenergic receptor (human placenta), **141**, 411
 - binding sites in spleen, bone marrow, and leukemia cells, differences, detection by cytolysis with melittin (mouse), **139**, 222
 - boundary, in endocytic vesicles, hydrophobic domain, close association with clathrin (bovine brain), **141**, 878
 - bound Na^+ , K^+ -ATPase, solubilization and purification (porcine kidney), **135**, 1008
 - cardiac, identification of substrates for cholera toxin- and pertussis toxin-catalyzed ADP-ribosylation (rat), **138**, 1355
 - cation permeability, α -latrotoxin effects (rat PC12 neurosecretory cells), **134**, 320
 - chromaffin granule, $\text{p60}^{\text{c-src}}$ activity detection (bovine), **134**, 736
 - cyanobacterial, a_3 -type cytochrome oxidase immunological identification (*Anacystis nidulans*), **136**, 235
 - erythrocyte
 - bicarbonate exchange kinetics at equilibrium, ^{13}C NMR (human), **136**, 266
 - corticotropin-releasing factor binding sites, identification and modulation by chronic ethanol treatment (human, rat), **136**, 137
 - outer bilayer, phosphatidylserine localization (newborn human), **136**, 914
 - ozone-induced oxidation, uric acid protective effects (bovine), **137**, 536
 - phosphatidic acid metabolism, activation by perfringolysin O (human), **137**, 23
 - phosphorylation-induced Ca^{2+} release (human), **136**, 288
 - protein band 3, dephosphorylation by phosphotyrosine protein phosphatase (human), **134**, 557
 - proteins and/or glycoproteins, functional group labeling with biotin derivatives and ^{125}I -labeled streptavidin (human), **136**, 80
 - sickle cell, hematorporphyrin-sensitized photohemolysis, inhibition by lawsone (human), **139**, 780
 - fusion, synergistic effects of micromolar Zn^{2+} and Ca^{2+} levels (bovine brain), **137**, 101
 - glomerular basement, degradation by glomerular neutral metalloproteinase (rat kidney), **141**, 898
 - Golgi, detection of putative opioid-peptide processing activity (rat), **138**, 356
 - hepatocyte
 - low-density lipoprotein receptors, binding activity, inhibition by serum phosphorylcholine-binding protein (rat), **141**, 151
 - transferrin receptor recovery after perfusion with collagenase (rat), **134**, 711
 - inner mitochondrial
 - association with creatine kinase in physiological salt solution, relative importance of salt composition and ionic strength (rat heart), **139**, 1262
 - ATP synthase subunit 8, localization of hydrophilic C terminus (*Saccharomyces cerevisiae*), **138**, 78
 - intestinal basolateral
 - Ca^{2+} -ATPase and Ca^{2+} pump activity, comparison (rat), **141**, 979
 - Na^+ - H^+ exchange, detection by fluorescence quenching (rabbit), **134**, 827
 - lens, major intrinsic polypeptide covalent change during senile cataractogenesis (human), **135**, 965
 - leukocyte, penetration by arachidonic acid, enhancement by ethanol (human), **140**, 782
 - lipid bilayer
 - pH-dependent insertion of diphtheria toxin B fragment, conformational analysis, **136**, 160
 - transmembrane K^+ currents, alteration induced by concanavalin A-polysaccharide interaction, **141**, 912
 - lymphocyte, calcium transport by ionophorous peptides, detection by quin-2 fluorescence (dog, human), **134**, 731
 - mitochondrial, potential, safranin as probe: side effects on Ca^{2+} transport (rat liver), **135**, 189
 - neuroblastoma cell, degradation of dynorphins, role of cysteine protease (mouse), **140**, 335
 - neutrophil

Subject Index

- binding of platelet-activating factor, inhibition by GTP (human), **141**, 353
- fluidity changes associated with chemotactic peptide-induced stimulation (human), **141**, 1151
- stimulus-dependent movement of protein kinase C and phorbol myristate acetate receptor from cytosol (human), **134**, 587
- nuclear
- receptor role in γ -interferon transport into cell nucleus (mouse), **138**, 254
 - structure and function, effect of dietary lipid content (murine liver), **135**, 655
- outer
- action as barrier against disaccharide penetration (*Pseudomonas aeruginosa*), **134**, 106
 - permeability (*Bacteroides fragilis*), **141**, 292
- ovarian
- hydroxyurea uptake by diffusion mechanism (Chinese hamster), **134**, 1254
 - identification of gonadotropin-inducible high-density lipoprotein receptors (rat), **134**, 1006
- pancreatic acinar, damage induced by toxic agents, role of Ca^{2+} (rat), **139**, 530
- permeability changes induced by *Clostridium perfringens* type A enterotoxin in absence of extracellular Ca^{2+} (HeLa, Vero cells), **141**, 704
- peroxisomal, associated polypeptides, synthesis *in vitro* (rat liver), **137**, 626
- phospholipid bilayer, solvent-free, incorporated Na^+ , K^+ -ATPase, ion-channel activity, voltage and ATP effects (canine renal microsomes), **136**, 767
- photoreceptor
- inositide and phosphatidic acid metabolism, effect of cationic amphiphilic drugs (bovine rod outer segments), **134**, 378
 - K^+ currents, Ca^{2+} -dependent inactivation, protein kinase C effects (*Hermisenda crassicornis*), **134**, 1245; *erratum*, **140**, 774
- placental, identification of atrial natriuretic factor receptors (human), **135**, 480
- plant and animal, [^3H]verapamil binding, effect of 3,4,5-triiodobenzoic acid, **138**, 1269
- plasma, *see* Plasma membranes
- platelet
- Ca^{2+} -ATPase activity, effect of activating hormones (human), **138**, 215
 - phosphoinositide hydrolysis, stimulation by GTP and cytosol (human), **137**, 801
- protein kinase C activation, inhibition by exogenous phospholipase C (murine epidermal HEL-37 cells), **136**, 1083
- pulmonary, detection of high-affinity atriopeptin-III receptors (rabbit), **140**, 302
- renal, amiloride-sensitive sodium channels, interaction with phenamil, characterization (swine), **135**, 25
- renal basolateral
- binding of human α -atrial natriuretic factor, hypertension effects (rat), **137**, 1079
 - Na^+/K^+ -ATPase activity, effect of insulin (non-insulin-dependent diabetic rat), **139**, 1313
- sarcoplasmic reticulum, transmembrane pH equilibration rate, measurement (rabbit skeletal muscle), **140**, 934
- skeletal and heart muscle, formation of phosphatidylinositol 4,5-bisphosphate (rabbit), **138**, 1395
- striatal, D-1 dopamine receptor extraction (rat), **137**, 943
- 3T6 cell, permeabilization by external ATP, **134**, 453
- vesicles, GABA-stimulated chloride uptake, effect of benzodiazepines and β -carbolines (rat cerebral cortex), **141**, 1
- Mepyramine, *see* Pyrilamine
- 7-Mercaptoheptanoylthreonine phosphate
- binding to component C of methylcoenzyme M methyl-reductase system, demonstration (*Methanobacterium thermoautotrophicum*), **139**, 889
- Meromyosin
- heavy, interaction with actin, analysis, **141**, 541
- Metal ions
- role in metallodaunomycin-induced DNA degradation, **136**, 1
- Metallodaunomycin
- induced DNA degradation, role of metal ions, **136**, 1
- Metalloproteinase
- neutral, glomerular, degradation of glomerular basement membrane (rat kidney), **141**, 898
- Methane
- synthesizing complex, detection of tetrahydromethanopterin methyltransferase (*Methanobacterium thermoautotrophicum*), **136**, 542
- Methanobacterium thermoautotrophicum*
- occurrence of cAMP, **136**, 1078
- Methiocarb
- oxidation by sulfoxidase (soybean cotyledonal microsomes), **135**, 922
- Methionine
- formation from 5'-deoxy-5'-methylthioadenosine and L-homocysteine, role in cell growth (BHK, CCL 39, Raji cells), **135**, 131
 - in ubiquitin, role in conformation and activity (human erythrocytes), **138**, 437
- Methionine-enkephalin
- crystal structure, determination with X rays, **134**, 698
 - detection in intraglandular colloid of intermediate lobe origin (bovine pituitary), **139**, 480
 - effect on atrial natriuretic factor release by atrial slices (rat), **135**, 1090
- Methionyl growth hormone
- human recombinant DNA-derived, 20-kDa variant, early insulin-like effects (hypophysectomized rat), **138**, 342
- Methionyl interleukin 2
- and recombinant human IL-2, produced in *Escherichia coli*, separation, **135**, 837
- Methotrexate
- effect on uracil incorporation into DNA (HeLa cells, human lymphoblastoid cells), **135**, 886
 - induction of DNA lesions, inhibition by leucovorine (human colon adenocarcinoma cells), **138**, 544
- Methoxatin
- pulse radiolysis, **139**, 955
- Methylamine dehydrogenase
- covalent cofactor, structure and linkage (*Methylobacterium* sp.), **141**, 562
- Methylation
- c-myc proto-oncogene, changes during aging process (mouse), **139**, 1299

Subject Index

- plasmid containing cell cycle regulatory sequence, analysis after transient transfection into fibroblasts (hamster), **135**, 942
- thyroglobulin gene, relationship to gene expression (human), **134**, 1109
- 3-Methyl-6-chloro-9-azido-1H-2,3,4,5-tetrahydro-3-benzazepine
- ³H-labeled, photoaffinity labeling of α_2 -adrenoceptors (human platelets, rabbit kidney), **137**, 606
- 3-Methylcholanthrene
- effect on glutathione transferase expression in seminal vesicles and pituitary gland (rat), **136**, 1057
- treated liver, aflatoxin B₁-specific cytochrome P-450 purification and characterization (hamster), **139**, 466
- 5-Methylchrysene
- K-regions, identification as sites of oxidative metabolism (rat hepatic microsomes), **141**, 734
- Methyl cobalamine
- inactivation by butyl nitrite, NMR analysis, **135**, 201
- Methylcoenzyme M methylreductase system
- component C, demonstration of binding to 7-mercaptoheptanoylthreonine phosphate (*Methanobacterium thermoautotrophicum*), **139**, 889
- 1-Methylcyclohexanecarboxylic acid
- effects on cellular and mitochondrial respiration (murine cerebral cortex, neuroblastoma), **140**, 789
- 7-Methylguanosine
- and L-phenylalanine and 5'-monophosphate, crystal structure, interaction model, **136**, 294
- 1-Methyl-3-isobutyl-xanthine
- effect on cAMP metabolism by thrombopathic platelets (dog), **140**, 595
- N-Methyl-N'-nitro-N-nitrosoguanidine
- exposed mammalian cells, poly(ADP-ribose) biosynthesis and suicidal NAD⁺ depletion, relationship (mouse, rat), **138**, 1051
- 1-Methyl-4-phenylpyridine
- induction of ATP depletion in isolated hepatocytes (rat), **137**, 310
- mitochondrial uptake, relationship to inhibition of NAD⁺-linked substrate oxidation (rat liver), **134**, 743
- and paraquat, cytotoxic effects on isolated hepatocytes, comparison (rat), **137**, 303
- production of superoxide and hydroxyl radicals in presence of NADPH-cytochrome P-450 reductase and oxygen, **135**, 583
- 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine
- induction of ATP depletion in isolated hepatocytes (rat), **137**, 310
- 5-Methyl tetrahydrofolate
- inactivation by butyl nitrite, NMR analysis, **135**, 201
- 1-Methyl tetrahydroisoquinoline
- endogenous, detection in brain (rat), **140**, 448
- Mevinolin
- effect on acetoacetyl-CoA thiolase activity in fibroblasts (CHO-K1 cells), **138**, 953
- Mezerein
- inactivation of glycogen synthase by phosphorylation (rat hepatocytes), **134**, 113
- inhibition of chondroitin sulfate sulfation (human promyelocytic leukemia cells), **135**, 909
- Micelles
- reverse, application in protein extraction, **135**, 629
- Microciona prolifera*
- cell aggregation, induction by Ca²⁺ pulses, ionophores, and phorbol esters without extracellular Ca²⁺, **134**, 1319
- Micrococcal nuclease
- conversion of adenylylated glutamine synthetase to phosphoryl form (*Mycobacterium smegmatis*), **137**, 716
- Microfilaments
- role in vitamin D metabolism (chicken renal tubules), **134**, 218
- β_2 -Microglobulin
- fragments, detection in amyloid kidney stones from uremic patients, **136**, 665
- inhibition of *in vitro* calcification (murine osteoblastic cells), **141**, 360
- Microsomes
- hepatic, reduction of vanadate (rat), **141**, 171
- oxidation of methiocarb (soybean cotyledons), **135**, 922
- seminal vesicle, hydroperoxide-dependent epoxidation of 3,4-dihydroxy-3,4-dihydrobenzo[a]anthracene (sheep), **140**, 181
- Microtubule-associated proteins
- type 2, related muscle proteins, identification as substrates for soluble insulin-stimulatable serine/threonine kinase (3T3-L1 adipocytes), **134**, 565
- Microtubules
- assembly inhibition, effect on transferrin receptor cycling by lymphoid cells (human), **138**, 1216
- associated protein τ , self-assembly into filaments resembling Alzheimer's disease-associated filaments (porcine brain), **141**, 790
- elongation, inhibition by GDP, **137**, 351
- and τ proteins, association with retinal heat shock protein, analysis (rabbit), **139**, 974
- role in vitamin D metabolism (chicken kidney), **134**, 218
- Microvilli
- fusion and pinching-off, induction by concanavalin A (rat colon), **140**, 766
- Milk
- detection of immunoreactive growth hormone-releasing factor (human), **135**, 1084
- Mineralocorticoid receptors
- antagonist-specific, identification in kidney (rat), **134**, 261
- Mitochondria
- cardiac, ATP-ADP translocase and creatine kinase levels, determination with enzyme inhibitors (chicken, dog, rabbit, rat), **134**, 359
- S-(2-chloro-1,1,2-trifluoroethyl)-L-cysteine-induced toxicity, role of hydrogen sulfide (rat kidney), **138**, 707
- cytochrome and polypeptide composition, effect of mitochondrial myopathy and renal dysfunction (human), **137**, 911
- energized, internalization of CAT₁₂ (rat liver), **136**, 941
- energy-linked processes, antagonistic effects of halothane and L-carnitine, role of long-chain acyl-CoA (rat liver), **139**, 303
- fluxes of K⁺ and Mg²⁺, quinine effects (rat liver), **135**, 830
- heme excess generation, inhibition of L-alanine:4,5-dioxovalerate transaminase (rat liver), **138**, 751
- inner membrane
- association with creatine kinase in physiological salt so-

Subject Index

- lution, relative importance of salt composition and ionic strength (rat heart), **139**, 1262
- ATP synthase subunit 8, localization of hydrophilic C terminus (*Saccharomyces cerevisiae*), **138**, 78
- localization of enzyme for heme attachment to apocytochrome *c* (yeast), **141**, 1145
- membrane potential, safranin as probe: side effects on Ca^{2+} transport (rat liver), **135**, 189
- 1-methyl-4-phenylpyridine uptake, relationship to inhibition of NAD^{+} -linked substrate oxidation (rat liver), **134**, 743
- oxidative phosphorylation, effects of prostaglandin B_1 oligomeric derivatives (rat liver), **137**, 215
- photosensitization *in vitro* by hematoporphyrin derivative: impairment of anion translocation (rat liver), **141**, 584
- production of free radicals, induction by hydroperoxide (rat liver), **141**, 1123
- protein phosphorylation, effect of estrone treatment (genetically diabetic mouse), **139**, 56
- protein synthesis, inhibition by rhodamine 123 (rat liver, tumor cells), **137**, 759
- respiration, effects of 1-methylcyclohexanecarboxylic acid (murine cerebral cortex, neuroblastoma), **140**, 789
- subparticles, cyanide-resistant duroquinol oxidase activity, comparison with NADH and succinate oxidation (*Arum maculatum*), **135**, 677
- superoxide formation, role of ubiquinone and cytochrome *b566* (rat heart), **138**, 533
- thiamine uptake, analysis *in vitro* (rat liver), **141**, 466
- translation products *in vitro*, contamination with bacterial proteins (*Drosophila melanogaster*), **136**, 797
- ubiquinone levels, thyroxine-induced changes (rat liver), **138**, 381
- uptake and processing of ornithine carbamyl transferase precursor *in vitro* after production in *Escherichia coli* (rat heart, liver), **134**, 21
- Mitomycin C**
- induction of phage-like particles in pathogenic *Vibrio parahaemolyticus* strain, **141**, 1249
- Mitoxantrone**
- effect on topoisomerase I and II (human breast cancer cells), **136**, 521
- inhibition of angiogenesis (rat avascular cornea), **140**, 901
- Moloney murine sarcoma virus**
- ts110 mutant-transformed cells, immunodetection of transformation-associated protein (rat kidney), **134**, 1223
- Molybdenum**
- cofactor, xanthine oxidase, dephosphorylated oxidation product, identification as 2-(1,2-dihydroxyethyl)thieno[3,2-*g*]pterin, **135**, 256
- detection in tetrathionate reductase (*Salmonella typhimurium*), **136**, 577
- iron cofactor of *Azotobacter vinelandii* nitrogenase, solubilization in dimethylformamide and acetonitrile, **139**, 740
- Monensin**
- effect on
- intracellular Ca^{2+} levels and contractile activity in isolated perfused heart (rabbit), **139**, 816
- protein synthesis inhibition by bovine cell-surface sialoglycopeptide (murine 3T3 cells), **136**, 976
- Monoamine oxidase**, *see* Amine oxidase (flavin-containing)
- Monoclonal antibodies**
- binding sites
- flexible opiate conformations, ^1H transferred nuclear Overhauser effect measurements, **141**, 1267
- in F_1 γ -subunit, detection (*Escherichia coli*), **137**, 468
- to cell-surface nerve growth factor receptor, in identification of chromatin-associated nerve growth factor receptors (human melanoma cell), **140**, 174
- to cytomegalovirus, generation of producing hybridomas (human), **137**, 273
- demonstration of high-molecular-weight forms of inhibin in follicular fluid (bovine), **136**, 1103
- to herpes simplex virus, producing hybridomas, generation after stimulation *in vitro* (human), **135**, 495
- OKT-3, alteration of accessibility of surface thiols (human lymphocytes), **140**, 999
- to pyruvate oxidase, characterization: enzyme conformation effects on antibody-induced inhibition (*Escherichia coli*), **137**, 884
- recognizing superficial tubulin-like antigen, identification in *Trypanosoma cruzi*-infected mice, **139**, 1176
- to recombinant human tumor necrosis factor, production and characterization, **137**, 847
- RNA polymerase II, in analysis of subunit functional properties (bovine), **136**, 281
- to sarcoma DNA polymerase α , isolation and characterization (mouse), **141**, 60
- for species distinction of type I insulin-like growth factor receptors, identification, **138**, 1341
- surface antigen-directed, coordinate inhibition of NADPH oxidase activation and phosphorylation of membrane proteins (human neutrophils), **140**, 1121
- to transcortin, in phylogenetic analysis of epitopes (human), **138**, 1037
- Monocytes**
- cAMP levels, synergistic action of adenosine and formyl-methionyl-leucyl-phenylalanine (human), **138**, 1376
- effects of κ elastin (human), **141**, 92
- phospholipase C, inhibition by aspirin, mechanism (human), **139**, 115
- Monooxygenase system**
- hepatic, reconstitution in solution from cytochrome *P*-450 and NADPH-specific flavoprotein monomers (rabbit), **139**, 883
- Monophenol monooxygenase**
- fungal, translation *in vitro*, **141**, 697
- 5'-Monophosphate**
- and L-phenylalanine and 7-methylguanosine, crystal structure, interaction model, **136**, 294
- Morphine**
- binding to anti-morphine antibodies and opiate receptors, inhibition by anti-paratypic antibody fraction (rabbit), **136**, 1177
- concentration estimation by comparative immunoremoval (bovine tissues), **137**, 316
- Morphine receptors**
- and enkephalin receptors, identification in neuroblastoma cells (human), **137**, 813
- Mouse hepatitis virus**
- nucleocapsid protein, phosphorylation sites, **141**, 7; *erratum*, **141**, 884

Subject Index

- Multiple sulfatase deficiency
associated rapid degradation of steroid sulfatase (human), **135**, 189
- Murabutide
-synthetic pre-S viral peptide-toxoid carrier conjugates, in production of antibodies recognizing hepatitis B virus surface antigens, **140**, 557
- Muramyltripeptides
lipophilic MTP-PE, activation of protein kinase C (porcine brain), **140**, 1043
- Murine leukemia virus
reverse transcriptase, inhibition by 2',3'-dideoxythymidine 5'-triphosphate and derivations modified on 3' position, **140**, 498
- Murine mammary tumor virus
cytoplasmic RNA, destabilization by heat shock, prevention by cycloheximide, **137**, 1028
promoter-linked *v-ras*^H oncogene, expression, cAMP effect, **134**, 436
- Muscarinic receptors
binding activity in neuroblastoma cells, effects of phorbol esters (human), **140**, 160
coupling to calcium channels and polyphosphoinositide hydrolysis, induction by point-mutated *p21^{ras}* (murine fibroblasts), **141**, 591
in pancreatic acini, chloroquine binding, effect on secretory responses (rat), **137**, 664
quinuclidinyl benzilate binding, effect of guanine nucleotides (rat heart), **134**, 172
stimulation of inositol tetrakisphosphate accumulation in cerebral cortical slices, enhancement by K⁺-induced depolarization (rat), **141**, 1130
- Muscimol
³H-labeled, photoaffinity labeling of γ -aminobutyric acid receptor binding site, comparison with [³H]flunitrazepam binding (bovine brain), **138**, 1308
- Muscles
skeletal
accumulation of myosin light chain 3 during development, densitometric measurement (quail embryo), **135**, 778
chemically skinned fibers, contractile mechanism, sensitivity to Ca²⁺, enhancement by inositol 1,4,5-trisphosphate (rabbit), **135**, 662
types 1, 2A, and 2B myosin heavy chains, electrophoretic analysis (rat), **138**, 981
smooth
cerebrovascular, glycogen metabolism, regulatory mechanism, comparison with endothelium (rat), **134**, 484
contraction
and α_1 -adrenergic receptor-stimulated phosphoinositide hydrolysis, inhibition by phorbol esters (rat aorta), **134**, 970
effect of myosin light chain phosphorylation (norepinephrine-stimulated canine veins), **135**, 735
3,4,5-triiodobenzoic acid effects (rabbit), **138**, 1269
vascular, cytoplasmic Ca²⁺ levels, measurement by laser microfluorospectrometry (spontaneously hypertensive rat), **141**, 340
- Muscular dysgenesis
effect on electrophysiological expression of Ca²⁺ channels and apamin-sensitive Ca²⁺-activated K⁺ channels in skeletal muscle (mouse), **136**, 935
- Muscular dystrophy
associated increase in calmodulin gene expression (chicken), **137**, 507
induced alteration of protein kinase C distribution in satellite cells, reversal by liposome-delivered phospholipids (mouse), **137**, 752
inherited, antioxidant enzyme activities in muscle, liver, and lung (chicken), **134**, 550
- Mutagenesis
benzo[a]pyrene *in vitro*, inhibition by resorufin (rat hepatic microsomes), **136**, 555
frameshift, by reversible DNA intercalators, effect of mutation in *uvrB* gene (*Salmonella typhimurium*), **138**, 505
- Mutants
cell division cycle 35-1, adenylate cyclase alterations (yeast), **140**, 723
protease-secreting, isolation and characterization (*Bacillus subtilis*), **138**, 526
temperature-sensitive, ATP synthesis, characterization (*Escherichia coli*), **138**, 72
- Mutations
in IVS 1 donor site consensus sequence inducing β thalassemia, identification (human), **139**, 709
point, c-K-ras codon 12 GGT-CGT, occurrence in lung cancer (human), **140**, 699
polA, enhancement of instability of IncFII plasmid basic replicon, **139**, 918
random single-point, induced destabilization of protein spatial structures, protection by codon bias in genes, **139**, 1094
uvrB gene, effect on frameshift mutations induced by reversible DNA intercalators (*Salmonella typhimurium*), **138**, 505
- Myelin
D-aspartic acid content (bovine, human brain), **135**, 683
- Myelin basic protein
D-aspartic acid content (bovine, human brain), **135**, 683
induction of structural transition of lysophosphatidylcholine micelles into multilaminar vesicles (bovine brain), **134**, 313
inhibition of formyl-peptide-induced neutrophil chemotaxis (human), **141**, 524
interaction with tubulin, effect on tubulin carboxypeptidase, **136**, 482
phosphoinositide covalent linkage, *in vivo* occurrence and effect of allergic encephalomyelitis *in vitro* (rabbit, guinea pig), **136**, 426
synthetic peptide analogs, protein kinase C inhibition (bovine), **134**, 78
- Myeloperoxidase
oxidation of lysine side chains in chicken embryonic elastin (dog), **135**, 451
- Myocytes
cardiac, synthesis of atrial natriuretic factor, serum-mediated enhancement (rat), **140**, 151
myosin isozyme pattern, changes during long-term culture (rat heart), **137**, 855
potassium levels, determination by null-point technique (rat heart), **137**, 573

Subject Index

- Myoglobin
geminate recombination of nitric oxide, analysis on picosecond time scale (elephant, whale), **140**, 962
- Myometrial cells
sodium channel detection with neurotoxins (rat), **137**, 675
- Myopathy
mitochondrial, effect on mitochondrial cytochrome and polypeptide composition (human), **137**, 911
- Myosin
amino acid sequence, homology with group A streptococcal Pep M5 protein, **140**, 684
degradation, calcium effects (rat), **134**, 1269
heavy chains, in types 1, 2A, and 2B skeletal muscle fibers, electrophoretic analysis (rat), **138**, 981
isozyme pattern in cardiac ventricular myocytes, changes during long-term culture (rat), **137**, 855
light chains
alkali, developmentally regulated accumulation in breast and limb muscles, densitometric measurement (quail embryo), **135**, 778
phosphorylation, role in platelet activation (human), **140**, 280
reactivity with Ca^{2+} -dependent and -independent protein kinases, **134**, 240
20-kDa, phosphorylation, effect on smooth muscle contraction (norepinephrine-stimulated canine veins), **135**, 735
- Myosin ATPase
smooth muscle, regulation by thin filament-associated proteins, **136**, 962
- N**
- NAD^+ (oxidized nicotinamide adenine dinucleotide)
-linked substrates, mitochondrial oxidation, inhibition by 1-methyl-4-phenylpyridine, relationship to uptake (rat liver), **134**, 743
suicidal depletion, and poly(ADP-ribose) biosynthesis in mammalian cells following carcinogen exposure, relationship (mouse, rat), **138**, 1051
- NAD^+ ADP-ribosyltransferase
activation by DNA damage, induction of alterations in glucose metabolism (murine L1210 cells), **134**, 227
role in altered deoxynucleoside triphosphate metabolism after DNA damage (murine L1210 cells), **137**, 1153
- NADH (reduced nicotinamide adenine dinucleotide)
cyanide-resistant oxidation in submitochondrial particles, comparison with duroquinol oxidase activity (*Arum maculatum*), **135**, 677
ferricyanide- or transferrin-stimulated oxidation (HeLa cells), **135**, 110
- NADH dehydrogenase
mitochondrial
deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
inhibition by pyridine derivatives, relationship to experimental and idiopathic parkinsonism, **135**, 269
- NADH dehydrogenase (ubiquinone)
mitochondrial, purification of ubiquinone-binding protein (bovine heart), **138**, 1237
subunits in cardiac mitochondria, deficiency in patient with mitochondrial encephalomyopathy and cardiomyopathy, **140**, 88
- NADH -ubiquinone oxidoreductase, *see* NADH dehydrogenase (ubiquinone)
- Nadolol
effect on intracellular Ca^{2+} levels and contractile activity in isolated perfused heart (rabbit), **139**, 816
- NADP (oxidized nicotinamide adenine dinucleotide phosphate)
improvement of cholera toxin-catalyzed ADP-ribosylation efficiency (canine cardiac, rat hepatic membranes), **137**, 328
- NADPH (reduced nicotinamide adenine dinucleotide phosphate)
-specific flavoprotein monomers, and cytochrome *P*-450 in solution, reconstitution of hepatic monooxygenase system (rabbit), **139**, 883
- NADPH -cytochrome *P*-450 reductase
and aromatase, reconstituted system, stereochemistry of estrogen biosynthesis (human placenta), **136**, 454
binary complex formation with cytochrome *P*-450 LM_2 and cytochrome b_5 , mechanism (rabbit hepatic microsomes), **134**, 519
catalysis of superoxide and hydroxyl radical formation from 1-methyl-4-phenylpyridinium ion under aerobic conditions, **135**, 583
cysteine residues, localization and role in enzyme function, assessment by fluorescent titration (rat hepatic microsomes), **134**, 393
interaction with cytochrome *P*-450 isozymes, spectrophotometric analysis (rabbit, rat), **137**, 437
- NAD(P)H dehydrogenase (quinone)
inhibition by lapachol (rat liver), **137**, 781
- NADPH oxidase
activation
by arachidonic acid, role of phospholipase A_2 in intact neutrophils and cell-free system, comparison (human), **138**, 1099
and phosphorylation of membrane proteins, coordinate inhibition by surface antigen-directed monoclonal antibody (human neutrophils), **140**, 1121
in Ca^{2+} -depleted neutrophils, activity restoration by formyl-methionyl-leucyl-phenylalanine and concanavalin A stimulation (human), **140**, 1
and phosphoinositide turnover, independent activation in Ca^{2+} -depleted phorbol ester-primed neutrophils by formyl-methionyl-leucyl-phenylalanine (human), **135**, 785
and protein kinase C, coactivation in plasma membrane (bovine neutrophils), **134**, 305
thyroidal, solubilization and characterization, **141**, 839
- Nalidixic acid
effect on expression of related promoter genes (*Escherichia coli*), **135**, 411
inhibition of EDTA-induced autolysis (*Escherichia coli*), **134**, 338
- Nalorphine
conformation at monoclonal antibody binding sites, ^1H transferred nuclear Overhauser effect measurements, **141**, 1267
- β -Naphthoflavone
enhancement of cardiac prostaglandin release *in vitro* (chick embryo), **136**, 582

Subject Index

- induction of cytochrome *P*-450c in hepatocytes and hepatic hematopoietic cells (fetal rat), **141**, 299
- Neomycin
effect on herpes simplex virus infection (BHK cells), **141**, 198
- Neopterin
transient accumulation in human T cells and various cell lines, induction by phorbol ester, **141**, 494
- Nerve growth factor
binding to high-affinity type-1 receptors, inhibition by phenylmethylsulfonyl fluoride (chicken), **134**, 1000
calcium-dependent activation of glycogen phosphorylase in pheochromocytoma PC12 cells (rat), **138**, 917
effect on protein kinase C substrate B-50 levels in pheochromocytoma PC12 cells (rat), **139**, 644
gene cloning: mRNA distribution in developing and adult tissues (chicken), **141**, 1116
mRNA, localization by *in situ* hybridization (murine brain), **138**, 813
synthesis and secretion by cultured astroglial cells (mouse), **136**, 57
- Nerve growth factor receptors
in melanoma cell chromatin, identification with monoclonal antibodies (human), **140**, 174
- Nerves
optic, unilateral crush, subsequent expression of proteins in retina (goldfish), **138**, 1177
- Nervous system
cyclohexadenosine receptor phylogenetic distribution, **137**, 182
- Neurocyprenin
apoforn, inhibition of dopamine β -monooxygenase (bovine brain), **138**, 1
- Neurohypophysis
-hypothalamus complex in culture, arginine vasopressin release, effect of synthetic atrial natriuretic factor (rat), **134**, 652
- Neuromedin U
-like immunoreactivity, characterization in tissue extracts (guinea pig, human, rat, swine), **140**, 1127
- Neurons
cortical, somatomedin receptors, occurrence and interactions with insulin and insulin-like growth factors (mouse), **135**, 228
spinal lower motor, excitation, relationship to thyrotropin-releasing hormone receptor binding, hypotheses, **138**, 1184
- Neuropeptide Y
homologous peptide, amino acid sequence analysis (coho salmon endocrine pancreas), **141**, 1084
- Neurophysin
-pro-octocin, cleavage at paired basic residues by endopeptidase associated with neurohypophyseal secretory granules (bovine), **134**, 1190
- Neurotensin
amino acid sequence, homology with kinetensin (human), **136**, 983
- Neutral endopeptidase, *see* Membrane metallo-endopeptidase
- Neutrophils
activated
cytosolic protein kinase C translocation to plasma membrane, mechanism (human), **136**, 228
oxidation of lysine side chains in chicken embryonic elastin (human), **135**, 451
activation, effects of protein kinase C inhibitors H-7 and H-9 (human), **135**, 749
 Ca^{2+} -depleted phorbol ester-primed, phosphoinositide turnover and NADPH oxidase, independent activation by formyl-methionyl-leucyl-phenylalanine (human), **135**, 785
chemotactic factor, generation by ethanol-metabolizing hepatocytes (human), **137**, 688
chemotaxis
formyl-peptide-induced, inhibition by myelin basic protein (human), **141**, 524
leukotriene B₄-induced, inhibition by SM-9064 (rat), **138**, 540
formyl-methionyl-leucyl-phenylalanine-induced inositol phospholipid metabolism and Ca^{2+} mobilization, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
12-hydroperoxyeicosatetraenoic acid-stimulated, leukotriene B₄ synthesis, requirement of free arachidonic acid (human), **138**, 589
ionophore A23187-stimulated, arachidonic acid release and leukotriene synthesis, enhancement by phorbol ester (human), **141**, 399
leukotriene B₄ production, comparison with A23187- or zymosan-induced production in whole blood (human), **137**, 334
leukotriene C₄ synthesis, selective inhibition by ethacrynic acid (human), **137**, 195
metabolism of 6-*trans*-leukotriene B₄, pathways (human), **136**, 707
NADPH oxidase activation and phosphorylation of membrane proteins, coordinate inhibition by surface antigen-directed monoclonal antibody (human), **140**, 1121
oxidation of chemoattractants, effect on efficacy (human), **138**, 66
oxidative metabolism, enhancement by interferon γ (human), **138**, 1276
phagocytic degranulation, stimulation by aggregated IgG, requirement for arachidonic acid lipooxygenation (human), **136**, 310
platelet-activating factor binding to membranes, inhibition by GTP (human), **141**, 353
production of superoxide anions, induction by tumor necrosis factor (human), **137**, 1094
protein kinase C and phorbol myristate acetate receptor, stimulus-dependent movement from cytosol to membranes (human), **134**, 587
respiratory burst, phosphoinositide hydrolysis, and Ca^{2+} mobilization induced by formyl-methionyl-leucyl-phenylalanine, phorbol ester effects (human), **135**, 556
responses to chemoattractants, phorbol ester-mediated inhibition, reversal by protein kinase inhibitor H7 (rabbit), **137**, 50
stimulated with chemotactic peptide formyl-methionyl-leucyl-phenylalanine, protein kinase C translocation, features (guinea pig), **139**, 1169
stimulation by chemotactic peptides, associated changes in membrane fluidity (human), **141**, 1151
superoxide production, induction by 1-oleoyl-2-acetylgllycerol

Subject Index

- enhancement by extracellular Ca^{2+} and cytochalasin B (human), **134**, 690
- after incorporation into pH-sensitive liposomes (guinea pig), **136**, 242
- superoxide radical-generating protein, photolabeling with arylazido NADP^+ analog (bovine blood), **139**, 85
- synthesis of 9,10-epoxy-12-octadecenoate (dog, guinea pig, human), **137**, 424
- Nicotine
- induction of damage in pancreatic acinar membranes, role of calcium (rat), **139**, 530
- Nitrendipine receptors
- affinity resin preparation with carboxy-nifedipine, **134**, 922
- detection in nonneuronal cells derived from cultured fetal spinal cord (mouse), **136**, 783
- Nitric oxide
- geminate recombination to myoglobin, analysis on picosecond time scale (elephant, whale), **140**, 962
- Nitrile hydratase
- purification and characterization (*Brevibacterium* R312), **139**, 1305
- 4-Nitrobenzyl mercaptan S-sulfate
- covalent binding to cytosolic proteins, inhibition by glutathione (rat liver), **137**, 1055
- Nitrogenase
- iron-molybdenum cofactor, solubilization in dimethylformamide and acetonitrile (*Azotobacter vinelandii*), **139**, 740
- repression by ammonia (*Anabaena cylindrica*), **134**, 835
- Nitrogen fixation
- inhibition by proline (*Anabaena* 7120), **139**, 940
- p-Nitrophenylphosphate
- hydrolysis
- by μ -opioid receptors (bovine brain), **140**, 660
- quantitation in homogenates of temporal regions of Alzheimer's brain (human), **139**, 315
- p-Nitrophenylphosphorylcholine
- hydrolysis, quantitation in homogenates of temporal regions of Alzheimer's brain (human), **139**, 315
- 4-Nitroquinoline 1-oxide
- adducts, lack of excision from DNA by cell extracts capable of excising pyrimidine dimers, **140**, 775
- induction of 8-hydroxyguanine residue formation in cellular DNA after exposure (Ehrlich ascites cells), **139**, 626
- Noradrenaline, *see* Norepinephrine
- Nordihydroguaiaretic acid
- inhibition of agonist evoked aggregation, ATP secretion, and cytosolic Ca^{2+} increase in aspirinated platelets (human), **139**, 509
- Norepinephrine
- brain levels, effect of sustained exercise (rat), **137**, 149
- effect on
- adenylate cyclase activity and [^3H]clonidine specific binding in brown adipose tissue (fetal rat), **138**, 1390
- intracellular Ca^{2+} levels and contractile activity in isolated perfused heart (rabbit), **139**, 816
- induction of
- increase in cytosolic Ca^{2+} levels in cultured aortic smooth muscle cells (rat), **135**, 9
- iodothyronine 5'-deiodinase activity in brown adipose tissue, reduction during pregnancy and lactation (rat), **138**, 1315
- release by cultured chromaffin cells, inhibition by potassium (bovine adrenal gland), **134**, 1
- in smooth muscle cell monolayers containing fluorescent probes, effect on cytosolic Ca^{2+} levels (hamster, rabbit, rat), **136**, 927
- Novobiocin
- inhibition of EDTA-induced autolysis (*Escherichia coli*), **134**, 338
- NTPase, *see* Nucleoside-triphosphatase
- Nuclear magnetic resonance
- ^{13}C
- bicarbonate exchange kinetics across erythrocyte membranes at equilibrium (human), **136**, 266
- and ^{15}N , characterization of peptidoglycan stem lengths (*Aerococcus viridans*), **137**, 736
- solid-state, dermatan sulfate L-iduronate ring conformation (porcine skin), **137**, 87
- deuterium, in monitoring metabolic processes *in vivo*, evaluation (mouse), **137**, 579
- ^1H
- assessment of intact lens hexose monophosphate shunt activity (rabbit), **138**, 1068
- detection of sulfmyoglobin multiple forms, **135**, 309
- mastoparan binding by calmodulin, **137**, 1069
- transferred nuclear Overhauser effect measurements of flexible opiate conformations at monoclonal antibody binding sites, **141**, 1267
- ^{23}Na , spin-lattice relaxation times in various tissues, modification after carbon tetrachloride intoxication (rat), **134**, 1079
- ^{31}P
- cellular ATP levels after photodynamic therapy (murine mammary tumor), **140**, 273
- depth-selected EKG-gated, *in vivo* analysis of rhythmic changes in myocardial phosphate metabolite levels during cardiac cycle (rat), **135**, 808
- measurement of cytoplasmic pH with phenylphosphonate as probe (*Escherichia coli*), **138**, 1106
- observation of ATP-Mg^{2+} and uncomplexed ATP in slow exchange, **134**, 1379
- Rous or Kirsten sarcoma virus-transformed fibroblasts (chicken, rat), **138**, 699
- peptide conformation analyses, computer applications, **134**, 907
- two-dimensional, observation of loop structure in d-GAATTCGCCGAATTC, **139**, 1224
- Nuclear matrix
- association of
- acute phase protein genes during inflammation (rat liver), **137**, 633
- glucocorticoid receptor complexes, evaluation (rat liver), **137**, 640
- relationship to androgen-dependent genes, basis for prediction of transcriptional activity (rat prostate), **138**, 1334
- Nuclear Overhauser effect
- ^1H transferred, flexible opiate conformations at monoclonal antibody binding sites, **141**, 1267
- Nuclease
- DNA cleavage rates at different sites, calculation from partial digestion of end-labeled molecule, method reevaluation, **138**, 778
- Nuclei
- γ -interferon transport into, role of nuclear membrane recep-

Subject Index

- tors (mouse), **138**, 254
- membrane structure and function, effect of dietary lipid content (murine liver), **135**, 655
- neuronal, topoisomerase II and DNA polymerase α levels, independent regulation (rat), **138**, 1116
- Nucleosides
- sodium-dependent concentrative transport in cultured intestinal epithelial cells (rat), **140**, 1028
- Nucleoside-triphosphatase
- nuclear membrane-associated, effect of dietary lipid content (murine liver), **135**, 655
- Nucleosomes
- c-fos*, relaxation during differentiation (HL-60 cells), **141**, 213
 - c-myc*, reassembly during differentiation (HL-60 cells), **141**, 213
 - preferential placement on plasmid pBR322 DNA restriction fragments (*Escherichia coli*), **138**, 110
- 5'-Nucleotidase
- activation by diadenosine tetraphosphate (*Artemia* embryo), **138**, 261
- Nucleotides
- affinity for active sites of detergent-solubilized Ca^{2+} -ATPase from sarcoplasmic reticulum (rabbit muscle), **138**, 652
 - binding to ADP/ATP carrier, analysis with 6'-*O*-dansyl- γ -aminobutyryl atractyloside (bovine cardiac mitochondria), **134**, 266
 - exchange from actin, effect of erythro-9-[3-(2-hydroxynonyl)]adenine, **137**, 1181
- Nucleotide sequence
- atrial natriuretic factor precursor gene (bovine), **136**, 396
 - C1 inhibitor-encoding cDNA (human), **137**, 620
 - cDNA to α_{2u} -globulin mRNA induced by phenobarbital (rat liver), **134**, 1182
 - cDNA partial clone for heparin cofactor II (human), **137**, 431
 - creatine kinase B mRNA: coding and 3' noncoding region homology among species (canine brain), **138**, 392
 - creatine kinase M subunit cDNA (human), **140**, 981
 - cytochrome *P*-450-like pseudogene (rabbit liver), **134**, 499
 - galactosyltransferase cDNA (human milk), **139**, 163
 - gene encoding U3 RNA, homology with vertebrate U3 RNA (*Drosophila melanogaster*), **138**, 512
 - glutathione *S*-transferase H_a subunit I cDNA (human liver), **141**, 229
 - hepatitis B virus subtype *adw* pre-S2 region, comparison with other subtypes, **135**, 382
 - insulin and insulin receptor mRNAs, occurrence of short complementary sequences, **139**, 94
 - interferon- α gene: homology with enhancer elements in gene regulatory region (mouse), **138**, 826
 - interleukin-2 cDNA and homologous genomic DNA fragments (human placenta), **138**, 966
 - lipase gene (*Pseudomonas fragi*), **141**, 185
 - mitochondrial ADP/ATP-carrier protein mRNA (bovine heart), **138**, 850
 - mRNA of DNA polymerase β in phage λ gt11 via cDNA cloning (human), **136**, 341
 - postmeiotically expressed cDNA clone encoding lactate dehydrogenase (mouse), **136**, 760
 - reovirus serotype 1 Lang strain mRNA encoding major capsid surface polypeptide $\sigma 3$, **136**, 183
 - minor capsid polypeptide $\sigma 1a$ and nonstructural polypeptide $\sigma 1bNS$, **140**, 508
 - nonstructural σNS polypeptide, **139**, 845
 - ribosomal gene spacer, polymorphisms (*Triploneustes gratilla*), **137**, 834
 - RNA of 14-kDa β -galactoside-binding lectin (chicken embryo), **134**, 51
 - rRNA, 5 S (*Thermococcus celer*), **134**, 1167
 - sphingolipid activator protein cDNA (human), **134**, 989
 - substance P precursor-encoding mRNA (rat), **139**, 1040
 - 3' terminus of 16 S RNA (*Streptomyces aureofaciens*), **137**, 80
 - U3 small nucleolar RNA gene (human), **137**, 1133
- O**
- Obtusifolius
- 14 α -demethylation by cytochrome *P*-450 monooxygenase (corn microsomes), **140**, 1064
- Octadeoxynucleotides
- d(A-T)₄ and d(T-A)₄ forms, different behavior at high concentrations of cesium fluoride, **139**, 1158
- Octyl glucoside
- induction of high Mg^{2+} -ATPase activity of coupling factor 1 after preactivation with heat and dithiothreitol (spinach chloroplasts), **141**, 636
- Oestradiol-17 β , *see* Estradiol
- 1-Oleoyl-2-acetylgllycerol
- differential effects on carbamylcholine- and high K^{+} -induced elevation of intracellular free Ca^{2+} levels in cultured chromaffin cells (bovine adrenal), **139**, 903
 - incorporated into pH-sensitive liposomes, stimulation of superoxide release from neutrophils (guinea pig), **136**, 242
 - induction of
 - sodium-hydrogen exchange (human platelets), **141**, 13
 - superoxide production, enhancement by extracellular Ca^{2+} and cytochalasin B (human neutrophils), **134**, 690
 - stimulation of phosphate accumulation in cultured renal tubular cells, synergistic effect of A23187 (mouse), **136**, 177
- 2',5'-Oligoadenylate
- binding protein, and 2-5A synthetase, induction by interferon, age-related differences (guinea pig macrophages), **140**, 455
- 2',5'-Oligoadenylate synthetase
- and 2-5A binding protein, induction by interferon, age-related differences (guinea pig macrophages), **140**, 455
 - accumulation in spleen, liver, kidney, and brain, induction by misaligned dsRNA (mouse), **138**, 40
 - induction by multiple molecular forms of interferon, comparison (MDBK cells), **141**, 1042
- Oligonucleotides
- degradation by 2',5'-phosphodiesterase: role of 3'-hydroxyl moiety in penultimate position of substrate, **141**, 555
 - d-GAATTCGCCGAATTC, loop structure, observation by two-dimensional NMR, **139**, 1224
 - synthetic, as probes for identification of cDNA clones for ligninase

Subject Index

- (*Phanerochaete chrysosporium*), **137**, 649
 screening recombinant DNA junctions, **138**, 945
 Tp(AP)pT, synthesis, **134**, 1204
- Oligopeptides
 intestinal transport in intact form (rabbit), **137**, 682
- Oligosaccharides
 -ceramide linkage in glycosphingolipids, identification and characterization of cleaving enzyme (leech), **141**, 346
 chains, role in glucocerebrosidase routing to lysosomes (U937 cells), **141**, 452
 lipid-linked, acetylated and pyruvylated, synthesis *in vitro* (*Rhizobium trifolii*), **134**, 205
 N-linked, endo-glycosidase F- and peptide:N-glycosidase F-mediated release from mammalian cells, applications, **140**, 967
 sequences, position of O-acetylated residues, mass spectrometric determination, **135**, 1126
 triantennary complex-type, precipitation of D-galactose-specific lectin (*Erythrina indica*), **141**, 963
- Oocytes
 expression of *Agrobacterium tumefaciens* T-DNA gene 7 (*Xenopus laevis*), **141**, 1242
- Opioid receptors
 binding of
 morphine, inhibition by anti-paratypic antibody fraction (rabbit), **136**, 1177
 [D-Pro¹⁰]dynorphin(1-11) mono- and diiodinated derivatives, comparison, **134**, 1142
 type μ , associated phosphatase activity (bovine brain), **140**, 660
 types μ and δ , selective binding of Tyr¹-substituted fluorescent Pya¹-enkephalins (rat brain), **136**, 1170
- Opsin
 incorporation of *myo*-inositol *in vivo* (frog), **136**, 815
- Ornithine carbamoyltransferase
 precursor produced in *Escherichia coli*, uptake and processing *in vitro* by rat cardiac and hepatic mitochondria, **134**, 21
- Ornithine carbamyl transferase, *see* Ornithine carbamoyltransferase
- Ornithine decarboxylase
 activation in 3T3 fibroblasts, relationship to phorbol ester-induced prostaglandin biosynthesis (BALB/c mouse), **139**, 186
 -antizyme complex, interaction with α -difluoromethylornithine, analysis (rat liver), **137**, 1101
 induction by glucocorticoid *in vivo*, enhancement by diacylglycerol (adrenalectomized rat), **138**, 275
 mRNA levels in pulmonary tumor cells, relationship to sensitivity to difluoromethylornithine (human), **134**, 572
- Osmolality
 effect on angiotensin-stimulated aldosterone production (bovine adrenal glomerulosa cells), **139**, 46
- Osteoblasts
 -like MC3T3-E1 cells
 calcification, inhibition by β_2 -microglobulin (mouse), **141**, 360
 production of macrophage colony-stimulating activity and prostaglandin E₂, stimulation by recombinant human interleukin-1 α and β (mouse), **141**, 285
 synthesis of colony-stimulating and differentiation-inducing factors (mouse), **134**, 400
- Ouabain
 displacement from binding sites by enterolactone (guinea pig, human heart), **134**, 1064
 effect on
 basal and stimulated α -aminoisobutyric acid transport by Na⁺,K⁺-ATPase (rat skeletal muscle), **134**, 1342
 metabolic cooperation between normal, preneoplastic, or neoplastic mammary cells and drug-resistant mammary tumor cells *in vitro* (mouse), **140**, 955
 -like substance
 detection in plasma of patients with primary aldosteronism, **135**, 41
 purification and characterization (bovine plasma), **135**, 1
 sensitivity, induction by Kirsten murine sarcoma virus-mediated transformation (human HOS cells), **136**, 807
- Ovalbumin
 gene, specific binding of estrogen receptor (chicken), **139**, 396
- Oviduct
 isthmus, 5 S RNA accumulation and specific cleavage (laying hen), **138**, 1405
- N-Oxalylcysteine
 quantitative analysis in various tissues (rat), **141**, 1051
- Oxalyl thioesters
 inhibition of phosphorylase phosphatase catalytic subunit, role in insulin action mechanism, **134**, 93
 quantitative analysis in various tissues (rat), **141**, 1051
- Oxidase
 superoxide radical-generating, photolabeling with arylazido NADP⁺ analog (bovine neutrophils), **139**, 85
- Oxidation
 biomimetic, lignin model compounds by simple inorganic complexes, analysis, **139**, 1104
 chemoattractant, by responding cells, effect on efficacy (human neutrophils), **138**, 66
 cyanide-resistant, NADH and succinate in submitochondrial particles, comparison with duroquinol oxidase activity (*Arum maculatum*), **135**, 677
 iron chromophore of purple acid phosphatase during interaction with phosphate (bovine spleen), **136**, 490
 lysine side chains in elastin by canine myeloperoxidase and stimulated human neutrophils (chicken embryonic aorta), **135**, 451
 NADH, stimulation by ferricyanide or transferrin (HeLa cells), **135**, 110
 NAD⁺-linked substrates in mitochondria, inhibition by 1-methyl-4-phenylpyridine, relationship to uptake (rat liver), **134**, 743
 ozone-induced, bovine erythrocyte membranes and linolenic acid, uric acid protective effects, **137**, 536
 polysaccharides by H₂O₂, role in cell wall breakdown (larch, orange), **141**, 238
 selective, imidazole ring in histidine residues by ascorbic acid-copper ion system, **138**, 659
- 2,3-Oxidosqualene cycloartenol
 inhibition by *N*-[(1,5,9)-trimethyl-decyl]-4 α ,10-dimethyl-8-aza-*trans*-decal-3 β -ol, **138**, 764
- 19-Oxoandrostenedione
 cytochrome P-450_{11 β} -catalyzed conversion to estrone, **140**, 530
 cytochrome P-450_{11 β} -dependent formation from 19-hy-

Subject Index

- droxyandrostenedione (bovine), **136**, 369
- Oxotremorine
effect on atrial natriuretic factor release by atrial slices (rat), **135**, 1090
- Oxygen
active, detection with lucigenin (rat hepatocytes), **140**, 468
consumption by monocytes, effect of κ elastin (human), **141**, 92
 ^{17}O , hyperfine structure for hydroxyl and superoxide radical adducts of spin traps, **141**, 622
peroxidase-dependent transfer from hydrogen peroxide to thioanisoles, **135**, 166
singlet
inactivation of firefly luciferase, **140**, 888
production in hydrogen peroxide-hypochlorous acid reaction, enhancement by 1,4-diazabicyclo[2.2.2]octane, **134**, 777
- Oxygen radicals
free, stimulation of tyrosine-specific protein phosphorylation in plasma membrane (rat liver), **139**, 439
reduction at epidermal surface, role of thioredoxin reductase (guinea pig, human), **136**, 630
- Ozone
induced oxidation of bovine erythrocyte membranes and linolenic acid, protective effects of uric acid, **137**, 536
- P**
- Pancreas
insulin secretion
lysophosphatidylinositol effects (rat), **138**, 720
polymyxin B effects (rat), **136**, 1001
neonatal, high peptide α -amidating activity (rat), **138**, 179
secretory responses, effect of chloroquine binding to muscarinic receptors (rat), **137**, 664
- Papain
domain boundaries, prediction from amino acid sequence, **139**, 11
- Paraquat
and 1-methyl-4-phenylpyridine, cytotoxic effects on isolated hepatocytes, comparison (rat), **137**, 303
- Parathyroid hormone
C-terminal peptide, detection after biliary ligation: implications for diagnosis of hepatobiliary disease (rat), **139**, 633
fragment 1-84, degradation in rat osteoblast-like osteosarcoma cells (human), **141**, 762
secretion by parathyroid cells
electroporabilized, effects of phorbol ester, cAMP, and changes in free calcium (bovine), **139**, 1233
regulation by calcium, effect of pertussis toxin treatment (bovine), **138**, 960
- Parkinsonism
experimental and idiopathic, relationship to mitochondrial NADH dehydrogenase inhibition by pyridine derivatives, **135**, 269
- Pectin
oxidation by hydrogen peroxide, role in cell wall breakdown (orange), **141**, 238
- Penicillamine selenotrisulfides
isolation, characterization, and thiol exchange reaction, **135**, 183
- Pentachlorophenol
catabolism by *Flavobacterium* sp., analysis, **141**, 825
- Pentafluorobenzyl oxime
derivatives, in determination of 4-hydroxyalkenal tissue levels by gas chromatography-mass spectrometry (rat), **139**, 144
- Pentagastrin
binding in gastric glands, characterization (guinea pig), **134**, 895
- Pentapeptides
content in elastin, measurement for protein quantitation (swine), **136**, 672
- Pentobarbital
effect on glutathione transferase expression in seminal vesicles and pituitary gland (rat), **136**, 1057
inhibition of adenylate cyclase, relationship to functional coupling of guanine nucleotide-binding stimulatory protein and catalytic unit (rat brain), **140**, 237
- Pepsinogen
carbachol-induced secretion from isolated gastric glands, effects of pirenzepine and atropine, comparison (guinea pig), **136**, 193
secretion from gastric chief cells, effects of A23187 and ionomycin (guinea pig), **137**, 281
type A, chromosomal genes, close linkage (human), **138**, 289
- Peptide B
immunoreactive, distribution in brain (rat), **139**, 1024
- Peptide:N-glycosidase F
deglycosylation of thyroid-stimulating hormone (bovine), **138**, 230
and endo-glycosidase F, release of N-linked oligosaccharides from mammalian cells, applications, **140**, 967
- Peptides
 α -amidation
analysis in neonatal pancreas (rat), **138**, 179
purification and characterization of catalyzing enzyme (*Xenopus laevis* skin), **137**, 984
tissue distribution and characterization of catalyzing enzyme (rat), **140**, 230
 α -amylase, signal and promoter, from *Bacillus subtilis*, secretion activities in *Escherichia coli*, **134**, 624
arginine-containing, vasodilating properties, dependence on endothelium (rat aorta), **141**, 446
associated with apolipoprotein B-100 surface-exposed segments in low-density lipoprotein particles, isolation and amino acid sequence (human), **140**, 250
atrial natriuretic factor (99-126 and 101-126), comparative biological activities (bovine, chicken, rabbit, rat), **135**, 987
atrial natriuretic factor-like, detection in parasympathetic ganglia (rat), **134**, 279
bombesin-related, specific binding and mitogenic effects, inhibition by substance P antagonist (Swiss 3T3 cells), **137**, 135
calcitonin gene-related
partial homology with N-terminal sequence of amyloid fibril protein from insulin-producing pancreatic tumor (human), **140**, 827
structure-activity relationship (human), **134**, 1306
captopril analogs, inhibition of captopril binding to angiotensin-converting enzyme (rat lung), **135**, 770

Subject Index

- chemotactic, stimulation of neutrophils, associated changes in membrane fluidity (human), **141**, 1151
- conformations constrained by NMR data, computer-aided search, **134**, 907
- crystalline delta endotoxin, 26-kDa, role in larvicidal and hemolytic activities (*Bacillus thuringiensis* var. *israelensis*), **136**, 1142
- C-terminal, parathyroid hormone, detection after biliary ligation: implications for diagnosis of hepatobiliary disease (rat), **139**, 633
- derived from VIP precursor protein, characterization (cat, guinea pig, human, rat, swine), **139**, 1142
- diphtheria toxin B fragment, pH-dependent insertion into lipid membrane, conformational analysis, **136**, 160
- hemolytic, 28-kDa, removal from *Bacillus thuringiensis* var. *israelensis* solubilized crystal protein, effect on mosquito toxicity, **141**, 937
- high-sensitivity mass spectrometric assay: direct analysis of aqueous solutions, **141**, 1058
- homologous to neuropeptide Y, amino acid sequence analysis (coho salmon endocrine pancreas), **141**, 1084
- hydroxamic acids, inhibition of skin collagenase (human), **136**, 390
- ionophorous, Ca^{2+} -transporting, detection in lymphocytes by quin-2 fluorescence (dog, human), **134**, 731
- myelin basic protein synthetic analogs, protein kinase C inhibition (bovine), **134**, 78
- nascent, in gastric mucus glycoprotein, cotranslational attachment of fatty acids (rat), **141**, 387
- opioid, putative processing activity in Golgi-enriched fraction from brain (rat), **138**, 356
- from pancreatic juice, growth stimulating activity on 3T3 fibroblasts (rat), **139**, 545
- pro-opiomelanocortin-derived, secretion from anterior and intermediate pituitary cells *in vitro*, suppression by atrial natriuretic factor (rat), **135**, 1035
- pseudo-, cyclic and linear enkephalin analogs, synthesis and biological activity *in vitro*, **136**, 730
- stimulating pancreatic enzyme secretion, intestinal administration, elevation of plasma cholecystokinin levels, radioimmunoassay (rat), **136**, 701
- synthetic
- containing D- or L-threonine residues, in analysis of protein kinase C substrate specificity (rat brain), **134**, 1358
 - corresponding to 17- and 28-residue sequences of Alzheimer's disease β -protein, *in vitro* formation of amyloid fibrils (human), **141**, 782
 - pre-S hepatitis B virus-murabutide-toxoid carrier conjugates, in production of antibodies recognizing hepatitis B virus surface antigens, **140**, 557
 - tyrosine phosphorylation by casein kinase A (human erythrocytes), **139**, 855
- Peptidoglycan receptors
- and β -1,3-glucan receptors, identification in phenoloxidase-activating system (silkworm), **141**, 1177
- Peptidoglycans
- peptide chain length, characterization by solid-state ^{13}C and ^{15}N NMR (*Aerococcus viridans*), **137**, 736
- Peptidyl dipeptidase
- angiotensin I conversion to type II (bovine endothelial cells), **134**, 770
- Perfringolysin O
- activation of phosphatidic acid metabolism in erythrocyte membranes (human), **137**, 23
- Peroxidase
- associated activity of polyethylene glycol-modified hemin in organic solvents, **138**, 283
 - catalysis of hydrogen peroxide metabolism and irreversible inactivation, role of superoxide radicals, **139**, 327
 - dependence of oxygen transfer from hydrogen peroxide to thioanisoles, **135**, 166
- Peroxidation
- etoposide and teniposide: free radical formation, **135**, 215
- lipids
- effects of anthrapyrazoles and aminoacridine derivative (rabbit hepatic microsomes), **140**, 797
 - enhancement by polyriboinosinic acid-polyribocytidylic acid treatment (mouse, rat), **134**, 632
 - formation of hydrogen atoms and hydroxyl radicals, ESR spectroscopy, **141**, 381
 - in hepatic microsomes, hepatocytes, and granulosa cells, relationship to aldrin epoxidation (rat), **138**, 24
 - inhibition by anisylidithiolthione (rat hepatic microsomes), **135**, 1015
- Peroxisomes
- conjugation of cholic acid with taurine and glycine (rat liver), **138**, 167
 - matrix, localization of coenzyme A (rat liver), **139**, 1195
 - membrane polypeptides, synthesis *in vitro* (rat liver), **137**, 626
 - protein phosphorylation, analysis (rat liver), **140**, 188
- Peroxydisulfate
- copper system, biomimetic oxidation of lignin model compounds, **139**, 1104
- Pertussis toxin
- ADP-ribosylated guanine nucleotide-binding protein, detection in spermatozoa (invertebrates, vertebrates), **138**, 728; *addendum*, **140**, 773
 - catalyzed ADP-ribosylation, substrate identification on cardiac cell membranes (rat), **138**, 1355
 - at different concentrations, effects on agonist-induced prostaglandin E_2 synthesis (rat mesangial cells), **141**, 39
 - effect on
 - α -adrenergic-stimulated responses in brown adipocytes (hamster), **135**, 823
 - calcium regulation of parathyroid hormone secretion (bovine parathyroid cells), **138**, 960 - inhibition of Ca^{2+} transients in cultured smooth muscle cells (rat aorta), **140**, 195
 - sensitive G proteins, role in α_1 -adrenergic and thyrotropin receptor-mediated activation of membrane phospholipases and iodide fluxes, analysis (FRTL-5 cells), **141**, 1000
 - sensitive N protein, mediation of chemoattractant receptor signal transduction (human), **138**, 887
- Pesticides
- phototoxicity, evaluation using *Paramecium aurelia*, **139**, 79
- pH
- cytoplasmic, measurements by ^{31}P NMR with phenylphosphonate as probe (*Escherichia coli*), **138**, 1106
 - dependence of K^+/H^+ antiporter, desensitization by N-ethylmaleimide (*Vibrio alginolyticus*), **136**, 1030

Subject Index

- effect on diphtheria toxin B fragment insertion into lipid membrane, conformational analysis, **136**, 160
- transmembrane equilibration rate in isolated sarcoplasmic reticulum vesicles, measurement (rabbit skeletal muscle), **140**, 934
- Phanerochaete chrysosporium*
- particulate fractions, H₂O₂-dependent decolorization of poly R-481, **136**, 220
- Phenamil
- interaction with amiloride-sensitive Na⁺ channels, characterization (porcine renal membranes), **135**, 25
- Phenobarbital
- effect on
- epidermal growth factor and insulin receptor expression after chronic treatment (rat liver), **135**, 501
- glutathione transferase expression in seminal vesicles and pituitary gland (rat), **136**, 1057
- induction of
- 2-carboxybenzaldehyde reductase (rat liver), **141**, 488
- α_{2u} -globulin mRNA: characterization of cDNA clone (rat liver), **134**, 1182
- polycyclic hydrocarbon-inducible cytochrome P-450 (rat hepatoma cells), **137**, 120
- Phenol red, *see* Phenolsulfonphthalein
- Phenolsulfonphthalein
- estrogenic activity in cultured anterior pituitary cells (rat), **141**, 885
- Phentolamine
- effect on
- β -adrenergic ligand binding to pineal particulate fractions (rabbit), **138**, 1196; *erratum*, **139**, 1320
- intracellular Ca²⁺ levels and contractile activity in isolated perfused heart (rabbit), **139**, 816
- Phenylalanine
- glycyl derivative, effect on IgG binding to bacterial Fc receptors, **135**, 1105
- and 7-methylguanosine and 5'-monophosphate, crystal structure, interaction model, **136**, 294
- Phenylalanine ammonia-lyase
- entrapment in silk fibroin for protection from proteolysis (*Rhodotorula glutinis*), **141**, 165
- Phenylalanine-tRNA ligase
- subunits, purification of cytoplasmic precursors (yeast mitochondria), **137**, 1119
- Phenylalanyl-tRNA synthetase, *see* Phenylalanine-tRNA ligase
- 2-Phenyl-1-2-benzisoxselenazol-3(2H)-one
- prevention of doxorubicin-induced killing of breast cancer cells (human), **135**, 330
- Phenyl-n-butylborinic acid
- as probe for mechanism and inhibition of cholesterol esterase- and lipoprotein lipase-catalyzed hydrolyses, **134**, 386
- 1-Phenylcyclopropylamine
- monoamine oxidase inactivation: identification of cysteine residue involved in labile adduct formation, **135**, 154
- Phenylephrine
- effect on
- atrial natriuretic factor release by atrial slices (rat), **135**, 1090
- cardiac and plasma atrial natriuretic factor levels after chronic administration (rat), **136**, 510
- cytoplasmic free Ca²⁺ levels, comparison with angiotensin II and vasopressin (rat hepatocytes), **140**, 107
- phosphoinositide metabolism (rat muscle), **136**, 863
- enhancement of β -adrenergic ligand binding to pineal particulate fractions (rabbit), **138**, 1196; *erratum*, **139**, 1320
- Phenylhydrazine
- induction of L-alanine:4,5-dioxovalerate transaminase, role of heme level (rat kidney, liver), **138**, 742
- Phenylmethylsulfonyl fluoride
- inhibition of nerve growth factor binding to high-affinity type-1 receptors (chicken), **134**, 1000
- Phenylphosphonate
- as probe for cytoplasmic pH measurements by ³¹P NMR (*Escherichia coli*), **138**, 1106
- Phlebotomy
- induced changes in hemoglobin proportions, effect of acetylsalicylic acid (adult rat), **137**, 69
- Phorbol
- effect on
- carbachol-induced inositol 1-phosphate accumulation (NCB-20 cells), **136**, 622
- phosphoprotein B-50 phosphorylation in native synaptic plasma membranes (rat brain), **136**, 1007
- Phorbol 12,13-dibutyrate
- effect on
- α_1 -adrenergic receptor-stimulated phosphoinositide hydrolysis and aortic contraction (rat), **134**, 970
- carbachol-induced inositol 1-phosphate accumulation (NCB-20 cells), **136**, 622
- phosphoprotein B-50 phosphorylation in native synaptic plasma membranes (rat brain), **136**, 1007
- phosphorylation of 80K protein (3T3 cells), **140**, 379
- induced contraction of thoracic aorta, mechanism (rabbit), **138**, 1362
- Phorbol 12,13-dibutyrate receptors
- affinity states, effect of ionophore A23187 (human B cells), **138**, 554
- Phorbol 12,13-didecanoate
- effect on
- chondroitin sulfate sulfation (human promyelocytic leukemia cells), **135**, 909
- muscarinic receptor binding and polyphosphoinositide breakdown (human neuroblastoma cells), **140**, 160
- stimulation of phosphate accumulation in cultured renal tubular cells, synergistic effect of A23187 (mouse), **136**, 177
- Phorbol 12-myristate 13-acetate
- with diisopropyl fluorophospho-thrombin, stimulation of DNA synthesis (NIL cells), **141**, 650
- effect on
- α_1 -adrenergic receptor-stimulated phosphoinositide hydrolysis and aortic contraction (rat), **134**, 970
- arachidonic acid release and leukotriene synthesis in A23187-stimulated neutrophils (human), **141**, 399
- basal and stimulated adenylate cyclase activity in 235-1 pituitary cells, **135**, 276
- calcium channel function (PC12 cells), **136**, 1049
- carbachol-induced inositol 1-phosphate accumulation (NCB-20 cells), **136**, 622
- compound 48/80-induced histamine secretion by peritoneal mast cells (rat), **137**, 1112

Subject Index

- formyl-methionyl-leucyl-phenylalanine-induced respiratory burst, phosphoinositide hydrolysis, and Ca^{2+} mobilization (human neutrophils), **135**, 556
- muscarinic receptor binding and polyphosphoinositide breakdown (human neuroblastoma cells), **140**, 160
- protein kinase C activity and cAMP production (rat brain), **139**, 251
- induction of
- protein kinase C translocation from soluble to particulate cell fraction (rat hepatocytes), **138**, 1328
 - sponge cell aggregation without extracellular calcium (*Microciona prolifera*), **134**, 1319
- inhibition of
- neutrophil responses to chemoattractants, reversal by protein kinase inhibitor H7 (rabbit), **137**, 50
 - polyphosphoinositide metabolism and cytosolic Ca^{2+} mobilization, reversal by protein kinase C inhibitor (thrombin-activated human platelets), **134**, 868
 - and insulin, effects on cytosolic protein kinase C (rat adipocytes), **135**, 1119
 - and ionomycin, effects on fatty acid synthesis by isolated hepatocytes (rat), **139**, 403
 - primed Ca^{2+} -depleted neutrophils, phosphoinositide turnover and NADPH oxidase, independent activation by formyl-methionyl-leucyl-phenylalanine (human), **135**, 785
 - stimulation of 2-deoxyglucose uptake by fibroblasts, inhibition by sphingolipids (mouse), **138**, 463; *erratum*, **139**, 852
- Phorbol 12-myristate 13-acetate receptors
- stimulus-dependent movement from cytosol to membranes (human neutrophils), **134**, 587
- Phosphate
- accumulation in cultured renal tubular cells, synergistic effects of phorbol esters and A23187 (mouse), **136**, 177
 - hepatic levels, arsenite-induced changes, ^{31}P NMR study *in vivo* (rat), **139**, 228
 - inorganic
 - inhibition of adenylate cyclase, dependence on III^{glc} protein of phosphoenolpyruvate: glyoxylate phosphotransferase system (*Escherichia coli*), **141**, 1138
 - and sodium, cotransport in renal epithelium, inhibition by phosphonoformic acid (human), **139**, 679
 - interaction with purple acid phosphatase, associated oxidation of iron chromophore (bovine spleen), **136**, 490
 - loss from guanylate cyclase during fertilization, stoichiometry (*Arbacia punctulata* sperm), **137**, 1148
 - metabolites, myocardial levels, rhythmic changes during cardiac cycle, depth-selected EKG-gated ^{31}P NMR analysis *in vivo* (rat), **135**, 808
- Phosphatidic acid
- incorporation of ^{32}P , effect of α_1 -adrenergic receptor stimulation (rabbit, rat aorta), **141**, 1213
 - metabolism in
 - erythrocyte membranes, activation by perfringolysin O (human), **137**, 23
 - neutrophils, formyl-methionyl-leucyl-phenylalanine-induced, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
 - photoreceptor membranes, effect of cationic amphiphilic drugs (bovine), **134**, 378
- Phosphatidylcholine
- formation in dimethyl sulfoxide-treated HL-60 cells, **140**, 515
 - incorporation of ^{32}P , effect of α_1 -adrenergic receptor stimulation (rabbit, rat aorta), **141**, 1213
 - synthesis in plasma membranes, stimulation by insulin and ATP (rat adipocytes), **137**, 1
- Phosphatidylethanolamine
- immunoliposomes, destabilization by antigen binding: rapid assay for virus, **138**, 931
 - transmethylation in HL-60 cells, induction by dimethyl sulfoxide, **140**, 515
- Phosphatidylinositol
- containing liposomes
 - interaction with spermine, **135**, 58
 - opsonized by concanavalin A, stimulation of phosphatidylinositol uptake by macrophages, **138**, 1090
 - dependent activation of DNA polymerase α (murine sarcoma cells), **135**, 880
 - incorporation of ^{32}P , effect of α_1 -adrenergic receptor stimulation (rabbit, rat aorta), **141**, 1213
 - metabolism in
 - B cells, role in antigen-induced activation, **140**, 1079
 - neutrophils, formyl-methionyl-leucyl-phenylalanine-induced, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
 - platelets
 - modification by platelet-derived growth factor (human), **135**, 52
 - role in reversible shape changes (human), **141**, 673
 - phosphorylated forms, detection in suspension-cultured cells, role in growth regulation (*Catharanthus roseus*), **134**, 1175
 - specific phospholipase C, immunologically distinct forms, purification and characterization (bovine brain), **141**, 137
 - turnover in thymocytes, stimulation by concanavalin A, effect on activation (mouse), **141**, 657
- Phosphatidylinositol 4,5-bisphosphate
- containing liposomes, interaction with spermine, **135**, 58
 - formation in skeletal and heart muscle membranes (rabbit), **138**, 1395
 - ^3H -labeled, hydrolysis by phospholipase C, stimulation by GTP and GDP (human platelets), **140**, 728
 - incorporation of ^{32}P , effect of α_1 -adrenergic receptor stimulation (rabbit, rat aorta), **141**, 1213
 - metabolism in
 - neutrophils, formyl-methionyl-leucyl-phenylalanine-induced, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
 - platelets, modification by platelet-derived growth factor (human), **135**, 52
- 1-Phosphatidylinositol-4,5-bisphosphate phosphodiesterase
- stimulation by L-alanine or GTP (catfish olfactory cilia), **137**, 36
- Phosphatidylinositol kinase
- associated activity of $\text{p60}^{\text{v-src}}$ -related tyrosine kinase from acetylcholine receptor-rich membranes (*Narke japonica*), **139**, 473
- Phosphatidylinositol 4-phosphate
- containing liposomes, interaction with spermine, **135**, 58
 - incorporation of ^{32}P , effect of α_1 -adrenergic receptor stimu-

Subject Index

- lation (rabbit, rat aorta), **141**, 1213
- metabolism in
- neutrophils, formyl-methionyl-leucyl-phenylalanine-induced, effects of cAMP-increasing agents and phorbol ester (guinea pig), **139**, 1272
 - platelets, modification by platelet-derived growth factor (human), **135**, 52
- Phosphatidylserine
- activation, role in antipolytic effect of insulin (human adipocytes), **141**, 1157
 - adenylate cyclase system in thrombopathic and normal platelets, regulatory control (dog), **140**, 589
 - cAMP-specific, purification by affinity chromatography on immobilized Rolipram (rat heart), **138**, 205
 - containing liposomes, Ca^{2+} -induced phase separation, potentiation by micromolar levels of Zn^{2+} (bovine brain), **136**, 476
 - localization in erythrocyte outer membrane bilayer (new-born human), **136**, 914
- 2',5'-Phosphodiesterase
- degradation of oligonucleotides: role of 3'-hydroxyl moiety in penultimate position of substrate, **141**, 555
- 6-Phosphofructokinase
- activity in brown and white adipose tissue (rat), **138**, 687
 - assay in brown adipose tissue during aging (rat), **140**, 419
 - cDNA, isolation (human muscle), **135**, 615
 - effects of arabinose- and ribose 1,5-bisphosphate (rat liver), **138**, 159
 - and fructose-bisphosphatase, interaction at physiological concentrations, effect on kinetic behavior (rabbit muscle), **135**, 852
 - genomic clones, isolation and characterization (human liver), **141**, 374
 - phosphorylation by protein kinase C, inhibition by phosphofructokinase stabilizers (rabbit, rat muscle), **136**, 445
 - reversible activation and inactivation by tissue-homologous protein-phosphorylating and -dephosphorylating enzymes (*Ascaris suum*), **139**, 215
- 6-Phosphofructo-2-kinase
- effects of
 - diabetes (rat liver), **136**, 498
 - tolbutamide (rat liver), **139**, 687
- Phosphogluconate dehydrogenase (decarboxylating)
- effects of epidermal growth factor, growth hormone, and thyroxine after chronic administration (hypophysectomized rat), **141**, 78
- 6-Phosphogluconic dehydrogenase, *see* Phosphogluconate dehydrogenase (decarboxylating)
- Phosphoglycerate mutase
- wheat germ, metalloenzymatic nature, **136**, 336
- Phosphoinositides
- α -adrenergic-stimulated breakdown in brown adipocytes, mediatory role of guanyl nucleotide-binding protein, evaluation with pertussis toxin (hamster), **135**, 823
 - covalent linkage to myelin basic protein, *in vitro* effect of allergic encephalomyelitis (rabbit, guinea pig), **136**, 426
 - dependent activation of sarcoma DNA polymerase α , blocking monoclonal antibody, isolation and characterization (mouse), **141**, 60
 - hydrolysis
 - α_1 -adrenergic receptor-stimulated, and aortic contraction, inhibition by phorbol esters (rat), **134**, 970
 - induction by formyl-methionyl-leucyl-phenylalanine, inhibition by phorbol ester (human neutrophils), **135**, 556
 - in isolated platelet membranes, stimulation by GTP and cytosol (human), **137**, 801
 - by platelet-derived growth factor receptor, effect of protein kinase C activation (human skin, murine 3T3 fibroblasts), **137**, 343
 - incorporation of inositol, effects of cationic amphiphilic drugs (rat glioma cells), **135**, 465
- metabolism
- effects of electrical stimulation and phenylephrine (rat muscle), **136**, 863
 - mediation by guanine nucleotide-binding protein (GH_3 cells), **138**, 1383
- phosphorylation, relationship to mammary tumor growth (rat), **134**, 876
- turnover
- in Ca^{2+} -depleted neutrophils, effect of formyl-methionyl-leucyl-phenylalanine and concanavalin A (human), **140**, 1
 - and NADPH oxidase, independent activation in Ca^{2+} -depleted phorbol ester-primed neutrophils by formyl-methionyl-leucyl-phenylalanine (human), **135**, 785
- Phospholamban
- structural unit, amino acid sequencing and electrophoretic analysis (canine cardiac sarcoplasmic reticulum), **138**, 1044
- Phospholipase A_2
- effect of 47-kDa protein phosphorylation in γ -thrombin-stimulated platelets (human), **141**, 459
 - inhibition by protein I (bovine pancreas), **139**, 455
 - membrane-associated, activation mediated by α_1 -adrenergic and thyrotropin receptors, role of pertussis toxin-sensitive G proteins, analysis (FRTL-5 cells), **141**, 1000
 - reversible inhibition by 1-stearyl-2-stearoylaminodeoxyphosphatidylcholine (cobra venom), **137**, 587
- role in
- activation of NADPH oxidase by arachidonic acid in intact neutrophils and cell-free system, comparison (human), **138**, 1099
 - insulin release from pancreatic islets (rat), **138**, 720
- Phospholipase C
- from *Bacillus cereus*, antibodies, cross-reaction with phosphatidylcholine-preferring phospholipase C in human monocytic cells, **140**, 114
 - binding to substance P receptors, stimulation by guanine nucleotide-dependent regulatory protein (rat parotid gland), **136**, 362
 - diglyceride lipase system, role in arachidonic acid release in macrophages (mouse), **141**, 367
 - exogenous, inhibition of protein kinase C activation in HEL-37 cells, **136**, 1083
 - hydrolysis of [^3H]phosphatidylinositol 4,5-bisphosphate, stimulation by GTP and GDP (human platelets), **140**, 728
 - inhibition by aspirin, mechanism (human monocytes), **139**, 115
 - membrane-associated, activation mediated by α_1 -adrener-

Subject Index

- gic and thyrotropin receptors, role of pertussis toxin-sensitive G proteins, analysis (FRTL-5 cells), **141**, 1000
- partial purification and characterization (human platelets), **136**, 713
- phosphatidylcholine-preferring, cross-reaction with *Bacillus cereus* phospholipase C antibodies (human monocytic cells), **140**, 114
- phosphatidylinositol-specific, immunologically distinct forms, purification and characterization (bovine brain), **141**, 137
- zymosan-stimulated activation, intracellular Ca^{2+} requirement (murine peritoneal macrophages), **134**, 367
- Phospholipid kinase**
associated activity of estrogen receptor (MCF-7 cells), **135**, 597
- Phospholipid methyltransferase**
inhibition of adenylate cyclase (pigeon erythrocyte ghosts), **138**, 596
- Phospholipids**
bilayers, solvent-free, incorporated Na^+ , K^+ -ATPase, ion-channel activity, voltage and ATP effects (canine renal microsomes), **136**, 767
- binding regions of apolipoproteins, proline-containing, magnitude of helical hydrophobic moment, **139**, 202
- liposome-delivered, reversal of altered protein kinase C distribution in dystrophic muscle cells (mouse), **137**, 752
- plasma, arachidonate and docosahexaenoate content, reduction in Usher's syndrome (human), **141**, 600
- polar head group manipulation, effect on low-density lipoprotein metabolism in cultured fibroblasts (human), **137**, 43
- supported bilayers, for induction of two-dimensional antibody crystallization, **134**, 819
- N*-(Phosphonacetyl)-L-aspartate
binding to aspartate carbamoyltransferase, mechanism (*Escherichia coli*), **136**, 822
- Phosphonoformic acid**
inhibition of renal epithelial Na^+/P_i cotransport (human), **139**, 679
- Phosphopeptides**
phosphoseryl and phosphotyrosyl, distinct specificities of repressible acid phosphatase (yeast), **139**, 1202
- Phosphoprotein phosphatase**
activation by polyamines (HeLa cell nuclei), **134**, 1372
- associated with asparaginase complex, characterization (*Leptosphaeria michotii*), **140**, 1135
- ATP, Mg-dependent, 70-kDa modulator complexes, isolation and characterization (rabbit skeletal muscle), **135**, 367
- calmodulin-dependent, phosphorylation by protein kinase C (bovine brain), **138**, 783
- and protein kinase, tissue-homologous, reversible activation and inactivation of phosphofructokinase (*Ascaris suum*), **139**, 215
- Phosphoproteins**
B-50
phosphorylation in native synaptic plasma membranes, enhancement by 1,2-dioctanoylglycerol and phorbol diester (rat brain), **136**, 1007
- as protein kinase C substrate, levels in PC12 cells, effect of nerve growth factor (rat), **139**, 644
- from free polyribosomes, uv light-induced crosslinking with poly(A)⁺ RNA, effects of transcription and translation inhibitors (rat), **136**, 356
- P150, phosphorylation in adriamycin-resistant HL60 cells, effects of dimethylsulfoxide, retinoic acid, and phorbol ester, **138**, 9
- thylakoid 9-kDa, amino acid composition (pea), **138**, 146
- Phosphoenolpyruvate**
-glycose phosphotransferase system, III^{glc} protein, role in adenylate cyclase inhibition by P_i (*Escherichia coli*), **141**, 1138
- Phosphoenolpyruvate carboxykinase**
gene expression and regulation by insulin during preadipose cell differentiation, **138**, 468
- in glutamine utilization pathway, intracellular distribution in lymphocytes (rat), **138**, 318
- Phosphoenolpyruvate carboxylase**
diurnal regulatory changes, role of phosphorylation-dephosphorylation processes (CAM plants), **136**, 151
- Phosphoriboisomerase**, see Ribose-5-phosphate isomerase
- Phosphoribulokinase**
and phosphoriboisomerase, association with ribulose-1,5-bisphosphate carboxylase (pea), **139**, 947
- Phosphorus**
³²P, labeling of membrane phospholipids, effect of α_1 -adrenergic receptor stimulation (rabbit, rat aorta), **141**, 1213
- Phosphorylase phosphatase**
ATP, Mg-dependent, activation by kinase F_A , role of Mg^{2+} , **136**, 16
- catalytic subunit, inhibition by oxalyl thioesters, role in insulin action mechanism (bovine heart), **134**, 93
- Phosphorylation**, see also Protein kinase
acidic protein, effect of phorbol 12,13-dibutyrate, bombesin, and vasopressin (3T3 cells), **140**, 379
- auto-, see Autophosphorylation
- calpains I and II *in vivo*, analysis (K-562 cells), **136**, 1090
- dephosphorylation processes, role in phosphoenolpyruvate carboxylase diurnal regulatory changes (CAM plants), **136**, 151
- erythrocyte membranes, induction of Ca^{2+} release (human), **136**, 288
- glycogen synthase, effect on kinetic properties (rabbit, rat skeletal muscle), **134**, 1129
- glycoprotein hormone α -subunits secreted by tumor cell lines (human), **140**, 143
- growth hormone in pituitary glands *in vitro* (rat), **137**, 806
- heat shock proteins, 26- and 27-kDa, induction by hyperthermic or ecdysone treatment (*Drosophila melanogaster*), **141**, 426
- initiation factor eIF-2 α , role in inhibition of protein synthesis under heat shock conditions (HeLa cells), **138**, 519
- insulin receptor, analysis in skeletal muscle plasma membranes (normal, obese-diabetic mice), **137**, 286
- link proteins, analysis (human cartilage), **137**, 599
- mediation of glycogen synthase inactivation by mezerein and phorbol esters (rat hepatocytes), **134**, 113
- membrane proteins**
and NADPH oxidase activation, coordinate inhibition by surface antigen-directed monoclonal antibody (human neutrophils), **140**, 1121

Subject Index

- placental, atrial natriuretic factor-induced, comparison with guanosine 3':5'-cyclic monophosphate effect (human), **139**, 431
- mitochondrial proteins
- effect of *ras* oncogene-encoded proteins (rat liver), **135**, 316
 - estrone treatment effects (genetically diabetic mouse), **139**, 56
- myosin light chain
- role in platelet activation (human), **140**, 280
 - 20-kDa, effect on smooth muscle contraction (norepinephrine-stimulated canine veins), **135**, 735
- nucleocapsid protein of mouse hepatitis virus, analysis, **141**, 7; *erratum*, **141**, 884
- oxidative, effect of prostaglandin B₁ oligomeric derivatives (rat hepatic mitochondria), **137**, 215
- p82, stimulation by phorbol ester (murine epidermis), **137**, 766
- phosphoinositides, relationship to mammary tumor growth (rat), **134**, 876
- phosphoprotein B-50 in native synaptic plasma membranes, enhancement by 1,2-dioctanoylglycerol and phorbol diester (rat brain), **136**, 1007
- pp67, stimulation by interleukin 2 or phorbol ester (NK-7 cells), **135**, 487
- proteins
- angiotensin II-stimulated, inhibition by atrial natriuretic peptide (bovine adrenal glomerulosa cells), **140**, 814
 - 47-kDa, stimulation by γ -thrombin, effect on phospholipase A₂ (human platelets), **141**, 459
 - 48-kDa, intracellular, stimulation by phorbol ester in plasminogen activator-secreting melanoma cells (human), **137**, 263
 - in inner and outer adrenocortical zones (guinea pig), **138**, 895
 - in peroxisomes (rat liver), **140**, 188
 - tyrosine-specific, in plasma membrane, stimulation by oxygen radicals (rat liver), **139**, 439
- recombinant interferon γ , site identification (human), **136**, 679
- regulation of fodrin-binding proteins, role in structure of neuronal postsynaptic density (rat brain), **138**, 59
- retinoid-binding proteins, effect of protein kinase C *in vitro* (murine brain), **138**, 330
- rhodopsin, reduction in retinal dystrophy (dog), **135**, 1022
- sites on β -casein, localization by fast atom bombardment mass spectrometry (buffalo), **140**, 28
- tyrosine on soluble 66-kDa protein, induction by dimethyl sulfoxide (human T cells), **140**, 743
- Phosphorylcholine**
- binding protein, inhibition of low-density lipoprotein binding to hepatic membrane receptors (rat serum), **141**, 151
 - hydrolysis, quantitation in homogenates of temporal regions of Alzheimer's brain (human), **139**, 315
- Phosphorylcreatine shuttle**
- mitochondrial component, enhancement during perinatal cardiac development (rat), **141**, 319
- Phosphotyrosine**
- content in proteins, regulation by tyrosine kinase and phosphotyrosine phosphatase during induced granulocytic and monocytic differentiation (HL-60 cells), **140**, 440
 - Phosphotyrosine phosphatase, *see* Protein-tyrosine-phosphatase
 - Phosphotyrosyl-protein phosphatase
 - calcineurin-associated, regulation by calmodulin and protease treatment (bovine brain), **140**, 320
 - magnesium-dependent, identification and characterization (rat liver), **140**, 212
 - partial purification and characterization (human erythrocytes), **137**, 566
 - Photinus*-luciferin 4-monooxygenase (ATP-hydrolyzing)
 - inactivation by singlet oxygen, **140**, 888
 - Photoaffinity labeling
 - thromboxane A₂-prostaglandin H₂ antagonist binding site (human platelets), **140**, 128
 - Photofrin II
 - photosensitization of isolated mitochondria: impairment of anion translocation (rat liver), **141**, 584
 - Photoreceptors
 - membranes, inositol and phosphatidic acid metabolism, effect of cationic amphiphilic drugs (bovine), **134**, 378
 - Photosystem II
 - binding sites for 3-(3',4'-dichlorophenyl)-1,1-dimethylurea, characterization (spinach), **141**, 682
 - oxygen-evolving preparations, signal II_s alteration by calcium chloride and trypsin (spinach chloroplasts), **134**, 272
 - Phototoxicity
 - chemically-induced, evaluation using *Paramecium aurelia*, **139**, 79
 - hematoporphyrin derivative, enhancement by selenium (murine bladder tumor cells), **140**, 1086
 - insulin receptor-mediated: killing of activated lymphocytes by insulin-conjugated psoralen derivative, **141**, 502
 - Phytohemagglutinin
 - stimulated T cells, concomitant binding and degradation of and pyruvate dehydrogenase activation by insulin (human), **134**, 412
 - stimulation of cytosolic and particulate protein kinase C (human T cells), **137**, 128
 - Pineal gland
 - detection of type-II thyroxine 5'-deiodinase (rat), **137**, 863
 - Pinocytosis
 - fluid-phase, kinetics in *Dictyostelium discoideum* amoebae, **138**, 1146
 - Pirenzepine
 - effect on carbachol-induced pepsinogen secretion from isolated gastric glands, comparison with atropine (guinea pig), **136**, 193
 - Pituitary
 - cultured cells
 - desensitization induced by gonadotropin-releasing hormone, effect of wheat germ agglutinin (rat), **140**, 750
 - GH3, K⁺ conductance response to thyrotropin-releasing hormone, decrease after patch recording, reversal by GH3 cell extracts, **137**, 388
 - luteinizing hormone secretion stimulated by gonadotropin-releasing hormone, arachidonic acid and protein kinase C coordinate actions (rat), **134**, 134
 - desensitization, induction by copper (rat), **136**, 417
 - detection of glycosylated growth hormone with lectin-binding immunoassay (human), **140**, 491
 - expression of glutathione transferase, effect of xenobiotics

Subject Index

- (rat), **136**, 1057
- secretion of
- ACTH and prostaglandin E₂, induction by phorbol ester, role of Ca²⁺; effect of dexamethasone (rat), **138**, 323
- pro-opiomelanocortin-derived peptides and growth hormone *in vitro*, suppression by atrial natriuretic factor (rat), **135**, 1035
- synthesis and secretion of phosphorylated growth hormone *in vitro* (rat), **137**, 806
- Placenta
- first-trimester normal noncultured, analysis of human chorionic gonadotropin intracellular forms, **137**, 443
- as source of inhibin (human), **140**, 485
- Plantacyanin
- cobalt(II)-substituted, preparation and spectroscopic analysis (cucumber), **139**, 961
- Plasma
- antioxidant levels, effect of *Plasmodium vinckei* infection (mouse), **134**, 152
- aromatic and branched-chain amino acid levels, effect of sustained exercise (rat), **137**, 149
- atrial natriuretic factor levels
- effect of chronically administered adrenergic and cholinergic agonists (rat), **136**, 510
- increase during hypertension (salt-sensitive rat), **137**, 876
- increase in spontaneously hypertensive and spontaneously hypertensive stroke-prone rats, **135**, 74
- measurement in salt-sensitive Dahl rats, **136**, 411
- radioimmunoassay (hyper-, hypothyroid rats), **134**, 178
- atriopeptin levels, effect of dietary salt and water intake manipulations (rat), **137**, 922
- cholecystokinin levels, increase after intestinal administration of pancreatic enzyme secretion-stimulating peptide, radioimmunoassay (rat), **136**, 701
- detection of ouabain-like immunoreactivity in primary aldosteronism (human), **135**, 41
- pooled, as secondary standard in apolipoprotein E radioimmunoassay (human), **138**, 455
- seminal, *see* Seminal plasma
- Plasma membranes
- binding of
- apolipoprotein A-IV-phospholipid complexes (rat liver), **139**, 122
- copper ceruloplasmin, and uptake (rat tissues), **139**, 822
- cyclo(His-Pro) (rat liver), **134**, 443
- electron transport, role in diferric transferrin reductase system (rat liver), **139**, 1117
- fusion with pituitary secretion vesicles, inhibition by somatostatin (rat), **139**, 673
- identification of calmodulin-dependent Ca²⁺-activated K⁺ channel (rat adipocytes), **135**, 934
- interaction with calmodulin (rat liver), **135**, 323
- isolated, phosphorylation by casein kinase 1 (AH-66 hepatoma ascites cells), **138**, 95
- native synaptic, phosphoprotein B-50 phosphorylation, enhancement by 1,2-dioctanoylglycerol and phorbol diester (rat brain), **136**, 1007
- prolactin binding sites, ethanol effects (rat liver, prostate), **134**, 1122
- in skeletal muscle, isolation, properties, and use in analysis of insulin binding and receptor phosphorylation (normal, obese-diabetic mice), **137**, 286
- synthesis of phosphatidylcholine, stimulation by insulin and ATP (rat adipocytes), **137**, 1
- translocation of cytosolic protein kinase C, mechanism (human neutrophils), **136**, 228
- vesicles, hexose transport, characterization (rat L6 myoblasts), **141**, 124
- Plasmids
- containing cell cycle regulatory sequence, methylation pattern, analysis after transient transfection into fibroblasts (hamster), **135**, 942
- IncFII, basic replicon, enhanced instability by *polA* mutation, **139**, 918
- NAH7, *nah* and *sal* operons, regulation by *nahR* region, **141**, 986
- pBR322, DNA restriction fragments, preferential nucleosome placement (*Escherichia coli*), **138**, 110
- pIJ702, *Streptomyces*, heterologous genes cloned in, structural stability, **140**, 372
- pP-450 PBc2, encoded protein, and cytochrome P-450 K from rabbit kidney, N-terminal amino acid sequences, comparison, **141**, 728
- pPyLT1, autonomous genetic elements, maintenance after DNA transfer in transgenic strains (mouse), **141**, 1162
- vector type and transcription terminator, effect on heterologous gene expression (yeast), **140**, 602
- yeast centromeric, as shuttle vectors between *Bacillus subtilis*, *Escherichia coli*, and *Saccharomyces cerevisiae*, **135**, 915
- α₂-Plasmin inhibitor
- domain structure, demonstration by calorimetry, **139**, 1062
- Plasminogen activator
- secreting melanoma cells, phorbol ester-stimulated synthesis and phosphorylation of 48-kDa intracellular protein (human), **137**, 263
- Plasminogen activator inhibitor
- endothelial cell levels, enhancement by interleukin 1 (human), **139**, 720
- Plasmodium*
- P. falciparum*, induced malaria, detection with cloned DNA probe, **138**, 363
- P. vinckei*, infection, effect on plasma antioxidant levels (mouse), **134**, 152
- Plastocyanin
- spectroscopic characterization (*Ceratophyllum demersum*), **136**, 610
- Platelet-activating factor
- binding to neutrophil membranes, inhibition by GTP (human), **141**, 353
- identification in normal uterus and characterization (rat), **138**, 1231
- induction of prostaglandin E₂ synthesis in mesangial cells, opposite effects of different concentrations of pertussis toxin (rat), **141**, 39
- Platelet-derived growth factor
- effect on platelet aggregation, phosphoinositide metabolism, and serotonin release (human), **135**, 52
- like polypeptide, induction of epidermal growth factor-potentiated phenotypic transformation of kidney cells (rat), **141**, 1229
- and polyoma virus middle T antigen, cooperation in transformation of secondary cultured embryo cells (rat), **136**, 921

Subject Index

- Platelet-derived growth factor receptors
 autophosphorylation and phosphoinositide hydrolysis, effect of protein kinase C activation (human skin, murine Swiss 3T3 fibroblasts), **137**, 343
- Platelet-derived growth inhibitors
 types α and β , for hepatocytes in primary culture, detection and properties (rat), **134**, 755
- Platelets
 activation, role of myosin light chain phosphorylation (human), **140**, 280
 aggregation, phosphoinositide metabolism, and serotonin release, platelet-derived growth factor effects (human), **135**, 52
 aspirinated, agonist evoked aggregation, ATP secretion, and cytosolic Ca^{2+} increase, inhibition by antioxidants (human), **139**, 509
 cytosolic Ca^{2+} levels
 effects of thrombin and ionomycin, analysis with fura-2 (human), **139**, 308
 and thromboxane B_2 generation, sodium fluoride-induced increase, inhibition by protein kinase C activation (human), **136**, 381
 cytosol, stimulation of phosphoinositide hydrolysis in isolated membranes (human), **137**, 801
 deficient leukotriene C_4 synthesis, role of abnormal glutathione metabolism (human), **140**, 857
 fibrinogen binding, inhibition by ajoene (human), **141**, 145
 fura-2-loaded, thrombin- and ADP-induced Ca^{2+} flux, kinetics (human), **136**, 1124
 glycolysis, basal- and thrombin-stimulated, regulatory role of fructose 2,6-bisphosphate (human), **138**, 666
 membrane Ca^{2+} -ATPase activity, effect of activating hormones (human), **138**, 215
 and psoriatic lesions, 12-hydroxy-5,8,10,14-eicosatetraenoic acid stereochemistry, comparison (human), **136**, 169
 reversible shape changes, regulation by phosphatidylinositol metabolism (human), **141**, 673
 serotonin release
 and clot retraction, effects of amiloride and cytochalasin B (human), **140**, 329
 and inositol trisphosphate production and Ca^{2+} mobilization, cGMP effects (human), **135**, 1099
 sodium-hydrogen exchange, induction by 1-oleoyl-2-acetyl-glycerol and phorbol ester (human), **141**, 13
 stimulus-response coupling, role of diacylglycerol stereospecificity (human), **140**, 1143
 storage of β -transforming growth factor as latent high-molecular-weight complex (human), **136**, 30
 thrombin-activated
 arachidonic acid release, role of diacylglycerol lipase pathway (human), **134**, 1327
 47-kDa protein phosphorylation, effect on phospholipase A_2 (human), **141**, 459
 polyphosphoinositide metabolism and cytosolic Ca^{2+} mobilization, phorbol ester-mediated suppression, reversal by protein kinase C inhibitor (human), **134**, 868
 thrombopathic
 adenylate cyclase-phosphodiesterase system, regulatory control (dog), **140**, 589
 cAMP metabolism, effects of forskolin and 1-methyl-3-isobutyl-xanthine (dog), **140**, 595
- Polidocanol
 selective inhibition of microsomal hepatic cholesterol epoxide hydrolase (mouse), **140**, 632
- Polyadenylic acid
 irradiated, R- and S-8,5'-cycloadenosine yields, analysis by HPLC, **134**, 883
- Poly(ADP-ribose)
 biosynthesis, and suicidal NAD^+ depletion in mammalian cells following carcinogen exposure, relationship (mouse, rat), **138**, 1051
 effect on DNA-histone H4 interaction (rat liver), **139**, 523
- Poly(ADP-ribose) polymerase, *see* NAD^+ ADP-ribosyltransferase
- Poly(ADP-ribosylation)
 and DNA synthesis and mammary gland differentiation, interrelationships (mouse), **137**, 359
 inhibition of DNA polymerase β *in vitro* (bovine thymus), **140**, 666
- Polycyclic aromatic hydrocarbons
 nitro-, dihydrodiol derivatives, binding to DNA, catalysis by xanthine oxidase, **141**, 245
- Polyethylene glycol
 -modified hemin, peroxidase activity in organic solvents, **138**, 283
- Poly G
 bound to thymidylate synthase, identification (*Streptococcus faecium*), **139**, 461
- Polygalactosamine
 -containing glycoproteins, occurrence of $\text{Gal}\beta(1\rightarrow3)\text{GalNAc-Ser/Thr}$ in linkage region (*Cordyceps ophioglossoides*), **140**, 350
- Polygalacturonic acid
 oxidation by hydrogen peroxide, role in cell wall breakdown (orange), **141**, 238
- Polylysine
 modulation of multisubstrate Ca^{2+} - and cAMP-independent kinase (bovine aortic smooth muscle), **136**, 789
- Polymerization
 actin, effect of erythro-9-[3-(2-hydroxynonyl)]adenine, **137**, 1181
 tropomyosin, end-to-end, irreversible loss after exposure to urea or cyanate (rabbit muscle), **134**, 64
- Polymyxin B
 effect on
 insulin secretion from intact and permeabilised islets of Langerhans (rat), **136**, 1001
 protein synthesis (human T cells), **135**, 649
- Polyoma virus
 middle T antigen
 and mitogenic growth factors, cooperation in transformation of secondary cultured embryo cells (rat), **136**, 921
 transforming ability, enhancement by protein kinase C stimulation (rat F111 cells), **140**, 1106
 tyrosine residue phosphorylation, promotion by protein kinase C (murine py6, rat pyF cells), **136**, 995
 middle-T-only gene, transformed Ob17 cells, extracellular intermediates, promotion of Ob17 preadipocyte proliferation and differentiation, **137**, 378
- Polypeptides
 epidermal growth factor-like, synthesis and secretion by breast MCF-7 tumor cells (human), **136**, 300
 homologous with human insulin receptor precursor, cloning

Subject Index

- of encoding cDNA (*Drosophila* sp.), **141**, 474
 - major intrinsic, *see* Major intrinsic polypeptide
 - minor capsid σ 1a and nonstructural σ 1bNS, encoding mRNA from reovirus serotype 1 Lang strain, nucleotide sequence and molecular cDNA cloning, **140**, 508
- mitochondrial
 - composition, effect of mitochondrial myopathy and renal dysfunction (human), **137**, 911
 - synthesized *in vitro*, contamination with bacterial proteins (*Drosophila melanogaster*), **136**, 797
- nonstructural σ NS, encoding mRNA from reovirus serotype 1 Lang strain, cDNA cloning and nucleotide sequence (human), **139**, 845
- p31, hepatitis B virus, synthesis after gene modification, **141**, 942
- p36-associated, 10-kDa, homology with S100 protein subunits (porcine lymphocytes), **134**, 248
- peroxisomal membrane-associated, synthesis *in vitro* (rat liver), **137**, 626
- platelet-derived growth factor-like, induction of epidermal growth factor-potentiated phenotypic transformation of kidney cells (rat), **141**, 1229
- reovirus-specified, mRNA, molecular cloning and nucleotide sequence (*Escherichia coli*), **136**, 183
- synthesis induction by uv irradiation (*Saccharomyces cerevisiae*), **138**, 679
- Polyphosphoinositides
 - breakdown in neuroblastoma cells, effects of phorbol esters (human), **140**, 160
- hydrolysis
 - coupling to muscarinic receptors, induction by point-mutated p21^{ras} (murine fibroblasts), **141**, 591
 - enhancement by gonadotropin-releasing hormone (rat pituitary cells), **134**, 861
 - metabolism, phorbol ester-mediated suppression, reversal by protein kinase C inhibitor (thrombin-activated human platelets), **134**, 868
- Polyprenols
 - high-molecular-weight, covalently linked to proteins, detection in kidney (mouse), **139**, 209
- Poly R-481
 - decolorization, H₂O₂-dependent, by particulate fractions from *Phanerochaete chrysosporium*, **136**, 220
- Polyribocytidylic-polyriboinosinic acid
 - effect on
 - lipid peroxidation, xanthine oxidase activity, and hepatic cytochrome P-450 levels (mouse, rat), **134**, 632
 - xanthine oxidase and cytochrome P-450-mediated enzyme, role of H-2 haplotypes (mouse), **138**, 246
- Polysaccharides
 - in concanavalin A solutions, detection and quantitation with electrical oscillator, **137**, 978
- Poly sorbate 80
 - selective inhibition of microsomal hepatic cholesterol epoxide hydrolase (mouse), **140**, 632
- Porphyryns
 - meso-tetraaryl, bearing 1 to 4 positive charges, interaction with bovine thymic DNA, mechanism, **141**, 643
- Post-proline cleaving enzyme, *see* Prolyl endopeptidase
- Postsynaptic density
 - detection of protein kinase C subtype (rat brain), **140**, 691
 - neuronal structure, role of fodrin-binding proteins, regulation by phosphorylation (rat brain), **138**, 59
- Potassium
 - apamin-sensitive channels, and Ca²⁺ channels in muscular dysgenic skeletal muscle cells, action potentials and contractile activity (mouse), **136**, 935
 - calmodulin-dependent Ca²⁺-activated channels, identification in adipocyte plasma membranes (rat), **135**, 934
 - channels in lymphocytes, inhibition by retinoic acid (human), **138**, 560
 - currents through
 - lipid bilayer membranes, alteration induced by concanavalin A-polysaccharide interaction, **141**, 912
 - photoreceptor membranes, Ca²⁺-dependent inactivation, protein kinase C effects (*Hermisenda crassicornis*), **134**, 1245; *erratum*, **140**, 774
 - fluxes in mitochondria, quinine effect (rat liver), **135**, 830
 - GH3 cell conductance response to thyrotropin-releasing hormone, decrease after patch recording, reversal by GH3 cell extracts, **137**, 388
 - hydrogen antiporter, pH dependence, desensitization by N-ethylmaleimide (*Vibrio alginolyticus*), **136**, 1030
 - induction of
 - aldosterone secretion by glomerulosa cells, inhibition by diazepam (bovine adrenal), **135**, 994
 - catecholamine release from chromaffin cells, synergistic effect of Ca²⁺ and phorbol ester (bovine adrenal), **140**, 990
 - depolarization, enhancement of muscarinic receptor-stimulated inositol tetrakisphosphate accumulation in cerebral cortical slices (rat), **141**, 1130
 - elevation of intracellular free calcium levels in cultured chromaffin cells, differential effects of protein kinase C activators (bovine adrenal), **139**, 903
 - inhibition of calcium uptake and noradrenaline release by cultured chromaffin cells (bovine adrenal), **134**, 1
 - myocyte levels, determination by null-point technique (rat heart), **137**, 573
 - sodium-chloride cotransport system, biochemical and pharmacological properties (chicken cardiac cells), **134**, 326
 - sodium pump
 - associated antigens, binding to antibodies, enhancement by intracellular K⁺ (ovine erythrocytes), **134**, 982
 - in erythrocytes from cystic fibrosis patients and heterozygotes, decreased affinity for K⁺, **137**, 1021
- Potassium hydrogen persulfate
 - activation of bleomycin-iron III complex for DNA cleavage (bacteriophage λ), **136**, 1013
- Prazosin receptors
 - detection in skeletal muscle (rat), **136**, 1071
- Preadipocytes
 - growth arrest, coupling with expression of early markers during adipose conversion, **137**, 903
 - Ob17, proliferation and differentiation, promotion by extracellular intermediates from Ob17 cells transformed by polyoma virus middle-T-only gene, **137**, 378
 - phosphoenolpyruvate carboxykinase gene expression and regulation by insulin during differentiation, **138**, 468
- Pregnancy
 - effect on basal and noradrenaline-induced iodothyronine 5'-deiodinase activity in brown adipose tissue (rat), **138**, 1315

Subject Index

- Preproenkephalin
mRNA levels in neuroblastoma-glioma hybrid cells, synergistic regulation by glucocorticoids and cAMP, **139**, 1
- Preproinsulin
mRNA levels in pancreas, effect of glucose oral administration (rat), **141**, 1201
signal sequence removal, effect on signal recognition particle functions (carp), **134**, 790
- Preproparathyroid hormone
mRNA, role of 5'-terminal cap group and purine in 3' position in recognition of first AUG sequence as site of translation initiation, **140**, 288
- Primaquine
mediation of hydroxyl radical generation by *Trypanosoma cruzi* extracts, **135**, 1029
- Principal coordinate analysis
application to comparison of protein sequences, **140**, 808
- Proacrosin
mRNA, *in vitro* translation (porcine spermatozoa), **138**, 139
- Proadrenodoxin
adrenocortical, cell-free synthesis and characterization (human), **136**, 348
- Procaine
effect on hypoxia-, hyperkalemia-, and acidosis-induced changes in myocardial tissues (dog), **134**, 900
- Procollagen
type I and III, mRNA levels in cutaneous wounds treated with helium-neon laser (pig), **138**, 1123
type X disulfide-bonded, identification in embryonic sternum cartilage (chicken), **138**, 835
- [D-Pro¹⁰]dynorphin(1-11)
mono- and diiodinated derivatives, preparation and binding properties, **134**, 1142
- Profilin
effect on actin synthesis, fluorescence photobleaching recovery study (*Acanthamoeba castellanii*), **140**, 308
- Progesterone
accumulation and catabolism in cultured granulosa cells, effects of luteinizing hormone and human chorionic gonadotropin (rat), **141**, 1025
metabolism in granulosa cells, effects of prostaglandins E₂ and F_{2α} (rat), **135**, 764
- Progesterone receptors
interaction with progestin R 5020 and antiprogestin RU 38486, comparison (human breast cancer cells), **135**, 90
- Progestin, *see* Progesterone
- Prolactin
insulin-like action in adipose tissue *in vitro*, inhibition by glycosylation (rat), **134**, 764
related mRNA, detection in pituitary GH₄C₁ tumor cells (rat), **138**, 1015
release from pituitary tumor cells, induction by bacterial adenylate cyclase (rat), **136**, 463
stimulation of hepatic DNA synthesis (rat), **138**, 1138
- Prolactin receptors
in mammary gland, regulation by luteinizing hormone (rat), **135**, 1076
in plasma membrane and Golgi fractions, ethanol effects (rat liver, prostate), **134**, 1122
- Proline
-containing phospholipid-binding regions of apolipoproteins, magnitude of helical hydrophobic moment contents, **139**, 202
hydration state in water/dimethyl sulfoxide mixtures, ¹⁷O NMR, **135**, 363
inhibition of nitrogen fixation in *Anabaena* 7120, **139**, 940
X-ray crystal structure data, analysis, **140**, 895
- Prolyl endopeptidase
inhibition by *N*-benzyloxycarbonyl-Gly-Pro-diazomethyl ketone (bovine brain), **134**, 888
- Pro-opiomelanocortin
derived peptides, secretion from anterior and intermediate pituitary cells *in vitro*, suppression by atrial natriuretic factor (rat), **135**, 1035
- Pro-oxytocin
-neurophysin, cleavage at paired basic residues by endopeptidase associated with neurohypophyseal secretory granules (bovine), **134**, 1190
- Prophenoloxidase
activating system, identification of β-1,3-glucan and peptidoglycan receptors (silkworm plasma), **141**, 1177
- Propranolol
effect on
inositide and phosphatidic acid metabolism in photoreceptor membranes (bovine), **134**, 378
inositol incorporation into phosphoinositides and inositol phosphates (rat C6 glioma cells), **135**, 465
- Propylene
epoxidation by chloroperoxidase (*Caldariomyces fumago*), **136**, 778
- Propylinium
effect on calmodulin activity and cellular proliferation (rat astrocytoma cells), **135**, 574
- 7-α-Propyl spirolactone receptors
identification as anti-mineralocorticoid binding sites (rat kidney), **134**, 261
- Prostacyclin
high-density lipoprotein-induced synthesis, role of apoprotein and lipid constituents (rabbit heart), **139**, 416
synthesis in cultured vascular endothelial cells, stimulation by cachectin/tumor necrosis factor (human), **141**, 482
- Prostaglandin A₁
tumor cell biotransformation products, growth inhibitory effects (rat neuroblastoma, glioma cells), **141**, 1254
- Prostaglandin B₁
oligomeric derivatives, Ca²⁺ ionophoretic activity and effect on mitochondrial oxidative phosphorylation (rat), **137**, 215
- Prostaglandin E₂
effect on
formyl-methionyl-leucyl-phenylalanine-induced inositol phospholipid metabolism and Ca²⁺ mobilization (guinea pig neutrophils), **139**, 1272
β-migrating very-low-density lipoprotein metabolism (rat peritoneal macrophages), **137**, 461
progesterone metabolism (rat granulosa cells), **135**, 764
metabolic dehydration and cellular uptake, relationship to prostaglandin E₂-induced growth inhibition (murine leukemia cells), **139**, 808
regulation of tumor necrosis factor production by macrophages (mouse), **137**, 404
secretion by pituitary cells, phorbol ester-induced, role of Ca²⁺: effect of dexamethasone (rat), **138**, 323

Subject Index

- synthesis in
 bone, stimulation by recombinant human interleukin 1 α (fetal mouse), **138**, 618
 MC3T3-E1 cells, stimulation by recombinant human interleukin-1 α and β , **141**, 285
 mesangial cells, agonist-induced, opposite effects of different concentrations of pertussis toxin (rat), **141**, 39
 resting macrophages, stimulation by tumor necrosis factor (mouse), **136**, 94
- Prostaglandin E₂ receptors
 binding affinity, guanosine triphosphate effects (porcine brain), **135**, 282
- Prostaglandin F_{2 α}
 effect on
 cytosolic free Ca²⁺ levels in vascular smooth muscle cells (rat aorta), **136**, 247
 intracellular Ca²⁺ levels and Ca²⁺ translocation (canine thyroid cells), **141**, 569
 progesterone metabolism (rat granulosa cells), **135**, 764
- Prostaglandin H₂
 -thromboxane A₂ antagonist binding site, photoaffinity labeling (human platelets), **140**, 128
- Prostaglandin I₂
 effect on adenylate cyclase-phosphodiesterase system in thrombopathic and normal platelets (dog), **140**, 589
- Prostaglandin I₃
 production from eicosapentaenoic acid, enhancement by arachidonic acid (human endothelial cells), **135**, 403
- Prostaglandins
 cardiac release *in vitro*, enhancement by cytochrome P-448 inducers (chick embryo), **136**, 582
 glomerular release, effect of atrial natriuretic factor (rat kidney), **138**, 496
 synthesis in
 peritoneal macrophages, regulatory role of protein kinase C (mouse), **139**, 604
 3T3 fibroblasts, phorbol ester-induced, relationship to ornithine decarboxylase induction (BALB/c mouse), **139**, 186
- Prostatropin
 multiple forms, characterization (bovine brain), **136**, 1155
- α_1 -Protease inhibitor
 binding of trypsin and chymotrypsin, comparison, **138**, 925
- Proteases
 calcium-activated neutral
 autolytic activation (rabbit muscle), **138**, 638
 phosphorylation by rat cerebral protein kinase C (bovine heart), **137**, 559
 gene, human T-cell leukemia virus type I, identification (MT-2 cells), **139**, 129
 inactivation of *Bacillus thuringiensis* var. *israelensis* toxin (mosquito larvae), **135**, 902
 metallo-, from mitochondrial matrix fraction, processing of enzyme precursors (rat liver), **134**, 1151
 -secreting mutants, isolation and characterization (*Bacillus subtilis*), **138**, 526
- α_1 -Proteinase inhibitor
 without or with high-mannose, hybrid, or complex-type oligosaccharide side chains, clearance by isolated perfused liver (rat), **141**, 949
- Proteinases
 calcium-dependent neutral, erythrocyte levels and subcellular distribution, Ca²⁺-induced alterations (human), **138**, 87
- Protein-glutamine γ -glutamyltransferase
 calcium-dependent, catalyzed reactions, inhibition of adenylate cyclase (pigeon erythrocyte ghosts), **138**, 596
 involucrin action as substrate at multiple sites (human keratinocytes), **136**, 51
 substrate identification in inside-out vesicles by immunoblotting (human erythrocytes), **134**, 685
- Protein kinase, *see also* Phosphorylation
 associated activity of estrogen receptor (MCF-7 cells), **135**, 597
 associated with asparaginase complex, characterization (*Leptosphaeria michotii*), **140**, 1135
 calcium-activated calmodulin-dependent
 detection in bovine thyroid cytosol, **140**, 120
 phosphorylation of hepatic pyruvate kinase: characterization of two phosphorylation sites (rat), **139**, 1017
 calcium-activated phospholipid-dependent, *see* Protein kinase C
 calcium-dependent and -independent, reactivity with myosin light chains, **134**, 240
 cAMP-dependent
 activation of heparin-releasable lipoprotein lipase (rat heart), **135**, 196
 activity in
 H-ras-transformed NIH 3T3 cells, **139**, 787
 inner and outer adrenocortical zones (guinea pig), **138**, 895
 and casein kinase I, comparative effects on glycogen synthase behavior as substrate for hepatic synthase phosphatases, **139**, 1033
 catalytic subunit
 phosphorylation of microsome-bound cytochrome P-450 LM₂ (rabbit liver), **134**, 461
 and regulatory subunit, comparison using polyclonal antibodies (bovine heart, *Dictyostelium discoideum*), **136**, 651
 relationship to anticarcinogenic effect of retinoids on mammary tumor induction by 7,12-dimethylbenz[a]anthracene (rat), **135**, 116
 stimulation of cholesterol 7 α -hydroxylase (rat), **134**, 797
 insulin-stimulatable, identification of muscle protein substrates related to microtubule-associated protein 2 (3T3-L1 adipocytes), **134**, 565
 multisubstrate, Ca²⁺- and cAMP-independent, modulation by polycations (bovine aortic smooth muscle), **136**, 789
 phosphorylation of
 glycogen synthase, effect on kinetic properties (rabbit, rat skeletal muscle), **134**, 1129
 high-mobility-group proteins (Chinese hamster ovary cells), **134**, 617
 in plasma, effect of Fe²⁺ (human), **138**, 693
 and protein phosphatase, tissue-homologous, reversible activation and inactivation of phosphofructokinase (*Ascaris suum*), **139**, 215
 ribosomal S6, detection in insulin-treated 3T3-L1 cells and Rous sarcoma virus-transformed chicken embryo fibroblasts, **137**, 702
- Protein kinase C
 activation

Subject Index

- by dexamethasone and saccharin (rat hepatocytes), **135**, 33
- effect on platelet-derived growth factor receptor autophosphorylation and phosphoinositide hydrolysis (human skin, murine 3T3 fibroblasts), **137**, 343
- by ganglioside GM3 in presence of calcium and phorbol ester, **138**, 865
- inhibition by exogenous phospholipase C (HEL-37 cells), **136**, 1083
- inhibition of sodium fluoride-induced increase in platelet cytosolic free calcium and thromboxane B₂ generation (human), **136**, 381
- by lipophilic muramyltripeptide MTP-PE (porcine brain), **140**, 1043
- by phorbol ester, enhancement of polyoma virus middle T antigen transforming ability (rat F111 cells), **140**, 1106
- by tumor promoters, role of sulfatide (bovine brain), **138**, 153
- altered distribution in dystrophic muscle cells, reversal by liposome-delivered phospholipids (mouse), **137**, 752
- and arachidonic acid, coordinate actions in gonadotropin-releasing hormone-stimulated luteinizing hormone secretion (rat pituitary cells), **134**, 134
- assay in platelets (normotensive, spontaneously hypertensive rats), **141**, 769
- calcium-dependent, conversion to independent form of phorbol ester-binding protein by digestion with trypsin (rat brain), **139**, 320
- catalytic properties, phorbol ester-induced alterations in bovine adrenocortical cells, human lymphoid cells, and rat fibroblasts, **134**, 1031
- characterization (rat-1 cells), **138**, 1283
- cytosolic
 - effects of insulin and phorbol ester, comparison (rat adipocytes), **135**, 1119
 - and particulate, stimulation by phytohemagglutinin (human T cells), **137**, 128
 - phorbol ester-induced desensitization, effect on cell growth (human breast cancer), **135**, 65
 - translocation to activated plasma membranes, mechanism (human neutrophils), **136**, 228
- down-regulation, induction by phorbol esters (rat brain), **139**, 251
- effect on
 - calcium channel function (PC12 cells), **136**, 1049
 - calcium-dependent inactivation of K⁺ currents across photoreceptor membranes (*Hermisenda crassicornis*), **134**, 1245; *erratum*, **140**, 774
 - phosphorylation of retinoid-binding proteins *in vitro* (murine brain), **138**, 330
- effects of
 - arachidonic acid oxygenation products (human), **134**, 1215
 - bovine myelin basic protein synthetic peptide analogs, **134**, 78
 - staurosporine (rat brain), **135**, 397
 - 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (murine thymoma cells), **140**, 267
- identification and characterization in small intestinal epithelial cells (rabbit, rat), **139**, 875
- intracellular translocation and down-regulation, induction
 - by phorbol ester (MCF-7 cells), **137**, 1159
 - and NADPH oxidase, coactivation in plasma membrane (bovine neutrophil cytoplasts), **134**, 305
- phorbol ester-induced translocation
 - in promyelocytic leukemia cells (human), **135**, 1144
 - from soluble to particulate cell fraction (rat hepatocytes), **138**, 1328
- phosphorylation of
 - bovine cardiac calcium-activated neutral protease (rat brain), **137**, 559
 - caldesmon (chicken smooth muscle), **132**, 56; *erratum*, **134**, 1029
 - calmodulin-dependent protein phosphatase (bovine brain), **138**, 783
 - interleukin-2 receptor Tac antigen epitope (human T cells), **135**, 239
 - lipocortins *in vitro* (bovine), **141**, 547
 - middle T antigen tyrosine residues in polyoma virus-transformed cells (mouse, rat), **136**, 995
 - p36 *in vitro* (bovine lung), **137**, 397
 - 6-phosphofructokinase, inhibition by phosphofructokinase stabilizers (rabbit, rat muscle), **136**, 445
 - talin, **134**, 1276
- protein kinase M, characterization (guinea pig peritoneal macrophages), **141**, 605
- purification by HPLC (rat brain), **135**, 636; *erratum*, **136**, 844
- redistribution and proteolytic cleavage during B-cell mitogenesis (human), **135**, 146
- regulation of
 - arginine vasopressin-induced inositol trisphosphate accumulation in cultured aortic myocytes (rat), **140**, 1093
 - tumor necrosis factor binding to membrane receptors (human T cells), **141**, 855
- role in
 - calcium channel regulation (PC-12 cells), **134**, 1298
 - epidermal growth factor receptor phosphorylation in K-ras-transformed cells (rat kidney), **135**, 435
 - interleukin 1 production by peritoneal macrophages (mouse), **135**, 355
 - osmotic activation of Na⁺/H⁺ exchange (rat thymic lymphocytes), **134**, 8
 - prostaglandin synthesis by peritoneal macrophages (mouse), **139**, 604
 - stimulus-dependent movement from cytosol to membranes (human neutrophils), **134**, 587
 - substrate B-50 levels in PC12 cells, effect of nerve growth factor (rat), **139**, 644
 - substrate specificity, analysis with synthetic peptide containing D- or L-threonine residues (rat brain), **134**, 1358
 - subtype, detection in postsynaptic densities (rat brain), **140**, 691
 - translocation features in neutrophils stimulated with chemotactic peptide formyl-methionyl-leucyl-phenylalanine (guinea pig), **139**, 1169
- Protein kinase M
 - protein kinase C, characterization (guinea pig peritoneal macrophages), **141**, 605
- Protein kinase (tyrosine), *see* Protein-tyrosine kinase
- Protein phosphatase, *see* Phosphoprotein phosphatase
- Proteins
 - 2-5A-binding, and 2-5A synthetase, induction by interfer-

Subject Index

- on, age-related differences (guinea pig macrophages), **140**, 455
- acetaldehyde conjugates, detection in urine of alcoholics, **140**, 924
- acidic, phosphorylation, effect of phorbol 12,13-dibutyrate, bombesin, and vasopressin (Swiss 3T3 cells), **140**, 379
- acute phase, gene association with hepatic nuclear matrix during inflammation (rat), **137**, 633
- ADP/ATP-carrier
- mRNA, isolation and characterization (bovine cardiac mitochondria), **138**, 850
 - substrate and inhibitor binding, analysis with 6'-O-dansyl- γ -aminobutyryl atractyloside (bovine cardiac mitochondria), **134**, 266
- alveolyn-like, associated with murine type 2 pneumocytes, immunochemical characterization, **139**, 1135
- amino acid sequences, comparison by principal coordinate analysis, **140**, 808
- amyloid fibril, from insulin-producing pancreatic tumor, partial homology of N-terminal sequence with calcitonin gene-related peptide (human), **140**, 827
- androgen-binding, secretion by immature Sertoli cells, *in vitro* effects of germ cells (rat), **141**, 861
- bacterial, contamination of mitochondrial translation products (*Drosophila melanogaster*), **136**, 797
- band 3
- dephosphorylation by cytosolic acid phosphatase specific for phosphotyrosine (human erythrocytes), **134**, 557
 - as substrates for transglutaminase, identification in inside-out vesicles by immunoblotting (human erythrocytes), **134**, 685
 - tyrosine phosphorylation by casein kinase A (human erythrocytes), **139**, 855
- beta, Alzheimer's disease-associated, synthetic peptides corresponding to 17- and 28-residue sequences, *in vitro* formation of amyloid fibrils (human), **141**, 782
- calcium-binding
- identification in bovine heart, **139**, 596
 - vitamin D-dependent, activation of Ca^{2+} , Mg^{2+} -ATPase activity (rat kidney), **138**, 547
- calmodulin- and F-actin-binding, 34-kDa, isolation and characterization (chicken gizzard smooth muscle), **141**, 20
- calmodulin-like, detection in *Bacillus subtilis*, **134**, 212
- containing histidines and tyrosines, labeling with p-diazobenzoyl biocytin, **138**, 872
- core VII, adenovirus serotype 2, chromatin-like complex with DNA, transcription *in vitro*, **136**, 86
- crystal, from *Bacillus thuringiensis* var. *israelensis*, removal of 28-kDa hemolytic peptide from solubilized form, effect on mosquito toxicity, **141**, 937
- cytosolic, covalent binding of 4-nitrobenzyl mercaptan S-sulfate, inhibition by glutathione (rat liver), **137**, 1055
- enhancer-binding, prediction by informational spectrum method, **141**, 831
- envelope
- DNA and RNA plant viruses, sequence homology, **134**, 372
 - human T-lymphotropic virus type III, detection of hexapeptide homologous to interleukin-2 receptor binding region (murine spleen), **139**, 367
- eosinophil cationic, cytotoxic, ribonuclease activity, **139**, 1239
- erythrocyte, 4.1, related factor, role in caldesmon-actin interaction, **140**, 1051
- expression in retina after optic nerve crush (goldfish), **138**, 1177
- extraction, application of reverse micelles, **135**, 629
- fatty acid-binding, from heart, liver, and intestine, purification and tissue expression (rat), **137**, 929
- 47-kDa, phosphorylation, stimulation by γ -thrombin, effect on phospholipase A2 (human platelets), **141**, 459
- 48-kDa
- expressed in proliferating lymphocytes, isolation and characterization (human, mouse), **134**, 1238
 - intracellular, phorbol ester-stimulated production and phosphorylation in plasminogen activator-secreting melanoma cells (human), **137**, 263
 - glucose-regulated, induction in fibroblasts, effect of extracellular Ca^{2+} and temperature (hamster), **140**, 881
- group A streptococcal Pep M5, amino acid sequence homology with other coiled-coil proteins, **140**, 684
- growth hormone-binding, affinity purification and structural characterization (human serum), **139**, 150
- growth hormone-dependent insulin-like growth factor-binding, from plasma, comparison to amniotic fluid IGF-binding protein (human), **139**, 1256
- guanine nucleotide-binding
- ADP-ribosylated by pertussis toxin, detection in spermatozoa (invertebrates, vertebrates), **138**, 728; *addendum*, **140**, 773
 - mediation of thyrotropin-releasing hormone-induced GTPase activity and phosphoinositide metabolism (GH₃ cells), **138**, 1383
- pertussis toxin-sensitive, role in α_1 -adrenergic and thyrotropin receptor-mediated activation of membrane phospholipases and iodide fluxes, analysis (FRTL-5 cells), **141**, 1000
- role in
- formation of inositol phosphates in adrenal glomerulosa cells (rat), **140**, 941
 - mediation of α -adrenergic responses in brown adipocytes, evaluation with pertussis toxin (hamster), **135**, 823
- stimulatory
- effect on phospholipase C binding to substance P receptors (rat parotid gland), **136**, 362
 - functional coupling with adenylate cyclase catalytic unit, relationship to enzyme inhibition by pentobarbital (rat brain), **140**, 237
 - interaction with catalytic unit of adenylate cyclase system, forskolin stabilizing effect (rat erythrocytes), **137**, 190
- heat shock
- association with cytoskeletal elements and binding to calmodulin (rabbit retina), **139**, 974
 - hsp 90, relationship to steroid hormone receptors and tyrosine kinase oncogene products, **138**, 1298
 - induction in fibroblasts, effect of extracellular Ca^{2+} and temperature (hamster), **140**, 881
 - 26- and 27-kDa, phosphorylation induced by hyperthermic or ecdysone treatment (*Drosophila melanogaster*), **141**, 426
- high-mobility-group
- phosphorylation (Chinese hamster ovary cells), **134**, 617

Subject Index

- supercoil-dependent recognition of specific DNA sites (chicken), **138**, 910
- high-molecular-weight nonfunctional, with antigenic homology to dihydrofolate reductase, detection in brain (mouse), **141**, 258
- iron-sulfur and ubiquinone-binding, antibodies, interaction with mitochondrial complex III (bovine heart), **136**, 376
- larvicidal, purification (*Bacillus sphaericus* 1593), **141**, 756
- link, phosphorylation state (human cartilage), **137**, 599
- lipid transfer, isolation from hemolymph (*Manduca sexta*), **136**, 260
- masking, for β -transforming growth factor, partial purification and characterization (rat platelets), **141**, 176
- matrix, in purified peroxisomes, coenzyme A binding (rat liver), **139**, 1195
- membrane
- and/or glycoproteins, functional group labeling with biotin derivatives and ^{125}I -labeled streptavidin (human erythrocytes), **136**, 80
 - phosphorylation
 - atrial natriuretic factor-induced, comparison with guanosine 3':5'-cyclic monophosphate effect (human placenta), **139**, 431
 - effect of *ras* oncogene-encoded proteins (rat hepatic mitochondria), **135**, 316
 - and NADPH oxidase activation, coordinate inhibition by surface antigen-directed monoclonal antibody (human neutrophils), **140**, 1121
- MEP, secreted from transformed cells, protease specificity, comparison with cathepsin L (mouse), **139**, 156
- microtubule-associated, *see* Microtubule-associated proteins
- modulator complexes, 70-kDa, of ATP/Mg-dependent protein phosphatase, isolation and characterization (rabbit skeletal muscle), **135**, 367
- mono(ADP-ribosyl)ated, hydrolyzing enzyme, identification (rat skeletal muscle), **139**, 932
- muscle, related to microtubule-associated protein 2, identification as substrates for soluble insulin-stimulatable serine/threonine kinase (3T3-L1 adipocytes), **134**, 565
- myelin basic, *see* Myelin basic protein
- N-*ras*, GTPase and autophosphorylation activities *in vitro* (NIH 3T3 cells), **139**, 771
- nucleocapsid, mouse hepatitis virus, phosphorylation sites, **141**, 7; *erratum*, **141**, 884
- p21, Ha-*ras* oncogene-encoded
 - complexes with nucleotides, ^{31}P NMR, **135**, 549
 - point-mutated, induction of muscarinic receptor coupling to calcium channels and polyphosphoinositide hydrolysis (murine fibroblasts), **141**, 591
- p36
 - oligomeric, inhibition of phospholipase A₂ (bovine pancreas), **139**, 455
 - phosphorylation *in vitro* by protein kinase C (bovine lung), **137**, 397
- p60^{c-src}, kinase activity, detection in chromaffin granule membranes (bovine), **134**, 736
- p82, phosphorylation, stimulation by phorbol ester (murine epidermis), **137**, 766
- pertussis/cholera toxin-sensitive, mediation of chemoattractant receptor signal transduction (human), **138**, 887
- phosphoenolpyruvate:glycose phosphotransferase system III^{glc}, role in adenylate cyclase inhibition by P_i (*Escherichia coli*), **141**, 1138
- phosphorylation
 - angiotensin II-stimulated, inhibition by atrial natriuretic peptide (bovine adrenal glomerulosa cells), **140**, 814
 - in inner and outer adrenocortical zones (guinea pig), **138**, 895
 - in peroxisomes (rat liver), **140**, 188
 - tyrosine-specific, in plasma membrane, stimulation by oxygen radicals (rat liver), **139**, 439
- phosphorylcholine-binding, inhibition of low-density lipoprotein binding to hepatic membrane receptors (rat serum), **141**, 151
- phosphotyrosine content, regulation by tyrosine kinase and phosphotyrosine phosphatase during induced granulocytic and monocytic differentiation (HL-60 cells), **140**, 440
- plasma, effect of *Plasmodium vinckei* infection (mouse), **134**, 152
- pOb24, mRNA expression, coupling with growth arrest during adipose conversion of preadipocytes, **137**, 903
- pp67, phosphorylation, stimulation by interleukin 2 or phorbol ester (NK-7 cells), **135**, 487
- precursor, mitochondrial, processing by metalloprotease purified from matrix fraction (rat liver), **134**, 1151
- prenylated, detection in kidney (mouse), **139**, 209
- pulmonary surfactant-associated, chemical and biophysical properties (bovine), **135**, 527
- ras* oncogene-encoded, effect on phosphorylation of mitochondrial membrane proteins (rat liver), **135**, 316
- recA*, binding to double-stranded DNA, superhelicity effects (*Escherichia coli*), **136**, 404
- related to Pb-thionein, immunological analysis (rat liver), **136**, 535
- retinoid-binding
 - covalent binding of azidoretinoids, **135**, 124
 - phosphorylation, effect of protein kinase C *in vitro* (murine brain), **138**, 330
 - structural changes induced by retinol removal: molecular dynamics study (human), **139**, 564
- ribosomal, casein kinase II-phosphorylated, identification (cryptobiotic *Artemia* sp. gastrulae), **135**, 347
- RNA complexes, formation in 4-thiouridine-exposed and subsequently illuminated CV-1 cells, **141**, 847
- S-, and vitronectin: physicochemical, immunochemical, and functional comparison (human), **134**, 951
- S100, subunits, homology with lymphocytic p36-associated 10-kDa polypeptide (swine), **134**, 248
- serine-rich, gene expression, modulation by glucose (*Saccharomyces cerevisiae*), **138**, 297
- soluble 66-kDa, tyrosine phosphorylation, induction by dimethyl sulfoxide (human T cells), **140**, 743
- somatomedin-binding
 - from amniotic fluid, in insulin-like growth factor I purification by affinity chromatography (human), **136**, 253
 - high-molecular-weight, purification (human plasma), **135**, 178
 - identification of biologically active domain (human placenta), **141**, 263
- spatial structures, destabilization by random single-point

Subject Index

- mutations, protection by codon bias in genes, **139**, 1094
- sphingolipid activator type 1, encoding cDNA, isolation and characterization (human), **134**, 989
- spore coat, N-terminal amino acid sequence (*Dictyostelium discoideum*), **135**, 336
- src*, associated tyrosine kinase activity, phosphorylation of band 3 glycoprotein (human erythrocytes), **135**, 720
- ssDNA-binding, modulation of topoisomerase I relaxing activity (*Escherichia coli*), **137**, 795
- staphylococcal A, in apolipoprotein E radioimmunoassay, **138**, 455
- stress
- induction by auranofin (human, mouse, rat), **138**, 1074
 - role of glutathione in induction (fetal rat fibroblasts), **134**, 492
- sulfation, inhibition by chlorate (rat PC12 cells), **141**, 870
- tau, microtubule-associated, self-assembly into filaments resembling Alzheimer's disease-associated filaments (porcine brain), **141**, 790
- thin filament-associated, regulation of smooth muscle actomyosin ATPase, **136**, 962
- transformation-associated, immunodetection in cells transformed by ts110 mutant of Moloney murine sarcoma virus (rat kidney), **134**, 1223
- tyrosine sulfation sites, characterization and criteria for prediction of occurrence, **141**, 326
- ubiquinone-binding, purification from mitochondrial NADH-ubiquinone reductase (bovine heart), **138**, 1237
- VIP precursor, characterization of derived peptide (cat, guinea pig, human, rat, swine), **139**, 1142
- vitamin D-binding, inhibition of sequential binding of actin monomers to plasma gelsolin, **136**, 72
- v-rel* oncogene-encoded, identification in reticuloendotheliosis virus T-transformed bone marrow cells, expression in Cos1 cells (chicken), **134**, 716
- Protein synthesis
- cardiac, stress-induced suppression, role in subsequent hypertrophy (rat), **135**, 857
 - c-myc* gene product, role of *c-myc* first exon (human), **140**, 313
- heat shock proteins in
- HeLa cells at 42°C, **137**, 957
 - primary cell cultures and established cell lines, comparison (human), **141**, 46
- induction by uv irradiation (*Saccharomyces cerevisiae*), **138**, 679
- inhibition by
- bovine cell-surface sialoglycopeptide, effects of calcium and sodium ionophores (murine 3T3 cells), **136**, 976
 - bovine sialoglycopeptide, role of cell surface interaction (murine 3T3 cells), **134**, 1350
 - heat shock conditions, role of initiation factor eIF-2 α phosphorylation (HeLa cells), **138**, 519
 - heme deprivation, promotion of globin mRNA storage (rabbit), **134**, 1048
 - polymyxin B (human T cells), **135**, 649
- initiation, role of guanine nucleotide exchange factor (HeLa cells), **134**, 1160
- mitochondrial, inhibition by rhodamine 123 (rat liver, tumor cells), **137**, 759
- plasma proteins in cultured hepatocytes, comparative effects of insulin and insulin-like growth factors (chick embryo), **134**, 427
- 71-kDa stress protein, induction by heat shock (canine heart), **134**, 166
- ubiquinone-binding protein from mitochondrial cytochrome *bc₁* complex in cell-free system (rat liver), **138**, 1291
- Protein-tyrosine kinase
- detection in lymphocytes by renaturation in polyacrylamide gels, **134**, 963
 - epidermal growth factor receptor- and *src* protein-associated activity, phosphorylation of band 3 glycoprotein (human erythrocytes), **135**, 720
 - identification by peptide-specific antibodies (human T cells), **140**, 392
 - multiple forms in spleen, identification and characterization (rat), **139**, 910
 - src*-related, quantitation during fertilization (sea urchin egg), **138**, 349
- Protein-tyrosine-phosphatase
- regulation of protein phosphotyrosine content during induced granulocytic and monocytic differentiation (HL-60 cells), **140**, 440
 - role in dephosphorylation of membrane protein band 3 (human erythrocytes), **134**, 557
- Proteodermatan sulfate
- interaction with collagen, role of disulfide bridges in protein core (bovine skin), **138**, 1348
- Proteoglycans
- synthesis in cultured articular chondrocytes, enhancement by interleukin 1-like factor (human), **134**, 539
- Proteolysis
- calcium-binding proteins, calcium effects (rat), **134**, 1269
 - phenylalanine ammonia-lyase, protective effect of entrapment in silk fibroin (*Rhodotorula glutinis*), **141**, 165
- Prothrombin
- fragment 1, sequential Ca²⁺ and Mg²⁺ binding, associated spectral response (bovine), **141**, 1207
- Proton transfer
- thermodynamics of carboxylic acid-retinal Schiff base hydrogen bonds, **138**, 819
- Pseudomonas aeruginosa*
- outer membrane, action as barrier against disaccharide penetration, **134**, 106
- Psoralen
- phototoxicity evaluation using *Paramecium aurelia*, **139**, 79
- Psoriasis
- lesional scales, and platelets, 12-hydroxy-5,8,10,14-eicosatetraenoic acid stereochemistry, comparison (human), **136**, 169
- Psychosine
- inhibition of cytochrome-*c* oxidase (rat hepatic mitochondria), **137**, 323
- Pulse radiolysis
- methoxatin semiquinone, **139**, 955
- Purpurogallin
- inhibition of glutathione *S*-transferase isoenzymes from human and rat tissues, comparison, **141**, 1170
- Putrescine
- effect on protein phosphatase activity (HeLa cell nuclei), **134**, 1372

Subject Index

- role in mediation of androgen response in kidney (mouse), **138**, 771
- Pyridine**
 derivatives, inhibition of mitochondrial NADH dehydrogenase, relationship to experimental and idiopathic parkinsonism, **135**, 269
- D-3-Pyridinyl-alanine⁶
 -containing luteinizing hormone-releasing hormone antagonists, increased potency, **137**, 709
- α -(4-Pyridyl 1-oxide)-N-t-butyl nitro-
 hydroxyl and superoxide radical adducts, [¹⁷O]oxygen hypofluoride structure, **141**, 622
- Pyrilamine**
 binding to histamine H₁ receptors, characterization (bovine retina), **135**, 445
- Pyrimidine**
 moiety in thiamine, methyl carbon atom, origin during thiamine biosynthesis (*Salmonella typhimurium*), **134**, 1136
- Pyrimidine dimers**
 -dependent cleavage of ssDNA by T4 uv endonuclease, **141**, 204
 DNA, cell extracts capable of excision, effect on 4-nitroquinoline-1-oxide adducts, **140**, 775
- Pyrophosphate**
 role in sucrose metabolism in sucrose-importing plants, **141**, 440
- Pyroloquinoline quinone**
 9-carboxyl group, essential role as prosthetic group in formation of D-glucose dehydrogenase holoenzyme, **139**, 1279
 tricyclic, bound to methylamine dehydrogenase, structure and linkage (*Methylobacterium* sp.), **141**, 562
- Pyruvate carboxylase**
 in glutamine utilization pathway, intracellular distribution in lymphocytes (rat), **138**, 318
- Pyruvate dehydrogenase**
 activation by insulin concomitant with insulin binding and degradation in phytohemagglutinin-stimulated T cells (human), **134**, 412
 phosphorylation, effect of estrone treatment (genetically diabetic mouse), **139**, 56
- Pyruvate kinase**
 in glutamine utilization pathway, intracellular distribution in lymphocytes (rat), **138**, 318
 phosphorylation by Ca²⁺/calmodulin-dependent protein kinase: characterization of two phosphorylation sites (rat liver), **139**, 1017
 stimulation by insulin (rat hepatocytes), **134**, 292
- [Pyruvate kinase]-phosphatase
 stimulation by insulin (rat hepatocytes), **134**, 292
- Pyruvate oxidase**
 monoclonal antibodies, characterization: enzyme conformation effects on antibody-induced inhibition (*Escherichia coli*), **137**, 884
- Pyruvate transporter**
 α -cyanocinnamate-sensitive, extraction from mitochondria and reconstitution (castor bean), **140**, 1036
- 6-Pyruvoyl-tetrahydropterin synthase
 purification and characterization (human liver), **134**, 646
- Q**
- Quercetin**
 inhibition of glutathione S-transferase isoenzymes from human and rat tissues, comparison, **141**, 1170
- Quinine**
 effect on mitochondrial K⁺ and Mg²⁺ fluxes (rat liver), **135**, 830
- Quinuclidinyl benzilate**
 binding to muscarinic receptors, effect of guanine nucleotides (rat heart), **134**, 172
- R**
- R 5020, *see* 17,21-Dimethyl-19-nor-4,9-pregnadiene-3,20-dione
- Radioimmunoassay**
 apolipoprotein E, with solid-phase staphylococcal protein A, use of pooled plasma as secondary standard (human), **138**, 455
 atrial natriuretic factor
 in plasma (human), **139**, 287, 1215
 in plasma and urine (human), **137**, 231
 comparative immunoremoval step, in opiate concentration estimation (bovine tissues), **137**, 316
 peptide B distribution in brain (rat), **139**, 1024
- Reduction**
 L-ascorbate oxidase, effect on active site-associated Cu²⁺ (cucumber), **135**, 644
 vanadate by microsomal redox system (rat liver), **141**, 171
- Renin**
 inhibition by angiotensinogen peptide analogs, effect of peptide chain length (monkey), **139**, 982
 release by glomeruli, effect of atrial natriuretic factor (rat kidney), **138**, 496
 renal, purification and N-terminal sequence (human), **139**, 446
- Renin inhibitors**
 structure-activity relationship (human kidney), **134**, 71
- Reoviruses**
 serotype 1 Lang strain
 s1 mRNA encoding minor capsid polypeptides σ 1a and nonstructural σ 1bNS, molecular cDNA cloning and nucleotide sequence, **140**, 508
 s3 mRNA encoding σ polypeptide, cDNA cloning and nucleotide sequence (human), **139**, 845
 s4 mRNA molecular cloning and nucleotide sequence (*Escherichia coli*), **136**, 183
- Reperfusion injury**
 myocardial, allopurinol beneficial action, mechanism, **137**, 76
- Replication**
 autonomous, mitochondrial DNA fragment from *Lentivirus edodes* cloned in *Saccharomyces cerevisiae*, **138**, 1110
- Resorufin**
 inhibition of benzo[a]pyrene metabolism and mutagenesis *in vitro* (rat hepatic microsomes), **136**, 555
- Respiration**
 α -adrenergic-stimulated, in brown adipocytes, mediatory role of guanyl nucleotide-binding protein, evaluation

Subject Index

- with pertussis toxin (hamster), **135**, 823
- cellular and mitochondrial, effects of 1-methylcyclohexanecarboxylic acid (murine cerebral cortex, neuroblastoma), **140**, 789
- Restriction mapping
 - apolipoprotein E and CI gene sequences on chromosome 19 (human), **136**, 876
 - 21-hydroxylase genes (human), **136**, 722; *erratum*, **138**, 502
- Reticulocytes
 - adenylate cyclase system, β -adrenergic agonist stimulation, desensitization induced by tetradecanoyl phorbol acetate (rat), **138**, 125
- Reticuloendotheliosis virus
 - strain T-transformed bone marrow cells, identification of *v-rel* oncogene-encoded protein (chicken), **134**, 716
- Retina
 - dystrophy, associated reduction in rhodopsin phosphorylation (dog), **135**, 1022
 - protein expression after optic nerve crush (goldfish), **138**, 1177
 - rod outer segments, acyl transferase and fatty acid CoA synthetase identification (bovine), **138**, 988
- Retinoic acid
 - effect on
 - DNA synthesis in keratinocytes, role of extracellular Ca^{2+} (murine epidermis), **138**, 483
 - P150 phosphorylation in adriamycin-resistant HL60 cells, **138**, 9
 - induced teratocarcinoma cell differentiation, specific expression of homoeobox-containing genes, **137**, 520
 - inhibition of
 - interferon induction by transformed L-929 and WISH cells, **138**, 47
 - potassium channels in lymphocytes (human), **138**, 560
 - photolabile aromatic analogs, biological activity and covalent binding to cellular retinoic acid-binding protein, **135**, 124
- 13-*cis*-Retinoic acid
 - anticarcinogenic effect on mammary tumor induction by 7,12-dimethylbenz[*a*]anthracene, relationship to cAMP-dependent protein kinase (rat), **135**, 116
- Retinol
 - binding protein, structural changes induced by retinol removal: molecular dynamics study (human), **139**, 564
- Retinyl-palmitate hydrolase
 - quantitation by HPLC (bovine brain, eye, liver, muscle), **134**, 1209
- Retroviruses
 - endogenous A-type, *pol* gene, relationship to mRNA from colon tumor and surrounding mucosa (human), **139**, 1071
 - neurotropic, effect of chronic infection on PC-12 cell function, **140**, 706
 - purified, detection of calmodulin-like activity, **141**, 1077
- Reverse transcriptase, *see* RNA-directed DNA polymerase
- Rhodamine 123
 - inhibition of
 - calmodulin activity and cellular proliferation (rat astrocytoma cells), **135**, 574
 - mitochondrial protein synthesis (rat liver, tumor cells), **137**, 759
- Rhodopsin
 - like pigment, flash spectrophotometric identification in *Halobacterium halobium*, **139**, 389
 - phosphorylation, reduction in retinal dystrophy (dog), **135**, 1022
- Riboflavin
 - biosynthesis: 6,7-dimethyl-8-ribityllumazine formation by heavy riboflavin synthase (*Bacillus subtilis*), **139**, 1111
 - phototoxicity evaluation using *Paramecium aurelia*, **139**, 79
- Riboflavin synthase
 - heavy, formation of 6,7-dimethyl-8-ribityllumazine in riboflavin biosynthesis (*Bacillus subtilis*), **139**, 1111
- Ribonuclease
 - associated activity of cytotoxic eosinophil cationic protein, **139**, 1239
- Ribonuclease T₁
 - L-glutamic acid-58 modification to L-glutamine and L-aspartic acid (*Escherichia coli*), **138**, 789
- D-Ribose
 - inhibition of DNA repair synthesis in lymphocytes (human), **138**, 673
- Ribose 1,5-bisphosphate
 - effect on 6-phosphofructo-1-kinase and fructose-1,6-bisphosphatase (rat liver), **138**, 159
- Ribose-5-phosphate isomerase
 - and phosphoribulokinase, association with ribulose-1,5-bisphosphate carboxylase (pea), **139**, 947
- Ribulose bisphosphate
 - at high concentrations, effect on kinetic parameters of ribulose-1,5-bisphosphate carboxylase/oxygenase (spinach), **134**, 506
- Ribulose-bisphosphate carboxylase
 - association with phosphoriboisomerase and phosphoribulokinase (pea), **139**, 947
 - kinetic parameters at high protein concentrations (spinach), **134**, 506
 - subunit B, isolation with alkaline buffer (spinach leaf), **138**, 118
- Ribulose-bisphosphate carboxylase/oxygenase, *see* Ribulose-bisphosphate carboxylase
- Ricin A
 - IgG conjugate, mannose receptor-mediated uptake by hepatic nonparenchymal cells, measurement (rat), **137**, 892
- Rifampicin, *see* Rifampin
- Rifampin
 - inhibition of EDTA-induced autolysis (*Escherichia coli*), **134**, 338
- RNA
 - association with unactivated glucocorticoid receptor complex (human lymphoid cells), **140**, 204
 - cytoplasmic, complementary to pre-albumin intron V, detection (rat liver), **136**, 989
 - DNA mixed polymer, incorporation of cytidine diphosphate (permeabilized hamster cells), **135**, 340
 - hepatitis B virus, produced in HEP 3B cells, analysis (human), **138**, 131
 - infectious transcripts, enzymatic synthesis from cloned cDNAs of cucumber mosaic virus satellite, **135**, 290
 - messenger
 - albumin, hepatocyte levels, regulation by dexamethasone, **139**, 1071

Subject Index

- methasone and estrogen *in vitro* (*Xenopus laevis*), **136**, 8
- alcohol dehydrogenase subunits α , β , and γ , size heterogeneity and developmental changes (human), **140**, 1020
- aldolase A, expression in adult and fetal tissues and hepatoma (human), **134**, 1093
- δ -aminolevulinic synthase, hepatic levels, effects of testosterone and hemin (chicken embryo), **140**, 81
- apolipoprotein E, monocyte-macrophage levels, reduction in apolipoprotein E deficiency (human), **134**, 937
- atrial natriuretic peptide, regulation by glucocorticoids *in vivo* (rat heart), **139**, 1047
- c-myc, accumulation in cultured thyroid cells, enhancement by epidermal growth factor, forskolin, phorbol ester, and TSH (dog), **141**, 1066
- collagen type X
expression during development, spatiotemporal changes (embryonic chick sternum), **137**, 1125
translation *in vitro* in presence of pancreatic microsomes (chicken), **136**, 109
- from colon tumor and surrounding mucosa, relationship to *pol* gene of endogenous A-type retrovirus (human), **139**, 1071
- creatine kinase B from canine brain, nucleotide sequence: coding and 3' noncoding region homology among vertebrate species, **138**, 392
- cytochrome P-450, synthesis induction by *N*-alkanes (*Candida maltosa*), **136**, 1148
- DNA polymerase β , nucleotide sequencing via cDNA cloning (human), **136**, 341
- epidermal growth factor receptor, characterization of rodent EGF receptor transcripts (mouse), **141**, 1109
- globin, storage during protein synthesis inhibition by heme deprivation (rabbit), **134**, 1048
- α_2 -globulin, phenobarbital-induced: characterization of cDNA clone (rat liver), **134**, 1182
- glucoamylase, translation *in vitro* (rat intestine), **134**, 37
- histone, isolation and characterization (murine testis), **136**, 1116
- insulin and insulin receptor, nucleotide sequences: occurrence of short complementary sequences, **139**, 94
- interleukin 1α , detection in virus-transformed B and T cells (human), **139**, 353
- lectin, β -galactoside-binding, 14-kDa, nucleotide sequence (chicken embryo), **134**, 51
- mitochondrial ADP/ATP-carrier protein, isolation and characterization (bovine heart), **138**, 850
- mitochondrial glutamic-oxaloacetic transaminase, molecular cloning and expression regulation in regenerating liver (rat), **134**, 803
- nerve growth factor
distribution in developing and adult tissues (chicken), **141**, 1116
localization by *in situ* hybridization (murine brain), **138**, 813
- nonstructural σ NS polypeptide of reovirus serotype 1 Lang strain, cDNA cloning and nucleotide sequence (human), **139**, 845
- ornithine decarboxylase, levels in pulmonary tumor cells, relationship to sensitivity to difluoromethylornithine (human), **134**, 572
- phorbol ester-induced protein from melanoma cells, isolation (human), **136**, 122
- pOb24, expression, coupling with growth arrest during adipose conversion of preadipocytes, **137**, 903
- preproenkephalin, levels in neuroblastoma-glioma hybrid cells, synergistic regulation by glucocorticoids and cAMP, **139**, 1
- preproinsulin, pancreatic levels, effect of glucose oral administration (rat), **141**, 1201
- preproparathyroid hormone, role of 5'-terminal cap group and purine in 3' position in recognition of first AUG sequence as site of translation initiation, **140**, 288
- proacrosin, translation *in vitro* (porcine spermatozoa), **138**, 139
- procollagen type I and III, levels in cutaneous wounds treated with helium-neon laser (pig), **138**, 1123
- prolactin-related, detection in pituitary GH₄C₁ tumor cells (rat), **138**, 1015
- reovirus serotype 1 Lang strain
s1, encoding minor capsid polypeptides σ 1a and nonstructural σ 1bNS, molecular cDNA cloning and nucleotide sequence, **140**, 508
s4, molecular cloning and nucleotide sequence (*Escherichia coli*), **136**, 183
- 71-kDa stress protein, heat shock-induced synthesis (canine heart), **134**, 166
- substance P precursor, nucleotide sequence (rat), **139**, 1040
- sucrase-isomaltase, translation *in vitro* (rat intestine), **134**, 37
- 5'-terminal caps, role in translational dormancy (*Physarum polycephalum*), **134**, 469
- transferrin receptor-specific, regulation in differentiating HL-60 cells, **138**, 995
- 2.3kb, detection in diverse neuroendocrine tissues by chromogranin A cDNA (human), **137**, 418
- tyrosinase, translation *in vitro*, **141**, 697
- misaligned double-stranded poly(I) • poly(C₁₂,U), induction of 2',5'-oligoadenylate synthetase accumulation in spleen, liver, kidney, and brain (mouse), **138**, 40
- mouse mammary tumor virus, cytoplasmic, destabilization by heat shock, prevention by cycloheximide, **137**, 1028
- poly(A)⁺, uv light-induced crosslinking with phosphoproteins from free polyribosomes, effects of transcription and translation inhibitors, **136**, 356
- protein complexes, formation in 4-thiouridine-exposed and subsequently illuminated CV-1 cells, **141**, 847
- ribosomal
5 S
accumulation and specific cleavage in oviduct isthmus (laying hen), **138**, 1405
gene-specific transcription factor, modification of DNA linking number, **134**, 1086
nucleotide sequence (*Thermococcus celer*), **134**, 1167
gene number and organization (*Mycobacterium* sp.), **136**, 1136
- 16 S, 3'-terminal fragment, preparation and nucleotide sequence (*Streptomyces aureofaciens*), **137**, 80
- synthesis, effect of RU 38486 (rat thymocytes), **134**, 44
- transfer, initiator methionine, recognition by human autoimmune antibody (yeast), **139**, 275

Subject Index

- U3
 encoding gene, sequence homology with vertebrate U3 RNA (*Drosophila melanogaster*), **138**, 512
 small nucleolar, gene isolation and characterization (human), **137**, 1133
- RNA-directed DNA polymerase
 murine leukemia virus, inhibition by 2',3'-di-deoxythymidine 5'-triphosphate and 3'-modified derivatives, **140**, 498
- RNA polymerase
 type II, subunit functional properties, analysis with monoclonal antibodies (bovine), **136**, 281
- Rolipram
 in affinity chromatography of cAMP-specific phosphodiesterase (rat heart), **138**, 205
- Rous sarcoma virus
 transformation of fibroblasts
 induction of ribosomal protein S6 kinase (chicken embryo), **137**, 702
³¹P NMR spectra (chicken, rat), **138**, 699
- RU 38486, *see* 17β-Hydroxy-11β-(4-dimethylaminophenyl)-17α-(1-propynyl)-estra-4,9-diene-3-one
- Ryanodine receptors
 complex with Ca²⁺, solubilization and separation from Ca²⁺-ATPase, **139**, 235
- S**
- Saccharin
 stimulation of protein kinase C (rat hepatocytes), **135**, 33
- Saccharomyces cerevisiae*
 acid phosphatase gene expression, effects of vector type, host strain, and transcription terminator sequence, **140**, 602
 centromeric plasmids, application as shuttle vectors, **135**, 915
 fatty acid auxotroph grown in presence of oleic acid, lectin release, **136**, 596
- Safranin
 as probe for mitochondrial membrane potential, side effects on Ca²⁺ transport (rat liver), **135**, 189
- Salinomycin
 effect on intracellular Ca²⁺ levels and contractile activity in isolated perfused heart (rabbit), **139**, 816
- Salla disease
 associated defective lysosomal metabolism of sialic acid, **137**, 175
- Salmonella minnesota*
 interaction with lymphocytes, energy dependence and role of thiol groups, **137**, 222
- Salt solutions
 physiological, composition and ionic strength, relative importance to association of mitochondrial inner membrane surface with creatine kinase (rat heart), **139**, 1262
- Sarcolemma
 detection of α-adrenergic receptors (rat skeletal muscle), **136**, 1071
- Sarcoplasmic reticulum
 interaction with fura-2-am (rabbit skeletal muscle), **138**, 1153
- transmembrane pH equilibration rate in isolated vesicles, measurement (rabbit skeletal muscle), **140**, 934
- Satellite cells
 dystrophic, altered protein kinase C distribution, reversal by liposome-delivered phospholipids (mouse), **137**, 752
- Satellite tobacco necrosis virus
 and maize streak and wheat dwarf viruses, coat protein sequence homology, **134**, 372
- Schiff base
 retinal, -carboxylic acid hydrogen bonds, proton transfer thermodynamics, **138**, 819
- Secretin
 induced damage in pancreatic acinar membranes, role of calcium (rat), **139**, 530
- Selenium
 enhancement of hematoporphyrin derivative phototoxicity (murine bladder tumor cells), **140**, 1086
- Semen coagulum
 interaction with sialoglycoprotein from seminal vesicle secretions (rat), **136**, 753
- Seminal plasma
 identification of insulin-like growth factor II (human), **140**, 536
- Seminal vesicles
 glutathione transferase expression, effect of xenobiotics (rat), **136**, 1057
- Semiquinones
 methoxatin, pulse radiolysis, **139**, 955
- Serine
 occurrence in linkage region of polygalactosamine-containing glycoproteins (*Cordyceps ophioglossoides*), **140**, 350
 -rich proteins, gene expression, modulation by glucose (*Saccharomyces cerevisiae*), **138**, 297
- Serine hydroxymethyltransferase
 crystalline, purification and characterization (*Hyphomicrobium methylovorum*), **139**, 71
- Serine kinase
 insulin-stimulated, detection in extracts from IM-9 lymphocytes, **139**, 179
- Serine proteinase
 cleaving α and β chains of C3, purification from erythrocyte membranes and partial characterization (human), **140**, 1113
- Serotonin
 brain levels, effect of sustained exercise (rat), **137**, 149
 metabolism, role of hepatic alcohol dehydrogenases (human), **139**, 1009
 release from platelets
 amiloride and cytochalasin B effects (human), **140**, 329
 cGMP effects (human), **135**, 1099
 platelet-derived growth factor effects (human), **135**, 52
- Serotonin receptors
 effects of swainsonine and tunicamycin, comparison (NCB-20 cells), **134**, 1387
- Sertoli cells
 coculture with Leydig cells, effect on steroidogenesis (swine), **134**, 957
 detection of FSH-suppressible inhibiting factor of testicular steroidogenic activity (pig), **139**, 169
 immature, secretion of androgen-binding protein and es-

Subject Index

- tradiol-17 β , *in vitro* effects of germ cells (rat), **141**, 861
- Serum**
- effect on inhibition of DNA synthesis in leukemia cells by *cis*- and *trans*-diamminedichloroplatinum(II) (mouse), **138**, 631
 - masked antibodies, naturally occurring, recognition of mitotic spindle apparatus component (mouse), **136**, 145
 - mediation of increased atrial natriuretic factor synthesis by cardiac atrioocytes (rat), **140**, 151
- Serum albumin**
- amino acid sequence, homology with kinetensin (human), **136**, 983
 - and dinitrophenylated derivatives, ultrathin films, direct current conductivity measurements (bovine), **133**, 1152; *erratum*, **135**, 1150
 - effect on endothelium-dependent relaxation (rat aortic rings), **138**, 1211
 - tyrosine phosphorylation by human erythrocytic casein kinase A (bovine), **139**, 855
- Sialic acid**
- lysosomal metabolism, defect in Salla disease (rat liver), **137**, 175
- Sialoglycoconjugates**
- synthesized upon T-cell activation, selective covalent modification (human), **135**, 473
- Sialoglycopeptides**
- cell-surface, inhibition of 3T3 cell protein synthesis, effects of calcium and sodium ionophores (bovine), **136**, 976
 - inhibition of 3T3 cell protein synthesis, role of cell surface interaction (bovine), **134**, 1350
- Sialoglycoproteins**
- detection in seminal vesicle secretions and interaction with semen coagulum (rat), **136**, 753
- Signal recognition particle**
- functions, effect of signal sequence removal from preproinsulin (carp), **134**, 790
- Simian virus 40**
- DNA replication, inhibition by 1-*N*-acyl-3"-*N*-(trifluoroacetyl)kanamycin *in vitro* (CosI cells), **136**, 322
 - T antigen, purification by DNA-Sephrose affinity chromatography, **134**, 915
- Sinacalide**
- induced damage in pancreatic acinar membranes, role of calcium (rat), **139**, 530
 - sulfation by sulfotransferase (*Eubacterium* A-44), **140**, 38
- SKF 102229, *see* 3-Methyl-6-chloro-9-azido-1H-2,3,4,5-tetrahydro-3-benzazepine
- Skin**
- carcinogenesis, evaluation of cholera toxin as promoter (mouse), **137**, 486
 - human, grafted onto athymic nude mice, carcinogen metabolism: model system for analysis of carcinogenesis, **138**, 33
- SM-9064**
- pharmacological profile (mouse, rat), **138**, 540
- Sodium**
- amiloride-sensitive transport, inhibition by atrial natriuretic factor and cGMP (LLC-PK₁ cells), **134**, 852
 - dependent concentrative nucleoside transport in cultured intestinal epithelial cells (rat), **140**, 1028
 - high dietary levels, induced increase in vasopressin release, inhibitory role of hypothalamic atrial natriuretic polypeptide (rat), **139**, 1285
 - hydrogen exchange
 - in basolateral membranes, detection by fluorescence quenching (rabbit small intestine), **134**, 827
 - osmotic activation, protein kinase C role (rat thymic lymphocytes), **134**, 8
 - in platelets, induction by 1-oleoyl-2-acetyl glycerol and phorbol ester (human), **141**, 13
 - inorganic phosphate, cotransport in renal epithelium, inhibition by phosphonoformic acid (human), **139**, 679
 - ion pump inhibitor, purification (human urine), **131**, 1234; *erratum*, **134**, 452
 - potassium-chloride cotransport system, biochemical and pharmacological properties (chicken cardiac cells), **134**, 326
 - potassium pump
 - associated antigens, binding to antibodies, enhancement by intracellular K⁺ (ovine erythrocytes), **134**, 982
 - in erythrocytes from cystic fibrosis patients and heterozygotes, decreased affinity for K⁺, **137**, 1021
 - regulation of α_2 -adrenoreceptors, effect of 4,4'-diisothiocyano-2,2'-stilbene disulfonic acid (human platelets), **141**, 222
- Sodium benzoate**
- effect on ammonia toxicity (rat), **138**, 369
- Sodium butyrate**
- differential effects on cAMP phosphodiesterase activity of butyrate-sensitive and -resistant mastocytoma cells, **138**, 1030
- Sodium carboxylcellulose**
- oxidation by hydrogen peroxide, role in cell wall breakdown, **141**, 238
- Sodium channels**
- amiloride-sensitive, interaction with phenamil, characterization (porcine renal membranes), **135**, 25
 - detection in myometrial cells with neurotoxins (rat), **137**, 675
 - voltage-dependent, localization with acetylcholine receptor at neuromuscular junction (mouse), **139**, 196
- Sodium chloride**
- dietary intake, alteration, effect on atriopeptin levels in heart and plasma (rat), **137**, 922
 - effect on α -lactalbumin labeling with 3-(trifluoromethyl)-3-(*m*-[¹²⁵I]iodophenyl)diazirine (bovine), **138**, 604
 - modulation of multisubstrate Ca²⁺- and cAMP-independent kinase (bovine aortic smooth muscle), **136**, 789
 - sensitive Dahl rats, measurement of plasma atrial natriuretic factor levels, **136**, 411
- Sodium 3,5-dibromo-4-nitrosobenzenesulfonate**
- spin-trapping of superoxide radicals, **136**, 657
- Sodium fluoride**
- induced cytosolic free calcium increase and thromboxane B₂ generation, inhibition by protein kinase C activation (human platelets), **136**, 381
- Sodium hyaluronate**
- hydration, NMR analysis of oriented sample, **141**, 73
- Somatomedin**
- binding protein
 - from amniotic fluid, in insulin-like growth factor I purification by affinity chromatography (human), **136**, 253
 - high-molecular-weight, purification (human plasma), **135**, 178

Subject Index

- identification of biologically active domain (human placenta), **141**, 263
- growth hormone-induced production during development, relationship to delayed appearance of hepatic growth hormone receptors (rat), **136**, 38
- Somatomedin receptors
 - occurrence in fetal cortical neurons and interactions with insulin and insulin-like growth factors (mouse), **135**, 228
- Somatostatin
 - inhibition of pituitary secretion vesicle fusion with plasma membranes (rat), **139**, 673
 - octacosapeptide, solid-phase synthesis (anglerfish pancreatic islets), **140**, 565
 - pancreatic secretion, cysteamine effect (rat), **134**, 1291
 - surface-adsorbed, reaction products, mass spectrometry, **134**, 420
- Somatostatin receptors
 - chemical crosslinking to radioligands (rat adrenal cortex), **139**, 757
- Somatotropin
 - biosynthesis in neonatal pituitary, effects of growth hormone-releasing hormone and clonidine, comparison (rat), **138**, 1223
 - dependent insulin-like growth factor-binding protein, from plasma, comparison with amniotic fluid IGF-binding protein (human), **139**, 1256
 - effect on hepatic lipogenic enzymes after chronic administration (hypophysectomized rat), **141**, 78
 - glycosylated, detection in pituitary gland with lectin-binding immunoassay (human), **140**, 491
 - ¹²⁵I-labeled, for receptor studies, preparation by reverse-phase HPLC (human), **134**, 671
 - induced somatomedin production during development, relationship to delayed appearance of hepatic growth hormone receptors (rat), **136**, 38
 - phosphorylated, synthesis and secretion by pituitary glands *in vitro* (rat), **137**, 806
 - release from
 - anterior pituitary cells *in vitro*, suppression by atrial natriuretic factor (rat), **135**, 1035
 - pituitary tumor cells, induction by bacterial adenylate cyclase (rat), **136**, 463
 - specific binding protein in serum, affinity purification and structural characterization (human), **139**, 150
 - 20-kDa variant, receptor binding properties, effect on insulin-like and growth-promoting activities (human), **134**, 159
- Somatotropin receptors
 - hepatic, delayed appearance, relationship to growth hormone-induced somatomedin production during development (rat), **136**, 38
- Sorbinil
 - effect on hyperglycemia-induced sorbitol accumulation and *myo*-inositol depletion in cultured conceptus (rat), **140**, 974
- Sorbitol
 - accumulation in cultured conceptus, hyperglycemia-induced, effects of aldose reductase inhibitors (rat), **140**, 974
- Soybean flour
 - induction of cytochrome *P*-450 (*Streptomyces griseus*), **141**, 405
- Spectrin
 - binding proteins, role in structure of neuronal postsynaptic density, regulation by phosphorylation (rat brain), **138**, 59
- Spermatozoa
 - calcium levels before/after *in vitro* capacitation (rabbit), **134**, 664
 - detection of guanine nucleotide-binding protein ADP-ribosylated by pertussis toxin (invertebrates, vertebrates), **138**, 728; *addendum*, **140**, 773
- Spermidine
 - erythrocyte-associated, intracellular location (human), **134**, 1365
 - protein phosphatase activation (HeLa cell nuclei), **134**, 1372
- Spermidine synthase
 - N*-chlorosulfonyl dicyclohexylamine inhibitor, synthesis and effects on Molt4B cells, **141**, 718
- Spermine
 - erythrocyte-associated, intracellular location (human), **134**, 1365
 - interaction with polyphosphoinositide-containing liposomes and *myo*-inositol 1,4,5-triphosphate, **135**, 58
 - protein phosphatase activation (HeLa cell nuclei), **134**, 1372
- Sphingolipids
 - activator protein 1, encoding cDNA, isolation and characterization (human), **134**, 989
 - inhibition of insulin- and phorbol ester-stimulated uptake of 2-deoxyglucose by fibroblasts (mouse), **138**, 463; *erratum*, **139**, 852
- Spindle apparatus
 - mitotic, component recognition by naturally occurring masked antibodies in serum (mouse), **136**, 145
- Spiperone receptors
 - effects of swainsonine and tunicamycin, comparison (NCB-20 cells), **134**, 1387
- Spironolactone
 - inactivation of hepatic cytochrome *P*-450 (rat), **136**, 1162
- Splenocytes
 - lipopolysaccharide-stimulated, interleukin 3-like activity detection in supernatants (mouse), **138**, 1322
- Squalene synthetase
 - hepatic microsomal, stimulation by cholestyramine, effect of diurnal cycle (rat), **138**, 335
- Stallimycin
 - synthetic analog, DNA-binding characteristics, **140**, 626
- Starch
 - in concanavalin A solutions, detection and quantitation with electrical oscillator, **137**, 978
- Starch(bacterial glycogen) synthase
 - activation by glucose, role of glucose 6-phosphate (rat hepatocytes), **141**, 1195
 - behavior as substrate for hepatic synthase phosphatases, effect of phosphorylation by different protein kinases, **139**, 1033
 - phosphorylation and inactivation by mezerein and phorbol esters (rat hepatocytes), **134**, 113
- Statil
 - effect on hyperglycemia-induced sorbitol accumulation and *myo*-inositol depletion in cultured conceptus (rat), **140**, 974

Subject Index

- Status epilepticus
bicuculline-induced, associated enhancement of inositol turnover (rat brain), **136**, 827
- Staurosporine
inhibition of protein kinase C (rat brain), **135**, 397
- 1-Stearyl-2-stearoylaminodeoxy phosphatidylcholine
reversible inhibition of phospholipase A₂ (cobra venom), **137**, 587
- Steroid hormone receptors
relationship to heat shock protein hsp 90, **138**, 1298
- Steroidogenesis
in bovine adrenocortical cells, effect of human α -atrial natriuretic factor, **137**, 657
in Leydig cells
effect of coculture with Sertoli cells (swine), **134**, 957
regulation by atrial natriuretic factor (mouse), **138**, 399
stimulation by insulin-like growth factor I (rat), **137**, 950
testicular, FSH-suppressible inhibiting factor, detection in extract from Sertoli cells (pig), **139**, 169
- Steroids
binding sites on
estrogen and glucocorticoid receptors, computer-based search, **139**, 281
glucocorticoid receptors and triamcinolone acetonide antibodies, comparison (rat), **136**, 470
- Steroid sulfatase, *see* Steryl-sulfatase
- Sterol 14-reductase
role in lanosterol 14 α -demethylation (yeast), **134**, 659
- Sterols
biosynthesis, ketoconazole effects (*Trypanosoma cruzi* epimastigotes), **136**, 851
C-24 transalkylation, regulation, role in development (*Gibberella fujikuroi*), **139**, 410
- Steryl-sulfatase
rapid degradation in multiple sulfatase deficiency (human), **135**, 389
- trans*-Stilbene oxide
effect on glutathione transferase expression in seminal vesicles and pituitary gland (rat), **136**, 1057
- Stomach
antrum, production of active and inactive forms of β -endorphin (rat), **134**, 14; *erratum*, **138**, 502
- Streptavidin
¹²⁵I-labeled, selective labeling of functional groups on erythrocyte membrane proteins and/or glycoproteins (human), **136**, 80
- Streptomyces lividans*
cloning and expression of α -amylase inhibitor gene from *S. griseosporus*, **141**, 1099
- Streptomycin
effect on EDTA-induced autolysis (*Escherichia coli*), **134**, 338
- Streptozocin
induction of diabetes
associated decreased expression of epidermal growth factor receptors (rat), **137**, 258
associated insulin receptor autophosphorylation and kinase activity, effect of short fast (rat), **140**, 850
- Streptozotocin, *see* Streptozocin
- Stress
induced suppression of cardiac protein synthesis, role in subsequent hypertrophy (rat), **135**, 857
- Styrene
epoxidation by chloroperoxidase (*Caldariomyces fumago*), **136**, 778
- Substance K
induced DNA synthesis, correlation with *myc*-gene transcript levels (quail fibroblasts), **137**, 167
- Substance P
antagonist, inhibition of specific binding and mitogenic effects of vasopressin and bombesin-related peptides (3T3 cells), **137**, 135
precursor, mRNA nucleotide sequence (rat), **139**, 1040
- Substance P receptors
binding to phospholipase C, stimulation by guanine nucleotide-dependent regulatory protein (rat parotid gland), **136**, 362
- Subtilisin
effect on microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
- Succinate
cyanide-resistant oxidation in submitochondrial particles, comparison with duroquinol oxidase activity (*Arum maculatum*), **135**, 677
- Succinate dehydrogenase
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
purification and characterization (*Desulfovibrio multispirans*), **135**, 1000
- Succinyl-CoA synthetase
phosphorylation, effect of estrone treatment (genetically diabetic mouse), **139**, 56
- Sucrase-isomaltase
mRNA, translation *in vitro* (rat intestine), **134**, 37
- Sucrose
cellular location after hepatic endocytosis, analysis by isopycnic centrifugation in sucrose gradients (rat), **136**, 504
metabolism in sucrose-importing plants, role of pyrophosphate and activation by fructose 2,6-bisphosphate, **141**, 440
- Sulfatide
role in protein kinase C activation by tumor promoters (bovine brain), **138**, 153
- Sulfation
chondroitin sulfate, inhibition by protein kinase C activators (HL-60 cells), **135**, 909
proteins, inhibition by chlorate (rat PC12 cells), **141**, 870
tyrosine, sites in proteins, characterization and criteria for prediction of occurrence, **141**, 326
- Sulfite
detoxication, role of glutathione (bovine kidney), **136**, 1036
- Sulfite reductase
low-spin, isolation and characterization (*Desulfuromonas acetoxidans*, *Methanosarcina barkeri*), **141**, 1032
- Sulfmyoglobin
multiple forms, detection by ¹H NMR and chemical properties, **135**, 309
- Sulfolobus solfataricus*
occurrence of cAMP, **136**, 1078
- Sulfometuron methyl
cytochrome P-450-dependent metabolism, induction by sulfonyleurea (*Streptomyces griseolus*), **140**, 650

Subject Index

T

Sulfonylurea

induction of cytochrome *P*-450-dependent metabolism (*Streptomyces griseolus*), **140**, 650

Sulfotransferase

sulfation of tyrosine-containing peptides and proteins (*Eubacterium* A-44), **140**, 38

Sulfoxidase

oxidation of methiocarb (soybean cotyledonal microsomes), **135**, 922

Superoxide dismutase

activity in hepatic, muscular, and pulmonary tissues of chickens with inherited muscular dystrophy, **134**, 550

associated activity of pulsed cytochrome oxidase (bovine heart), **136**, 433

mimetic activity of cytokinin-copper(II) complexes, **137**, 372

plasma levels, effect of *Plasmodium vinckei* infection (mouse), **134**, 152

prevention of doxorubicin-induced killing of breast cancer cells (human), **135**, 330

Superoxide radicals

formation from 1-methyl-4-phenylpyridinium ion in presence of NADPH cytochrome *P*-450 reductase and oxygen, **135**, 583

-generating protein, photolabeling with arylazido NADP⁺ analog (bovine neutrophils), **139**, 85

3-hydroxykynurenine role as scavenger (*Aldrichina grahami*), **139**, 666

mitochondrial formation, role of ubiquinone and cytochrome *b*₅₆₆ (rat heart), **138**, 533

photogeneration by chlorophyll *a* in water/acetone, ESR studies (spinach), **139**, 1124

production by neutrophils, induction by

1-oleoyl-2-acetyl glycerol

enhancement by extracellular Ca²⁺ and cytochalasin B (human), **134**, 690

after incorporation into pH-sensitive liposomes (guinea pig), **136**, 242

tumor necrosis factor (human), **137**, 1094

role in lactoperoxidase-catalyzed hydrogen peroxide metabolism and irreversible enzyme inactivation, **139**, 327

spin-trapping

[¹⁷O]oxygen hyperfine structure, **141**, 622

by sodium 3,5-dibromo-4-nitrosobenzenesulfonate, **136**, 657

Surfactants

pulmonary, chemical and biophysical properties of associated proteins (bovine), **135**, 527

Swainsonine

and tunicamycin, comparative effects on various neuroreceptors (NCB-20 cells), **134**, 1387

Synaptosomes

calcium levels, kyotorphin-induced increase (rat brain stem), **137**, 897

magnesium uptake and release, characterization and properties (guinea pig cerebral cortex), **136**, 906

Synthase phosphatase

hepatic, components S and G, behavior of glycogen synthase as substrate, effect of different protein kinases, **139**, 1033

Talin

phosphorylation by protein kinase C, **134**, 1276

Tamoxifen

antiestrogenic and antiproliferative effects, identification of mediating growth-promoting histamine receptor as antiestrogen binding site (rat liver), **134**, 601

and buserelin and estradiol, combined effects on breast tumor growth *in vitro* (human), **140**, 550

Taurine

conjugation with cholic acid by hepatic peroxisomes (rat), **138**, 167

protection against experimental arterial calcinosis (mouse), **140**, 679

TCDD. *see* 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin

Teleocidin

activation of protein kinase C, role of sulfatide (bovine brain), **138**, 153

Temperature

effect on

elastin and polypentapeptide length (bovine ligamentum nuchae), **141**, 749

F₁-ATPase kinetics and structure (bovine cardiac mitochondria), **136**, 891

induction of heat shock and glucose-regulated proteins (hamster fibroblasts), **140**, 881

steady-state cultures of *Escherichia coli*, **135**, 708

high, role in induction of high Mg²⁺-ATPase activity of coupling factor 1 by octyl glucoside (spinach chloroplasts), **141**, 636

increase, as cell cycle monitoring parameter (murine breast cancer cells), **140**, 414

Teniposide

peroxidative free radical formation and *O*-demethylation, **135**, 215

stimulation of topoisomerase II-induced cleavage sites in *c-myc* proto-oncogene *in vivo* (N417 cells), **137**, 154

Testosterone

developmental regulation of glutathione *S*-transferase (mouse), **140**, 581

effect on δ -aminolevulinate synthase mRNA levels (embryonic chicken liver), **140**, 81

production in Leydig cells, regulation by atrial natriuretic factor *in vitro* (mouse), **138**, 399

renal response, role of putrescine (mouse), **138**, 771

Tetanus toxin

inhibition of macrophage secretion, role of cytosolic Ca²⁺ homeostasis (human), **135**, 16

-synthetic pre-S viral peptide-murabutide conjugates, in production of antibodies recognizing hepatitis B virus surface antigens, **140**, 557

3,4,3',4'-Tetrachlorobiphenyl

enhancement of cardiac prostaglandin release *in vitro* (chick embryo), **136**, 582

2,3,7,8-Tetrachlorodibenzo-*p*-dioxin

effect on protein kinase C (murine thymoma cells), **140**, 267

enhancement of cardiac prostaglandin release *in vitro* (chick embryo), **136**, 582

induced myelotoxicity, antagonism by 1-amino-3,7,8-trichlorodibenzo-*p*-dioxin (murine bone marrow), **139**, 747

Subject Index

- 12-*O*-Tetradecanoylphorbol-13-acetate
and A23187, synergistic stimulation of lipoxygenase-mediated arachidonic acid metabolism (murine macrophages), **140**, 861
- activation of protein kinase C
in presence of calcium and ganglioside GM3, **138**, 865
role of sulfatide (bovine brain), **138**, 153
- alteration of protein kinase C catalytic properties
in bovine adrenocortical cells, human lymphoid cells, and rat fibroblasts, **134**, 1031
characterization (rat-1 cells), **138**, 1283
- and calcium, synergistic effect on K⁺-induced catecholamine release from chromaffin cells (bovine adrenal gland), **140**, 990
- effect on
carbamylcholine- and high K⁺-induced elevation of intracellular free calcium levels in cultured chromaffin cells, comparison (bovine adrenal), **139**, 903
formyl-methionyl-leucyl-phenylalanine-induced inositol phospholipid metabolism and Ca²⁺ mobilization (guinea pig neutrophils), **139**, 1272
- 25-hydroxyvitamin D₃ metabolism in kidney (chicken), **139**, 495
- P150 phosphorylation in adriamycin-resistant HL60 cells, **138**, 9
- parathyroid hormone secretion in electroporated parathyroid cells (bovine), **139**, 1233
- enhancement of
c-myc mRNA accumulation in cultured thyroid cells (dog), **141**, 1066
malignant transformation of embryonic fibroblasts, inhibition by 3-aminobenzamide (mouse), **134**, 1334
- inactivation of glycogen synthase by phosphorylation (rat hepatocytes), **134**, 113
- induction of
adenylate cyclase system desensitization to β -adrenergic agonist stimulation (rat reticulocytes), **138**, 125
- adrenocorticotropin and prostaglandin E₂ secretion from pituitary, role of Ca²⁺: effect of dexamethasone (rat), **138**, 323
- atrial natriuretic peptide secretion, regulation by forskolin, Bay k 8644, and 8-bromo-cyclic GMP (rat heart), **139**, 266
- cytosolic protein kinase C desensitization, effect on cell growth (human breast cancer), **135**, 65
- 8-hydroxydeoxyguanosine accumulation in DNA of exposed granulocytes (human), **137**, 841
- prostaglandin biosynthesis in 3T3 fibroblasts, relationship to ornithine decarboxylase induction (mouse), **139**, 186
- protein kinase C translocation
and down-regulation (MCF-7 cells), **137**, 1159
in HL-60 cells, **135**, 1144
- protein from melanoma cells, mRNA isolation (human), **136**, 122
- sodium-hydrogen exchange in platelets (human), **141**, 13
- transient bipterin and neopterin accumulation in human T cells and various cell lines, **141**, 494
- inhibition of chondroitin sulfate sulfation (HL-60 cells), **135**, 909
- stimulation of
intracellular 48-kDa protein synthesis and phosphorylation in plasminogen activator-secreting melanoma cells (human), **137**, 263
- p82 phosphorylation (murine epidermis), **137**, 766
- phosphate accumulation in cultured renal tubular cells, synergistic effect of A23187 (mouse), **136**, 177
- pp67 phosphorylation (NK-7 cells), **135**, 487
- protein kinase C, enhancement of polyoma virus middle T antigen transforming ability (F111 cells), **140**, 1106
- Tetrahydrobiopterin
cofactor activity for indoleamine 2,3-dioxygenase (murine epididymis), **137**, 1106
- Tetrahydroisoquinoline
endogenous, detection in brain (rat), **140**, 448
- Tetrahydromethanopterin methyltransferase
detection in methane-synthesizing complex (*Methanobacterium thermoautotrophicum*), **136**, 542
- Tetraphenylporphyrin
-iron system, generated hemoprotein-peroxide complex, characterization, **135**, 972
- Tetrathionate reductase
molybdenum detection (*Salmonella typhimurium*), **136**, 577
- Tetrodotoxin
³H-labeled, binding in neuronal and nonneuronal spinal cord cultures (mouse), **138**, 1250
- β -Thalassemia
associated mutation in IVS 1 sequence donor site consensus sequence, identification (human), **139**, 709
- Thebaine
concentration estimation by comparative immunoremoval (bovine tissues), **137**, 316
- Theophylline
inhibition of erythroleukemic cell differentiation (mouse), **134**, 845
- Thiamine
biosynthesis, origin of pyrimidine moiety methyl carbon atom (*Salmonella typhimurium*), **134**, 1136
uptake by hepatic mitochondria, analysis *in vitro* (rat), **141**, 466
- Thioanisoles
peroxidase-dependent oxygen transfer from hydrogen peroxide, **135**, 166
- Thiolate
-hemin complexes, scission of plasmid DNA strands, **136**, 645
- Thiol groups
exchange reaction: penicillamine selenotrisulfides, **135**, 183
role in lymphocyte interaction with *Salmonella minnesota*, **137**, 222
surface, accessibility, alteration by ionophores or OKT-3 antibodies (human lymphocytes), **140**, 999
- Thionein
lead-containing, immunological analysis of related protein (rat liver), **136**, 535
- Thioredoxin reductase (NADPH)
role in reduction of free radicals at epidermal surface (guinea pig, human), **136**, 630
- 4-Thiouridine
-exposed and subsequently illuminated CV-1 cells, detection of RNA-protein complexes, **141**, 847
- Threonine
D- or L-enantiomer-containing peptides, in analysis of pro-

Subject Index

- tein kinase C substrate specificity (rat brain), **134**, 1358
 occurrence in linkage region of polygalactosamine-containing glycoproteins (*Cordyceps ophioglossoides*), **140**, 350
- Thrombin**
 effect on platelet cytosolic Ca^{2+} levels, analysis with fura-2 (human), **139**, 308
 induction of Ca^{2+} flux in fura-2-loaded platelets, kinetics (human), **136**, 1124
 inhibition by benzoxazinones, effects of electron withdrawal and 5-substitution (human), **140**, 928
 -stimulated glycolysis, regulatory role of fructose 2,6-bisphosphate (human platelets), **138**, 666
 type α , diisopropyl fluorophosphate derivative, with dioc-tanoyl glycerol or phorbol myristate acetate, stimulation of DNA synthesis (NIL cells), **141**, 650
 type γ , stimulation of 47-kDa protein phosphorylation, effect on phospholipase A_2 (human platelets), **141**, 459
- Thromboxane A_2**
 -prostaglandin H_2 antagonist binding site, photoaffinity labeling (human platelets), **140**, 128
- Thromboxane B_2**
 sodium fluoride-induced generation, inhibition by protein kinase C activation (human platelets), **136**, 381
- Thymidylate synthase**
 poly G-bound, identification (*Streptococcus faecium*), **139**, 461
- Thymine**
 oxidized residues, specific DNA glycosylase, identification in *Drosophila melanogaster* embryos, **134**, 201
- Thyroglobulin**
 and acetylcholinesterase, sequence and structure homology: functional and clinical significance, **137**, 142
 gene
 activation by thyrotropin and cAMP in hormonally depleted FRTL-5 cells, **141**, 1261
 expression, relationship to gene methylation pattern (human), **134**, 1109
- Thyroid**
 cells in primary culture, c-myc mRNA accumulation, enhancement by epidermal growth factor, forskolin, phorbol ester, and TSH (dog), **141**, 1066
 cellular calmodulin levels and calcium uptake, thyrotropin-stimulated increases (swine), **136**, 306
 cytosol, detection of Ca^{2+} -activated calmodulin-dependent protein kinase (bovine), **140**, 120
- Thyrotropin**
 deglycosylation with peptide:N-glycosidase F and endo- β -N-acetylglucosaminidase F (bovine), **138**, 230
 enhancement of c-myc mRNA accumulation in cultured thyroid cells (dog), **141**, 1066
 induction of
 increases in calcium uptake and calmodulin levels in thyroid cells (swine), **136**, 306
 thyroglobulin gene in hormonally depleted FRTL-5 cells, **141**, 1261
 α -subunit secreted by tumor cell lines, phosphorylation (human), **140**, 143
 β -subunit gene, chromosomal assignment by chromosome sorting with velocity sedimentation and Southern hybridization (human), **134**, 477
- Thyrotropin receptors**
 low- and high-affinity binding components, characterization (porcine thyroid), **137**, 61
 -mediated activation of membrane phospholipases and iodide fluxes, role of pertussis toxin-sensitive G proteins, analysis (FRTL-5 cells), **141**, 1000
 physicochemical properties (guinea pig adipocytes), **135**, 208
- Thyrotropin-releasing hormone**
 analogs, relationship of receptor binding to net excitation of spinal lower motoneurons, hypotheses, **138**, 1184
 effect on Ca^{2+} intracellular levels and translocation (canine thyroid cells), **141**, 569
 GH3 cell K^+ conductance response, decrease after patch recording, reversal by GH3 cell extracts, **137**, 388
 induced GTPase activity, mediation by guanine nucleotide-binding protein (GH3 cells), **138**, 1383
 metabolite cyclo(His-Pro), specific binding to hepatic membranes (rat), **134**, 443
- Thyroxine**
 effect on
 hepatic lipogenic enzymes after chronic administration (hypophysectomized rat), **141**, 78
 mitochondrial ubiquinone levels (rat liver), **138**, 381
 role in atrial natriuretic factor release, analysis by measurement of cardiac and plasma levels in hyper- and hypothyroid rats, **134**, 178
- Thyroxine 5'-deiodinase**
 type II, detection in pineal gland (rat), **137**, 863
- Tight junctions**
 structural models: Forssman glycolipid flow between MDCK epithelial cells, **138**, 1163
- Tin oxide**
 electrode, analysis of electron transfer reaction of cytochrome-c peroxidase, **139**, 1003
- Tissue distribution**
 free D-aspartic acid, measurement (human, rat), **141**, 27
- Tolazoline**
 enhancement of β -adrenergic ligand binding to pineal particulate fractions (rabbit), **138**, 1196; *erratum*, **139**, 1320
- Tolbutamide**
 effect on fructose-6-phosphate 2-kinase and fructose-2,6-bisphosphatase activities (rat liver), **139**, 687
- Topoisomerase I**
 relaxing activity, modulation by proteins binding ssDNA (*Escherichia coli*), **137**, 795
 stimulation by mitoxantrone (human breast cancer cells), **136**, 521
- Topoisomerase II**
 and DNA polymerase α levels in developing neuronal nuclei, independent regulation (rat brain), **138**, 1116
 induced cleavage sites in c-myc proto-oncogene, *in vivo* stimulation by antitumor drugs (N417 cells), **137**, 154
 inhibition by mitoxantrone (human breast cancer cells), **136**, 521
- Toromycin**
 binding to DNA, analysis, **136**, 885
- Toxins, see also specific toxins**
Bacillus sphaericus 1593, larvicidal, purification, **141**, 756
Bacillus thuringiensis var. *israelensis*
 crystalline delta endotoxin, larvicidal activity, role of

Subject Index

- 26-kDa peptide, **136**, 1142
 inactivation by proteases from mosquito larvae, **135**, 902
Clostridium botulinum C₂, component I, ADP-ribosylation of avian nonmuscle actin, **136**, 802
Clostridium difficile B, ADP-ribosylation activity, analysis in fibroblasts (human), **139**, 64
Escherichia coli 0157:H7, Vero cell cytotoxic, purification and physicochemical properties, **139**, 424
 sea anemone, in detection of Na⁺ channels in myometrial cells (rat), **137**, 675
Tityus serrulatus toxin VII, pharmacologic properties, comparison with β -toxin and insect toxins from scorpion venoms, **139**, 296
 tricho-, hexapeptide fragment
 C-terminal, crystal structure, **139**, 590
 crystal structure, **136**, 870
- Transalkylation
 sterol C-24, regulation, role in development (*Gibberella fujikuroi*), **139**, 410
- Transcortin
 epitopes, phylogenetic analysis with monoclonal antibodies (human), **138**, 1037
- Transcription
 activity prediction from relationship of androgen-dependent genes to nuclear matrix (rat prostate), **138**, 1334
 casein gene, hydrocortisone role (rat mammary explants), **134**, 812
 chromatin-like complex of adenovirus serotype 2 major core protein VII and DNA *in vitro*, **136**, 86
 control sequence, detection in cauliflower mosaic virus DNA, **138**, 17
- Transcription factors
 5 S RNA gene-specific, modification of DNA linking number, **134**, 1086
- Transferrin
 stimulation of NADH oxidation (HeLa cells), **135**, 110
- Transferrin receptors
 cycling by lymphoid cells, effect of microtubule assembly inhibition (human), **138**, 1216
 hepatocyte membrane-associated, recovery after perfusion with collagenase (rat), **134**, 711
 transcriptional regulation in differentiating HL-60 cells, **138**, 995
- Transferrin reductase
 diferric, associated activity of hepatocyte plasma membrane (rat), **139**, 1117
- Transformation
 fibroblasts by Rous sarcoma virus, induction of ribosomal protein S6 kinase (chicken embryo), **137**, 702
 Kirsten murine sarcoma virus-mediated
 associated loss of epidermal growth factor binding, mechanism (rat renal cells), **135**, 435
 induction of ouabain sensitivity (HOS cells), **136**, 807
 malignant, radiation-induced and phorbol ester-enhanced, inhibition by 3-aminobenzamide (murine embryonic fibroblasts), **134**, 1334
- Transforming growth factor
 and polyoma virus middle T antigen, effect on secondary cultured embryo cell transformation (rat), **136**, 921
 type β
 associated activity of bovine brain-derived growth factor (NRK cells), **139**, 619
 effect on
 differentiated adrenocortical cell functions (bovine), **139**, 693
 luteinizing hormone receptor expression in ovarian granulosa cells (rat), **139**, 800
 ¹²⁵I-labeled, preparation and binding analysis (pig), **138**, 714
 inhibition of
 alkaline phosphatase proliferation and expression in murine osteoblast-like cells (porcine platelets), **140**, 56
 endothelial cell proliferation, interaction with acidic and basic fibroblast growth factors (bovine aorta), **138**, 476
 proliferation of endothelial cells *in vitro* (bovine aorta), **137**, 295
 masking protein, partial purification and characterization (rat platelets), **141**, 176
 modulation of FSH-induced aromatase activity in granulosa cells, comparison with inhibin (rat), **136**, 969
 role in phenotypic transformation of kidney cells induced by platelet-derived growth factor-like polypeptide (rat), **141**, 1229
 stimulation of
 collagen production and processing (human fibroblasts), **138**, 974
 follicle-stimulating hormone secretion in pituitary monolayer system (rat), **135**, 950
 storage in platelets as latent high-molecular-weight complex (human), **136**, 30
- Transglutaminase, *see* Protein-glutamine γ -glutamyltransferase
- Translation
 collagen type X mRNA in presence of pancreatic microsomes (chicken), **136**, 109
 dormancy, role of mRNA 5'-terminal caps (*Physarum polycephalum*), **134**, 469
 fungal tyrosinase mRNA *in vitro*, **141**, 697
 glucoamylase and sucrase-isomaltase mRNAs *in vitro* (rat intestine), **134**, 37
in vitro systems, efficient and prolonged, detection in etioplasts (cucumber), **135**, 248
- Transthyretin
 variant produced by transgenic mouse, as model of familial amyloidotic polyneuropathy (human), **139**, 794
- TRH, *see* Thyrotropin-releasing hormone
- Triacetyl glycerol
 and water, hydrogen bond formation, detection by high-pressure infrared spectrometry, **134**, 140
- Triacylglycerol lipase
 gene, molecular cloning and nucleotide sequence analysis (*Pseudomonas fragi*), **141**, 185
- Triamcinolone acetonide
 antibodies, and glucocorticoid receptors, steroid-binding domains, comparison (rat), **136**, 470
- Triethylenediamine
 enhancement of singlet oxygen production in hydrogen peroxide-hypochlorous acid reaction, **134**, 777
- 3-(Trifluoromethyl)-3-(*m*-iodophenyl) diazirine
¹²⁵I-labeled, α -lactalbumin labeling, effects of protein and sodium chloride levels (bovine), **138**, 604

Subject Index

- Triglycerides
 synthesis and secretion in isolated hepatocytes, epinephrine effects (rat), **141**, 191
- 3,4,5-Triiodobenzoic acid
 effect on smooth muscle contraction and [³H]verapamil binding to plant and animal membrane fractions, **138**, 1269
- Triiodothyronine
 binding to nuclear membranes, effect of dietary lipid content (murine liver), **135**, 655
 neogenesis in lean and obese LA/N-*cp* rats, **140**, 134
- N*-[(1,5,9)-Trimethyl-decyl]-4 α ,10-dimethyl-8-aza-*trans*-decal-3 β -ol
 inhibition of 2,3-oxidosqualene cycloartenol and lanosterol cyclases, **138**, 764
- 2,3,5-Trimethyl-6-(12-hydroxy-5,10-dodecadiynyl)-1,4-benzoquinone, *see* AA861
- 1,3,4-Trisphosphate
 formation from inositol 1,4,5-trisphosphate, calcium sensitivity and ATP dependence (bovine adrenal glomerulosa cells), **139**, 259
- Triton WR-1339, *see* Tylloxapol
- Tropomyosin
 amino acid sequence, homology with group A streptococcal Pep M5 protein, **140**, 684
 noncrosslinked, anomalous concentration dependence of equilibrium thermal unfolding curve, simulation by theoretical model, **141**, 394
 polymerization, end-to-end, irreversible loss after exposure to urea or cyanate (rabbit muscle), **134**, 64
- Troponin C
 degradation, calcium effects (rat), **134**, 1269
- Tropylum tetrafluoroborate
 as substrate for aldehyde oxidase (rabbit liver), **140**, 609
- Trypanosoma cruzi*
 epimastigotes, sterol biosynthesis, ketoconazole effects, **136**, 851
 extracts, primaquine-mediated hydroxyl radical generation, **135**, 1029
 infection, identification of monoclonal antibody recognizing superficial tubulin-like antigen (mouse), **139**, 1176
- Trypsin
 alteration of signal II_s in oxygen-evolving photosystem II preparations (spinach chloroplasts), **134**, 272
 binding by α_1 -protease inhibitor, comparison with chymotrypsin, **138**, 925
 conversion of Ca²⁺-dependent protein kinase C to independent form of phorbol ester-binding protein (rat brain), **139**, 320
 effect on microsomal *trans*-2-enoyl CoA reductase component of fatty acid chain elongation system (rat liver), **140**, 74
 induction of destabilization of dioleoylphosphatidylethanolamine- and glycoporphin-containing liposomes, **141**, 973
- Trypsin inhibitor
 sulfation by sulfotransferase (*Eubacterium* A-44), **140**, 38
- α_1 -Trypsin inhibitor
 and peptide aldehydes, as elastase inhibitors in treatment of emphysema, comparison (human), **139**, 896
- Tryptophan
 plasma levels, effect of sustained exercise (rat), **137**, 149
- Tryptophanase
 tyrosyl residue, catalytic function (*Escherichia coli*), **137**, 964
- Tubulin
 interaction with myelin basic protein, effect on tubulin carboxypeptidase, **136**, 482
 -like superficial antigen, recognition by *Trypanosoma cruzi* monoclonal antibody (mouse), **139**, 1176
- β subunit
 exchangeable GTP-binding domain, localization (bovine brain), **135**, 1135
 gene detection in recombinant phages constructed from mung bean genomic DNA, **137**, 788
 synthesis induction by 20-hydroxyecdysone (*Drosophila melanogaster*), **134**, 191
 -vincristine complexes, formation and stability in tumor cytosols, GTP effect (human), **135**, 695
- Tubulin carboxypeptidase
 effect of myelin basic protein interaction with tubulin (bovine brain), **136**, 482
- Tumor cells, *see also* Cell lines
 astrocytoma, proliferation, effects of dequalinium, propylinium, and rhodamine (rat), **135**, 574
- bladder
 clonogenic growth, inhibition by 1,25-dihydroxyvitamin D₃, role of receptors (human), **139**, 136
 phototoxicity of hematoporphyrin derivative, enhancement by selenium (mouse), **140**, 1086
- brain, in primary cultures, heat shock response, comparison with established cell lines (human), **141**, 46
- breast
 doxorubicin-induced killing, prevention by iron chelators and oxygen radical scavengers (human), **135**, 330
 drug-resistant, metabolic cooperation with normal, preneoplastic, or neoplastic mammary cells *in vitro*, ouabain effects (mouse), **140**, 955
 estrogen-induced glycoprotein, identification as lysosomal acidic protease (human), **138**, 102
- growth
 clonogenic, inhibition by 1,25-dihydroxyvitamin D₃, role of receptors (human), **139**, 136
 combined effects of buserelin, estradiol, and tamoxifen *in vitro* (human), **140**, 550
 effect of phorbol ester-induced desensitization of cytosolic protein kinase C (human), **135**, 65
 relationship to phosphoinositide phosphorylation (rat), **134**, 876
 induction by 7,12-dimethylbenz[a]anthracene, anticarcinogenic effect of retinoids, relationship to cAMP-dependent protein kinase (rat), **135**, 116
 phorbol ester-induced protein kinase C intracellular translocation and down-regulation (human), **137**, 1159
 progesterone receptors, interaction with progesterin R 5020 and antiprogesterin RU 38486, comparison (human), **135**, 90
 proliferation, effect of high-affinity hydroxylated antiestrogens (human), **140**, 523
 synthesis and secretion of epidermal growth factor-like polypeptide (human), **136**, 300
- buccal pouch, regression, association with β -carotene (hamster), **136**, 1130
- colonic adenocarcinoma, mRNA, relationship to *pol* gene

Subject Index

- of endogenous A-type retrovirus (human), **139**, 1071
- Ehrlich ascites
- glucose effect on self-transport (mouse), **134**, 1231
- growth, role of *N*-isopentenyl adenine (mouse), **137**, 208
- embryonal carcinoma, differentiation, induction by α -difluoromethylornithine *in vitro* (human), **140**, 66
- Engelbreth-Holm-Swarm, amyloid P component isolation (mouse), **134**, 995
- epidermoid carcinoma, terminal cellular differentiation, relationship to epidermal growth factor receptors (human), **140**, 837
- erythroleukemia
- antiproliferative effects of AA861 and cirsiolol (human), **140**, 832
- differentiation, inhibition by theophylline and dibutyl cAMP (mouse), **134**, 845
- proto-oncogene sequential expression during differentiation (mouse), **135**, 1112
- gastric carcinoma, associated GA733 antigen, isolation and characterization (human), **135**, 297
- glioma C6
- growth inhibition by prostaglandin A₁ biotransformation products (rat), **141**, 1254
- inositol incorporation into phosphoinositides and inositol phosphates, effects of cationic amphiphilic drugs (rat), **135**, 465
- HeLa
- (ADP ribose)_n degradation rates, **139**, 37
- detection of (ADP-ribose)_n glycohydrolase I and II, **135**, 979
- extracts capable of excising pyrimidine dimers from DNA, effect on 4-nitroquinoline-1-oxide adducts, **140**, 775
- membrane permeability changes induced by *Clostridium perfringens* type A enterotoxin in absence of extracellular Ca²⁺, **141**, 704
- synthesis of specific protein after exposure to heat shock at 42°C, **137**, 957
- transient accumulation of pteridines, induction by phorbol ester, **141**, 494
- hepatoma
- aflatoxin B₁-induced, expression of *c-myc*, *c-Ha-ras* and *N-ras* genes (rat), **138**, 858
- ascites, isolated plasma membranes, phosphorylation by casein kinase I, **138**, 95
- expression of aldolase A mRNA (human), **134**, 1093
- glucose-6-phosphatase induction by dexamethasone *in vitro*, **139**, 1130
- oncogene expression, Northern blot analyses (human), **135**, 262
- polycyclic hydrocarbon-inducible cytochrome *P*-450, induction by phenobarbital (rat), **137**, 120
- processing and secretion of complement components C3 and C4, effects of acidotropic compounds (human), **136**, 603
- stationary-phase, dipyrindamole sensitivity, restoration by amphotericin B (rat), **140**, 434
- HEP 3B, produced hepatitis B virus transcripts, analysis (human), **138**, 131
- increased actin nucleating activity, **134**, 1284
- leukemia
- DNA synthesis inhibition by *cis*- and *trans*-diamminedichloroplatinum(II), effect of serum (mouse), **138**, 631
- sensitivity to amsacrine, correlation with magnitude of drug-induced DNA cleavage (human), **134**, 638
- lung
- interferon- β binding, enhancement by lanthanide ions (human), **137**, 480
- occurrence of *c-K-ras* codon 12 GGT-CGT point mutations (human), **140**, 699
- sensitivity to difluoromethylornithine, relationship to ornithine decarboxylase mRNA levels (human), **134**, 572
- lymphoid leukemia
- antiproliferative effects of AA861 and cirsiolol (human), **140**, 832
- cytolysis by melittin, relationship to membrane structure (mouse), **139**, 222
- DNA breakage induction by clomiphene (human), **138**, 1203
- effects of *N*-chlorosulfonyl dicyclohexylamine (human), **141**, 718
- growth characteristics, effect of vibriobactin (mouse), **136**, 273; *erratum*, **137**, 585
- prostaglandin E₂-induced growth inhibition, relationship to metabolic dehydration of prostaglandin E₂ and cellular uptake (mouse), **139**, 808
- transferrin receptor cycling, effect of microtubule assembly inhibition (human), **138**, 1216
- lymphoma, and nonmalignant lymphoid tissues, regulatory mechanisms of ferritin synthesis (human), **139**, 652
- mastocytoma
- antitumor potential of activated macrophages, receptor-mediated inhibition by dexamethasone (mouse), **136**, 130
- butyrate-sensitive and resistant, cAMP phosphodiesterase activity, differential effects of sodium butyrate, **138**, 1030
- permeabilized, Ca²⁺ release, induction by inositol 1,4,5-trisphosphate (mouse), **135**, 46
- medulloblastoma, isolation of heparin-binding growth factors related to bovine brain acidic fibroblast growth factor from lysates (human), **139**, 861
- melanoma
- adhesion to defined extracellular matrix components (mouse), **139**, 515
- Bowes, plasminogen activator-secreting, phorbol ester-stimulated synthesis and phosphorylation of 48-kDa intracellular protein (human), **137**, 263
- clonogenic growth, inhibition by 1,25-dihydroxyvitamin D₃, role of receptors (human), **139**, 136
- detection of warfarin-sensitive vitamin K₁ 2,3-epoxide reductase (mouse), **137**, 1015
- differentiation and adhesion, 120- and 80-kDa detergent-insoluble glycoproteins as markers, **140**, 638
- phorbol ester-induced protein, mRNA isolation (human), **136**, 122
- proliferation, role of platelet-derived acid-sensitive growth factor (human), **138**, 795
- neuroblastoma
- acetylcholine-operated ion channel and α -bungarotoxin receptor, localization on different molecules (human), **137**, 1141

Subject Index

- cellular and mitochondrial respiration, effects of 1-methylcyclohexanecarboxylic acid (mouse), **140**, 789
- glioma hybrid, preproenkephalin mRNA levels, synergistic regulation by cAMP and glucocorticoids, **139**, 1
- growth
 - clonogenic, inhibition by 1,25-dihydroxyvitamin D₃, role of receptors (human), **139**, 136
 - inhibition by prostaglandin A₁ biotransformation products (rat), **141**, 1254
- identification of enkephalin and morphine receptors (human), **137**, 813
- inositol trisphosphate release, effects of D-Ala²-D-Leu⁵ enkephalin and bradykinin, **135**, 507
- neurohybridoma, carbachol-induced inositol 1-phosphate production, effects of lithium and phorbol esters, **136**, 622
- osteosarcoma
 - FBJ virus-induced, type V collagen staining by concanavalin A (mouse), **136**, 857
- growth
 - clonogenic, inhibition by 1,25-dihydroxyvitamin D₃, role of receptors (rat), **139**, 136
 - enhancement by 1,25-dihydroxyvitamin D₃ after inoculation with receptor-rich cells (athymic nude mouse), **139**, 1292
 - osteoblast-like, degradation of human parathyroid hormone (1–84) (rat), **141**, 762
- pancreatic
 - conditioned medium, purification of macrophage colony-stimulating factor (human), **138**, 238
 - insulin-producing, purified amyloid fibril protein, partial homology of N-terminal sequence with calcitonin gene-related peptide (human), **140**, 827
 - point mutation-induced amplification of *c-myc* and activated *c-Ki-ras* genes (human), **140**, 167
- pheochromocytoma PC12
 - calcium channels
 - phorbol ester and protein kinase C effects, **136**, 1049
 - protein kinase C effects, **134**, 1298
 - calcium-dependent activation of glycogen phosphorylase by nerve growth factor (rat), **138**, 917
 - cellular functions, effect of chronic retrovirus infection, **140**, 706
 - levels of protein kinase C substrate B-50, effect of nerve growth factor (rat), **139**, 644
- pituitary
 - cAMP accumulation and hormone release, induction by bacterial adenylate cyclase (rat), **136**, 463
 - detection of prolactin-related mRNA (rat), **138**, 1015
 - insulin-like growth factor II receptor purification and antibody generation (human), **138**, 304
- promyelocytic leukemia HL60
 - adriamycin-resistant, P150 phosphorylation, effects of dimethylsulfoxide, retinoic acid, and phorbol ester (human), **138**, 9
 - antiproliferative effects of AA861 and cirsiolol (human), **140**, 832
 - 3-deazaneplanocin effects (human), **135**, 688
- differentiation
 - granulocytic and monocytic, protein phosphotyrosine content regulation by tyrosine kinase and phosphotyrosine phosphatase (human), **140**, 440
 - induced by interferon γ or tumor necrosis factor, appearance of membrane-bound tyrosine kinase (human), **140**, 908
 - monocytic, expression of colony-stimulating factor (human), **141**, 924
 - reassembly of *c-myc* and relaxation of *c-fos* nucleosomes (human), **141**, 213
 - transferrin receptor transcriptional regulation (human), **138**, 995
- induction of arachidonic acid release by dimethyl sulfoxide (human), **140**, 515
- resistant to differentiation induction, 5-aza-2'-deoxycytidine growth inhibitory effects (human), **141**, 629
- transient accumulation of pteridines, induction by phorbol ester (human), **141**, 494
- rhabdomyosarcoma
 - isolation of heparin-binding growth factors related to bovine brain acidic fibroblast growth factor from lysates (human), **139**, 861
 - vincristine–tubulin complex formation and stability, GTP effects (human), **135**, 695
- salivary duct, subclones, alkaline phosphatase expression (human), **138**, 625
- sarcoma, associated DNA polymerase α , reactive monoclonal antibody, purification and characterization (mouse), **141**, 60
- teratocarcinoma
 - drug-induced differentiation, specific expression of homoeobox-containing genes, **137**, 520
 - undifferentiated, *c-mos* proto-oncogene expression, **140**, 477
- teratoma, secreted insulin-related factor, characterization (mouse), **135**, 533
- Walker 256, DNA interstrand crosslinking by CB 1954 (rat), **140**, 1073
- Tumor necrosis factor
 - cachectin, stimulation of prostacyclin synthesis in cultured vascular endothelial cells (human), **141**, 482
- human recombinant
 - cytotoxicity and cachectin-like activity, analysis with monoclonal antibodies, **137**, 847
 - depression of cytochrome P-450-dependent microsomal drug metabolism (mouse), **136**, 316
 - effects on circulating leukocytes (mouse), **141**, 818
- induction of
 - HL-60 cell differentiation, associated appearance of membrane-bound tyrosine kinase, **140**, 908
 - superoxide anion production by neutrophils (human), **137**, 1094
- production by macrophages
 - association with β -carotene (hamster), **136**, 1130
 - regulation by prostaglandin E₂ (mouse), **137**, 404
 - stimulation of interleukin 1 and prostaglandin E₂ production in resting macrophages (mouse), **136**, 94
- Tumor necrosis factor receptors
 - binding capacity, regulation by protein kinase C (human T cells), **141**, 855
- Tunicamycin
 - effect on hexose transport reversal and insulin binding in glucose-starved fibroblasts after glucose refeeding (human), **140**, 844

Subject Index

- and swainsonine, comparative effects on various neuroreceptors (NCB-20 cells), **134**, 1387
- Tween 80, *see* Polysorbate 80
- Tyloxapol
selective inhibition of hepatic microsomal cholesterol epoxide hydrolase (mouse), **140**, 632
- Tyrosinase, *see* Monophenol monooxygenase
- Tyrosine
glycyl derivative, effect on IgG binding to bacterial Fc receptors, **135**, 1105
labeling with *p*-diazobenzoyl biocytin, **138**, 872
phosphorylation
by casein kinase A (human erythrocytes), **139**, 855
in plasma membrane proteins, stimulation by oxygen radicals (rat liver), **139**, 439
in soluble 66-kDa protein, induction by dimethyl sulfide (human T cells), **140**, 743
plasma levels, effect of sustained exercise (rat), **137**, 149
sulfation sites in proteins, characterization and criteria for prediction of occurrence, **141**, 326
in tryptophanase, catalytic function (*Escherichia coli*), **137**, 964
- Tyrosine aminotransferase
induction by glucocorticoid *in vivo*, enhancement by diacylglycerol (adrenalectomized rat), **138**, 275
- Tyrosine hydroxylase
activation by calpains I and II (bovine adrenal medulla), **134**, 749
- Tyrosine kinase
associated activity of
insulin and insulin-like growth factor I receptors (rat brain), **134**, 532
insulin-like growth factor I receptors, effect of human polyclonal antibody (human placenta, rat liver), **138**, 1023; *erratum*, **140**, 1152
detection in plasma, serum, and erythrocyte extracts by pp60^{src} antiserum (human), **135**, 426
membrane-bound, appearance during interferon γ - or tumor necrosis factor-induced differentiation of HL-60 cells, **140**, 908
oncogene products, relationship to heat shock protein hsp 90, **138**, 1298
p60^{v-src}-related, isolation from acetylcholine receptor-rich membranes: association and dissociation of phosphatidylinositol kinase activity (*Narke japonica*), **139**, 473
regulation of protein phosphotyrosine content during induced granulocytic and monocytic differentiation (HL-60 cells), **140**, 440
- Tyrosine-protein kinase, *see* Protein-tyrosine kinase
- U**
- Ubiquinol-cytochrome-c oxidoreductase
mitochondrial, deficiency in mitochondrial myopathy and renal dysfunction (human), **137**, 911
- Ubiquinone
-binding protein
antibodies, interaction with mitochondrial complex III (bovine heart), **136**, 376
from mitochondrial cytochrome *bc*₁ complex, cell-free synthesis (rat liver), **138**, 1291
purification from mitochondrial NADH-ubiquinone reductase (bovine heart), **138**, 1237
mitochondrial, thyroxine-induced changes (rat liver), **138**, 381
role in mitochondrial superoxide formation (rat heart), **138**, 533
- Ubiquitin
conformation and activity, role of methionine-1 (human erythrocytes), **138**, 437
- UDPGalactose-glucose galactosyltransferase, *see* Lactose synthase
- UDPGalactose:lactose galactosyltransferase
multiple forms, partial purification and characterization (human renal microsomes), **141**, 84
- UDPGalactose:lactosylceramide galactosyltransferase
multiple forms, partial purification and characterization (human renal microsomes), **141**, 84
- UDPGlucose glucosyltransferase
regulation by amylose-precipitating factor (*Neurospora crassa*), **135**, 701
- Ultraviolet irradiation
induction of polypeptide synthesis (*Saccharomyces cerevisiae*), **138**, 679
- Unspecific monooxygenase
inhibition by recombinant tumor necrosis factor (mouse), **136**, 316
- Uracil
incorporation into DNA, methotrexate effect (HeLa cells, human lymphoblastoid cells), **135**, 886
- Urea
effect on tropomyosin end-to-end polymerization (rabbit muscle), **134**, 64
formation by periportal and perivenous hepatocytes, zonal characteristics, retention in primary culture (rat), **139**, 1055
induced transformation of native estrogen receptor: evidence for separate DNA- and ATP-binding sites (immature bovine uterus), **139**, 1250
- Uremia
associated amyloid kidney stones, detection of β_2 -microglobulin fragments, **136**, 665
detection of 2-amino-3,8-dimethylimidazo [4,5-f] quinoxaline in dialysate (human), **138**, 1084
- Uric acid
plasma levels, effect of *Plasmodium vinckei* infection (mouse), **134**, 152
protection of bovine erythrocyte membranes and linolenic acid from ozone-induced oxidation, **137**, 536
- Uridine
2',3'-*seco* nucleosides and nucleotides, behavior in various enzymatic systems, **137**, 453
- Usher's syndrome
associated reduction of arachidonate and docosahexaenoate content in plasma phospholipids (human), **141**, 600
- Uterus
normal, identification and characterization of platelet-activating factor (rat), **138**, 1231

Subject Index

V

- Valinomycin
alteration of accessibility of surface thiols (human lymphocytes), **140**, 999
- Vanadate
reduction by microsomal redox system (rat liver), **141**, 171
stimulation of insulin-like growth factor binding to adipocytes (rat), **138**, 174
- Vanadium
-binding substance, purification from tunicate blood cells, **141**, 251
- Vas deferens
twitch response, inhibition by clonidine-displacing substance (rat), **134**, 184
- Vasoactive intestinal polypeptide
precursor protein, characterization of derived peptide (cat, guinea pig, human, rat, swine), **139**, 1142
- Vasopressin
arginine, *see* Arginine vasopressin
effect on
cytoplasmic free Ca^{2+} levels, comparison with angiotensin II and phenylephrine effects (rat hepatocytes), **140**, 107
phosphorylation of 80K protein (Swiss 3T3 cells), **140**, 379
release after sodium loading, inhibitory role of hypothalamic atrial natriuretic polypeptide (rat), **139**, 1285
in smooth muscle cell monolayers containing fluorescent probes, effect on cytosolic Ca^{2+} levels (hamster, rabbit, rat), **136**, 927
specific binding and mitogenic effects, inhibition by substance P antagonist (Swiss 3T3 cells), **137**, 135
- Venoms
scorpion, β -toxin and insect toxins, pharmacological properties, comparison with *Tityus serrulatus* toxin VII, **139**, 296
snake, induced damage in pancreatic acinar membranes, role of calcium (rat), **139**, 530
- Verapamil
 ^3H -labeled, binding to plant and animal membrane fractions, effect of 3,4,5-triiodobenzoic acid, **138**, 1269
stimulation of low-density lipoprotein receptor synthesis (human skin fibroblasts), **136**, 845
- Veratridine
in detection of Na^+ channels in myometrial cells (rat), **137**, 675
- Veratryl alcohol
as lignin model compound, biomimetic oxidation by simple inorganic complexes, **139**, 1104
- Veratrylglycerol- β -guaiacyl ether
as lignin model compound, biomimetic oxidation by simple inorganic complexes, **139**, 1104
- Vesicles
endocytic, hydrophobic domain of boundary membranes, close association with clathrin (bovine brain), **141**, 878
inside-out, transglutaminase substrate identification by immunoblotting (human erythrocytes), **134**, 685
lysophosphatidylcholine/cholesterol, structural analysis by ^{31}P NMR, **139**, 25
plasma membrane, hexose transport, characterization (rat myoblasts), **141**, 124
secretion, fusion with plasma membranes, inhibition by somatostatin (rat pituitary), **139**, 673
- Vibriobactin
effect on growth characteristics of L1210 cells (mouse), **136**, 273; *erratum*, **137**, 585
- Vibrio parahaemolyticus*
pathogenic strain WP1, phage-like particles, induction by mitomycin C, **141**, 1249
- Vincristine
-tubulin complexes, formation and stability in tumor cytosols, GTP effects (human), **135**, 695
- VIP, *see* Vasoactive intestinal polypeptide
- Vitamin C, *see* Ascorbic acid
- Vitamin D
-binding protein, inhibition of sequential binding of actin monomers to plasma gelsolin, **136**, 72
-dependent calcium-binding protein, activation of human erythrocyte Ca^{2+} , Mg^{2+} -ATPase (rat kidney), **138**, 547
metabolism, cytoskeleton role (chicken renal tubules), **134**, 218
- Vitamin E
plasma levels, effect of *Plasmodium vinckei* infection (mouse), **134**, 152
- Vitamin K_2
cyclic, structural determination (*Nocardia* spp.), **140**, 916
- Vitamin K_1 2,3-epoxide reductase
warfarin-sensitive, detection in B16 melanoma cells (mouse), **137**, 1015
- Vitronectin
and S-protein: physicochemical, immunochemical, and functional comparison (human), **134**, 951
- Voltage
effect on ion-channel activity of Na^+ , K^+ -ATPase incorporated into solvent-free phospholipid bilayers (canine renal microsomes), **136**, 767

W

- Warfarin
-sensitive vitamin K_1 2,3-epoxide reductase, detection in B16 melanoma cells (mouse), **137**, 1015
- Water
-acetone, chlorophyll *a* photogeneration of superoxide ion in, ESR study (spinach), **139**, 1124
-dimethyl sulfoxide mixtures, hydration states of L-alanine and L-proline, ^{17}O NMR, **135**, 363
intake alteration, effect on atriopeptin levels in heart and plasma (rat), **137**, 922
and triacetyl glycerol, hydrogen bond formation, detection by high-pressure infrared spectrometry, **134**, 140
- Wheat dwarf virus
and satellite tobacco necrosis virus, coat protein sequence homology, **134**, 372
- Wheat germ agglutinin
effect on gonadotropin-releasing hormone-stimulated luteinizing hormone release and gonadotrope desensitization (rat pituitary), **140**, 750
- Wound healing
enhancement with helium-neon laser, mechanism (pig), **138**, 1123

Subject Index

X

- Xanthine dehydrogenase
absence in rabbit myocardium, **141**, 1104
- Xanthine oxidase
absence in rabbit myocardium, **141**, 1104
catalysis of DNA binding to dihydrodiol-derivatized nitro-
polycyclic aromatic hydrocarbons, **141**, 245
induction by polyriboinosinic-polyribocytidylic acid
analysis (mouse, rat), **134**, 632
role of H-2 haplotypes (mouse), **138**, 246
molybdenum cofactor, dephosphorylated oxidation prod-
uct, identification as 2-(1,2-dihydroxyethyl)thieno-
[3,2-*g*]pterin, **135**, 256
- X-ray diffraction
solid-state, dermatan sulfate L-iduronate ring conformation
(porcine skin), **137**, 87
- X rays
induced malignant transformation of embryonic fibroblasts,
inhibition by 3-aminobenzamide (mouse), **134**, 1334
- Xylan
oxidation by hydrogen peroxide, role in cell wall breakdown
(larch), **141**, 238
- Xylem sap
identification of nonproteic amino acid (*Pisum sativum*),
139, 383

Y

- Ytterbium chloride
enhancement of interferon- β binding to human A549 and
murine L cells, **137**, 480

Z

- Zinc
and calcium, micromolar levels, synergistic effects on
membrane fusion (bovine brain), **137**, 101
content in cytochrome-*c* oxidases, analysis by proton-in-
duced X-ray emission (bovine, rat, swine), **140**, 1007
dietary deficiency, effect on fructose-1,6-bisphosphatase
(rat liver), **134**, 944
micromolar levels, potentiation of Ca^{2+} -induced phase sep-
aration of phosphatidylserine-containing liposomes
(bovine brain), **136**, 476
- Zymosan
serum-treated, induction of leukotriene B_4 production in
whole blood, comparison with production by isolated
neutrophils (human), **137**, 334
-stimulated phospholipase C activation, intracellular Ca^{2+}
requirement (murine peritoneal macrophages), **134**,
367

10589