

PURIFICATION OF THE YELLOW FLUORESCENT PROTEIN FROM *VIBRIO FISCHERI*  
AND IDENTITY OF THE FLAVIN CHROMOPHORE

P. Macheroux, K.U. Schmidt, P. Steinerstauch, and S. Ghisla  
Faculty of Biology, University of Konstanz, POB 5560,  
D-7750 Konstanz, FRG

P. Colepiccolo, R. Buntic, and J.W. Hastings,  
Department of Cellular and Developmental Biology,  
Harvard University, Cambridge, MA 02138

Received May 1, 1987

---

A low molecular weight protein (~ 25,000 D) exhibiting a yellow fluorescence emission peaking at ~ 540 nm was isolated from *Vibrio fischeri* (strain Y-1) and purified to apparent homogeneity. FMN is the chromophore, but it exhibits marked red shifts in both the absorption ( $\lambda_{\max}$  = 380, 460 nm) and the fluorescence emission. When added to purified luciferase from the same strain, which itself catalyzes an emission of blue-green light ( $\lambda_{\max}$  ~ 495 nm), this protein induces a bright yellow luminescence ( $\lambda_{\max}$  ~ 540 nm); this corresponds to the emission of the Y-1 strain *in vivo*. This yellow bioluminescence emission is thus ascribed to the interaction of these two proteins, and to the excitation of the singlet FMN bound to this fluorescent protein. © 1987 Academic Press, Inc.

---

The light emission of bioluminescent bacteria is characteristically centered in the blue-green region, with a maximum at about 495 nm (1). About ten years ago, however, Ruby and Nealson (2) reported a strain of *Photobacterium* (formerly *Vibrio*) *fischeri* isolated from sea water off San Diego that emitted yellow light. Spectrally, two maxima were observed: one at the usual 495 nm and a second in the yellow about 540 nm.

This yellow emission was later identified with a yellow fluorescent protein, distinct from luciferase, but capable of generating the yellow bioluminescence when included in a luciferase *in vitro* reaction system otherwise emitting blue light (3). The chromophore appeared to be a flavin, but neither its identity nor the purification and properties of this novel protein were reported. In view of the keen interest in the mechanism of populating excited states in chemi- and bioluminescence (4), as well as the identity of the emitter(s) in the bacterial luciferase system (1,5-9), we undertook a detailed study of this yellow fluorescent protein (YFP).

#### MATERIALS AND METHODS

Instrumentation and Assay: Bioluminescence was measured with a calibrated photomultiplier photometer (10). Fluorescence excitation and emission

spectra were recorded with a Perkin Elmer MPF4 spectrophotofluorimeter; the spectral composition of bioluminescence was measured in the same instrument without the exciting light. Bioluminescence and fluorescence emission spectra were corrected for wavelength dependent differences in sensitivity. Absorption spectra were measured with the Kontron Uvicon 820 spectrophotometer. SDS-PAGE was carried out according to Laemmli (11) with protein molecular weight standards from Pharmacia. The standard buffer contained 10 mM K/Na phosphate, pH 7.0, with 1 mM dithiothreitol, 1 mM EDTA and 1  $\mu$ M FMN (Sigma). All chemicals used were of analytical (or best available) grade and not further purified.

The activity of luciferase alone, without regard for the color of the light emitted, was measured photometrically at 22° in the "dithionite" assay (14), in which luciferase and FMN are first reduced with dithionite and the reaction is initiated by the rapid mixing with long chain aldehyde and excess oxygen. In this reaction the half time of the light emission is quite short, perhaps 10-20 s, too short for easy measurement of its spectral distribution. The activity of the YFP was estimated by measuring the intensity of the yellow (~ 540 nm) band relative to the blue in a luciferase assay at 12°C. To do this the coupled reductase-luciferase assay (12) was used, in which FMN is reduced by NADH (Sigma) via added flavin reductase (13); with excess NADH assay light emission may be caused to continue much longer. The reaction mixture (150  $\mu$ l final volume) included 10  $\mu$ l of a purified luciferase (~ 50  $\mu$ M); 10  $\mu$ l of purified reductase (~ 50  $\mu$ M); 10  $\mu$ l FMN (0.4 mM); 10  $\mu$ l decanal suspension (0.01% v/v); 100  $\mu$ l of the sample in standard buffer; and 10  $\mu$ l of NADH (0.1 M) to initiate the reaction in a microcuvette thermostatted at 12° in the fluorimeter. Under these conditions the light emission was steady for up to 10 min.

Cell Growth and Harvesting: *Vibrio fischeri*, strain Y-1 (ATCC #33715) was obtained from Dr. K.H. Nealson and grown at 16° in a 250 l New Brunswick fermenter, in a complete sea water medium containing peptone and yeast extract (14). Cell density (by OD<sub>660 nm</sub>) and bioluminescence (photometrically) were monitored during growth; at peak luminescence (about 18 hr) the culture was rapidly cooled to 4°C and cells were harvested with a Braun continuous centrifuge. The resulting paste, about 500 g (wet weight) per 180 l culture, was frozen and stored at -20°.

## RESULTS AND DISCUSSION

Cells were lysed at 4°C in standard buffer (600 ml/100 g cells) with stirring for 1 hr. The lysate was centrifuged at 27,500 x g for 60 min and the precipitate discarded. Solid ammonium sulfate was added to the supernatant and the fraction precipitating between 40 and 75% saturation was dissolved in a minimal amount (~ 100 ml) of standard buffer and dialyzed 3 times against the same buffer. This solution was combined with a DEAE-Sephacel suspension (80 ml); after one hour the solid with the adsorbed protein was separated by centrifugation, packed into a column (18 x 2.5 cm) and eluted with a gradient of the standard buffer with the phosphate concentration ranging from 10 to 300 mM. The fractions were assayed for the different activities; those containing the YFP, as detected by the shift in fluorescence emission to longer wavelengths (see below), were combined and concentrated by ultrafiltration (Amicon PM 10) to 20 ml. This was loaded on a Sephacryl S-200 column (90 x 6 cm) and the fractions containing YFP were eluted with standard buffer and applied to a


HA-Ultrogel (hydroxyl-apatite, BioRad) column (27 x 2.5 cm). The standard buffer without EDTA with a phosphate gradient (10-50 mM) was used to elute the YFP activity; active fractions were pooled, and concentrated by dialysis against polyethylene glycol to a volume of about 20 ml. YFP concentrations were measured in terms of flavin absorbance assuming one flavin per protein molecule.

For luciferase and flavin reductase, the material obtained from the 40-75%  $(\text{NH}_4)_2\text{SO}_4$  fractionation was dialyzed and subjected to gel filtration over Sephacryl S-200; the two activities were well separated and were pooled separately. Luciferase was further purified according to the procedure of Holzman and Baldwin (15); FMN reductase was concentrated and used without further purification.


Several difficulties were encountered in the purification of YFP, notably the apparent denaturation of the protein. Considering that this might be due to the dissociation of the flavin chromophore, FMN (1.0  $\mu\text{M}$ ) was added to all buffers used in the purification procedures. This appeared to be successful and the YFP protein was obtained in a purity estimated to be >95% by SDS-PAGE (Figure 1). Based on the standards, a molecular weight of about 25,000 Daltons was estimated. Amino acid analysis gave a similar value.

The purified holoprotein exhibits the two absorption bands in the visible and near UV typical of a flavoprotein (Figure 2), but differs in some respects. Whereas the 440 and 375 nm flavin bands are usually only slightly shifted in flavoenzymes, these bands are red-shifted in YFP, probably due to a decrease in the protein polarity around the flavin binding region (16,17). Moreover, the ratio of the peak absorbances of these two bands is 0.95, whereas in free FMN it is 1.2. The fluorescence emission maximum of the protein bound flavin is at 542 nm compared to about 525 nm for free flavin, but the fluorescence quantum yield is essentially unaltered.

The flavin chromophore associated with YFP was identified unambiguously as FMN by several independent criteria. Upon denaturation of the protein, either by heat or by the addition of SDS, the chromophore exhibits the same absorption and fluorescence values as authentic FMN. Its  $R_f$  value on thin layer chromatography (butanol:HAc:H<sub>2</sub>O; 12:3:5) and the retention time on HPLC (18) are identical with those of authentic FMN. This flavin can be used by bacterial luciferase, which is highly specific for FMN in the bioluminescence reaction (1,7). It is bound by the FMN specific apoflavodoxin from Megasphaera elsdenii (19), and also by riboflavin binding protein (which binds riboflavin and FMN, but not FAD) (20). Similar results were obtained in all of these tests utilizing YFP isolated and purified in the absence of added FMN.



①



②

Figure 1. SDS-PAGE with 15% polyacrylamide. Lanes 1 and 2, standards, with approximate molecular weights indicated in kilodaltons:  $\alpha$ -lactalbumin (14,000 D), soybean trypsin inhibitor (20,000 D), carbonic anhydrase (30,000 D), Ovalbumin (43,000 D), bovine serum albumin (67,000 D), and phosphorylase b (96,000 D). Lane 2: crude extract after lysis and dialysis. Lane 3 after DEAE-Sephacel. Lane 4, after Sephacryl S-200 gel filtration. Lanes 5 and 6, after ultragel hydroxyl-apatite chromatography.

Figure 2. Absorption spectrum of purified YFP (best fractions after ultragel hydroxyl-apatite chromatography), 5.6  $\mu$ M in 10 mM Tris-HCl, pH 7.0 containing 1.0  $\mu$ M FMN, (left, solid lines). The sample was dialyzed extensively against this buffer and the spectrum was recorded with buffer in the reference. For the U.V. band (240-300 nm) the scale is 5 times greater. The arrows indicate the absorption maxima obtained after addition of 0.05% SDS; these correspond to those of free FMN. Corrected fluorescence emission spectrum of YFP (dashed line, right) measured using the same sample as above with excitation at 470 nm (slits, 4 mm). The fluorescence emission maximum observed after addition of SDS is shown by the arrow. The fluorescence excitation spectrum for YFP (not shown) corresponds to its absorption spectrum, and after denaturation to that of the free flavin. Fluorescence in arbitrary units.

The effect of pure YFP on the bioluminescence emission of bacterial luciferase is demonstrated in Fig. 3, where the appearance of a new emission band with a  $\lambda_{\max}$  at about 540 nm is evident. The similarity of this emission to that observed in vivo indicates that the yellow fluorescent protein is involved in the generation of yellow light in the Y-1 strain of Vibrio fischeri. Although energy transfer from a primary excited state has been suggested as a mechanism (2), the direct excitation of YFP by an electron exchange mechanism has also been proposed (1,8).

It seems likely that a second chromophore may be involved in bacterial bioluminescence in cases where the emission in vivo differs from that of the pure luciferase in vitro (21,22). Indeed a low molecular weight protein with a tightly bound chromophore identified as 8-ribityllumazine has been isolated




Figure 3. Comparison of the corrected bioluminescence emission spectra of the *in vitro* reactions with *Vibrio fischeri* Y-1 luciferase in the absence (solid line, left) and in the presence of 5  $\mu$ M YFP at 7°. Luminescence in arbitrary units.

from *Photobacterium phosphoreum* and *Photobacterium leiognathi* (23) and shown to cause light emission to peak at 476 nm, compared to ~ 495 nm with luciferase alone. This is phenomenologically similar to the YFP protein: similar mechanisms for the population of the excited states may be involved.

#### ACKNOWLEDGEMENTS

This work was supported by a grant from the Deutsche Forschungsgemeinschaft to S.G. (DFG Gh 2/4-5,6) and from the U.S. National Science Foundation to J.W.H. (DMB-8676522). P.C. was supported by a Brazilian National Research Council Fellowship (CNPq). We thank Dr. K.H. Nealson for providing the bacterial strain and for helpful suggestions and discussion.

#### REFERENCES

1. Hastings, J.W., Potrikus, C.J., Gupta, S., Kurfürst, M., Makemson, J.C. (1985) *Adv. Micro. Physiol.* **26**, 235-291.
2. Ruby, E.G. and Nealson, K.H. (1977) *Science* **196**, 432-434.
3. Leisman, G. and Nealson, K.H. (1982) in *Flavins and Flavoproteins* (Massey, V., and Williams, C.H., eds.), Elsevier, New York, pp. 383-386.
4. Wilson, T. (1985) in *Singlet Oxygen* (A.A. Frimer, ed.), Vol. II, CRC Press, Inc., Boca Raton, FL, pp. 37-65.
5. Macheroux, P. and Ghisla, S. (1985) *Nachr. Chem. Tech. Lab.* **33**, 785-790.
6. Lee, J. (1985) *Chemil- and Bioluminescence* (Burr, J.G., ed.), Plenum, New York, pp. 401-437.
7. Ziegler, M.M. and Baldwin, T.O. (1981) *Curr. Top. Bioenerg.* **12**, 65-113.

8. Hastings, J.W. (1986) in Light Emission by Plants and Bacteria (Govindjee, J. Ames and D.C. Fork, eds.), Academic Press, New York, pp. 363-398.
9. Kurfürst, M., Ghisla, S. and Hastings, J.W. (1984) *Proc. Nat. Acad. Sci. USA* 81, 2990-2994.
10. Mitchell, G. and Hastings, J.W. (1971) *Anal. Biochem.* 39, 243-250.
11. Laemmli, U.K. (1970) *Nature* 227, 680-685.
12. Hastings, J.W., Baldwin, T.O. and Nicoli, M.Z. (1978) in Methods in Enzymology (Kaplan, N.O. and Colowick, S.P., eds.), Vol. 57 (M. DeLuca, ed.), Academic Press, New York, pp. 135-152.
13. Watanabe, H. and Hastings, J.W. (1982) *Mol. Cell Biochem.* 44, 181-187.
14. Nealson, K.H. (1978) in Methods in Enzymology (Kaplan, N.O. and Colowick, S.P., eds.), Vol. 57 (M. DeLuca, ed.), Academic Press, New York, pp. 153-166.
15. Holzman, T.F. and Baldwin, T.O. (1982) *Biochemistry* 21, 6194-6201.
16. Ghisla, S., Massey, V., Lhoste, J.M. and Mayhew, S.G. (1974) *Biochemistry* 13, 589-597.
17. Hellis, P.F. (1982) *Chem. Soc. Rev.* 11, 15-39.
18. Ghisla, S., Massey, V. and Yagi, K. (1986) *Biochemistry* 25, 3282-3289.
19. Wassink, J.H. and Mayhew, S.G. (1975) *Analytical Biochemistry* 68, 609-616.
20. Becvar, J. and Palmer, G. (1982) *J. Biol. Chem.* 257, 5607-5617.
21. Seliger, H.H. and Morton, R.A. (1968) in Photophysiology (A.C. Giese, ed.), Vol. 4, Academic Press, NY pp. 253-314.
22. Matheson, I.B.C., Lee, J. and Müller, F. (1981) *Proc. Natl. Acad. Sci. USA* 78, 948-952.
23. O'Kane, D.J., Karle, V.A. and Lee, J. (1985) *Biochemistry* 24, 1461-1467.