

TABLE I
Analysis of the 340 nm Fluorescence Intensity Decay of Annexin V in the Absence
and in the Presence of (PC/PS) Liposomes of Various Ratios

	τ_i (ns)	α_i	f_i (%)
Annexin V + EDTA	5.4 ± 0.1	0.08 ± 0.01	36 ± 1
	1.30 ± 0.03	0.44 ± 0.03	48 ± 1
	0.40 ± 0.03	0.48 ± 0.04	16 ± 1
Annexin V + (PC/PS=10)liposomes (pCa = 2.5)	7.2 ± 0.1	0.58 ± 0.03	82 ± 3
	2.2 ± 0.1	0.42 ± 0.03	18 ± 3
Annexin V + (PC/PS= 40)liposomes (pCa = 2)	6.9 ± 0.1	0.63 ± 0.05	85 ± 4
	2.0 ± 0.1	0.37 ± 0.05	15 ± 4
Annexin V + (PC/PS=200)liposomes (pCa = 1.9)	7.0 ± 0.2	0.60 ± 0.05	84 ± 5
	2.0 ± 0.1	0.40 ± 0.05	16 ± 5
Annexin V + PC liposomes (pCa = 2)	5.9 ± 0.1	0.20 ± 0.04	59 ± 2
	1.57 ± 0.05	0.42 ± 0.01	34 ± 2
	0.98 ± 0.02	0.38 ± 0.04	7 ± 1

Note. Lifetime components τ_i , normalized preexponential terms α_i and fractional intensities f_i were expressed as means (\pm S.E.M.) for at least three independent experiments, corresponding each to 20 decays. The experiments have been carried out with solutions containing 2.9 μ M annexin V in buffer A (see Materials and Methods) with Ca^{2+} concentrations corresponding to maximal annexin V binding. The excitation wavelength was set at 295 nm.

Volume **235**, Number 1 (1997), in Article No. RC976735, "Identification of Glycyrrhizin as a Thrombin Inhibitor," by Ivo Mauricio B. Francischetti, Robson Q. Monteiro, and Jorge A. Guimarães, pages 259–263: On page 259, in the author line, the name of the first author was inadvertently divided into two names. For the reader's convenience, the correct author line is printed here.

Ivo Mauricio B. Francischetti, Robson Q. Monteiro,
and Jorge A. Guimarães

This erratum is Article No. RC977093.