

ERRATUM

Volume **254**, Number 3 (1999), in Article ID bbrc.1998.9992, "Validation of Methods for CYP2C9 Genotyping: Frequencies of Mutant Alleles in a Swedish Population," by Ümit Yasar, Erik Eliasson, Marja-Liisa Dahl, Inger Johansson, Magnus Ingelman-Sundberg, and Folke Sjöqvist, pages 628–631: On page 629, Table 1 was incorrect as printed. The corrected Table 1, incorporating the corrected primer sets used by Nasu *et al.* for detection of CYP2C9*2 and *3 and the correct sizes (bp) of cleavage products, is reprinted below.

This erratum is Article ID bbrc.1999.0540.

TABLE 1
 Primers (5' → 3') Used in the Different Methods and the Lengths of the DNA Fragments
 after Possible Endonuclease Cleavage

Method	Primer Ref.	Sequence	PCR product (bp)	Endonuclease	Positive cleavage (allele)	Cleavage products (bp)
CYP2C9*2 analysis						
Wang	23	GGATATGAAGCAGTGAAGGAA	420	<i>Ava</i> II	CYP2C9*1 or *3	363 + 57
	23	GGCCTTGTTTTCTCAACTC*				
Sullivan-Klose	11	TACAAATACAATGAAAATATCATG	691	<i>Ava</i> II	CYP2C9*1 or *3	527 + 164
	11	CTAACAACCAGACTCATAATG*				
Nasu	24	GTATTTTGGCCTGAAACCCATA	454	<i>Ava</i> II	CYP2C9*1 or *3	397 + 57
	24	GGCCTTGTTTTCTCAACTC*				
CYP2C9*3 analysis						
Wang	23	TGCACGAGGTCCAGAGGTAC	131	<i>Kpn</i> I	CYP2C9*3	111 + 20
	23	TGCACGAGGTCCAGAGATGC	131	<i>Nsi</i> I	CYP2C9*1 or *2	110 + 21
	23	AAACATGGAGTTGCAGTGTA*				
Sullivan-Klose	11	AATAATAATATGCACGAGGTCCAGAGGTAC	141	<i>Kpn</i> I	CYP2C9*3	111 + 30
	11	AATAATAATATGCACGAGGTCCAGAGATGC	141	<i>Nsi</i> I	CYP2C9*1 or *2	110 + 31
	11	GATACTATGAATTTGGGACTTC*				
Nasu	23	TGCACGAGGTCCAGAGGTAC	105	<i>Kpn</i> I	CYP2C9*3	85 + 20
	23	TGCACGAGGTCCAGAGATGC	105	<i>Nsi</i> I	CYP2C9*1 or *2	84 + 21
	24	ACAAACTTACCTTGGGAATGAGA*				
* Reverse primer						