Volume **260**, Number 2 (1999), in Article ID bbrc.1999.0925, "APOBEC-2, a Cardiac- and Skeletal Muscle-Specific Member of the Cytidine Deaminase Supergene Family," by Wei Liao, Seung-Ho Hong, Benny Hung-Junn Chan, Frederick B. Rudolph, Sandra C. Clark, and Lawrence Chan, pages 398–404: On page 398, in the author line, the third author's name was misspelled. It should be "Benny Hung-Junn Chang" instead of "Benny Hung-Junn Chan" as printed. For the reader's convenience, the complete author line is printed below. Also, on page 400, Fig. 2 contained some errors. The revised Fig. 2 and revised legend are printed below.

This erratum is Article ID bbrc.1999.1337.

Wei Liao, Seung-Ho Hong, Benny Hung-Junn Chang, Frederick B. Rudolph, Sandra C. Clark, and Lawrence Chan




FIG. 2. (Top) Map figure from The Jackson BSS backcross showing part of chromosome 17. The map is depicted with the centromere toward the top. A 3 cM scale bar is shown to the right of the chromosome bar. Loci mapping to the same position are listed in alphabetical order. Human syntenic genes are indicated in boxes to the far right. To the left of the chromosome bar are offset numbers and other loci of interest from the Mouse Genome Database at http://www.informatics.jax.org. Missing typings were inferred from surrounding data where assignment was unambiguous. Raw data from The Jackson Laboratory were obtained from the World Wide Web address http://www.jax.org/resources/ documents/cmdata. (Bottom) Haplotype figure from the Jackson BSS backcross showing part of chromosome 17 with loci linked to *Apobec-2*. Loci are listed in order with the most proximal at the top. The black boxes represent the C57BL6/JEi allele and the white boxes the SPRET/Ei allele. The number of animals with each haplotype is given at the bottom of each column of boxes. The percent recombination (R) between adjacent loci is given to the right of the figure, with the standard error (SE) for each R.