

The involvement of Gab1 and PI 3-kinase in β 1 integrin signaling in keratinocytes

Yoshihiro Kuwano ^{a,b}, Manabu Fujimoto ^{c,*}, Rei Watanabe ^a, Nobuko Ishiura ^a,
Hiroko Nakashima ^a, Mayumi Komine ^a, Tatsuo S. Hamazaki ^b,
Kunihiko Tamaki ^a, Hitoshi Okochi ^b

^a Department of Dermatology, Faculty of Medicine, University of Tokyo, Tokyo, Japan

^b Department of Regenerative Medicine, Research Institute, International Medical Center of Japan, Tokyo, Japan

^c Department of Dermatology, Kanazawa University Graduate School of Medical Science, 13-1 Takaramachi, Kanazawa, Ishikawa 920-8641, Japan

Received 4 July 2007

Available online 16 July 2007

Abstract

The control of the stem cell compartment in epidermis is closely linked to the regulation of keratinocyte proliferation and differentiation. β 1 integrins are expressed 2-fold higher by stem cells than transit-amplifying cells. Signaling from these β 1 integrins is critical for the regulation of the epidermal stem cell compartment. To clarify the functional relevance of this differential expression of β 1 integrins, we established HaCaT cells with high β 1 integrin expression by repeated flow cytometric sorting of this population from the parental cell line. In these obtained cells expressing β 1 integrins by 5-fold, MAPK activation was markedly increased. Regarding the upstream of MAPK, Gab1 phosphorylation was also higher with high β 1 integrin expression, while Shc phosphorylation was not altered. In addition, enhanced phosphatidylinositol 3-kinase activation was also observed. These observations suggest that Gab1 and phosphatidylinositol 3-kinase play pivotal roles in the β 1 integrin-mediated regulation of the epidermal stem cell compartment.

© 2007 Elsevier Inc. All rights reserved.

Keywords: HaCaT cell; Keratinocyte; Stem cell; β 1 Integrin; Gab1; Phosphatidylinositol 3-kinase; Mitogen-activated protein kinase

The epidermis is a stratified, rapidly renewing tissue in which terminally differentiated cells are continuously lost from the skin surface and replaced by proliferation in the innermost or basal layer adjacent to the basement membrane. Cell regeneration is thought to be achieved by a hierarchy of proliferative cells within the basal layer consisting of stem cells and transient amplifying cells [1]. In the early stages of epidermal differentiation, basal cells detach from the basement membrane and migrate into the suprabasal layers, where they progress through an intricate series of stage-specific morphological and biochemical changes involving the synthesis of the differentiation-specific keratins K1 and K10, as well as involucrin, filagrin, and loricrin. Hence, the adhesive properties of basal kerat-

inocytes are closely linked to the regulation of keratinocyte proliferation and differentiation. The anchorage of keratinocytes to extracellular matrix negatively regulates their differentiation [2,3].

Epidermal cells adhere to the basement membrane via adhesion molecules known as integrins, which comprise a superfamily of heterodimeric receptors that consist of one α and one β subunit. The two subunits collaborate to bind to ligands, which are extracellular matrix proteins or counter-receptors of the Ig superfamily [4]. Watt and colleagues have established that a major cell-surface marker that distinguishes epidermal stem cells from transit-amplifying cells is β 1 integrins, which are expressed 2-fold higher on stem cells than transit-amplifying cells [5,6]. Exit from the stem cell compartment is accompanied by a decrease in surface β 1 integrin levels [7].

* Corresponding author. Fax: +81 76 234 4270.

E-mail address: fujimoto-m@umin.ac.jp (M. Fujimoto).

Whereas $\beta 1$ integrins have been proved to be a useful epidermal stem cell marker, the functional relevance of this increased cell-surface density of $\beta 1$ integrins is on the way to the resolution. High expression of $\beta 1$ integrins helps to maintain the patterned distribution of stem cells, since stem cells are less motile than transit-amplifying cells and thus tend to remain clustered within the epidermal basal layer [8]. Integrins can also mediate signals provided by neighboring cells and the extracellular milieu [4], and $\beta 1$ integrin-mediated signaling maintains the epidermal stem cell compartment via mitogen-activated protein kinase (MAPK) activation [9]. In addition to the epidermal stem cells, Janes et al. suggest that the variation of the integrin expression influences the development of the squamous cell carcinoma (SCC) [10]. Whereas Khavari and colleagues have demonstrated the involvement of Gab1 in Ras/MAPK activation from epidermal growth factor receptor [11], their roles in $\beta 1$ integrin-mediated signaling remained unclear.

Here, we have described further roles of $\beta 1$ integrin expression for the signaling transduction pathway in keratinocytes. We obtained HaCaT cells showing high expression levels of $\beta 1$ integrins, by a repetition of selecting and culturing cells expressing relatively high levels of $\beta 1$ integrins. HaCaT cells with high $\beta 1$ integrin expression showed increased activation of the MAPK cascade. Those accompanied augmented phosphorylation of Gab1, but not of Shc. Also, those cells showed enhanced phosphatidylinositol 3-kinase (PI3K) activation and the amplified association between PI3K and $\beta 1$ integrin. Thus, increased $\beta 1$ integrin expression, which is the marker of the epidermal stem cells, regulates the MAPK signal transduction via Gab1 and PI3K signal transduction in keratinocytes.

Materials and methods

Cells and immunofluorescence analysis. A human keratinocyte cell line, HaCaT, was kindly provided by Dr. Toshio Kuroki (Institute of Molecular Oncology, Showa University, Tokyo, Japan). The HaCaT cells were cultured in DMEM supplemented with 2 mM glutamine, 10% FCS, and 15 mM HEPES (Invitrogen, Carlsbad, CA).

HaCaT cells were grown to confluency and dispersed by trypsin treatment. Cells were washed and stained at 4 °C using mouse anti-human $\beta 1$ integrin antibody (HUTS-4, Chemicon, Temecula, CA) for 20 min, followed by labeling with FITC-conjugated secondary antibody (Southern Biotech, Birmingham, AL). After the washing, the population of HaCaT cells of which $\beta 1$ integrin expression was in the highest 3–5% was collected using an Epics Altra flow cytometer (Beckman Coulter, Miami, FL) and this population was passaged. When cultured, these cells expressed higher levels of $\beta 1$ integrins than their parental cells, and the sorting was repeated. After repeated six times, acquired cells ($\beta 1$ integrin^{high} cells) were used for the experiment. HaCaT cells with median levels of $\beta 1$ integrin expression were also sorted six times and used as controls ($\beta 1$ integrin^{normal} cells). Immunofluorescence analysis was performed as described [12]. Briefly, cells were stained as described above and then analyzed with fluorescence intensity shown as 50% log density plots. Unreactive isotype-matched antibodies (Beckman Coulter) were used as controls for background staining.

Immunoprecipitations and Western blot analysis. Immunoprecipitations and immunoblotting were performed as described previously [13].

Cells were lysed in buffer containing 1% Nonidet P-40, 150 mM NaCl, 50 mM Tris-HCl (pH 8.0), 1 mM Na orthovanadate, 2 mM EDTA, 50 mM NaF, and protease inhibitors. Then, the cell lysates were pre-cleared twice by incubation with appropriate control antibodies plus protein G-Sepharose beads (Amersham Pharmacia Biotech, Buckinghamshire, UK), followed by incubation with protein G-beads plus antibodies to proteins of interest for 2 h at 4 °C. The antibodies used for immunoprecipitation included anti-ERK2, anti-Gab1, anti-Shc (Santa Cruz Biotech, Santa Cruz, CA) and anti-PI3K antibodies (Upstate Biotech, Lake Placid, NY). Immunoprecipitated proteins were subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) and transferred onto nitrocellulose membranes for immunoblotting. These membranes were incubated with HRP-conjugated anti-phosphotyrosine antibody (4G10, Upstate Biotech) to detect protein tyrosine phosphorylation, or were incubated with Anti-ACTIVE[®] MAPK antibodies (Promega, Madison, WI), followed by incubation with HRP-conjugated anti-rabbit IgG antibodies (Jackson ImmunoResearch Laboratories, West Grove, PA). The blots were developed using an enhanced chemiluminescence kit (Pierce, Rockford, IL) and detected using Fluoro-S imager (Bio-Rad Laboratories, Hercules, CA). The blots were stripped and reprobed with antibodies specific for the protein of interest to verify equivalent amounts of protein in each lane. Band intensity was quantified using Quantity One software (Bio-Rad Laboratories). For the inhibition of PI3K activity, cells were cultured with or without 20 μ M LY294002 (Calbiochem, San Diego, CA) for 30 min before the lysis of cells.

PI3K activity assay. PI3K activity was measured in vitro using a competitive ELISA (Echelon Biosciences Inc., Salt Lake City, UT). PI3K was immunoprecipitated from an equal amount of protein lysate from HaCaT cells, followed by incubation with phosphatidylinositol (4,5)-biphosphate [PI(4,5)P₂] substrate (100 pmol) with ATP for 2 h at room temperature. The supernatant was then incubated with a PI(3,4,5)P₃ detector protein for 1 h at room temperature, and the reaction mixtures were transferred to PI(3,4,5)P₃-coated detection plates for 1 h at room temperature. After washing, secondary detection reagent and developing solution (supplied with the kit) were added. PI(3,4,5)P₃ detector protein binding to the plate was determined by measuring the absorbance at 450 nm.

Statistical Analysis. All data are shown as mean values \pm SEM. Comparisons between groups were made using the *t* test.

Results

Generation of HaCaT cells expressing high levels of $\beta 1$ integrins

HaCaT cells expressing high levels of $\beta 1$ integrins were generated by repeated sorting and culture of cells expressing relatively high $\beta 1$ integrin levels (see Materials and methods). Expressions of high $\beta 1$ integrin levels were confirmed by flow cytometry. $\beta 1$ integrin^{high} cells showed higher expression of $\beta 1$ integrins by approximately 5-fold compared with $\beta 1$ integrin^{normal} cells (Fig. 1). $\beta 1$ integrin^{normal} cells expressed $\beta 1$ integrins at the identical levels as the original cells (data not shown). In addition, these $\beta 1$ integrin^{high} and $\beta 1$ integrin^{normal} cells expressed all the other examined surface proteins including epidermal growth factor receptor and intracellular signaling molecules at identical levels (data not shown). Thus, this system was considered to be suitable for assessing how $\beta 1$ integrin levels influence signaling outcomes since $\beta 1$ integrin expression was considered physiological in these cells.




Fig. 1. Cell surface expression of $\beta 1$ integrins by $\beta 1$ integrin^{normal} HaCaT cells and $\beta 1$ integrin^{high} cells. $\beta 1$ Integrin^{high} cells were generated by repeatedly sorting the portion with relatively higher levels of $\beta 1$ integrins. These cells were examined by immunofluorescence staining with flow cytometry analysis. These results are representative of those obtained in four similar experiments.

Increased ERK activity in $\beta 1$ integrin^{high} cells

The direct signaling event mediated by integrins is the activation of the MAPK cascade [9,11,14]. Therefore, ERK activation was first determined in $\beta 1$ integrin^{high} cells and $\beta 1$ integrin^{normal} cells. Cell lysates were resolved by SDS-PAGE and transferred to nitrocellulose membranes. After those were probed with anti-phosphoERK antibodies, band intensity was quantified. Total protein amount of ERK in $\beta 1$ integrin^{high} cells was equivalent to that in $\beta 1$ integrin^{normal} cells (data not shown). However, $\beta 1$ integrin^{high} cells showed 80% increase of ERK activity compared to $\beta 1$ integrin^{normal} cells ($182.5 \pm 5.6\%$; Fig. 2A). ERK activity in $\beta 1$ integrin^{normal} cells was identical to that in the original HaCaT cells (data not shown). We also examined the activation of the EGF receptor, and $\beta 1$ integrin^{high} cells, $\beta 1$ integrin^{normal} cells, and original HaCaT cells had the same tyrosine phosphorylation levels of the EGF receptor (data not shown). Thus, the elevation of $\beta 1$ integrin expression in HaCaT cells amplifies MAPK signal transduction pathway similarly to the previous study using the normal keratinocyte [9].

Increased Gab1 activation in $\beta 1$ integrin^{high} cells

While $\beta 1$ integrins control MAPK cascade and regulate the proliferation and differentiation in keratinocytes [9], the upstream of Ras in $\beta 1$ integrin signaling cascade remains unresolved. In other integrins, the stimulation from $\alpha 6\beta 4$ integrins in keratinocytes causes the Ras/MAPK activation via Shc [15]. In the case of the stimulation from epidermal growth factor receptor, Gab1 facilitates Ras/MAPK signaling [11]. Therefore, we examined whether the activation




Fig. 2. The signaling mediated by $\beta 1$ integrins. (A) Phosphorylation of ERK in $\beta 1$ integrin^{normal} HaCaT cells and $\beta 1$ integrin^{high} cells. Cell lysates were incubated with protein-G beads and anti-ERK Abs. Immunoprecipitated proteins were subjected to SDS-PAGE and transferred onto membranes for immunoblotting. Membranes were incubated with Anti-ACTIVE[®]MAPK antibodies and detected using Fluoro-S imager. The blots were subsequently stripped and reprobed with anti-ERK antibodies to verify equivalent loading of proteins between samples. (B) Shc and (C) Gab1 activation in $\beta 1$ integrin^{normal} HaCaT cells and $\beta 1$ integrin^{high} cells. Cell lysates were subjected to SDS-PAGE and transferred to nitrocellulose membranes for subsequent immunoblotting with phospho-specific antibodies for the activated forms of Shc and Gab1. Membranes are stripped and reprobed with anti-Shc and anti-Gab1 Abs. These results are representative of those obtained in four similar experiments. (D) PI3K activity and (E) $\beta 1$ integrin association with PI3K in $\beta 1$ integrin^{normal} HaCaT cells and $\beta 1$ integrin^{high} cells. (D) PI3K activity was determined with competitive ELISA. (E) Cell lysates were incubated with protein-G beads and anti-PI3K Ab. Immunoprecipitated proteins were subjected to SDS-PAGE and transferred onto membranes. The coprecipitations of $\beta 1$ integrins were determined by Western blot analysis using anti- $\beta 1$ integrin antibody. (A,D,E) Values represent the relative mean optical density (\pm SEM; $n = 4$) of band intensities determined by scanning densitometry from four independent immunoblotting experiments. An asterisk indicates a significantly different sample mean from $\beta 1$ integrin^{normal} HaCaT cells; ** $p < 0.001$ using the t test. * $p < 0.05$ using the t test.

of Shc or Gab1 was located upstream of the MAPK activation in $\beta 1$ integrin^{high} cells. $\beta 1$ integrin density did not influence the expression of Gab1 or Shc since total amount of Shc and Gab1 was not altered between $\beta 1$ integrin^{high} cells and $\beta 1$ integrin^{normal} cells (Fig. 2B and C). Shc was equally

tyrosine phosphorylated in $\beta 1$ integrin^{high} cells compared with $\beta 1$ integrin^{normal} cells (Fig. 2B). By contrast, Gab1 tyrosine phosphorylation was markedly stronger in $\beta 1$ integrin^{high} cells than in $\beta 1$ integrin^{normal} cells (Fig. 2C). Thus, $\beta 1$ integrin upregulation resulted in enhanced tyrosine phosphorylation of Gab1.

Enhanced PI3K activity and association with $\beta 1$ integrins in $\beta 1$ integrin^{high} cells

One of the signaling pathways that protects cells from anoikis is activation of AKT through upstream activation of PI3K [16]. PI3K is a central molecule which regulates cell survival pathways, gene expression and cell metabolism, and cytoskeletal rearrangements [17]. In addition, PI3K could serve as a downstream factor mediating signals from activation of $\beta 1$ integrins in keratinocytes [18]. Therefore, we investigated the relationship between the increased expression of $\beta 1$ integrins and PI3K activity. Assessed by competitive ELISA, PI3K activity was significantly enhanced in $\beta 1$ integrin^{high} cells ($156.8 \pm 10.6\%$) compared with $\beta 1$ integrin^{normal} cells (Fig. 2D). PI3K activity in $\beta 1$ integrin^{normal} cells was identical to that in original HaCaT cells (data not shown). PI3K expression levels were not altered in $\beta 1$ integrin^{high} cells compared with $\beta 1$ integrin^{normal} cells or the original HaCaT cells (data not shown).

Then, to clarify further the relation of increased $\beta 1$ integrins to the enhanced PI3K activity, we examined the association between $\beta 1$ integrins and PI3K (Fig. 2E). $\beta 1$ Integrin coprecipitates were augmented in $\beta 1$ integrin^{high} cells ($216.5 \pm 35.1\%$) compared to $\beta 1$ integrin^{normal} cells. Hence, amplified $\beta 1$ integrin expression causes augmented PI3K association and leads to enhanced PI3K activity.

Effects of inhibited PI3K on Gab1 and ERK activities

Increase of $\beta 1$ integrin density led to the increased Gab1 and ERK activities and enhanced PI3K activity. Since increase of $\beta 1$ integrins also caused enhanced PI3K association, PI3K seems to be an upstream factor of Gab1 and ERK. Therefore, we tested whether a PI3K inhibitor, LY294002, can block the increased activities of Gab1 and ERK induced by increased $\beta 1$ integrins (Fig. 3). Cells were cultured with or without 20 μ M LY294002 for 30 min before the lysis of cells. LY294002 blocked the increase of the Gab1 phosphorylation in $\beta 1$ integrin^{high} cells (Fig. 3A). The increase of ERK activity was also blocked by LY294002 (Fig. 3B). Therefore, Gab1 and ERK are the downstream factor of PI3K in $\beta 1$ integrin^{high} cells, suggesting that increased $\beta 1$ integrin enhanced the Gab1 and ERK activity via PI3K.

Discussion

Epidermal stem cells express 2-fold higher $\beta 1$ integrins than transit-amplifying cells [5,6] and suprabasal keratinocytes do not normally express integrins [5,19]. While




Fig. 3. The inhibition of (A) Gab1 activity and (B) ERK activity by LY294002 in $\beta 1$ integrin^{normal} cells and $\beta 1$ integrin^{high} cells. Cells were cultured with or without 20 μ M LY294002 for 30 min before the lysis of cells. Cell lysates were incubated with protein-G beads and anti-Gab1 or anti-ERK Abs. Immunoprecipitated proteins were subjected to SDS-PAGE and transferred onto membranes for immunoblotting. These membranes were incubated with HRP-conjugated anti-phosphotyrosine antibody or Anti-ACTIVE^(R)MAPK antibodies and detected using Fluoro-S imager. The blots were subsequently stripped and reprobed with anti-Gab1 or anti-ERK antibodies to verify equivalent loading of proteins between samples. Values represent the relative mean optical density (\pm SEM; $n = 4$) of band intensities determined by scanning densitometry from four independent immunoblotting experiments. An asterisk indicates a significantly different sample mean from cells cultured without LY294002; $*p < 0.05$ using the t test.

the introduction of a dominant-negative $\beta 1$ integrin mutant into cultured human keratinocytes reduces MAPK activation and stimulates the exit from the stem cell compartment [9], the further detailed mechanisms for the regulation by the increased $\beta 1$ integrins remain unresolved. The current study supports this previous observation about the functional role of highly expressed $\beta 1$ integrins in keratinocytes. $\beta 1$ Integrin^{high} cells showed increased ERK activity (Fig. 2A). Then, this increased activity was accompanied with amplified Gab1 tyrosine phosphorylation (Fig. 2C), while Shc tyrosine phosphorylation showed no apparent change in $\beta 1$ integrin^{high} cells (Fig. 2B). Furthermore, $\beta 1$ integrin^{high} cells also showed enhanced PI3K activity (Fig. 2D) and augmented PI3K association with $\beta 1$ integrins (Fig. 2E). Then, the inhibition of PI3K led to the inhibition of increased Gab1 and ERK activities (Fig. 3), suggesting that Gab1 and ERK are the downstream factors of PI3K. PI3K is also reported as the downstream factor of Ras [20]. Thus, PI3K may have dual roles in this pathway by facilitating Gab1 phosphorylation and by serving as downstream effector of Ras (Fig. 4). This intrinsic loop may sustain the activity of this signaling pathway, which may be important in epidermal stem cell fate. These changes in $\beta 1$ integrin^{high} cells suggest the further detailed functional relevance of increased expression of $\beta 1$ integrins in epidermal stem cells [5,6] (Fig. 4).




Fig. 4. Proposed model of the involvement of Gab1 and PI3K in $\beta 1$ integrin signaling in keratinocytes. Augmented $\beta 1$ integrin expression increases the association with PI3K, which enhances PI3K activity. PI3K produces phosphatidylinositol-3,4,5-trisphosphates, which recruits Gab1 to the membrane, where they are activated. Then, Gab1 regulates Ras activation, which leads to ERK activation. PI3K also works as the downstream effector of Ras. This intrinsic loop may sustain the activity of this signaling pathway, which may be important in epidermal stem cell fate.

When sections of SCCs are examined by immunohistochemistry, considerable variation in integrin expression is observed and associated with the prognosis. An increase in surface integrin levels by only several folds is unlikely to be detected by conventional immunohistochemical staining and thought to exist more frequently in SCCs. Then, it has been supposed that these small changes could profoundly influence the proliferation and differentiation [10]. Our observations of enhanced PI3K and MAPK activity in $\beta 1$ integrin^{high} HaCaT cells also support this hypothesis and suggest that $\beta 1$ integrins and/or those downstream molecules are reasonable target molecules for the treatment of SCC.

It has been suggested that Ras acts within epidermis to support proliferative capacity and oppose terminal differentiation [11,21]. While Ras/MAPK activation in $\beta 1$ integrin-mediated signal transduction is independent of FAK [9], little is known about the upstream of Ras/MAPK activation except FAK. In $\alpha 6\beta 4$ integrins, Shc mediates the Ras/MAPK activation [15]. In the signal transduction from growth factor, MAPK activation can proceed normally in Shc^{-/-} cells [22] and is impaired in Gab1^{-/-} cells [23,24]. Gab1 overexpression prolongs the persistence of active Ras/MAPK in response to epidermal growth factor stimulation [11]. Our observations of increased tyrosine phosphorylation of Gab1, but not Shc, in $\beta 1$ integrin^{high} cells, are consonant with these prior observations about the epidermal growth factor stimulation. The difference of the signal transduction pathway between $\alpha 6\beta 4$ integrins and $\beta 1$ integrins may make intricate regulation of differentiation and proliferation by these integrins possible.

Relationship between Gab1 and Ras/MAPK signal transduction pathway and PI3K remains unresolved. The lipid product of PI3K, phosphatidylinositol-3,4,5-trisphosphate, recruits a subset of signaling proteins with pleckstrin homology domains including Gab1 to the membrane, where they are activated [17]. This is consistent with the result suggesting that Gab1 and ERK are the downstream factors of PI3K (Fig. 3). Besides, PI3K is one of the major downstream effector cascades of Ras [20]. Nevertheless, both Gab1/Ras/MAPK and PI3K formed the intricate signal transduction pathway from $\beta 1$ integrins (Fig. 4). In $\beta 1$ integrin^{normal} cells, the inhibition of Gab1 and ERK phosphorylation by LY294002 was limited compared with the

inhibition in $\beta 1$ integrin^{high} cells (Fig. 3). It implies that various pathways or surface molecules play roles for the Gab1 and ERK phosphorylation and the engagement of PI3K and $\beta 1$ integrin is partial in $\beta 1$ integrin^{normal} cells. Then, the marked inhibition in $\beta 1$ integrin^{high} cells suggests the critical role of PI3K in the activation of Gab1 and ERK by the increase of $\beta 1$ integrins. Whereas we used HaCaT cells for acquisition of enough abundant and uniform cells to determine the precise extent of tyrosine phosphorylation of signaling molecules, further investigations in primary keratinocytes may be needed for the confirmation of the role in vivo.

In conclusion, the results of this study have provided a novel mechanism of signaling alteration by increased expression of $\beta 1$ integrins, which leads to increased MAPK activation via Gab1 and enhanced PI3K activity via augmented association of $\beta 1$ integrins and PI3K, indicating the functional relevance in keratinocyte stem cells or malignant cells. Especially, tonic signaling from high surface density of $\beta 1$ integrins may be important to establish constitutive activation in keratinocyte stem cells. These signal transduction pathways may be worth intense explorations for understanding the regulation of the stem cell compartment as well as malignancies of the keratinocyte.

Acknowledgments

This work was supported by grants from the Ministry of Education, Science and Culture of Japan and from the Ministry of Health, Labour and Welfare of Japan.

References

- [1] P.A. Hall, F.M. Watt, Stem cells: the generation and maintenance of cellular diversity, *Development* 106 (1989) 619–633.
- [2] F.M. Watt, M.D. Kubler, N.A. Hotchin, L.J. Nicholson, J.C. Adams, Regulation of keratinocyte terminal differentiation by integrin-extracellular matrix interactions, *J. Cell Sci.* 106 (Pt 1) (1993) 175–182.
- [3] L. Levy, S. Broad, D. Diekmann, R.D. Evans, F.M. Watt, Beta1 integrins regulate keratinocyte adhesion and differentiation by distinct mechanisms, *Mol. Biol. Cell* 11 (2000) 453–466.
- [4] F.M. Watt, Role of integrins in regulating epidermal adhesion, growth and differentiation, *Embo J.* 21 (2002) 3919–3926.
- [5] P.H. Jones, S. Harper, F.M. Watt, Stem cell patterning and fate in human epidermis, *Cell* 80 (1995) 83–93.
- [6] P.H. Jones, F.M. Watt, Separation of human epidermal stem cells from transit amplifying cells on the basis of differences in integrin function and expression, *Cell* 73 (1993) 713–724.
- [7] A. Gandarillas, F.M. Watt, c-Myc promotes differentiation of human epidermal stem cells, *Genes Dev.* 11 (1997) 2869–2882.
- [8] U.B. Jensen, S. Lowell, F.M. Watt, The spatial relationship between stem cells and their progeny in the basal layer of human epidermis: a new view based on whole-mount labelling and lineage analysis, *Development* 126 (1999) 2409–2418.
- [9] A.J. Zhu, I. Haase, F.M. Watt, Signaling via beta1 integrins and mitogen-activated protein kinase determines human epidermal stem cell fate in vitro, *Proc. Natl. Acad. Sci. USA* 96 (1999) 6728–6733.
- [10] S.M. Janes, F.M. Watt, New roles for integrins in squamous-cell carcinoma, *Nat. Rev. Cancer* 6 (2006) 175–183.

- [11] T. Cai, K. Nishida, T. Hirano, P.A. Khavari, Gab1 and SHP-2 promote Ras/MAPK regulation of epidermal growth and differentiation, *J. Cell Biol.* 159 (2002) 103–112.
- [12] M. Fujimoto, A.P. Bradney, J.C. Poe, D.A. Steeber, T.F. Tedder, Modulation of B lymphocyte antigen receptor signal transduction by a CD19/CD22 regulatory loop, *Immunity* 11 (1999) 191–200.
- [13] M. Fujimoto, Y. Kuwano, R. Watanabe, N. Asashima, H. Nakashima, S. Yoshitake, H. Okochi, K. Tamaki, J.C. Poe, T.F. Tedder, S. Sato, B cell antigen receptor and CD40 differentially regulate CD22 tyrosine phosphorylation, *J. Immunol* 176 (2006) 873–879.
- [14] A. Howe, A.E. Aplin, S.K. Alahari, R.L. Juliano, Integrin signaling and cell growth control, *Curr. Opin. Cell Biol.* 10 (1998) 220–231.
- [15] F. Mainiero, C. Murgia, K.K. Wary, A.M. Curatola, A. Pepe, M. Blumberg, J.K. Westwick, C.J. Der, F.G. Giancotti, The coupling of $\alpha 6 \beta 4$ integrin to Ras-MAP kinase pathways mediated by Shc controls keratinocyte proliferation, *Embo J.* 16 (1997) 2365–2375.
- [16] S.M. Frisch, R.A. Screaton, Anoikis mechanisms, *Curr. Opin. Cell Biol.* 13 (2001) 555–562.
- [17] L.C. Cantley, The phosphoinositide 3-kinase pathway, *Science* 296 (2002) 1655–1657.
- [18] H.W. Jeong, I.S. Kim, TGF- $\beta 1$ enhances β 1-integrin-mediated keratinocyte cell migration through the $\alpha 3 \beta 1$ integrin and PI3K, *J. Cell. Biochem.* 92 (2004) 770–780.
- [19] J.C. Adams, F.M. Watt, Changes in keratinocyte adhesion during terminal differentiation: reduction in fibronectin binding precedes $\alpha 5 \beta 1$ integrin loss from the cell surface, *Cell* 63 (1990) 425–435.
- [20] J.M. Shields, K. Pruitt, A. McFall, A. Shaub, C.J. Der, Understanding Ras: ‘it ain’t over ‘til it’s over’, *Trends Cell Biol.* 10 (2000) 147–154.
- [21] M. Dajee, M. Tarutani, H. Deng, T. Cai, P.A. Khavari, Epidermal Ras blockade demonstrates spatially localized Ras promotion of proliferation and inhibition of differentiation, *Oncogene* 21 (2002) 1527–1538.
- [22] K.M. Lai, T. Pawson, The ShcA phosphotyrosine docking protein sensitizes cardiovascular signaling in the mouse embryo, *Genes Dev.* 14 (2000) 1132–1145.
- [23] M. Itoh, Y. Yoshida, K. Nishida, M. Narimatsu, M. Hibi, T. Hirano, Role of Gab1 in heart, placenta, and skin development and growth factor- and cytokine-induced extracellular signal-regulated kinase mitogen-activated protein kinase activation, *Mol. Cell. Biol.* 20 (2000) 3695–7304.
- [24] M. Sachs, H. Brohmann, D. Zechner, T. Muller, J. Hulsken, I. Walther, U. Schaeper, C. Birchmeier, W. Birchmeier, Essential role of Gab1 for signaling by the c-Met receptor in vivo, *J. Cell Biol.* 150 (2000) 1375–1384.