

## RAP80 interacts with the SUMO-conjugating enzyme UBC9 and is a novel target for sumoylation

Jun Yan, Xiao-Ping Yang, Yong-Sik Kim, Joung Hyuck Joo, Anton M. Jetten \*

*Cell Biology Section, Division of Intramural Research, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, NC 27709, USA*

Received 26 July 2007  
Available online 8 August 2007

### Abstract

RAP80, a nuclear protein with two functional ubiquitin-interaction motifs (UIMs) at its N-terminus, plays a critical role in the regulation of estrogen receptor alpha and DNA damage response signaling. A yeast two-hybrid screen identified the SUMO-conjugating enzyme UBC9 as a protein interacting with RAP80. The interaction of RAP80 with UBC9 was confirmed by co-immunoprecipitation and GST pull-down analyses. The region between aa 122–204 was critical for the interaction of RAP80 with UBC9. In addition, we demonstrate that RAP80 is a target for SUMO-1 modification in intact cells. Expression of UBC9 enhanced RAP80 mono-sumoylation and also induced multi-sumoylation of RAP80. In addition to SUMO-1, RAP80 was efficiently conjugated to SUMO-3 but was only a weak substrate for SUMO-2 conjugation. These findings suggest that sumoylation plays a role in the regulation of RAP80 functions. Published by Elsevier Inc.

**Keywords:** RAP80; UBC9; SUMO-1; Sumoylation

Post-translational modifications are important in controlling the conformation, function, and activity of proteins and, as such, play a critical role in the regulation of many cellular processes. Ubiquitination and sumoylation represent a unique type of post-translational modification, in which ubiquitin or ubiquitin-like proteins are covalently conjugated to lysine residues in target proteins [1–6]. Poly-ubiquitination plays a key role in guiding the proteolytic degradation of proteins by the ubiquitin-proteasome system while monoubiquitination has been implicated in several processes, including the regulation of endocytosis, DNA-repair, and transcription [1,2,7].

Members of the small ubiquitin-related modifier (SUMO) family are structurally related to ubiquitin [2–5]. SUMO-1 shows ~50% amino acid sequence identity to SUMO-2 and

SUMO-3 while SUMO-2 and SUMO-3 exhibit ~95% amino acid sequence identity to each other. The pathway of sumoylation is analogous to that of ubiquitination, but involves a different set of enzymes. SUMO is activated by an E1-activating enzyme consisting of an AOS1/UBA2 heterodimer, transferred to the E2-conjugating enzyme, UBC9, and is subsequently (with the aid of various E3 ligases) attached to the  $\epsilon$ -amino group of specific lysines in target proteins [2–5]. Sumoylation can be reversed by SUMO-cleaving enzymes. In some cases, UBC9 can catalyze sumoylation without the help of E3 ligases, but its efficiency and specificity is increased by E3 ligases. Sumoylation is emerging as a critical modifier of the conformation and activity of many, mostly nuclear, proteins with functions in many cellular processes, including transcriptional regulation, DNA damage repair, histone modification, and apoptosis [6,8–11].

Receptor-associated protein 80 (RAP80) (HUGO nomenclature: ubiquitin-interaction motif containing 1 or UIMC1) is a nuclear protein identified in our laboratory [12,13]. It contains two putative Cys-X<sub>2</sub>-Cys-X<sub>11</sub>-His-X<sub>3</sub>-Cys zinc finger-like motifs at its carboxyl-terminus and two

*Abbreviations:* GST, glutathione sulfotransferase; UIM, ubiquitin-interaction motif; RAP80, receptor-associated protein 80; ER $\alpha$ , estrogen receptor  $\alpha$ ; IR, ionizing radiation.

\* Corresponding author. Fax: +1 919 541 4133.

E-mail address: [jetten@niehs.nih.gov](mailto:jetten@niehs.nih.gov) (A.M. Jetten).

ubiquitin-interacting motifs (UIMs) near its amino-terminus. Recently, we demonstrated these UIMs are able to bind (poly)ubiquitin [12]. In addition, we reported that RAP80 can interact with the estrogen receptor  $\alpha$  (ER $\alpha$ ) in an agonist-dependent manner and regulate its transcriptional activity. Moreover, RAP80 interacts with the tumor suppressor BRCA1 and is essential for the translocation of BRCA1 to DNA damage foci after irradiation [14–17]. The UIMs play a critical role in these activities of RAP80 [12,17].

To obtain further insights into the functions of RAP80, we performed a yeast two-hybrid screen to identify potential RAP80-interacting proteins. In this study, we report that the SUMO E2-conjugating enzyme UBC9 was identified as a novel RAP80 binding partner. The interaction was confirmed by co-immunoprecipitation and GST pull-down assays. In addition, we demonstrate that RAP80 is a target of sumoylation in intact cells and that UBC9 overexpression promotes multi-sumoylation of RAP80. RAP80 sumoylation was independent of its UIMs. Ionizing radiation (IR) increases RAP80 ubiquitination but not sumoylation. Both ubiquitination and sumoylation may play an important role in regulating the activity and function of RAP80.

## Materials and methods

**Plasmids.** pLXIN-3 $\times$ FLAG-RAP80, pLXIN-3 $\times$ FLAG-RAP80 $\Delta$ UIM,  $\Delta$ C582,  $\Delta$ C504,  $\Delta$ C404,  $\Delta$ C304,  $\Delta$ C204, and  $\Delta$ C122 were described previously [12,13]. Details on other RAP80, UBC9, SUMO-1, 2, and 3 plasmids used in this study are provided in [Supplementary data](#).

**Yeast two-hybrid screening.** The Gal4 yeast two-hybrid system was purchased from BD Biosciences. Library screening was conducted according to the manufacturer's instructions and details on it are provided in [Supplementary data](#).

**GST pull-down assay.** The methods of purifying GST or GST-UBC9 fusion proteins and their binding to [ $^{35}$ S]-methionine-labeled RAP80 were previously described [12].

**Co-immunoprecipitation assay.** HeLa or HEK293T cells were transiently transfected with pLXIN-3 $\times$ FLAG-RAP80 (full-length or mutants) and pCMV-Myc-UBC9 using Fugene 6 transfection reagent (Roche, Indianapolis, IN). Forty-eight hours after transfection, cells were processed as previously described [17].

**Sumoylation analysis.** HEK293T cells were transiently transfected with pLXIN-3 $\times$ FLAG-RAP80 (full-length or mutants), pCMV-HA-SUMO-1, and pCMV-Myc-UBC9 as indicated. Forty-eight hours later, cells were harvested and lysed for 1 h in modified RIPA buffer (50 mM Tris/HCl, pH 7.8, 150 mM NaCl, 5 mM EDTA, 15 mM MgCl<sub>2</sub>, 0.5% NP-40, 0.3% Triton X-100, 0.5% sodium deoxycholate, 1 mM dithiothreitol, 10 mM *N*-ethylmaleimide, and 0.1% SDS) containing protease inhibitor cocktails. The cell lysates were centrifuged for 10 min at 14,000g and 4 °C. The supernatants were incubated overnight with anti-HA antibody (Roche) and protein-G agarose (Sigma) to pull-down sumoylated proteins. The agarose was then washed three times with RIPA buffer. The bound proteins were solubilized in sample buffer and analyzed by Western blot analysis using anti-FLAG M2 antibody.

## Results


### Identification of UBC9 as a RAP80-interacting protein

To identify proteins that interact with RAP80 or modulate RAP80 function, a yeast two-hybrid screen was per-

formed using full-length RAP80 as bait and an E17 mouse embryo library as prey. From the  $1.5 \times 10^6$  cDNA clones screened, several positive clones were isolated. One of the positive clones encoded the full-length region of the SUMO-conjugating enzyme UBC9.

The interaction between RAP80 and UBC9 was validated by co-immunoprecipitation analysis. Myc-UBC9 and FLAG-RAP80 were transiently expressed in HeLa cells and cell lysates incubated with anti-FLAG M2 affinity resin to isolate FLAG-RAP80 protein complexes. FLAG-RAP80 protein complexes were then examined by Western blot analysis. As shown in [Fig. 1A](#), UBC9 was co-immunoprecipitated with FLAG-RAP80 in a specific manner. These data indicate that RAP80 and UBC9 interact with each other consistent with results of the yeast two-hybrid analysis.

We next examined the interaction of RAP80 with UBC9 by *in vitro* pull-down analysis using GST-UBC9 fusion protein and *in vitro* translated, [ $^{35}$ S]-methionine-labeled RAP80. GST-UBC9 protein complexes were isolated with glutathione-Sepharose beads and analyzed by SDS-PAGE. As shown in [Fig. 1B](#), GST-UBC9 fusion protein effectively pulled down RAP80 ([Fig. 1B](#)). Very little



**Fig. 1.** RAP80 interacts with UBC9. (A) HeLa cells were co-transfected with pCMV-Myc-UBC9 and pLXIN-3 $\times$ FLAG-RAP80 expression plasmids as indicated. Cells were harvested 48 h later and cell lysates prepared. Five percent of each cell lysate was used directly for Western blot analysis while the remaining was incubated with anti-FLAG M2 affinity resin to isolate FLAG-RAP80 protein complexes. Protein complexes were then examined by Western blot analysis using anti-Myc and anti-FLAG antibodies. (B) GST and GST-UBC9 fusion protein were bound to glutathione-Sepharose 4B beads and then incubated with [ $^{35}$ S]-methionine-labeled RAP80. After 1.5 h incubation, beads were washed extensively and bound proteins solubilized. Radiolabeled proteins were analyzed by SDS-PAGE and visualized by autoradiography. The first lane showed 10% of the input of radiolabeled RAP80. GST and GST-RAP80 proteins are shown in the lower panel.

RAP80 was bound to GST. These observations suggest that RAP80 interacts directly with UBC9.

#### Mapping the RAP80 region required for the interaction with UBC9

RAP80 contains two UIMs and two putative zinc fingers. The UIMs are critical for RAP80 function [12,17]. To determine which region of RAP80 was required for its interaction with UBC9, a series of RAP80 deletion mutants (Fig. 2A) were examined by co-immunoprecipitation analysis for their ability to bind UBC9. The results demonstrated that the UIMs of RAP80 were not required for its interaction with UBC9 (Fig. 2B). Carboxyl-terminal deletions up to Ser<sup>204</sup> had little effect on the ability of RAP80 to pull-down UBC9; however, RAP80 $\Delta$ C122 was unable to interact with UBC9 (Fig. 2C). These data indicated that the region between Arg<sup>122</sup> and Ser<sup>204</sup> is crucial for the interaction of RAP80 with UBC9 and that the 2 zinc finger-like motifs located at the carboxyl-terminus are not required for binding.

#### RAP80 is a novel target of sumoylation

Since UBC9 functions as an E2 SUMO-conjugating enzyme, we determined whether RAP80 was a target of sumoylation. To investigate this, HEK293T cells were transfected with pLXIN-3 $\times$ FLAG-RAP80, pCMV-HA-

SUMO-1 in the presence or absence of pCMV-Myc-UBC9. Forty-eight hours later cell extracts were prepared and sumoylated proteins immunoprecipitated with an anti-HA antibody. Analysis of immunoprecipitated proteins by Western blot analysis with an anti-FLAG antibody showed that RAP80 was sumoylated. SUMO-1 has a calculated mass of 11.5 kDa and migrates in SDS-PAGE at about 17 kDa while RAP80 migrates at 105 kDa [12,18]. Sumoylated RAP80 migrated at about 120 kDa suggesting that it is mono-sumoylated (Fig. 3A). Often sumoylated RAP80 appeared as 2–3 bands migrating closely together what might represent mono-sumoylation of different RAP80 molecules at distinct sites resulting in slightly different electrophoretic mobilities.

Co-expression with Myc-UBC9 greatly enhanced mono-sumoylation of RAP80. Moreover, several additional, higher molecular weight RAP80 species were observed suggesting that multiple lysines in RAP80 were sumoylated (Fig. 3A). These results suggested that UBC9 expression leads to multi-sumoylation of RAP80. Recent studies have reported that proteins can be poly-sumoylated *in vitro* [19,20]. To determine whether the higher molecular weight, sumoylated RAP80 species represents multi-mono-sumoylation or poly-sumoylation, we examined sumoylation of RAP80 by SUMO-1(KO) in which all lysines were mutated. SUMO-1(KO) cannot form a poly-SUMO-1 chain but can still mono-sumoylate proteins. As shown in Supplementary Fig. 1, RAP80 is still multi-sumoylated




Fig. 2. Mapping of the UBC9-interacting region in RAP80. (A) Schematic view of the different RAP80 deletion mutants. UIM and ZF indicate the two ubiquitin-interacting motifs and putative zinc fingers, respectively. (B, C) Effect of different mutation on the interaction of RAP80 with UBC9. HeLa cells were co-transfected with pCMV-Myc-UBC9 and pLXIN-3 $\times$ FLAG-RAP80 or pLXIN-3 $\times$ FLAG-RAP80 mutants shown in (A). After 48 h incubation cells were collected and processed as described in the legend to Fig. 1A.




Fig. 3. RAP80 is a target for *in vivo* sumoylation. (A) HEK293T cells were transfected with pLXIN-3×FLAG-RAP80, pCMV-HA-SUMO-1, and pCMV-Myc-UBC9 as indicated. Forty-eight hours later cell extracts were prepared and sumoylated proteins immunoprecipitated with an anti-HA antibody. Isolated proteins were subsequently examined by Western blot analysis with an anti-FLAG antibody. (B) HEK293T cells were transfected with pLXIN-3×FLAG-RAP80, pCMV-HA-SUMO-1, pcDNA3-HA-SUMO-2 or pcDNA3-HA-SUMO-3 as indicated. Forty-eight hours later cell extracts were prepared and sumoylated proteins immunoprecipitated with anti-HA antibody. Proteins were then examined by Western blot analysis with an anti-FLAG antibody. (C) HEK293T cells were transfected with pLXIN-3×FLAG-RAP80, pCMV-HA-SUMO-1, or pCMV-HA-Ub as indicated. Forty-eight hours later cells were irradiated (10 Gy) and collected after 3 h incubation. Cells extracts were immunoprecipitated with anti-HA antibody. Proteins were then examined by Western blot analysis with an anti-FLAG antibody.

when co-expressed with UBC9 and SUMO-1(KO) supporting the conclusion that RAP80 is multi-mono-sumoylated on distinct lysines.

To determine whether RAP80 could be covalently modified by SUMO-2 and SUMO-3, HEK293T cells were transfected with pLXIN-3×FLAG-RAP80 and expression vectors encoding HA-tagged SUMO-1, -2, or -3 and 48 h later the level of sumoylated RAP80 analyzed. As shown in Fig. 3B, RAP80 was efficiently conjugated to SUMO-1 and -3 but was only a weak substrate for SUMO-2 conjugation.

IR induces translocation of RAP80 to DNA damage foci [14–17]. We, therefore, examined whether IR had an effect on RAP80 sumoylation. As shown in Fig. 3C, IR did not influence the extent of RAP80 sumoylation. In contrast to sumoylation, IR enhanced RAP80 ubiquitination.

#### The amino-terminus of RAP80 is sumoylated

Previously, it was reported that the UIMs of RAP80 can bind (poly)ubiquitin and are of critical importance for several RAP80 functions [12,14–17]. We therefore examined whether the UIMs had a role in the sumoylation of RAP80. As shown in Supplementary Fig. 2, as wild-type RAP80, FLAG-RAP80ΔUIM was efficiently mono-sumoylated. Deletion of the UIMs also did not affect the multi-sumoylation of RAP80 induced by exogenous

UBC9 expression. These results suggest that the UIMs are not required for RAP80 sumoylation.

To obtain greater insight into the regions within RAP80 that are targeted for sumoylation, the effect of a series of carboxyl-terminal truncations on the sumoylation of RAP80 was examined. As was often observed for full-length RAP80, sumoylated RAP80ΔC582 appeared as doublet what might be due to mono-sumoylation of different RAP80 molecules. RAP80ΔC122, the shortest deletion mutant tested, was still sumoylated (Fig. 4A). These observations indicated that the amino-terminus of RAP80 contains lysine(s) that are targeted for sumoylation.

As shown in Fig. 4A, the amino-terminus of RAP80 contains 17 lysine residues, some of which are part of a nuclear localization signal [13]. Sequence analysis for potential sumoylation consensus motifs (<http://www.abgent.com/doc/sumoplot> and <http://bioinformatics.lcd-ustc.org/sumosp/prediction.php>) revealed that this region does not contain any high-probability sumoylation sites but has five low-probability sumoylation sites at K9, K19, K31, K52, and K61. Because K52 received the highest score, we first examined the effect of the K52A single mutation on RAP80 sumoylation. The result in Fig. 4B shows that this mutation did not affect RAP80 sumoylation. Next, we examined the effects of the K9A, K19A, K31A, and K61A single mutations and the K19,52A double mutation on the sumoylation of RAP80ΔC122. None of these






Fig. 4. RAP80 is sumoylated at its amino-terminus. (A) HEK293T cells were transfected with pCMV-HA-SUMO-1 and a series of carboxyl-terminal deletion mutants of pLXIN-3×FLAG-RAP80. Forty-eight hours later sumoylated proteins were immunoprecipitated with an anti-HA antibody and examined by Western blot analysis with an anti-FLAG antibody. (\*) The migration of the sumoylated mutant RAP80 proteins. (B) Effect of several mutations on RAP80 sumoylation. The amino acid sequence of the N-terminal region of RAP80 (aa 1–122) is shown; potential sumoylation sites are underlined. HEK293T cells were transfected with pCMV-HA-SUMO-1, wild-type pLXIN-3×FLAG-RAP80 or its K52A mutant. The cells were collected 48 h later and processed as described under (A). (C) HEK293T cells were transfected with pCMV-HA-SUMO-1 and wild-type pLXIN-3×FLAG-RAP80 $\Delta$ C122 or RAP80 $\Delta$ C122 mutants as indicated. The cells were collected 48 h later and processed as described under (A). 5K/A indicates the RAP80 $\Delta$ C122 (5K/A) mutant in which all five potential sites are mutated.

mutations affected sumoylation (Fig. 4C). Lastly, we generated a RAP80 $\Delta$ C122(5K/A) mutant in which all 5 potential sites were mutated. Fig. 4C shows that RAP80 $\Delta$ C122(5K/A) was still sumoylated.

To determine whether RAP80 $\Delta$ C122 was multi-sumoylated when Myc-UBC9 was co-expressed, HEK293T cells were transfected with pLXIN-3×FLAG-RAP80 $\Delta$ C122, pCMV-HA-SUMO-1, and pCMV-Myc-UBC9. Forty-eight hours later cells were collected and sumoylation of RAP80 $\Delta$ C122 analyzed. The data show that UBC9 effectively promoted multi-sumoylation of RAP80 $\Delta$ C122 (Supplementary Fig. 3).

## Discussion

Previously, we reported that RAP80 interacts with ER $\alpha$  in an agonist-dependent manner and regulates its stability and transcriptional activity [12]. Recently, we and others demonstrated that RAP80 interacts with the tumor suppressor BRCA1 and has a critical role in DNA damage repair response [14–17]. The UIMs which bind (poly)ubiquitin, play a critical role in these activities of RAP80. In this study, we identified using a yeast two-hybrid screen UBC9 as a novel interacting partner of RAP80. This interaction was confirmed by immunoprecipitation and *in vitro* GST-pull-down analysis.

Our study also demonstrates that RAP80 is a novel target of sumoylation. RAP80 was effectively conjugated

to SUMO-1 and SUMO-3 and to a much lesser extent to SUMO-2. This is in agreement with other studies showing that proteins can exhibit different specificities for SUMO-1–3 conjugation [21]. Based on the size of the sumoylated RAP80, it was concluded that RAP80 is mostly mono-sumoylated. Deletion analysis indicated that the amino-terminus (aa 1–122) of RAP80 can be modified by sumoylation. Although many proteins are sumoylated at the  $\Psi$ KXD/E consensus sequence, several studies have reported that proteins can be sumoylated at sites that do not conform to the  $\Psi$ KXD/E consensus sequence. For example, PML and Smad4 are sumoylated at lysines in AKCP and VKYC, respectively, while Mdm2 and CREB are sumoylated and do not contain a  $\Psi$ KXD/E sequence [4,22,23]. How these sites are recognized has not yet been established. Although the amino-terminus contains 17 lysine residues none of them are part of a high-probability  $\Psi$ KXD/E sumoylation motif. Five low-probability motifs were identified; however, mutation of all five sites did not have any effect on the sumoylation of RAP80 suggesting that another lysine can be a target for sumoylation. Sumoylation of many proteins involves binding of UBC9 to the consensus  $\Psi$ KXD/E sumoylation motif; however, in the case of other proteins, including the transcription factor AP2, the co-repressor N-COR, and Sovar, UBC9 interacts with a region distant to the sumoylation site [24–26]. The amino-terminus of RAP80 is sumoylated but does

not bind UBC9. Possibly, this sumoylation does not require a direct interaction between RAP80 and UBC9, and sumoylation is mediated by E3 ligases. Deletion analysis showed that the region of RAP80 between Arg<sup>122</sup> and Ser<sup>204</sup> is important for the interaction of RAP80 with UBC9; however, this region does not contain a high-probability ΨKXD/E sumoylation motif. The binding of UBC9 to this region may involve different interaction motif and be implicated in a different activity of RAP80. Since RAP80 is part of a larger protein complex, it may recruit UBC9 to the complex thereby promoting the sumoylation of other proteins in the complex.

Sumoylation has been reported to be a critical modifier of the conformation and activity of many, mostly nuclear, proteins with functions in a variety of cellular processes [3,6,8–11]. For example, sumoylation plays an important role in the regulation of the NF-κB signaling pathway. After sumoylation NEMO, the regulatory subunit of IκB kinase, is translocated from the cytoplasm to the nucleus [27]. In IκBα sumoylation targets the same lysines as ubiquitination thereby stabilizing the protein by preventing its ubiquitination and destruction by the proteasome [28]. Recently, we and others reported that RAP80 interacts with the tumor suppressor BRCA1 and translocates with BRCA1 to DNA damage foci after irradiation suggesting a role for RAP80 in DNA-damage repair signaling [14–17]. The UIMs play a critical role in the transfer of RAP80 to DNA damage foci, and as a consequence also in the transfer of BRCA1. Interestingly, ubiquitination and sumoylation have multiple functions in DNA damage repair signaling [6,8,9] and a number of proteins involved in DNA repair are subject to sumoylation. For example, Lys<sup>164</sup> in proliferating cell nuclear antigen (PCNA) is a target for ubiquitination as well as sumoylation [29]. By competing with ubiquitin for attachment to this lysine SUMO can block ubiquitination and inhibit damage-induced DNA repair. Rad51 and Rad52, which play a key role in homologous recombination, interact with UBC9 and are targets of sumoylation [30,31]. The DNA repair factor XPC becomes sumoylated after irradiation [32]. Sumoylation of DNA repair proteins can regulate their stability, protein–protein interactions, localization, and activity. Sumoylation did not affect the stability of RAP80 or its nuclear localization (Yan, J., unpublished observations). We hypothesize that this post-translational modification plays an important role in regulating the activity of RAP80 and its function in DNA damage response and possibly ERα signaling.

## Acknowledgments

We thank Drs. T. Wang and B. Cuthbertson for their comments on the manuscript. This research was supported by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.bbrc.2007.07.158](https://doi.org/10.1016/j.bbrc.2007.07.158).

## References

- [1] S. Fang, A.M. Weissman, A field guide to ubiquitylation, *Cell. Mol. Life Sci.* 61 (2004) 1546–1561.
- [2] G. Gill, SUMO and ubiquitin in the nucleus: different functions, similar mechanisms? *Genes Dev.* 18 (2004) 2046–2059.
- [3] R.T. Hay, SUMO: a history of modification, *Mol. Cell.* 18 (2005) 1–12.
- [4] E.S. Johnson, Protein modification by SUMO, *Annu. Rev. Biochem.* 73 (2004) 355–382.
- [5] O. Kerscher, R. Felberbaum, M. Hochstrasser, Modification of proteins by ubiquitin and ubiquitin-like proteins, *Annu. Rev. Cell. Dev. Biol.* 22 (2006) 159–180.
- [6] S. Muller, A. Ledl, D. Schmidt, SUMO: a regulator of gene expression and genome integrity, *Oncogene* 23 (2004) 1998–2008.
- [7] D. Hoeller, N. Crosetto, B. Blagoev, C. Raiborg, R. Tikkanen, S. Wagner, K. Kowanzetz, R. Breitling, M. Mann, H. Stenmark, I. Dikic, Regulation of ubiquitin-binding proteins by monoubiquitination, *Nat. Cell Biol.* 8 (2006) 163–169.
- [8] H.L. Klein, A SUMOry of DNA replication: synthesis, damage, and repair, *Cell* 127 (2006) 455–457.
- [9] S.J. Moschos, Y.Y. Mo, Role of SUMO/Ubc9 in DNA damage repair and tumorigenesis, *J. Mol. Histol.* 37 (2006) 309–319.
- [10] J.S. Seeler, O. Bischof, K. Nacerddine, A. Dejean, SUMO, the three Rs and cancer, *Curr. Top. Microbiol. Immunol.* 313 (2007) 49–71.
- [11] A. Verger, J. Perdomo, M. Crossley, Modification with SUMO. A role in transcriptional regulation, *EMBO Rep.* 4 (2003) 137–142.
- [12] J. Yan, S.-K. Kim, X.-P. Yang, M. Albers, M. Koegl, A.M. Jetten, Ubiquitin interacting motifs of RAP80 are critical in its regulation of Estrogen Receptor α, *Nucleic Acids Res.* 35 (2007) 1673–1686.
- [13] Z. Yan, Y.S. Kim, A.M. Jetten, RAP80, a novel nuclear protein that interacts with the retinoid-related testis-associated receptor, *J. Biol. Chem.* 277 (2002) 32379–32388.
- [14] H. Kim, J. Chen, X. Yu, Ubiquitin-binding protein RAP80 mediates BRCA1-dependent DNA damage response, *Science* 316 (2007) 1202–1205.
- [15] B. Sobhian, G. Shao, D.R. Lilli, A.C. Culhane, L.A. Moreau, B. Xia, D.M. Livingston, R.A. Greenberg, RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites, *Science* 316 (2007) 1198–1202.
- [16] B. Wang, S. Matsuoka, B.A. Ballif, D. Zhang, A. Smogorzewska, S.P. Gygi, S.J. Elledge, Abraxas and RAP80 form a BRCA1 protein complex required for the DNA damage response, *Science* 316 (2007) 1194–1198.
- [17] J. Yan, Y.S. Kim, X.P. Yang, L.P. Li, G. Liao, F. Xia, A.M. Jetten, The ubiquitin-interacting motif containing protein RAP80 interacts with BRCA1 and functions in DNA damage repair response, *Cancer Res.* 67 (2007) 6647–6656.
- [18] M.J. Matunis, J. Wu, G. Blobel, SUMO-1 modification and its role in targeting the Ran GTPase-activating protein, RanGAP1, to the nuclear pore complex, *J. Cell Biol.* 140 (1998) 499–509.
- [19] C.B. Gocke, H. Yu, J. Kang, Systematic identification and analysis of mammalian small ubiquitin-like modifier substrates, *J. Biol. Chem.* 280 (2005) 5004–5012.
- [20] M. Yang, C.T. Hsu, C.Y. Ting, L.F. Liu, J. Hwang, Assembly of a polymeric chain of SUMO1 on human topoisomerase I in vitro, *J. Biol. Chem.* 281 (2006) 8264–8274.
- [21] Z. Ji, C. Degerny, N. Vintonenko, J. Deheuninck, B. Foveau, C. Leroy, J. Coll, D. Tulasne, J.L. Baert, V. Fafeur, Regulation of the Ets-1 transcription factor by sumoylation and ubiquitylation, *Oncogene* 26 (2007) 395–406.

- [22] K.M. Comerford, M.O. Leonard, J. Karhausen, R. Carey, S.P. Colgan, C.T. Taylor, Small ubiquitin-related modifier-1 modification mediates resolution of CREB-dependent responses to hypoxia, *Proc. Natl. Acad. Sci. USA* 100 (2003) 986–991.
- [23] X. Lin, M. Liang, Y.Y. Liang, F.C. Brunicardi, X.H. Feng, SUMO-1/Ubc9 promotes nuclear accumulation and metabolic stability of tumor suppressor Smad4, *J. Biol. Chem.* 278 (2003) 31043–31048.
- [24] J.J. Eloranta, H.C. Hurst, Transcription factor AP-2 interacts with the SUMO-conjugating enzyme UBC9 and is sumolated in vivo, *J. Biol. Chem.* 277 (2002) 30798–30804.
- [25] J. Tiefenbach, N. Novac, M. Ducasse, M. Eck, F. Melchior, T. Heinzel, SUMOylation of the corepressor N-CoR modulates its capacity to repress transcription, *Mol. Biol. Cell* 17 (2006) 1643–1651.
- [26] Z. Fan, Y. Zhuo, X. Tan, Z. Zhou, J. Yuan, B. Qiang, J. Yan, X. Peng, G.F. Gao, SARS-CoV nucleocapsid protein binds to hUbc9, a ubiquitin conjugating enzyme of the sumoylation system, *J. Med. Virol.* 78 (2006) 1365–1373.
- [27] T.T. Huang, S.M. Wuerzberger-Davis, Z.H. Wu, S. Miyamoto, Sequential modification of NEMO/IKKgamma by SUMO-1 and ubiquitin mediates NF-kappaB activation by genotoxic stress, *Cell* 115 (2003) 565–576.
- [28] J.M. Desterro, M.S. Rodriguez, R.T. Hay, SUMO-1 modification of IkappaBalpha inhibits NF-kappaB activation, *Mol. Cell* 2 (1998) 233–239.
- [29] C. Hoege, B. Pfander, G.L. Moldovan, G. Pyrowolakis, S. Jentsch, RAD6-dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO, *Nature* 419 (2002) 135–141.
- [30] O.V. Kovalenko, A.W. Plug, T. Haaf, D.K. Gonda, T. Ashley, D.C. Ward, C.M. Radding, E.I. Golub, Mammalian ubiquitin-conjugating enzyme Ubc9 interacts with Rad51 recombination protein and localizes in synaptonemal complexes, *Proc. Natl. Acad. Sci. USA* 93 (1996) 2958–2963.
- [31] M. Sacher, B. Pfander, C. Hoege, S. Jentsch, Control of Rad52 recombination activity by double-strand break-induced SUMO modification, *Nat. Cell Biol.* 8 (2006) 1284–1290.
- [32] Q.E. Wang, Q. Zhu, G. Wani, M.A. El-Mahdy, J. Li, A.A. Wani, DNA repair factor XPC is modified by SUMO-1 and ubiquitin following UV irradiation, *Nucleic Acids Res.* 33 (2005) 4023–4034.