



Synergistic action of gastrin and ghrelin on gastric acid secretion in rats

Kaori Fukumoto^a, Keiko Nakahara^a, Tetsuro Katayama^b, Mikiya Miyazatao^c, Kenji Kangawa^c, Noboru Murakami^{a,*}

^a Department of Veterinary Physiology, Faculty of Agriculture, University of Miyazaki, Miyazaki 889-2192, Japan

^b Frontier Science Research Center, Kibana Campus, University of Miyazaki, Miyazaki 889-2192, Japan

^c Department of Biochemistry, National Cardiovascular Center Research Institute, Fujishirodai 5-7-1, Suita, Osaka 565-8565, Japan

ARTICLE INFO

Article history:

Received 16 June 2008

Available online 9 July 2008

Keywords:

Gastric acid
Gastrin
Ghrelin

ABSTRACT

Gastrin and ghrelin are secreted from G cells and X/A-like cells in the stomach, respectively, and respective hormones stimulate gastric acid secretion by acting through histamine and the vagus nerve. In this study, we examined the relationship between gastrin, ghrelin and gastric acid secretion in rats. Intravenous (iv) administration of 3 and 10 nmol of gastrin induced transient increases of ghrelin levels within 10 min in a dose-dependent manner. Double immunostaining for ghrelin and gastrin receptor revealed that a proportion of ghrelin cells possess gastrin receptors. Although (iv) administration of gastrin or ghrelin induced significant gastric acid secretion, simultaneous treatment with both hormones resulted in a synergistic, rather than additive, increase of gastric acid secretion. This synergistic increase was not observed in vagotomized rats.

These results suggest that gastrin may directly stimulate ghrelin release from the stomach, and that both hormones may increase gastric acid secretion synergistically.

© 2008 Elsevier Inc. All rights reserved.

Ghrelin, recently purified from rat and human stomachs as an endogenous ligand for the growth hormone (GH) secretagogue receptor (GHS-R), is a 28-amino-acid peptide with an *n*-octanoylation modification at the Ser3 residue [1]. This octanoylation is essential for stimulation of GH secretion from the pituitary gland [1]. Although cells showing immunostaining for ghrelin are distributed widely in the stomach, hypothalamus, pituitary gland, liver, kidney, pancreas and placenta, the main source of circulating ghrelin has been considered to be the gastrointestinal tract, especially in X/A-like cells of fundic glands [2–6]. Several studies on the physiological function of ghrelin have demonstrated that, in addition to stimulating GH secretion, ghrelin also stimulates food intake, body weight gain, stomach motility, insulin release, cell proliferation and gastric acid secretion [7–13].

It is well known that ghrelin secretion is stimulated by starvation, and that subsequently this increase of ghrelin induces food intake [7–9,14]. Certainly, central and peripheral treatment with ghrelin causes initiation of food intake within 5 min in rats [9,14]. Therefore, ghrelin has been considered an appetite-promoting hormone. In addition to regulation of ghrelin secretion by the autonomic nervous system and nutrients, some previous studies have demonstrated that ghrelin secretion is influenced by many gastrointestinal hormones, such as glucagon, insulin, cholecystokinin, glucagon-like peptide, gastrin and somatostatin

[15–21]. However, the relationship between ghrelin secretion influenced by these gastrointestinal hormones and the physiological role of secreted ghrelin is still largely unknown. Circulating ghrelin levels exhibit a diurnal pattern, with bimodal peaks occurring before the dark and the light periods, respectively [14]. These two peaks are consistent with the minimum (emptiness) and maximum (satiety) volumes of the gastric content, respectively. This indicates that ghrelin secretion increases not only when the stomach is empty but also when it is full. Therefore, it has been assumed that the function of the ghrelin peak observed at stomach emptiness is to stimulate food intake, whereas that of the satiety peak is to stimulate gastric acid secretion [14]. If this hypothesis is correct, it is important to clarify what stimulates ghrelin secretion under satiety conditions. We have suspected that gastrin may be involved in this ghrelin increase at satiety, since gastrin is released at this time and stimulates gastric acid secretion, whereas somatostatin and cholecystokinin inhibit gastric acid secretion [22].

In this study using rats, therefore, we examined whether intravenous injection of gastrin affects ghrelin release from the stomach, or whether ghrelin and gastrin stimulate gastric acid secretion independently or cooperatively.

Materials and methods

Animals. Male Wistar rats weighing 400–500 g were used in all experiments. The rats were maintained under a 12:12-h light/dark

* Corresponding author. Fax: +81 985 58 7265.

E-mail address: a0d201u@cc.miyazaki-u.ac.jp (N. Murakami).

cycle (lights on at 07.00 h) and a room temperature of $23 \pm 1^\circ\text{C}$ with standard laboratory food and water provided *ad libitum*. All procedures were performed in accordance with the Japanese Physiological Society's guidelines for animal care.

Gastrin administration. Rats were anesthetized with an intraperitoneal (ip) injection of pentobarbital. To exclude the possibility that several treatments, such as implantation of a polyethylene tube into a stomach vein, or injection of saline and gastrin, might affect ghrelin secretion by stimulating the endings of sensory neurons in the stomach or vagus nerve, rats were firstly vagotomized at the subphrenic esophagus level using the method described previously [23] and were then maintained at a body temperature at 37°C by a small-animal heat controller while monitoring the rectal temperature (Unique Medical Co., Ltd., Tokyo, Japan). Next, to analyze ghrelin secretion into the gastric vein, a heparin-treated PE10 tube (inner diameter 0.28 mm; Becton–Dickinson, NJ, USA) was inserted and fixed to the gastric vein [20]. Blood samples were collected at 10-min intervals. After the third blood sample collection, 0.2 ml of 3 or 10 nmol gastrin (Peptide Institute, Osaka, Japan), or saline as a control, was injected into the femoral vein. Thereafter, blood sampling was continued for 60 min.

Measurement of plasma ghrelin. Ghrelin concentration was measured by the method described previously [12]. Briefly, blood samples were collected into chilled polypropylene tubes containing a protease inhibitor, aprotinin (Sigma–Aldrich, St. Louis, USA), and EDTA-2Na, and immediately centrifuged at 14,000 rpm for 3 min. Plasma samples were acidified with a 10% volume of 1 N HCl and stored at -80°C until assay. Acyl-ghrelin was measured using an active ghrelin ELISA kit (Mitsubishi Kagaku Iatron, Tokyo, Japan).

Double immunostaining for ghrelin and gastrin receptor. A trimmed part of the glandular stomach was washed using saline and placed in 0.34% formalin for 4 days at 4°C , and then transferred to 0.1 M phosphate buffer containing 20% sucrose. Sections were cut at a thickness of $18\ \mu\text{m}$ with a cryostat at a temperature of -20°C . The sections were fixed with 40% paraformaldehyde for 20 min and blocked for 1 h in 5% normal donkey serum in PBST and then incubated overnight at 4°C with rabbit antiserum against rat ghrelin together with goat antiserum against CCKB-R C18 (Santa Cruz Biotechnology, Santa Cruz, CA). This anti-rat ghrelin antibody (antiserum #G606) specifically recognizes ghrelin with *n*-octanoylated Ser-3 and does not recognize des-acyl-ghrelin. After washing, the sections were incubated with a second antibody solution of Alexa-488-labeled anti-rabbit IgG antibody and Alexa-555-labeled donkey anti-goat IgG antibody solution for 30 min. The samples were observed using a fluorescence microscope (Axioskop 2plus; Zeiss, MA, USA). Digital images were contrasted and color-adjusted using Adobe Photoshop 7.0 for Windows.

Measurement of gastric acid secretion. Intact and vagotomized rats were used in this study. Vagotomy was performed at the subphrenic esophagus level under pentobarbital anesthesia as mentioned above. Four days after the operation, the rats were used for the following experiment. After anesthesia with urethane, two gastric perfusion cannulae were implanted in the pyloric and cardiac parts of the stomach by insertion from the duodenum and esophagus, respectively. The body temperature was maintained at 37°C until the end of sampling. Five milliliters of water warmed at 37°C was perfused from the cardiac side to the pyloric part side with an infusion pump at a flow rate of 2.5 ml/min at 10-min intervals. Acidity of the perfusate was recorded by the method reported previously [24]. After acidity had been stabilized at between pH5 and pH7, 3 nmol of either gastrin or ghrelin, or 3 nmol of both gastrin and ghrelin was injected into the femoral vein. Injection of saline was used as a control. After treatment with the agents, the perfusate was collected at 10-min intervals for 60 min.

Data analysis. Data are expressed as means \pm SEM for the groups. Comparisons between groups were performed using unpaired *t*-test or ANOVA followed by the Tukey–Kramer test for multiple comparisons. Changes in plasma ghrelin levels over time were analyzed using repeated measures ANOVA followed by Dunnett's test. Differences at $P < 0.05$ were considered significant.

Result and discussion

The effect of intravenous injection of gastrin on ghrelin secretion from the stomach in vagotomized rats is shown in Fig. 1C. The plasma concentration of ghrelin in the gastric vein at the start of the experimental period, at 0 min in Fig. 1C, was $44.7 \pm 5\ \text{fmol/ml}$. Saline administration did not cause significant changes in plasma ghrelin levels after treatment. On the other hand, gastrin induced a significant dose-dependent increase in ghrelin levels. The peak level of ghrelin was observed 10 min after treatment with 3 and 10 nmol of gastrin, thereafter falling rapidly to below the control level (Fig. 1C). This decrease of the ghrelin level to below the baseline may be due to exhaustion of stored ghrelin after secretion in quantity. These results indicate that gastrin may play an important role in ghrelin secretion, and that the effect is probably limited to stimulation of secretion, and not synthesis.

Double-staining immunohistochemistry for ghrelin and gastrin receptor in the lamina propria mucosae of the lower gastric corpus was performed to examine the possible presence of gastrin receptors on ghrelin cells. Cells immunoreactive for ghrelin and gastrin




Fig. 1. Effect of peripheral injection of gastrin on plasma ghrelin levels. (A) Photograph showing cannulation of stomach vein (small arrow) by a PE10 tube (large arrow). (B) Body temperature was maintained constantly using a thermostat. (C) Ghrelin levels after injection of saline, and with 3 nmol and 10 nmol gastrin into the femoral vein. Each symbol and vertical bar represent means \pm SEM ($n = 6$). Asterisks indicate significant differences (* $P < 0.01$, ** $P < 0.05$ vs saline).




Fig. 2. Double immunostaining for ghrelin and gastrin receptors in rat stomach. Distribution of ghrelin (A: stained green) and gastrin (B: stained red) receptors from the neck to the base of a rat oxyntic gland. The scale bars represent 100 μ m. (C) Co-localization of ghrelin and gastrin receptors is observed in a few cells. d,e: Higher magnification of cells stained for both ghrelin and gastrin receptors. The scale bars represent 5 μ m.

receptor were abundant from the neck to the base of the oxyntic glands (Fig. 2A and B). A few cells were reactive for both antibodies (Fig. 2C). At high magnification, gastrin receptors were detected as a spot on the border of ghrelin cells (Fig. 2D and E). This was probably because the antibody for gastrin receptor recognizes the extracellular domain of the receptor molecule, which has a scattered distribution on the cell membrane. These results suggested that a proportion of ghrelin cells express gastrin receptors on their plasma membrane, and that gastrin may stimulate ghrelin secretion by acting directly on ghrelin cells.

In intact rats, injection of 3 nmol of ghrelin or gastrin induced significant gastric acid secretion, gastrin showing the more potent effect (Fig. 3A). Simultaneous administration of ghrelin and gastrin caused an increase of gastric acid secretion, the effect being more potent and longer lasting than that of each agent administered alone. The increase of gastric acid secretion induced by simultaneous injection of ghrelin and gastrin suggested a synergistic rather than an additive effect. These results suggest that gastrin and ghrelin stimulate gastric acid secretion cooperatively. In vagotomized rats, injection of gastrin also increased gastric acid secretion, as seen in intact rats, but acidity at 20 min after treatment was significantly decreased in comparison with that of intact rats (pH 6.23 ± 0.22 in vagotomized rats vs pH 4.83 ± 0.25 in intact rats) (Fig. 3B). In addition, no significant increase was observed after treatment with ghrelin. This lack of ghrelin-induced gastric acid secretion in vagotomized rats is in agreement with a previous report by Masuda et al. [10]. Vagotomy also eliminated the synergistic increase of gastric acid stimulated by simultaneous treatment with gastrin and ghrelin (Fig. 3B). Therefore, the vagus nerve may be involved in the synergistic effect of gastrin and ghrelin on gastric acid secretion.




Fig. 3. Effect of peripheral injection of gastrin and ghrelin on gastric acid secretion in intact and vagotomized rats. Ghrelin and/or gastrin were injected into the femoral vein of intact (A) or vagotomized (B) rats under urethane anesthesia. Gastric acidity was determined by measurement the pH of the gastric perfusate. Vagotomy was performed 4 days before the experiment. A synergistic effect on gastric acid secretion was observed upon concomitant treatment with gastrin and ghrelin. Each symbol and vertical bar represent means \pm SEM ($n = 6$). Asterisks indicate significant differences (* $P < 0.01$, ** $P < 0.05$ vs saline).

It is well known that gastrin induces gastric acid secretion from parietal cells of the stomach by stimulation of histamine release [25]. The present study suggests that gastrin also induces gastric acid secretion by acting through ghrelin release. Gastrin induced ghrelin release in vagotomized rats. In addition, double immunohistochemistry revealed that the plasma membrane of some ghrelin cells expressed gastrin receptors. Therefore, gastrin-induced ghrelin secretion may be due to the direct action of gastrin on X/A-like ghrelin cells [3]. Lippl et al. [19] have demonstrated that gastrin stimulates ghrelin release in cultured rat stomach cells [19]. However, in that study, the possible involvement of hormones other than gastrin in ghrelin secretion could not be excluded because of the paracrine action of gastrin on other stomach cells, such as enterochromaffin cells.

In this study, we confirmed that ghrelin stimulates gastric acid secretion by acting through the vagus nerve, since vagotomy eliminated this effect of ghrelin. Our findings also indicated that gastric acid secretion induced by gastrin alone may partially involve ghrelin secretion, i.e., the increase of gastric acid secretion after single treatment with gastrin might be induced by both histamine and ghrelin. The diminution of acidity resulting from vagotomy may be attributable to gastrin-induced ghrelin.

When gastrin and ghrelin were administered simultaneously, gastric acid secretion was increased synergistically. Although the mechanism of this synergy was not clarified, total ghrelin including that administered exogenously and that induced endogenously by gastrin might reach a level that is optimal for stimulating the vagus nerve, thereafter leading to secretion of a large amount of gastric acid. Alternatively, gastrin might increase the expression of ghrelin receptors (GHSR-1a) in the nodose ganglion, since we have already reported the possible involvement of gastrointestinal hormone in regulation of GHSR-1a synthesis in the nodose ganglion and for axonal transport of GHSR-1a from the nodose ganglion to nerve endings [26]. The fact that vagotomy eliminated the synergistic effect of concomitant treatment with gastrin and ghrelin, may support these possibilities.

We have previously reported that the level of ghrelin exhibits a diurnal pattern, with bimodal peaks occurring before dark (06:00 h) and before light (15:00 h) [14]. These two peaks were consistent with the maximum and minimum volumes of gastric content, respectively, suggesting that ghrelin secretion is induced by both gastric emptying and filling. The former peak is coincident with the rise before a meal reported in many mammals, and acts to stimulate food intake [27–29]. The present study indicates that this later peak might occur through the action of gastrin.

In summary, the present study has shown that some ghrelin cells in gastric glands of the rat stomach possess gastrin receptors, whereby gastrin acts directly on ghrelin cells to stimulate ghrelin secretion. Our findings suggest that secreted ghrelin stimulates gastric acid secretion, and furthermore that acid secretion stimulated by gastrin may be partly mediated by ghrelin. The increase of ghrelin secretion stimulated by gastrin results in the ghrelin peak observed in the dark period, and is closely related to the diurnal cycle of ghrelin.

Acknowledgments

This study was supported by a Grant-in-Aid for Scientific Research (S: N.M.; C: K.N.) from the Ministry of Education, Science, Sports, and Culture of Japan.

References

- [1] M. Kojima, H. Hosoda, Y. Date, M. Nakazato, H. Matsuo, K. Kangawa, Ghrelin is a growth-hormone-releasing acylated peptide from stomach, *Nature* 402 (1999) 656–660.
- [2] K. Mori, A. Yoshimoto, K. Takaya, K. Hosoda, H. Ariyasu, K. Yahata, M. Mukoyama, A. Sugawara, H. Hosoda, M. Kojima, K. Kangawa, K. Nakao, Kidney produces a novel acylated peptide, ghrelin, *FEBS Lett.* 15 (2000) 213–216.
- [3] Y. Date, M. Kojima, H. Hosoda, A. Sawaguchi, M.S. Mondal, T. Suganuma, S. Matsukura, K. Kangawa, M. Nakazato, Ghrelin, a novel growth hormone-releasing acylated peptide, is synthesized in a distinct endocrine cell type in the gastrointestinal tracts of rats and humans, *Endocrinology* 141 (2000) 4255–4261.
- [4] O. Gualillo, J. Caminos, M. Blanco, T. Garcia-Caballero, M. Kojima, K. Kangawa, C. Dieguez, F. Casanueva, Ghrelin, a novel placental-derived hormone, *Endocrinology* 142 (2001) 788–794.
- [5] M. Korbonits, S.A. Bustin, M. Kojima, S. Jordan, E.F. Adams, D.G. Lowe, K. Kangawa, A.B. Grossman, The expression of the growth hormone secretagogue receptor ligand ghrelin in normal and abnormal human pituitary and other neuroendocrine tumors, *J. Clin. Endocrinol. Metab.* 86 (2001) 881–887.
- [6] Y. Date, M. Nakazato, S. Hashiguchi, K. Dezaki, M.S. Mondal, H. Hosoda, M. Kojima, K. Kangawa, T. Arima, H. Matsuo, T. Yada, S. Matsukura, A Ghrelin is present in pancreatic alpha-cells of humans and rats and stimulates insulin secretion, *Diabetes* 51 (2002) 124–129.
- [7] A.M. Wren, C.J. Small, H.L. Ward, K.G. Murphy, C.L. Dakin, S. Taheri, A.R. Kennedy, G.H. Roberts, D.G. Morgan, M.A. Ghatei, S.R. Bloom, The novel hypothalamic peptide ghrelin stimulates food intake and growth hormone secretion, *Endocrinology* 141 (2000) 4325–4328.
- [8] M. Tschöp, D.L. Smiley, M.L. Heiman, Ghrelin induces adiposity in rodents, *Nature* 407 (2000) 908–913.
- [9] M. Nakazato, N. Murakami, Y. Date, M. Kojima, H. Matsuo, K. Kangawa, S. Matsukura, A role for ghrelin in the central regulation of feeding, *Nature* 409 (2001) 194–198.
- [10] Y. Masuda, T. Tanaka, N. Inomata, N. Ohnuma, S. Tanaka, Z. Itoh, H. Hosoda, M. Kojima, K. Kangawa, Ghrelin stimulates gastric acid secretion and motility in rats, *Biochem. Biophys. Res. Commun.* 276 (2000) 905–908.
- [11] H.M. Lee, G. Wang, E.W. Englander, M. Kojima, J.G.H. Greeley, Ghrelin, a new gastrointestinal endocrine peptide that stimulates insulin secretion: enteric distribution, ontogeny, influence of endocrine and dietary manipulations, *Endocrinology* 143 (2002) 185–190.
- [12] K. Nakahara, M. Nakagawa, Y. Baba, M. Sato, K. Toshinai, Y. Date, M. Nakazato, M. Kojima, M. Miyazato, H. Kaiya, H. Hosoda, K. Kangawa, N. Murakami, Maternal ghrelin plays an important role in rat fetal development during pregnancy, *Endocrinology* 147 (2006) 1333–1342.
- [13] M. Sato, K. Nakahara, S. Goto, H. Kaiya, M. Miyazato, Y. Date, M. Nakazato, K. Kangawa, N. Murakami, Effects of ghrelin and des-acylghrelin on neurogenesis of the rat fetal spinal cord, *Biochem. Biophys. Res. Commun.* 350 (2006) 598–603.
- [14] N. Murakami, T. Hayashida, T. Kuroiwa, K. Nakahara, T. Ida, M.S. Mondal, M. Nakazato, M. Kojima, K. Kangawa, Role for central ghrelin in food intake and secretion profile of stomach ghrelin in rats, *J. Endocrinol.* 174 (2002) 283–288.
- [15] H. Hosoda, K. Kangawa, The autonomic nervous system regulates gastric ghrelin secretion in rats, *Regul. Pept.* 146 (2007) 12–18.
- [16] G. Gomez, E.W. Englander, J.G.H. Greeley, Nutrient inhibition of ghrelin secretion in the fasted rat, *Regul. Pept.* 117 (2004) 33–36.
- [17] F. Broglio, P. van Koetsveld, A. Benso, C. Gottero, F. Prodham, M. Papotti, G. Muccioli, C. Gauna, L. Hofland, R. Deghenghi, E. Arvat, A.J. van der Lely, E. Ghigo, Ghrelin secretion is inhibited by either somatostatin or cortistatin in humans, *J. Clin. Endocrinol. Metab.* 87 (2002) 4829–4832.
- [18] J. Kamegai, H. Tamura, T. Shimizu, S. Ishii, H. Sugihara, S. Oikawa, Effects of insulin, leptin, and glucagon on ghrelin secretion from isolated perfused rat stomach, *Regul. Pept.* 119 (2004) 77–82.
- [19] F. Lippl, F. Kircher, J. Erdmann, H.-D. Allescher, V. Schusdziarra, Effect of GIP, GLP-1, insulin and gastrin on ghrelin release in the isolated rat stomach, *Regul. Pept.* 119 (2004) 93–98.
- [20] T. Katayama, S. Shimamoto, H. Oda, K. Nakahara, K. Kangawa, N. Murakami, Glucagon receptor expression and glucagon stimulation of ghrelin secretion in rat stomach, *Biochem. Biophys. Res. Commun.* 357 (2007) 865–870.
- [21] M. Shimada, Y. Date, M.S. Mondal, K. Toshinai, T. Shimbara, K. Fukunaga, N. Murakami, M. Miyazato, K. Kangawa, H. Yoshimatsu, H. Matsuo, M. Nakazato, Somatostatin suppresses ghrelin secretion from the rat stomach, *Biochem. Biophys. Res. Commun.* 302 (2003) 520–525.
- [22] M.L. Schubert, Gastric secretion, *Curr. Opin. Gastroenterol.* 23 (2007) 595–601.
- [23] Y. Date, N. Murakami, A. Nijima, S. Matsukura, H. Matsuo, K. Kangawa, M. Nakazato, The role of gastric vagal afferent in feeding and growth hormone secretion effects of ghrelin, *Gastroenterology* 123 (2002) 1120–1128.
- [24] G. Dockray, R. Dimaline, A. Varro, Gastrin: old hormone, new functions, *Eur. J. Physiol.* 449 (2005) 344–355.
- [25] G. Cui, H.L. Waldum, Physiological and clinical significance of enterochromaffin-like cell activation in the regulation of gastric acid secretion, *World J. Gastroenterol.* 13 (2007) 493–496.
- [26] M. Sato, K. Nakahara, M. Kojima, M. Miyazato, K. Kangawa, N. Murakami, Regulation of GH secretagogue receptor (GHS-R) gene expression in the rat nodose ganglion, *J. Endocrinol.* 194 (2007) 1–7.
- [27] M. Arosio, C.L. Ronchi, P. Beck-Peccoz, C. Gebbia, C. Giavoli, V. Cappiello, D. Conte, M. Peracchi, Effects of modified sham feeding on ghrelin levels in healthy human subjects, *J. Clin. Endocrinol. Metab.* 89 (2004) 5101–5104.
- [28] M. Yokoyama, K. Nakahara, M. Kojima, H. Hosoda, K. Kangawa, N. Murakami, Influencing the between-feeding and endocrine responses of plasma ghrelin in healthy dogs, *Eur. J. Endocrinol.* 152 (2005) 155–160.
- [29] T. Sugino, J. Yamaura, M. Yamagishi, A. Ogura, R. Hayashi, Y. Kurose, M. Kojima, K. Kangawa, Y. Hasegawa, Y. Terashima, A transient surge of ghrelin secretion before feeding is modified by different feeding regimens in sheep, *Biochem. Biophys. Res. Commun.* 298 (2002) 785–788.