



Erratum

Erratum to “Adaptation of macrophages to exercise training improves innate immunity” [Biochem. Biophys. Res. Commun. 372 (2008) 152–156]

Takako Kizaki ^{a,*}, Tohru Takemasa ^b, Takuya Sakurai ^a, Tetsuya Izawa ^c, Tomoko Hanawa ^d, Shigeru Kamiya ^d, Shukoh Haga ^e, Kazuhiko Imaizumi ^f, Hideki Ohno ^a

^a Department of Molecular Predictive Medicine and Sport Science, Kyorin University, School of Medicine, 6-20-2 Shinkawa, Mitaka, Tokyo 181-8611, Japan

^b Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8573, Japan

^c Department of Sport and Health Science, Doshisha University, 1-3 Tatara Miyakodai, Kyotanabe, Kyoto 610-0394, Japan

^d Infectious Disease/Division of Medical Microbiology, Kyorin University, School of Medicine, Mitaka, Tokyo 181-8611, Japan

^e Institute of Health and Sport Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8574, Japan

^f Faculty of Human Sciences, Waseda University, Tokorozawa, Saitama 359-1192, Japan

The publisher regrets that the captions for Fig. 1 and Supplemental Fig. 1 are reversed. Each of these figures and its corresponding caption appear correctly here.

Supplementary material

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.ab.2006.07.003](https://doi.org/10.1016/j.ab.2006.07.003).




Fig. 1. Effects of exercise training on the intracellular growth of *L. monocytogenes* within peritoneal macrophages. Adherent cells from control ($n = 5$) or exercise-trained mice ($n = 5$) were pooled and infected with *L. monocytogenes* at the multiplicity of infection. Results were expressed as means \pm SEM from triplicate cultures. ●, control; ■, exercise trained. Significantly lower than 0 h ($P < 0.05$).

DOI of original article: [10.1016/j.bbrc.2008.05.005](https://doi.org/10.1016/j.bbrc.2008.05.005)

* Corresponding author.

E-mail address: kizaki@kyorin-u.ac.jp (T. Kizaki).