



Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc



Response to excess copper in the hyperthermophile *Sulfolobus solfataricus* strain 98/2

Aramis A. Villafane, Yekaterina Voskoboynik¹, Mariola Cuebas[†], Ilona Ruhl, Elisabetta Bini^{*}

Department of Biochemistry and Microbiology, and Biotechnology Center for Agriculture and the Environment, Rutgers, The State University of New Jersey, New Brunswick, NJ 08901-8525, USA

ARTICLE INFO

Article history:

Received 19 April 2009

Available online 8 May 2009

Keywords:

Archaea

Sulfolobus solfataricus

Copper resistance

Regulation of transcription

Metal toxicity

ABSTRACT

Copper is an essential micronutrient, but toxic in excess. *Sulfolobus solfataricus* cells have the ability to adapt to fluctuations of copper levels in their external environment. To better understand the molecular mechanism behind the organismal response to copper, the expression of the cluster of genes *copRTA*, which encodes the copper-responsive transcriptional regulator CopR, the copper-binding protein CopT, and CopA, has been investigated and the whole operon has been shown to be cotranscribed at low levels from the *copR* promoter under all conditions, whereas increased transcription from the *copTA* promoter occurs in the presence of excess copper. Furthermore, the expression of the copper-transporting ATPase CopA over a 27-h interval has been monitored by quantitative real-time RT-PCR and compared to the pattern of cellular copper accumulation, as determined in a parallel analysis by Inductively Coupled Plasma Optical Emission spectrometry (ICP-OES). The results provide the basis for a model of the molecular mechanisms of copper homeostasis in *Sulfolobus*, which relies on copper efflux and sequestration.

© 2009 Elsevier Inc. All rights reserved.

Introduction

Copper is a transition metal and an important trace element because of the essential role it plays in a range of biological processes. In contrast, the occurrence of copper levels beyond the physiological range causes serious damage to all molecular components. Studies on yeast have led to the proposition that virtually no free copper ions are present in the cell under normal conditions [1]. The response of cells to copper excess/deficiency is accomplished through the interplay of copper-binding proteins, copper-responsive regulators, transporters for the efflux and uptake of copper, and copper-requiring enzymes. Genetic determinants of copper homeostasis have been described for several bacterial species [2–7]. In particular, the various components of the *Enterococcus irae* and *Escherichia coli* copper-homeostasis systems, their regulation, and interactions have been thoroughly studied [2,8]. Many sequenced archaeal genomes encode homologs of Cu (I) and Cu (II) transporting ATPases [9]. However, investigations of the response in archaea to changes of copper levels are still limited. Structural studies of individual functional domains of the Cu (I)-transporting

ATPase CopA in *Archeoglobus fulgidus* have provided useful insights into its activities and functions [10–14]. *A. fulgidus* also possesses a Cu(II)-transporting ATPase, CopB, that has been biochemically characterized [15]. The transcriptional analysis of a *cop* locus responsible for survival in the presence of copper has been reported in the extreme acidophilic archaeon “*Ferroplasma acidarmatus*” strain Fer1, where cotranscription of genes encoding the copper-binding protein CopZ and the putative copper-transporting ATPase CopB was shown to increase in response to Cu (II) [16]. An interesting mechanism for copper detoxification has been described in *Sulfolobus metallicus*, which is based on sequestration by organic phosphate, possibly followed by active efflux of the metal-phosphate complex [17]. The *Sulfolobus solfataricus* genome encodes a *cop* locus, which includes the three open reading frames (ORFs) Sso2651, Sso2652, and Sso10823, encoding the CopA ATPase, a copper-responsive regulator, and a putative copper-binding protein, respectively [18]. Cotranscription of Sso2651 and Sso10823 has been reported to specifically increase in the presence of copper, while the copper-responsive regulator binds sequences surrounding the putative *copA* promoter in *S. solfataricus* strain P2 [19]. In this study, the response of *S. solfataricus* to copper has been further investigated in the strain 98/2. The selection of the genetically tractable strain 98/2 [20] will expand the scope of analyses aimed to the elucidation of archaeal interactions with copper. To gain better insights into the *Sulfolobus* response to copper levels, the transcription of the three genes of the *copRTA* operon has been examined under different conditions and in a time course experiment, and the changes in the amount of copper associated with

^{*} Corresponding author. Address: Department of Biochemistry and Microbiology, Rutgers University, 76 Lipman Drive, New Brunswick, NJ 08901-8525, USA. Fax: +1 732 932 8965.

E-mail address: bini@aesop.rutgers.edu (E. Bini).

¹ Present address: Department of Molecular Biology and Biochemistry, Rutgers University, USA.

[†] Deceased.

the cells have been monitored over time. Based on the data obtained, a preliminary model for the maintenance of copper homeostasis in *Sulfolobus* is proposed.

Materials and methods

Growth conditions. *Sulfolobus solfataricus* strains 98/2 or P2 (DSM 1617) were cultured at 80 °C in a defined standard medium (SM) as described in [20], the medium was supplemented with 0.2% sucrose as the carbon and energy source. Batch cultures were inoculated to obtain a density corresponding to an OD₅₄₀ of about 0.025, with aliquots withdrawn from mid-log phase cultures. Growth was monitored at a wavelength of 540 nm on a Beckman DU-520 spectrophotometer (Beckman Coulter, USA). All the experiments were carried out on the strain 98/2, unless otherwise stated.

RNA extraction. Total RNA was isolated from *S. solfataricus* cultures in their exponential phase of growth (OD₅₄₀ = 0.3–0.6). Before centrifugation at 3500g for 15 min, cells were mixed with two volumes of RNA Protect (Qiagen, USA). RNA was extracted from the cell pellets using the RNeasy Mini kit (Qiagen, USA) and treated with DNase (Ambion, USA), as recommended by the manufacturer. DNA contamination was excluded by PCR using primers targeting the 16S rRNA gene. The quantity and quality of the RNA obtained was evaluated both spectrophotometrically on a NanoDrop ND-1000 spectrophotometer (NanoDrop, USA) and by agarose gel electrophoresis [21].

Reverse-transcription PCR (RT-PCR) and real-time quantitative RT-PCR (qRT-PCR) analyses. For RT-PCR analysis, total RNA (0.5 µg) was analyzed in 25-µl reactions using the Enhanced Avian HS RT-PCR Kit (Sigma–Aldrich, USA). The amplification products were separated on a 1.2% agarose gel by electrophoresis, and the gel images were acquired using a GelLogic 440 Imaging System (Eastman Kodak, USA). Specific transcripts were quantified by qRT-PCR using the iScript One-Step RT-PCR kit with SYBR Green (Bio-Rad, USA) and the real-time detection system iCycler iQ (Bio-Rad, USA). Reactions, in triplicate, were assembled according to the protocol of the manufacturer, and using 0.2 µg of total RNA in a 25-µl reaction. Primers were designed using the OligoPerfect Designer software (Invitrogen, USA) to have a composition that was suitable for use in both RT-PCR and qRT-PCR (Table 1). Specificity of each pair of primers was confirmed by sequencing. The efficiency of the PCR amplifications was determined from the slopes of the dilution curves of the target RNA. The cycle threshold (C_t) values obtained were used in the “2^{−ΔΔC_t} Method” to calculate the relative changes in gene expression [22]. Expression of the target RNAs of interest was normalized to the level of the Sso0067 transcript, detected using the primers q0067-F and q0067-R (Table 1). Sso0067 encodes a ribosomal protein and its expression is not affected by copper exposure, as determined by microarray analysis (unpublished).

Analysis of copper content. Cell samples were harvested from exponentially growing *S. solfataricus* cultures. Cell pellets were washed with 10 mM EDTA to remove the copper adsorbed to the

cell wall, then rinsed with SM without added trace metals (Zn, Cu, Mo, V, and Co). Before centrifugation at 3500 g for 15 min, aliquots were removed from each sample for determination of protein concentration using the BCA Protein Assay (Pierce, USA). Cell pellets were resuspended in 50% nitric acid, and digested for 16 h at 25 °C. The total copper content was analyzed using a Vista Pro radial view Inductively Coupled Plasma Optical Emission Spectrometer (ICP-OES, Varian Inc., USA). Controls included non-inoculated SM medium and untreated cultures.

Results and discussion

Physiological response to copper

To establish the optimal concentration of copper to be used in this study, cells were exposed to CuCl₂ at concentrations ranging from 0 mM to 2.5 mM. The MIC, defined as the lowest concentration that completely inhibits cell growth immediately after exposure, was determined to be 1.5 mM (Fig. 1A). In response to copper concentrations that were equal or greater than 1.5-mM CuCl₂, a lag phase of variable duration was observed. The duration of the lag phase was directly proportional to the metal concentration, and growth resumed thereafter, indicating a slow adaptation to levels of copper that were above the MIC. To rule out the possibility that this effect was due to the appearance of copper-resistant mutants, cells adapted to 2-mM copper-containing medium were subcultured in fresh medium in the absence of the metal. After 4 cycles of cell divisions, the cells were re-inoculated into fresh medium containing 2 mM copper. A lag phase was observed again, indi-




Fig. 1. Effect of CuCl₂ on growth of *S. solfataricus*. (A) An exponentially growing culture of strain 98/2 was used as inoculum for 8 subcultures. Copper was added at the time indicated by the arrow, at the following final concentrations: (●), 0 mM; (○), 0.75 mM; (▼), 1 mM; (▽), 1.25 mM; (■), 1.5 mM; (□), 1.75 mM; (◆), 2 mM; (◇), 2.5 mM. (B) Growth of P2 (○) or 98/2 (●). Copper was added at the time of inoculum and cell growth is expressed as percentage of the untreated control. Best fit curves were obtained by nonlinear regression applied to both sets of data.

Table 1
Oligonucleotides used in this work.

ID	Sequence (5'–3')	Target
q2651-F	GAATAGTTGGGATGCATTGT	<i>copA</i>
q2651-R	ACTACCCCTTAACGTTTTC	
q2652-F	TTTATTGCCTTCGCCATTTC	<i>copR</i>
q2652-R	GTTGCGTGCAAAATTTTCCT	
2652-F	TGCAATTCCTTGCTTGTCTGG	<i>cop</i> operon (paired with q2651-R)
q10823-F	ATGATAATCGATCCGGTTTG	
q10823-R	ATTCTTAAATACTCTTCCGGA	<i>copT</i>
q0067-F	TACCAATGTGCGCTTTTGCT	
q0067-R	CAAATCACCATCTGGAGGAA	Reference transcript

cating that the response to copper concentrations above the MIC was the result of physiological adaptation (unpublished). To limit the manifestation of extensive stress responses that would overshadow the response to copper, only the sublethal concentration of 0.75 mM copper was used for transcript measurements. The level of tolerance to copper observed in *S. solfataricus* 98/2 was within the range observed for most microorganisms. It was previously observed that the carbon source affected the copper sensitivity of microbes [23], probably because of the binding of copper to the thiol groups of some amino acids and other molecules. Therefore, all experiments were carried out using cells grown on 0.2% sucrose as the sole carbon source. Interestingly, the *S. solfataricus* strain P2 showed a higher copper sensitivity than the strain 98/2, displaying a growth rate on 0.2 mM copper comparable to that observed for strain 98/2 grown in the presence of 2.5 mM copper (Fig. 1B). The two strains were originally isolated in Italy and USA, respectively. It was speculated that their physical separation might have led to different functionality of the copper responsive system. Thus, genetic diversity either at the level of the transcriptional regulator CopR or within the region of the putative *copTA* promoter was hypothesized to be responsible for the different copper sensitivities of the two strains. However, the sequencing of the segment of the *cop* operon that includes *copR*, *copT*, and their intergenic space in the strain 98/2 (GenBank Accession No. EU544670), showed that this region was identical to the corresponding sequence in strain P2, indicating that other factors probably contribute to these differences.

Cotranscription of *copRTA*

The *cop* operon consists of three genes oriented in the same direction, represented by ORF's Sso2652, Sso10823, and Sso2651. With the possible exception of the Cu (I) transporter CopA, and the Cu (II) transporter CopB, there are no unique identifiers for functional homologs of copper responsive sequences in prokaryotes. Herein, it is proposed the designation "CopR" for the product of ORF Sso2652, in agreement with the nomenclature applied to metal responsive transcriptional regulators: ArsR, NikR, MerR [24], and CopR [25,26]. "CopT" is suggested for Sso10823, which corresponds to a stand alone TRASH domain [27]. In an attempt to elucidate the molecular mechanisms behind the regulation of the *copRTA* operon (Fig. 2A), RT-PCR was used to test whether the genes *copR*, *copT*, and *copA* were cotranscribed. Using the primers pair 2652-F and q2651-R, designed to amplify across the genes *copR* and *copA*, a unique band of the expected size was obtained (Fig. 2A and B), sequencing of the amplicon confirmed its specificity and indicated the cotranscription of all the three genes. Cotranscription of the copper-binding protein and the ATPase-coding genes was previously reported to be detected by primer extension [19], but the cotranscription of the *copR* gene was not addressed in that analysis. By specifically targeting the region encompassing the three genes, the regulator-encoding *copR* was also shown to be a part of the transcript. This is an interesting finding because it provides additional insight into the regulation of copper homeostasis in *Sulfolobus*, as discussed later in this section. In the next step, the induction of individual genes was investigated by qRT-PCR. The primer pairs used for the detection of *copR* and *copT* are listed in Table 1. In cells treated with copper, the levels of *copT* mRNA exceeded the amounts of the same transcript in untreated cultures, displaying a greater than 20-fold induction after 120 min, thus behaving as *copA* (see section Transcription of *copA* occurs transiently in response to copper), however, the level of *copR* expression was unaffected by copper (Fig. 2C). This trend agreed with the data obtained by primer extension [19], and, moreover, this method provided an accurate quantitation of the fold induction. The fact that the genes of the operon *copRTA* are cotranscribed, that




Fig. 2. Co-transcription of *copR*, *copT*, and *copA*. (A) Diagram of the *cop* operon and location of the primers. (B) Agarose gel electrophoresis of the amplicons corresponding to the regions encompassing *copR/copA* (RA), *copT/copA* (TA) and targeting *copA* (AA), (no) negative control without reverse transcriptase. (C) Expression of *copR* and *copT* in the presence of copper (gray bars) relative to untreated controls (black bars). Results are reported as means \pm standard errors.

copR is constitutively expressed (Fig. 2C), and that *copTA* transcription is considerably affected by excess copper, suggests a model where, in the absence of excess copper, *copT* and *copA* are transcribed together with *copR*, constitutively, from the *copR* promoter, whereas the supplementary transcription of *copTA* in the presence of copper results from the induction of a second promoter upstream of *copTA* promoter. The constitutive expression of *copTA* probably provides a constant and low-level supply of the proteins CopT and CopA that maintain homeostasis, allowing the cell to adjust to small fluctuations in copper levels under normal conditions.

Transcription of *copA* occurs transiently in response to copper

To better understand the regulation of cell response, the expression of *copA* was monitored in a time course experiment. Total RNA samples, isolated at different times after treating cell cultures with 0.75 mM CuCl_2 were subjected to qRT-PCR analysis using primers targeting the *copA* transcript (Table 1). *Sulfolobus* responded to exposure to a sub-lethal copper excess by the transient active transcription of the *copA* gene. The *copA* transcript reached a peak 1 h after treatment, corresponding to approximately 35-fold the uninduced level, thereafter, the transcript level decreased until a steady-state level of 2–3-fold induction is reached in the following 16 h (Fig. 3). Moreover, in cells cultured for several generations in the presence of 0.75 mM copper, the amount of *copA* transcript was maintained at 2–3-fold the uninduced level (unpublished), indicating that its rate of expression was maintained constant during long-term exposures, provided the concentration of copper did not change. The above observation can be explained as follows. A high rate of CopA synthesis is necessary to reestablish homeostasis during early copper exposure. Thereafter, the drop in intracellular




Fig. 3. Time course analysis of copper content and *copA* induction in response to copper. (A) Total RNA was isolated at different times after treatment of the cultures with 0.75 mM copper. Samples were not collected at 180 and 1620 min. Results are expressed as means \pm standard errors. Bars represent the normalized ratio between amounts of transcript in copper treated and untreated cultures. For comparison, changes in copper content (\blacklozenge) after exposure to 0.75 mM copper are superimposed. (B) Copper content in cultures treated with 0.75 mM (white bars) or 1.25 mM copper (dark bars).

concentration of copper causes a decrease in *copA* transcription, which leads to the establishment of a rate of CopA synthesis sufficient to maintain the internal equilibrium at the new cytoplasmic copper concentration. It has been previously reported [19] that treating *S. solfataricus* P2 with 5 mM copper causes an accumulation of the *copA* transcript over a 2-h period. This might be explained by a higher demand for CopA, extended over time, in the presence of copper concentrations above the MIC. This observation is also consistent with the behavior of *S. solfataricus* 98/2 for copper levels above 0.75 mM (Fig. 1A). Furthermore, the fact that *copA* is transiently induced after copper challenge and that its transcription subsequently declines but is maintained at a low basal induced level for the duration of exposure might indicate an accumulation of the CopA transporter. The persistence of a stable CopA protein would reduce the demand for further *copA* transcription and translation.

Determination of copper content

To test if the pattern of expression of *copA* depended on internal fluctuations of copper, the copper associated with the cells was analyzed in a time course experiment by ICP-OES spectrometry, and was observed to slowly increase during the first 3 h of monitoring. This might be explained with the sequestration and consequent accumulation of intracellular excess copper by CopT. Alternatively, although the cell pellets were washed with EDTA, these measurements might reflect some residual copper adsorption to the cell surface. After prolonged exposure, the amount of copper returned to the level measured 1 h after treatment (Fig. 3A), probably due to efflux mediated by CopA. The amount

of copper in a 0.75 mM solution is $47.6 \times 10^3 \mu\text{g/L}$. However, the copper associated with the cells in a similar medium, 30 min after treatment, is three orders of magnitude lower, being approximately 11.6 $\mu\text{g/L}$ (or 183 nM). These data suggest that active efflux has a major role in maintenance of copper homeostasis, and that sequestration also contributes to it, possibly by fine tuning the system. Cells treated with inhibitory copper concentrations showed a similar pattern of copper accumulation, except for a large increase 60 min after exposure (Fig. 3B). Bioremediation studies have established that dead cells, i.e. algae, have a higher capacity of binding metal ions [28]. Therefore the large increase in copper content may indicate cell death.

Conclusions

When *S. solfataricus* cells are challenged with copper, their rate of growth slows down, and this decrease is proportional to the concentration of copper tested. Although immediately after copper challenge a MIC of 1.5 mM copper has been determined, cells become resistant to environs containing up to 2.5 mM of CuCl_2 after prolonged exposure. An increased level of the transcripts of genes directly involved in copper detoxification, namely *copA* and *copT*, which are co-transcribed, is observed after treatment with 0.75 mM copper. One hour after copper challenge, the level of *copA* transcript declines until a steady-state level corresponding to an approximately 2-fold induction is reached. Similarly, after treatment with excess copper, the amount of the metal associated with the cells is maintained constant, after a slight accumulation followed by a decrease. These results are consistent with the existence of a feedback-like mechanism of action, where accumulation of the CopA transporter, derived from the burst of *copA* transcription, causes a drop in the levels of cytoplasmic copper ions. Sequestration of the metal by CopT contributes to limit the availability of free copper. The lowered copper level results in a reduced effect on CopR and consequently a decline of *copA* transcription. The different copper sensitivity of the two *Sulfolobus* strains P2 and 98/2, both having identical *copR* and *copT* genes and *copTA* promoter sequences, supports the hypothesis that additional factors probably have a role in the response of the cells to copper concentrations, and that these factors might be different in the two strains studied. Further research is required to elucidate the steps involved and to identify the additional genetic determinants of the response of the cell to varying levels of copper in the surroundings.

Acknowledgments

This research was supported by Hatch Project NJ01136 and a SEBS Pre-Tenure Award to EB, a GAANN fellowship to AV, and NIH Bridge Award GM58389 to MC.

References

- [1] T.D. Rae, P.J. Schmidt, R.A. Pufahl, V.C. Culotta, T.V. O'Halloran, Undetectable intracellular free copper: the requirement of a copper chaperone for superoxide dismutase, *Science* 284 (1999) 805–808.
- [2] C. Rensing, G. Grass, *Escherichia coli* mechanisms of copper homeostasis in a changing environment, *FEMS Microbiol. Rev.* 27 (2003) 197–213.
- [3] S. Chintalapati, R. Al Kurdi, A.C. van Scheltinga, W. Kuhlbrandt, Membrane structure of CtrA3, a copper-transporting P-type-ATPase from *Aquifex aeolicus*, *J. Mol. Biol.* 378 (2008) 581–595.
- [4] S. Sitthithak, L. Knutsson, J.W. Webb, R.K. Jayaswal, Molecular characterization of the copper transport system in *Staphylococcus aureus*, *Microbiology* 153 (2007) 4274–4283.
- [5] D. Magnani, O. Barre, S.D. Gerber, M. Solioz, Characterization of the CopR regulon of *Lactococcus lactis* IL1403, *J. Bacteriol.* 190 (2008) 536–545.
- [6] G.T. Smaldone, J.D. Helmann, CsoR regulates the copper efflux operon *copZA* in *Bacillus subtilis*, *Microbiology* 153 (2007) 4123–4128.
- [7] M. Espariz, S.K. Checa, M.E. Audero, L.B. Pontel, F.C. Soncini, Dissecting the *Salmonella* response to copper, *Microbiology* 153 (2007) 2989–2997.

- [8] M. Solioz, J.V. Stoyanov, Copper homeostasis in *Enterococcus hirae*, FEMS Microbiol. Rev. 27 (2003) 183–195.
- [9] E. Bini, Archaea–metal interactions: metabolism and strategies of resistance, in: P. Blum (Ed.), Archaea: New Models for Prokaryotic Biology, Caister Academic Press, 2008, pp. 27–43.
- [10] M.H. Sazinsky, S. Agarwal, J.M. Arguello, A.C. Rosenzweig, Structure of the actuator domain from the *Archaeoglobus fulgidus* Cu⁺-ATPase, Biochemistry 45 (2006) 9949–9955.
- [11] M.H. Sazinsky, A.K. Mandal, J.M. Arguello, A.C. Rosenzweig, Structure of the ATP binding domain from the *Archaeoglobus fulgidus* Cu⁺-ATPase, J. Biol. Chem. 281 (2006) 11161–11166.
- [12] W.J. Rice, A. Kovalishin, D.L. Stokes, Role of metal-binding domains of the copper pump from *Archaeoglobus fulgidus*, Biochem. Biophys. Res. Commun. 348 (2006) 124–131.
- [13] A.K. Mandal, Y. Yang, T.M. Kertesz, J.M. Arguello, Identification of the transmembrane metal binding site in Cu⁺-transporting PIB-type ATPases, J. Biol. Chem. 279 (2004) 54802–54807.
- [14] D.I. Cattoni, F.L. Flecha, J.M. Arguello, Thermal stability of CopA, a polytopic membrane protein from the hyperthermophile *Archaeoglobus fulgidus*, Arch. Biochem. Biophys. 471 (2008) 198–206.
- [15] S. Mana-Capelli, A.K. Mandal, J.M. Arguello, *Archaeoglobus fulgidus* CopB is a thermophilic Cu²⁺-ATPase: functional role of its histidine-rich-N-terminal metal binding domain, J. Biol. Chem. 278 (2003) 40534–40541.
- [16] C. Baker-Austin, M. Dopson, M. Wexler, R.G. Sawers, P.L. Bond, Molecular insight into extreme copper resistance in the extremophilic archaeon *Ferroplasma acidarmanus* Fer1, Microbiology 151 (2005) 2637–2646.
- [17] F. Remonsellez, A. Orell, C.A. Jerez, Copper tolerance of the thermoacidophilic archaeon *Sulfolobus metallicus*: possible role of polyphosphate metabolism, Microbiology 152 (2006) 59–66.
- [18] Q. She, R.K. Singh, F. Confalonieri, Y. Zivanovic, G. Allard, M.J. Awayez, C.C. Chan-Weiher, I.G. Clausen, B.A. Curtis, A. De Moors, G. Erauso, C. Fletcher, P.M. Gordon, I. Heikamp-de Jong, A.C. Jeffries, C.J. Kozera, N. Medina, X. Peng, H.P. Thi-Ngoc, P. Redder, M.E. Schenk, C. Theriault, N. Tolstrup, R.L. Charlebois, W.F. Doolittle, M. Duguet, T. Gaasterl, R.A. Garrett, M.A. Ragan, C.W. Sensen, J. Van der Oost, The complete genome of the crenarchaeon *Sulfolobus solfataricus* P2, Proc. Natl. Acad. Sci. USA 98 (2001) 7835–7840.
- [19] T.J. Ettema, A.B. Brinkman, P.P. Lamers, N.G. Kornet, W.M. de Vos, J. van der Oost, Molecular characterization of a conserved archaeal copper resistance (cop) gene cluster and its copper-responsive regulator in *Sulfolobus solfataricus* P2, Microbiology 152 (2006) 1969–1979.
- [20] P. Worthington, V. Hoang, F. Perez-Pomares, P. Blum, Targeted disruption of the alpha-amylase gene in the hyperthermophilic archaeon *Sulfolobus solfataricus*, J. Bacteriol. 185 (2003) 482–488.
- [21] J. Sambrook, E.F. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989.
- [22] K.J. Livak, T.D. Schmittgen, Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method, Methods 25 (2001) 402–408.
- [23] K.W. Miller, S.S. Risanico, J.B. Risatti, Differential tolerance of *Sulfolobus* strains to transition metals, FEMS Microbiol. Lett. 93 (1992) 69–73.
- [24] M.A. Pennella, D.P. Giedroc, Structural determinants of metal selectivity in prokaryotic metal-responsive transcriptional regulators, Biometals 18 (2005) 413–428.
- [25] F. Cantini, L. Banci, M. Solioz, The copper-responsive repressor CopR of *Lactococcus lactis* is a 'winged helix' protein, Biochem. J. 417 (2009) 493–499.
- [26] S.D. Mills, C.K. Lim, D.A. Cooksey, Purification and characterization of CopR, a transcriptional activator protein that binds to a conserved domain (cop box) in copper-inducible promoters of *Pseudomonas syringae*, Mol. Gen. Genet. 244 (1994) 341–351.
- [27] T.J. Ettema, M.A. Huynen, W.M. de Vos, J. van der Oost, TRASH: a novel metal-binding domain predicted to be involved in heavy-metal sensing, trafficking and resistance, Trends Biochem. Sci. 28 (2003) 170–173.
- [28] J. Kadukova, E. Vircikova, Comparison of differences between copper bioaccumulation and biosorption, Environ. Int. 31 (2005) 227–232.