



Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc



Alix is involved in caspase 9 activation during calcium-induced apoptosis

Flavie Strappazon¹, Sakina Torch, Christine Chatellard-Causse, Anne Petiot, Chantal Thibert, Béatrice Blot, Jean-Marc Verna², Rémy Sadoul^{*}

Inserm, U836, Team 2, Grenoble F-38000, France
Université Joseph Fourier, Grenoble F-38000, France

ARTICLE INFO

Article history:

Received 9 May 2010

Available online 21 May 2010

Keywords:

ALG-2
Apoptosis
Cell death
Cerebellar granule neurons
Endoplasmic reticulum
Thapsigargin

ABSTRACT

The cytoplasmic protein Alix/AIP1 (ALG-2 interacting protein X) is involved in cell death through mechanisms which remain unclear but require its binding partner ALG-2 (apoptosis-linked gene-2). The latter was defined as a regulator of calcium-induced apoptosis following endoplasmic reticulum (ER) stress. We show here that Alix is also a critical component of caspase 9 activation and apoptosis triggered by calcium. Indeed, expression of Alix dominant-negative mutants or downregulation of Alix afford significant protection against cytosolic calcium elevation following thapsigargin (Tg) treatment. The function of Alix in this paradigm requires its interaction with ALG-2. In addition, we demonstrate that caspase 9 activation is necessary for apoptosis induced by Tg and that this activation is impaired by knocking down Alix. Altogether, our findings identify, for the first time, Alix as a crucial mediator of Ca^{2+} induced caspase 9 activation.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

Calcium plays a central role in regulating a variety of cellular responses, notably in neurons. In particular, it is now well established that alteration in intracellular calcium homeostasis is a common feature of many apoptotic processes, which are associated with local elevations of cytosolic Ca^{2+} released from endoplasmic reticulum (ER) stores [1–5]. Although cytosolic Ca^{2+} elevation induces caspase 9 activation as a necessary step for apoptosis, the way of activation of the caspase remains dubious. In particular, it is still unclear whether caspase 9 activation requires Apaf-1 or occurs in a separate ER based complex [6–8]. Using Apaf-1 $-/-$ cells, Rao et al. defined the apoptosis-linked gene-2 (ALG-2), a calcium-binding protein belonging to the penta-EF hand protein family, as a regulator of ER stress-induced death involving Ca^{2+} [8]. Indeed, knock-down of ALG-2 reduced cell death due to thapsigargin (Tg), which

elevates cytosolic calcium by blocking the ER Ca^{2+} -ATPase. They went on to suggest that ALG-2 is part of a high molecular weight complex containing caspases 9 and 12 and that knocking down ALG-2 inhibits Tg-induced caspase 9 activation.

Ca^{2+} binding to ALG-2 induces a conformational change which allows it to interact with several proteins [9,10], among which the multifunctional adaptor protein Alix/AIP1 [11,12]. Alix is a cytosolic protein which interacts with proteins involved in deforming membranes during various processes including endocytosis, endosomal sorting, virus budding and cytokinesis [13,14]. Up-regulation of Alix expression *in vivo* correlates with calcium-induced cell death, as observed in degenerating neurons of the rat hippocampus undergoing epileptic seizures [15]. We also found that over-expression of the full length protein (Alix-wt) in cultured neurons was sufficient to activate caspases and thereby apoptosis. In contrast expression of the C-terminal half of the protein encompassing the ALG-2 binding site blocked neuronal death both *in vivo* and *in vitro* [16,17]. The pro- and anti-apoptotic functions of Alix-wt and Alix-CT, respectively, were tightly dependent on their capacity to bind to ALG-2. Expression of Alix lacking its ALG-2 binding site could block naturally occurring death of motoneurons during development of the chick embryo [18,19]. Given the fact that ALG-2 needs to bind Ca^{2+} in order to interact with Alix, these observations suggest that ALG-2 and Alix cooperate in transducing Ca^{2+} -regulated signals during cell death. Here we report that knock-down of Alix protects cells from Tg-induced cell death which is mediated by caspase 9. Furthermore, we show that Alix

Abbreviations: ALG-2, apoptosis-linked gene-2; Alix, ALG-2 interacting protein X; Apaf-1, apoptotic protease activating factor-1; BHK, baby hamster kidney; DIV, days *in vitro*; ER, endoplasmic reticulum; RFP, red fluorescent protein; Tg, thapsigargin.

^{*} Corresponding author. Address: Grenoble Institut des Neurosciences, Bâtiment Edmond J. Safra des Neurosciences, Chemin Fortuné Ferrini, Site Santé, 38706 La Tronche cédex, France.

E-mail address: remy.sadoul@ujf-grenoble.fr (R. Sadoul).

¹ Present address: Laboratory of Molecular Neuroembryology, IRCCS Fondazione Santa Lucia, Via del Fosso di Fiorano 64/65, 00143 Rome, Italy.

² Present address: Service des Ressources Humaines, Délégation Alpes, CNRS – BP 166, 38042 Grenoble, France.

expression is required for caspase 9 activation induced by calcium elevation. These results suggest that Alix and ALG-2 act upstream of caspase 9 activation following cytosolic calcium elevation.

2. Material and methods

2.1. Constructs and antibodies

Polyclonal anti-Alix was previously described [16], rabbit anti-Flag (Sigma), monoclonal anti-HA (Cell Signaling), monoclonal anti-actin (Chemicon), alexa-488 conjugated goat anti-rabbit and alexa-488 conjugated goat anti-mouse (Molecular Probes), peroxidase-conjugated goat anti-mouse and peroxidase goat anti-rabbit (Jackson). Alix expression plasmids have been described previously [20]. pcDNA3 vectors coding for dominant-negative mutants of procaspase 8 (C360A) and 9 mutants (C287A) were a kind gift of P. Mehlen (Lyon).

2.2. Cell cultures

Primary cultures of cerebellar granule cells (CGN) were prepared from 6-day-old S/IOPS NMRI mice (Charles River Laboratories), as described previously [16]. Dissociated neurons were plated in Dulbecco's modified Eagle's medium (DMEM, Invitrogen) containing 25 mM KCl (K25 medium) supplemented with 10% fetal bovine serum. After 4 DIV, the medium was replaced with serum-free K25. Twenty-four hours later, thapsigargin (Invitrogen) (200 nM in 0.01% DMSO) or DMSO alone were added to K25 medium. Baby hamster kidney cells (BHK-21) were used as described previously [21]. Cells were treated with 1 μ M Tg 24 h after plating.

2.3. Cell transfections

Transient transfection of cultured CGN was performed using the calcium phosphate co-precipitation method as previously described [16]. BHK-21 cells were grown at 37 °C in Glasgow MEM (GMEM) supplemented with 5% fetal calf serum. Transient transfections of BHK-21 cells were performed using JetPEI (PolyPlus Transfection).

2.4. Establishment of an Alix-deficient BHK-21 cell line

An shRNA was cloned downstream of the human H1 promoter in the pSuperGFP vector (Oligoengine, Seattle, WA, USA). Forward and reverse sequences for the Alix shRNA construct were: 5'-GAT CCCGCGCTGGTGAAGTTCATCTTCAAGAGAGATGAACCTCACCAGC GGCTTTTGGAAA-3' and 5'-AGCTTTTCCAAAAGCCGCTGGTGAAGT TCATCTCTTGAAGATGAACCTCACCAGCGCGGG-3', respectively. The synthetic oligonucleotides were synthesised by Invitrogen. The annealed oligonucleotides were ligated into the BglII–HindIII site of the pSuperGFP vector.

BHK-21 cells were transfected with pSuper-Sh Alix or pSuper vectors as control by using JetSi transfection (Polyplus tranfection). Cells were selected by adding G418 (800 μ g/ml) to the culture medium. After 15 days, clones were isolated and screened by Western blot analysis for the best reduction in Alix expression. Permanent cell lines (pSuper/Sh Alix and control) were maintained with G418 (800 μ g/ml).

2.5. Determination of cell viability

MTT assay. The tetrazolium salt MTT [3-(4,5-dimethyl-2-thiazol-yl)-2,5-diphenyl-2H-tetrazoliumbromide] (Sigma) was added to cell cultures (1 mg/ml) and incubated for 30 min at 37 °C. Cells were then lysed in DMSO. Formazan production by living cells was assessed by measuring absorbance at 540 nm.

Hoechst staining. Cultures were fixed in 4% paraformaldehyde in PBS for 20 min at 4 °C and stained with the bisbenzimidazole (Hoechst 33342, Sigma; 4 μ g/ml) in PBS for 20 min at room temperature. Cells were washed three times with PBS, mounted in Mowiol (Calbiochem) and observed under an Axiovert microscope (Zeiss) connected to a CDD camera. For each condition, random images were captured and analyzed. Cell viability was then scored on the basis of nuclear morphology: cells containing condensed or fragmented nuclei were counted as dying or dead cells.

2.6. Determination of caspase 9 activation in cells

Caspase 9 Detection kit (Calbiochem) uses the fluorescent marker red LEHD-fmk which enters living cells and binds irreversibly to activated caspase 9. After 24 h treatment with 1 μ M Tg or vehicle (DMSO), cells were incubated for 1 h at 37 °C with the fluorescent marker according to the supplier's instructions and observed under an Axiovert microscope (Zeiss).




Fig. 1. Ectopic expression of the C-terminal half of Alix delays cell death of BHK-21 cells and CGN induced by an ER stress. BHK-21 cells grown for 24 h (A) or cerebellar granule neurons (CGN) cultured for 4 days (B) were transfected with vectors encoding either Flag-Alix-CT, Flag-Alix-CTΔALG-2 or RFP. One day later, cells were treated with Tg (1 μ M (a) 200 nM (B)) or vehicle (0.01% DMSO) for 24 h. Transfected cells were revealed using an anti-Flag antibody and cells with condensed/fragmented nuclei were scored as pyknotic. Results are expressed as the percentage of transfected cells displaying pyknotic nuclei. Values are given as a percentage (\pm SD) of cell death and statistical analysis was performed by analysis of variance (one-way ANOVA followed by Scheffe's post hoc test). * P < 0.05 versus RFP control.

2.7. Immunofluorescence

Cells were fixed in 4% paraformaldehyde in PBS for 10 min at 37 °C. After permeabilization and blocking with 3% goat pre-immune serum (Invitrogen), cells were incubated at room temperature with primary antibodies (monoclonal anti-HA: 1/100, polyclonal anti-Flag: 1/500) then with Alexa Fluor-conjugated secondary anti-mouse antibody (1/1000) or anti-rabbit antibody (1/1000) (Invitrogen). Cells were rinsed with PBS, and stained with Hoechst 33342 before examination.

2.8. Western blot analysis

Cells were washed twice with PBS before harvesting with lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.5% DOC (sodium deoxycholate), 1% NP40, 0.1% SDS (sodium dodecyl sulfate), 1 mM Na₃VO₄, 50 mM NaF, 10 mM β-glycerophosphate and 2× protease inhibitor cocktail (Complete™, Boehringer). Cell extracts were centrifuged at 15,000g for 15 min at 4 °C and 20 μg protein was separated by SDS-PAGE and transferred onto nylon membranes (Immobilon-P, Millipore). After incubation with the membranes primary antibodies were revealed using peroxidase-coupled secondary antibodies and the SuperSignal detection method (Pierce).

3. Results

3.1. Alix C-terminal half protects BHK-21 cells and cerebellar neurons from thapsigargin-induced cytotoxicity

We first used BHK-21 cells to analyze the effect of the C-terminal half of Alix (Alix-CT) on apoptosis following a Ca²⁺ increase due

to Tg treatment. 24 h after transfection, BHK-21 cells were treated with 1 μM Tg and viability estimated after a further 24 h culture period. Sixty percent of control RFP-expressing cells exhibited altered nuclear morphology indicative of apoptosis whereas 15% were dead or dying in untreated control cultures (Fig. 1A). In contrast, only 30% of Alix-CT-expressing cells died following the same treatment. Remarkably, Alix-CT deleted of its ALG-2-binding domain (Alix-CTΔALG-2) did not afford any protective effect against Tg-induced apoptosis (Fig. 1A).

Similar experiments using cerebellar granule neurons (CGN) gave comparable results. Indeed, while a 24 h incubation with 200 nM Tg increased death of RFP transfected CGN by 36%, the number of Alix-CT-expressing neurons showing a condensed or fragmented nucleus did not significantly differ from that of untreated cultures (Fig. 1B). Deletion of the ALG-2 binding site on Alix-CT abrogated its capacity to rescue neurons. Our findings indicate that expression of a dominant-negative form of Alix counteracts the deleterious effect of Tg in a way which depends on its capacity to interact with ALG-2.

3.2. Alix depletion reduces thapsigargin-induced cell death

To reinforce our suggestion of a possible role of Alix in mediating ER stress apoptosis, we established a BHK-21 cell line in which we down regulated Alix expression using a pSuper vector coding for a short hairpin RNA against Alix (*Sh Alix*). As illustrated in Fig. 2A, Alix content was reduced by at least 90% in cells stably expressing *Sh Alix* compared to control cells expressing an empty P-Super (*Sh control*). Exposure to 1 μM Tg did not significantly alter Alix expression in *Sh control* or *Sh Alix* cells (Fig. 2A). Using the MTT assay, we observed a time-dependent reduction in metabolic




Fig. 2. Reduction of Alix expression protects BHK-21 cells from Tg cytotoxicity. (A) Western blot analysis of Alix expression in *Sh control* or *Sh Alix* BHK-21 cells with or without Tg for 24 h. (B) *Sh control* or *Sh Alix* BHK-21 cells were treated with 1 μM Tg and the percentage of viable cells was estimated at various time intervals using the MTT assay. Results are expressed as a percentage of the metabolic activity at $t = 0$ in Tg-treated cultures relative to DMSO-treated-control cultures. Each point represents the mean \pm SD of triplicate wells from three independent experiments. (C) Alix expression restores sensitivity of *Sh Alix* cells to Tg cytotoxicity. *Sh control* or *Sh Alix* BHK-21 cells were transfected with a vector coding either for Flag-Alix, Flag-AlixΔALG-2 or RFP. Upper panel: WB analysis of cell extracts of *Sh control* and *Sh Alix* cells using an anti-Flag antibody demonstrating expression of Alix and AlixΔALG-2 in transfected cells. Bottom panel: 24 h after transfection, cells were exposed to Tg (1 μM) during 24 h and then immunostained with an anti-Flag antibody. Cells were counted in 10 random fields in two or three different wells and condensed/fragmented nuclei were scored as pyknotic. Results are expressed as the percentage of transfected cells displaying a pyknotic nucleus. Data are means \pm SD of three independent experiments. Statistical analysis was performed by analysis of variance (one-way ANOVA followed by Scheffe's post hoc test). * $P < 0.05$ versus RFP-overexpressing *Sh Alix* cells. # $P < 0.05$ versus *Sh control* cells.

activity of *Sh* control cells following Tg treatment reaching 70% at 72 h (Fig. 2B). In contrast, after an initial limited decline, MTT metabolism remained constant over until the end of the experiment in *Sh* Alix cells. These findings suggest that Alix-depleted cells are less sensitive to Tg cytotoxicity. To confirm this, cell viability was estimated using Hoechst nuclear staining 24 h after addition of Tg. After Tg treatment about 55% of *Sh* control cells exhibited condensed or fragmented nuclei, while only 15% of *Sh* Alix cells showed these apoptotic features (Fig. 2C). Therefore, reducing Alix expression greatly improves the survival of BHK-21 cells exposed to Tg.

To strengthen our finding, we restored Alix expression in *Sh* Alix BHK-21 cells using an expression vector coding for the protein (Fig. 2C, upper panel). Even though, Alix over-expression did not affect the sensitivity of *Sh* control cells towards Tg, it almost completely restored Tg-induced death in *Sh* Alix cells (Fig. 2C). In contrast, RFP expression did not influence viability in either *Sh* control or *Sh* Alix cells. This observation, together with the fact that Alix- Δ ALG-2 did not significantly restore Tg-induced cell death indicates that Alix mediates apoptosis following cytosolic Ca^{2+} elevation and that this activity requires its capacity to bind to ALG-2.




Fig. 3. Reduction of Alix expression impedes caspase 9 activation following Tg treatment in BHK-21 cells. (A) Expression of DN-caspase 9 protects cells from Tg-induced apoptosis. BHK-21 cells were transfected with vectors coding for RFP or dominant-negative forms of either caspase 8-HA or caspase 9-HA. Twenty-four hours later, cells were treated with 1 μ M Tg, 24 h. RFP transfected cells were treated or not with 100 μ M pan-caspase inhibitor BAF (Boc-aspartyl (OMe)-fluoromethylketone; Bio-Rad). Cells were then immunostained using a monoclonal anti-HA antibody and nuclei stained with Hoechst. Ten random fields from two or three different wells were counted and condensed/fragmented nuclei were scored as pyknotic. Results are expressed as the percentage of transfected cells displaying pyknotic nuclei. Data represent means \pm SD of three independent experiments. Statistical analysis was performed by analysis of variance (one-way ANOVA followed by Scheffé's post hoc test). * $P < 0.05$ versus DN-caspase 8 overexpressing BHK-21 cells (treated with DMSO). (B,C) Caspase 9 activation following Tg treatment requires Alix expression. *Sh* control or *Sh* Alix BHK-21 cells were treated with 1 μ M Tg or with DMSO vehicle. After 24 h cells were labeled for activated caspase 9 (red LETD-fmk) and stained with Hoechst. (B) Representative fields. Arrows indicate cells with pyknotic nuclei displaying activated caspase 9. (C) Quantification of cells displaying activated caspase 9. Results are expressed as the percentage of cells displaying activated caspases 9 could be observed between *Sh* Alix cells with or without Tg ($p = 0.14$).

3.3. Caspase 9 activation is required for Tg-induced apoptosis

Since Alix over-expression is sufficient to activate caspases in CGN [16], we hypothesized that Alix downregulation might affect activation of apical caspases following Tg treatment. Because some reports involved caspase 8 [22] and caspase 9 [7,23] in ER stress, we first determined which of these two caspases initiates Tg-induced death in BHK-21 cells. For this, we tested the effects on cell survival of catalytically dead mutants (DN-caspases) of caspases 8 and 9 which inhibit activation of their relevant endogenous counterparts. Twenty-four hours after transfection, BHK-21 cells were treated with 1 μ M Tg and further grown for 24 h (Fig. 3A). This treatment robustly increased the number of apoptotic cells among control cultures expressing RFP. Tg-induced apoptosis was significantly reduced by DN-caspase 9 expression as did the pan-caspase inhibitor BAF. In contrast expression of DN-caspase 8 did not afford protection but even significantly sensitized cells to Tg. Thus, activation of caspase 9 seems to be necessary for BHK-21 apoptosis induced by Tg.

3.4. Alix is involved in caspase 9 activation required for thapsigargin-induced apoptosis

We then tested the relationship between Alix and caspase 9 activation following Tg. Using a caspase 9 detection kit, activated caspase 9 could not be detected in either *Sh* control or *Sh* Alix cells (Fig. 3B). However, following a 24 h exposure to Tg, caspase 9 activation could be detected in 30% of *Sh* control cells. Most of these cells displayed condensed or fragmented nuclei typical of apoptosis. In sharp contrast, the number of cells containing activated caspase 9 did not increase significantly after Tg treatment in cells in which Alix was knocked down (Fig. 3C). This demonstrates that Alix is required for caspase 9 activation following Tg treatment.

4. Discussion

Alix controls some aspects of cell death, particularly neuronal apoptosis induced by various stimuli [15–17,24]. Here, we provide the first evidence for a role of Alix in mediating caspase 9 activation and apoptosis involving alteration in Ca^{2+} homeostasis. Expression of a dominant-negative form of Alix (Alix-CT) protects CGN and BHK-21 cells against apoptosis induced by Tg, which blocks the sarcoendoplasmic reticulum Ca^{2+} -ATPase (SERCA). Deletion of the ALG-2 binding site on Alix-CT abolished the capacity of the mutant to block cell death. Consistent with this, Alix knock-down in BHK-21 cells protected these cells against Tg whereas complementation with wt Alix (wild-type Alix) restored their susceptibility to Tg in a way depending on its capacity to bind ALG-2. Finally, we also show that Alix is necessary for caspase 9 activation, a process necessary for induction of apoptosis induced by Tg.

Our finding that, in BHK cells, caspase 9 is instrumental for apoptosis induced by cytosolic elevation is in good agreement with several reports showing that cells derived from caspase 9 knock-out animals or in which caspase 9 activation was blocked, are protected against death induced by alteration in ER calcium homeostasis [6–8]. More disputed is the way of activation of caspase 9, which was demonstrated to be both dependent and independent of the intrinsic/Apaf-1 pathway [6,8]. Rao et al. demonstrated that ALG-2 co-immunoprecipitates with vectorially expressed caspases 9 and 12. They also showed that knock-down of ALG-2 inhibits activation of procaspase 9 and suggested that activation by Tg stress features an Apaf-1-independent pathway that requires, among others, ATP, calcium, ALG-2, caspase 9, caspase 12 and additional factors present in microsomes [8]. Several studies have now shown that, in various situations and several types of cells [25], caspase 12 is not critical

for the ER stress-induced apoptotic cascade [6,22,23,26]. Our present results demonstrate that, like ALG-2, Alix expression is necessary for caspase 9 activation induced by Tg.

Another link between the ER and ALG-2 had been demonstrated by Draeby et al. [27] who reported that ALG-2 interacts with the ER-localized scotin protein which is a proapoptotic-p53-inducible protein. Noteworthy is that deletion of the PRD C-terminal region of scotin containing the ALG-2 binding site, abolished the ER localization and proapoptotic function of the protein. Furthermore, ALG-2 is temporarily and Ca^{2+} -dependently recruited by Sec31 of Cop-II at ER exit sites [28–30]. Alix also translocates to ER exit sites in response to a rise in cytoplasmic Ca^{2+} suggesting a possible function of the protein in this compartment [30]. In line with this, we have previously reported that Alix-CT expressed in HEK-293 cells concentrates on tubular vesicular structures stained with an antibody recognizing ER resident proteins bearing KDEL [20]. Thus, Alix and ALG-2 localization is compatible with a scenario in which changes in ER Ca^{2+} homeostasis induce the Ca^{2+} -dependent recruitment by ALG-2 of Alix and additional proteins required for the building of a platform allowing activation of caspase 9.

Acknowledgments

This study was partly supported by the Institut National de la Santé et de la Recherche Médicale (Inserm), the Université Joseph Fourier and by funds from the Association Française contre les Myopathies, the Association pour la recherche sur le cancer (ARC), the Association pour la Recherche sur la Sclérose Latérale Amyotrophique (ARS) and the Région Rhône-Alpes. F. Strappazzon is the recipient of a grant from the Ministère de l'Éducation Nationale, de l'Enseignement Supérieur et de la Recherche. The authors gratefully acknowledge P. Mehlen for caspase constructs, F.J. Hemming for revising the English manuscript, A.L. Mahul and S. Fraboulet for valuable suggestions throughout this work.

References

- [1] M.J. Berridge, M.D. Bootman, P. Lipp, Calcium – a life and death signal, *Nature* 395 (1998) 645–648.
- [2] M.J. Berridge, P. Lipp, M.D. Bootman, Signal transduction. The calcium entry pas de deux, *Science* 287 (2000) 1604–1605.
- [3] L.K. Nutt, A. Pataer, J. Pahler, B. Fang, J. Roth, D.J. McConkey, S.G. Swisher, Bax and Bak promote apoptosis by modulating endoplasmic reticular and mitochondrial Ca^{2+} stores, *J. Biol. Chem.* 277 (2002) 9219–9225.
- [4] B. Tombal, S.R. Denmeade, J.M. Gillis, J.T. Isaacs, A supramicromolar elevation of intracellular free calcium ($[\text{Ca}^{2+}]_i$) is consistently required to induce the execution phase of apoptosis, *Cell Death Differ.* 9 (2002) 561–573.
- [5] R. Rizzuto, P. Pinton, D. Ferrari, M. Chami, G. Szabadkai, P.J. Magalhaes, F. Di Virgilio, T. Pozzan, Calcium and apoptosis: facts and hypotheses, *Oncogene* 22 (2003) 8619–8627.
- [6] F. Di Sano, E. Ferraro, R. Tufi, T. Achsel, M. Piacentini, F. Cecconi, Endoplasmic reticulum stress induces apoptosis by an apoptosome-dependent but caspase 12-independent mechanism, *J. Biol. Chem.* 281 (2006) 2693–2700.
- [7] A. Masud, A. Mohapatra, S.A. Lakhani, A. Ferrandino, R. Hakem, R.A. Flavell, Endoplasmic reticulum stress-induced death of mouse embryonic fibroblasts requires the intrinsic pathway of apoptosis, *J. Biol. Chem.* 282 (2007) 14132–14139.
- [8] R.V. Rao, K.S. Poksay, S. Castro-Obregon, B. Schilling, G. del Rio, B.W. Gibson, H.M. Ellerby, D.E. Bredesen, R.H. Row, R.H. Row, Molecular components of a cell death pathway activated by endoplasmic reticulum stress, *J. Biol. Chem.* 279 (2004) 177–187.
- [9] H. Suzuki, M. Kawasaki, T. Inuzuka, M. Okumura, T. Kakiuchi, H. Shibata, S. Wakatsuki, M. Maki, Structural basis for Ca^{2+} -dependent formation of ALG-2/Alix peptide complex: Ca^{2+} /EF3-driven arginine switch mechanism, *Structure* 16 (2008) 1562–1573.
- [10] K. Katoh, H. Shibata, H. Suzuki, A. Nara, K. Ishidoh, E. Kominami, T. Yoshimori, M. Maki, The ALG-2-interacting protein Alix associates with CHMP4b, a human homologue of yeast Snf7 that is involved in multivesicular body sorting, *J. Biol. Chem.* 278 (2003) 39104–39113.
- [11] M. Missotten, A. Nichols, K. Rieger, R. Sadoul, Alix, a novel mouse protein undergoing calcium-dependent interaction with the apoptosis-linked-gene 2 (ALG-2) protein, *Cell Death Differ.* 6 (1999) 124–129.
- [12] P. Vito, L. Pellegrini, C. Guiet, L. D'Adamio, Cloning of AIP1, a novel protein that associates with the apoptosis-linked gene ALG-2 in a Ca^{2+} -dependent reaction, *J. Biol. Chem.* 274 (1999) 1533–1540.

- [13] R. Sadoul, Do Alix and ALG-2 really control endosomes for better or for worse, *Biol. Cell* 98 (2006) 69–77.
- [14] G. Odorizzi, The multiple personalities of Alix, *J. Cell Sci.* 119 (2006) 3025–3032.
- [15] F.J. Hemming, S. Fraboulet, B. Blot, R. Sadoul, Early increase of apoptosis-linked gene-2 interacting protein X in areas of kainate-induced neurodegeneration, *Neuroscience* 123 (2004) 887–895.
- [16] Y. Trioulier, S. Torch, B. Blot, N. Cristina, C. Chatellard-Causse, J.M. Verna, R. Sadoul, Alix, a protein regulating endosomal trafficking, is involved in neuronal death, *J. Biol. Chem.* 279 (2004) 2046–2052.
- [17] A.L. Mahul-Mellier, F.J. Hemming, B. Blot, S. Fraboulet, R. Sadoul, Alix, making a link between apoptosis-linked gene-2, the endosomal sorting complexes required for transport, and neuronal death in vivo, *J. Neurosci.* 26 (2006) 542–549.
- [18] A.L. Mahul-Mellier, F. Strappazzon, A. Petiot, C. Chatellard-Causse, S. Torch, B. Blot, K. Freeman, L. Kuhn, J. Garin, J.M. Verna, S. Fraboulet, R. Sadoul, F.J. Hemming, Alix and ALG-2 are involved in tumor necrosis factor receptor 1-induced cell death, *J. Biol. Chem.* 283 (2008) 34954–34965.
- [19] A.L. Mahul-Mellier, F. Strappazzon, C. Chatellard-Causse, B. Blot, D. Beal, S. Torch, F. Hemming, A. Petiot, J.M. Verna, S. Fraboulet, R. Sadoul, Alix and ALG-2 make a link between endosomes and neuronal death, *Biochem. Soc. Trans.* 37 (2009) 200–203.
- [20] C. Chatellard-Causse, B. Blot, N. Cristina, S. Torch, M. Missotten, R. Sadoul, Alix (ALG-2-interacting protein X), a protein involved in apoptosis, binds to endophilins and induces cytoplasmic vacuolization, *J. Biol. Chem.* 277 (2002) 29108–29115.
- [21] T. Tuomikoski, M.A. Felix, M. Doree, J. Gruenberg, Inhibition of endocytic vesicle fusion in vitro by the cell-cycle control protein kinase cdc2, *Nature* 342 (1989) 942–945.
- [22] A. Jimbo, E. Fujita, Y. Kouroku, J. Ohnishi, N. Inohara, K. Kuida, K. Sakamaki, S. Yonehara, T. Momoi, ER stress induces caspase-8 activation, stimulating cytochrome c release and caspase-9 activation, *Exp. Cell Res.* 283 (2003) 156–166.
- [23] E.A. Obeng, L.H. Boise, Caspase-12 and caspase-4 are not required for caspase-dependent endoplasmic reticulum stress-induced apoptosis, *J. Biol. Chem.* 280 (2005) 29578–29587.
- [24] D. Blum, F.J. Hemming, M.C. Galas, S. Torch, L. Cuvelier, S.N. Schiffmann, R. Sadoul, Increased Alix (apoptosis-linked gene-2 interacting protein X) immunoreactivity in the degenerating striatum of rats chronically treated by 3-nitropropionic acid, *Neurosci. Lett.* 368 (2004) 309–313.
- [25] Y. Kosuge, T. Sakikubo, K. Ishige, Y. Ito, Comparative study of endoplasmic reticulum stress-induced neuronal death in rat cultured hippocampal and cerebellar granule neurons, *Neurochem. Int.* 49 (2006) 285–293.
- [26] H. Shiraishi, H. Okamoto, A. Yoshimura, H. Yoshida, ER stress-induced apoptosis and caspase-12 activation occurs downstream of mitochondrial apoptosis involving Apaf-1, *J. Cell Sci.* 119 (2006) 3958–3966.
- [27] I. Draeby, Y.L. Woods, J.M. la Cour, J. Mollerup, J.C. Bourdon, M.W. Berchtold, The calcium binding protein ALG-2 binds and stabilizes Scotin, a p53-inducible gene product localized at the endoplasmic reticulum membrane, *Arch. Biochem. Biophys.* 467 (2007) 87–94.
- [28] J.M. la Cour, J. Mollerup, M.W. Berchtold, ALG-2 oscillates in subcellular localization, untemporally with calcium oscillations, *Biochem. Biophys. Res. Commun.* 353 (2007) 1063–1067.
- [29] H. Shibata, H. Suzuki, H. Yoshida, M. Maki, ALG-2 directly binds Sec31A and localizes at endoplasmic reticulum exit sites in a Ca^{2+} -dependent manner, *Biochem. Biophys. Res. Commun.* 353 (2007) 756–763.
- [30] A. Yamasaki, K. Tani, A. Yamamoto, N. Kitamura, M. Komada, The Ca^{2+} -binding protein ALG-2 is recruited to endoplasmic reticulum exit sites by Sec31A and stabilizes the localization of Sec31A, *Mol. Biol. Cell* 17 (2006) 4876–4887.