



## Visualization and quantification of endoplasmic reticulum $\text{Ca}^{2+}$ in renal cells using confocal microscopy and Fluo5F

Andre C. Eaddy<sup>a</sup>, Rick G. Schnellmann<sup>a,b,\*</sup>

<sup>a</sup> Center for Cell Death, Injury, and Regeneration, Department of Pharmaceutical and Biomedical Sciences, Medical University of South Carolina, Charleston, SC 29425, United States

<sup>b</sup> Ralph H. Johnson VA Medical Center Charleston, SC 29401, United States

### ARTICLE INFO

#### Article history:

Received 23 November 2010

Available online 3 December 2010

#### Keywords:

Calcium  
Endoplasmic reticulum  
Confocal microscopy  
Kidney  
Fluo5f

### ABSTRACT

Sarcoplasmic/endoplasmic reticulum (ER)  $\text{Ca}^{2+}$  is the most abundant store of intracellular  $\text{Ca}^{2+}$ , and its release is an important trigger of physiological and cell death pathways. Previous work in our laboratory revealed the importance of ER  $\text{Ca}^{2+}$  in toxicant-induced renal proximal tubular cell (RPTC) death. The purpose of this study was to evaluate the use of confocal microscopy and Fluo5F, a low affinity  $\text{Ca}^{2+}$  indicator, to directly monitor changes in RPTC ER  $\text{Ca}^{2+}$ . Fluo5F staining reflected ER  $\text{Ca}^{2+}$ , resolved ER structure, and showed no colocalization with tetramethyl rhodamine methyl ester (TMRM), a marker of mitochondrial membrane potential. Thapsigargin, an ER  $\text{Ca}^{2+}$  pump inhibitor, decreased ER fluorescence by 30% and 55% at 5 and 15 min, respectively, whereas A23187, a  $\text{Ca}^{2+}$  ionophore caused more rapid ER  $\text{Ca}^{2+}$  release (55% and 75% decrease in fluorescence at 5 and 15 min).

Carbonylcyanide-p-trifluoromethoxyphenylhydrazone (FCCP), a mitochondrial uncoupler, added at the end of the experiment, further decreased ER fluorescence after thapsigargin treatment, revealing that thapsigargin did not release all ER  $\text{Ca}^{2+}$ . In contrast, FCCP did not decrease ER fluorescence after A23187 treatment, suggesting complete ER  $\text{Ca}^{2+}$  release. ER  $\text{Ca}^{2+}$  release in response to A23187 or thapsigargin resulted in a modest but significant decrease in mitochondrial membrane potential. These data provide evidence that confocal microscopy and Fluo5F are useful and effective tools for directly monitoring ER  $\text{Ca}^{2+}$  in live cells.

© 2010 Elsevier Inc. All rights reserved.

### 1. Introduction

$\text{Ca}^{2+}$  acts as a universal second messenger and regulates numerous cellular functions including metabolism, motility, and transport [1]. Loss of  $\text{Ca}^{2+}$  homeostasis is critical to many disease processes and is a major component of cell death pathways including necrosis, apoptosis, and autophagy [2–8]. Sarcoplasmic/endoplasmic reticulum (ER)  $\text{Ca}^{2+}$  is the most abundant store of intracellular  $\text{Ca}^{2+}$  and its disruption often initiates the deleterious cascade of events leading to cell death and dysfunction [5–11].

Traditional methods of measuring of ER  $\text{Ca}^{2+}$  are indirect or require difficult probe loading techniques (i.e. membrane perme-

abilization, microinjection, or fused cell hybrids). For example, human embryonic kidney cells require 1 h of dye loading (2  $\mu\text{M}$  Fluo-3 at room temperature) and extracellular  $\text{Ca}^{2+}$  chelation using BAPTA or EGTA, just to monitor increases in cytosolic  $\text{Ca}^{2+}$  as an indirect measure of ER  $\text{Ca}^{2+}$  stores [12]. To monitor ER  $\text{Ca}^{2+}$  directly, Montero and Robert et al. engineered an ER-targeted aequorin chimera, a  $\text{Ca}^{2+}$ -sensitive photoprotein with a lower affinity for  $\text{Ca}^{2+}$ . However, in HeLa and skeletal muscle cells, the aequorin chimera was rapidly saturated by the high  $\text{Ca}^{2+}$  concentrations within the ER and required a non-physiological reduction of ER  $\text{Ca}^{2+}$  to detect changes in  $\text{Ca}^{2+}$  stores [13,14].

Primary cultures of renal proximal tubular cells (RPTC) are highly aerobic, gluconeogenic, and exhibit robust transport capacity, making them ideal for the study of kidney tubular function and injury [15,16]. Previous work in our laboratory revealed the importance of ER  $\text{Ca}^{2+}$  in toxicant-induced kidney injury; although, the mechanism by which  $\text{Ca}^{2+}$  plays such a pivotal role is not completely understood [5,7,9,10,17,18].

Confocal microscopy is a useful method for the visualization and quantification of fluorophores at the subcellular level in living cells, and is compatible with most  $\text{Ca}^{2+}$  indicators. Chemical fluorescent (UV and visible-wavelength excitation fluorescent indicators) and

**Abbreviations:** ER, endoplasmic reticulum; EGTA, ethylene glycol tetraacetic acid; RPTC, renal proximal tubular cell; TMRM, tetramethyl rhodamine methyl ester; Fluo5F, Fluo-5F-AM; LSCM, laser scanning confocal microscopy; FCCP, carbonylcyanide-p-trifluoromethoxyphenylhydrazone; SERCA, sarcoplasmic/endoplasmic reticulum ATPase; DAG, diacylglycerol; IP3, inositol triphosphate; DMSO, dimethyl sulfoxide.

\* Corresponding author at: Department of Pharmaceutical and Biomedical Sciences, Medical University of South Carolina, 280 Calhoun St, MSC 140, Charleston, SC 29425-1400, United States. Fax: +1 843 792 2620.

E-mail address: [schnell@muscc.edu](mailto:schnell@muscc.edu) (R.G. Schnellmann).

bioluminescent calcium indicators ( $\text{Ca}^{2+}$  binding photoproteins and GFP-based  $\text{Ca}^{2+}$  indicators) differ in their characteristics (loading requirements, excitation/emission spectrum, permeability, compartmentalization, relative brightness, and  $\text{Ca}^{2+}$  affinity) and have inherent drawbacks (i.e., dye leakage, cytotoxicity, bleaching, autofluorescence, intracellular buffering, and lack of ion specificity)[19]. Fluo5F is a chemical fluorescent  $\text{Ca}^{2+}$  indicator with a lower affinity for  $\text{Ca}^{2+}$  ( $K_d \sim 2.3 \mu\text{M}$ ), limited cytotoxicity and bleaching, and high  $\text{Ca}^{2+}$  specificity, making it ideal for studying ER  $\text{Ca}^{2+}$ . The purpose of this study was to evaluate the use of confocal microscopy and Fluo5F for directly monitoring changes in ER  $\text{Ca}^{2+}$  in RPTC.

## 2. Materials and methods

### 2.1. Materials

Female New Zealand White rabbits (1.5–2.0 kg) were purchased from Myrtle's Rabbitry (Thompson Station, TN). Tetramethyl rhodamine methyl ester (TMRM) and Fluo-5F-AM were purchased from Molecular Probes, Invitrogen (Carlsbad, CA). All other chemical and materials were obtained from Sigma Chemical (St. Louis, MO).

### 2.2. Isolation of rabbit RPTC and culture conditions

Rabbit RPTC were isolated using the iron oxide perfusion method and grown to confluence in 35 mm tissue culture dishes under improved conditions as previously described [15,16]. The culture medium was a 1:1 mixture of Dulbecco's modified Eagle's medium/Ham's F-12 medium (without glucose, phenol red, or sodium pyruvate) supplemented with 15 mM HEPES buffer, 2.5 mM L-glutamine, 1  $\mu\text{M}$  pyridoxine HCl, 15 mM sodium bicarbonate, and 6 mM lactate. Hydrocortisone (50 nM), selenium (5 ng/ml), human transferrin (5  $\mu\text{g}/\text{ml}$ ), bovine insulin (10 nM), and L-ascorbic acid-2-phosphate (50  $\mu\text{M}$ ) were added daily to fresh culture medium.

### 2.3. RPTC loading

RPTC were loaded with 2  $\mu\text{M}$  Fluo-5F-AM (Fluo5F) and 100 nM TMRM for 20 min at 37°C, washed twice with 37°C phosphate buffered saline, and media was replaced. Then, 1  $\mu\text{M}$  Fluo5F and 40 nM TMRM were added to the media to maintain dye equilibrium and incubated for 30 min at 37°C prior to imaging. The TMRM loading protocol was modified from Lemasters and Ramshesh [20].

### 2.4. Microscopic imaging and analysis

RPTC were imaged on a Leica Microsystems, TCS SP2 AOBS laser scanning confocal microscope (LSCM) using standardized pinhole,

gain, and black level settings. A  $63 \times 0.9$  NA water-immersion objective was used in an upright microscope configuration. All images were acquired at 8-bit resolution and at  $1024 \times 1024$  with a line averaging of two.

RPTC were monitored for 10 min prior to treatment to establish baseline. Then, RPTC were treated and monitored an additional 15 min prior to Carbonylcyanide-p-trifluoromethoxyphenylhydrazine (FCCP, 1  $\mu\text{M}$ ) addition to depolarize mitochondria. Images were acquired in 1 min intervals during the course of experiments. To quantify data, images were analyzed using Adobe Photoshop. Mean intensities of images in the green and red channel after background subtraction was interpreted as a quantitative measure of ER  $\text{Ca}^{2+}$  and mitochondrial membrane potential, respectively. Values are graphed as the percent mean intensity of each subsequent image vs. mean intensity of the image taken at time zero. Images comprise of 7–10 cells.


### 2.5. Statistical analysis

RPTC isolated from one animal represents one experiment ( $N = 1$ ). Data are presented as means  $\pm$  SE, and multiple means were compared using Student–Newman–Keuls test at each time point. Means with values  $p < 0.05$  were considered statistically significant.


## 3. Results and discussion

To investigate the use of confocal microscopy to monitor ER  $\text{Ca}^{2+}$  dynamics in RPTC, Fluo5F fluorescence was monitored as described above (Fig. 1A). Fluo5F fluorescence revealed a reticular pattern around nuclei void of staining. The peripheral granularity of staining and scarcity of staining at the cell margins provided evidence that the probe is not fluorescing in the cytosol. To confirm that Fluo5F was not in the mitochondria, RPTC were loaded with tetramethyl rhodamine methyl ester (TMRM), a marker of polarized mitochondria. TMRM staining (Fig. 1B) had a punctuate/tubular pattern unlike Fluo5F staining with no co-localization (Fig. 1C), confirming that Fluo5F does not reflect mitochondrial  $\text{Ca}^{2+}$ , the second most significant storage/buffering compartment for intracellular  $\text{Ca}^{2+}$  [4].


We investigated the use of Fluo5F to monitor real-time changes in RPTC ER  $\text{Ca}^{2+}$ . TMRM was used to measure mitochondrial membrane potential. In control cells, both Fluo5F and TMRM fluorescence intensity, morphology, and resolution were maintained over time (Fig. 2). Carbonylcyanide-p-trifluoromethoxyphenylhydrazine (FCCP), a mitochondrial uncoupler, was added at the end of each experiment to cause mitochondrial depolarization and indirectly ER  $\text{Ca}^{2+}$  release, due to ATP depletion and cessation of sarcoplasmic/ER  $\text{Ca}^{2+}$  ATPase (SERCA) activity.




**Fig. 1.** Morphology of Renal Proximal Tubule Cells (RPTC) co-loaded with Fluo5F and TMRM using confocal microscopy. RPTC were co-loaded with Fluo5F (2  $\mu\text{M}$ ) and TMRM (100 nM) as described in Section 2. Green fluorescence of Fluo5F and red fluorescence of TMRM were imaged by laser scanning confocal microscopy. The white bar represents 10  $\mu\text{m}$ . Fluo5F fluorescence (A) represents endoplasmic reticulum (ER)  $\text{Ca}^{2+}$ , TMRM fluorescence (B) represents polarized mitochondria, and there was no co-localization (C). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 2.** ER  $\text{Ca}^{2+}$  release RPTC exposed to A23187 and thapsigargin. RPTC were loaded with Fluo5F and TMRM (see in Fig. 1), monitored for 10 min to establish baseline, exposed to diluent (DMSO, Control), A23187 (10  $\mu\text{M}$ ), or thapsigargin (10  $\mu\text{M}$ ) for 15 min, and then exposed to FCCP (1  $\mu\text{M}$ ), a mitochondrial uncoupler. Open arrows represent time of A23187 or thapsigargin treatment and closed arrows represent the addition of FCCP. Green fluorescence (A) and red fluorescence (B) represents ER  $\text{Ca}^{2+}$  and mitochondrial polarization, respectively. Note the more rapid ER  $\text{Ca}^{2+}$  release (A) in response to A23187 exposure compared to thapsigargin and the rapid loss of mitochondrial polarization (B) in response to FCCP. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 3.** Quantification of A23187- and thapsigargin- induced ER  $\text{Ca}^{2+}$  release in RPTC. RPTC were co-loaded with Fluo5F and TMRM, treated as described in Fig. 2, and images acquired as described in Section 2. Values are mean intensities of images in the green channel  $\pm$  SE and loss of intensity represents ER  $\text{Ca}^{2+}$  release. The open arrow represents exposure to A23187, thapsigargin, or diluent control and the closed arrow represents the addition of FCCP. Note the more rapid ER  $\text{Ca}^{2+}$  release after A23187 exposure compared to thapsigargin and an additional loss of fluorescence with FCCP following thapsigargin exposure.



**Fig. 4.** Quantification of A23187- and thapsigargin-induced mitochondrial depolarization in RPTC. See above for experimental details. Values are mean intensities of images in the red channel  $\pm$  SE and loss of intensity represents mitochondrial depolarization. The open arrow represents exposure to A23187, thapsigargin, or diluent control and the closed arrow represents the addition of FCCP. Note the decrease in mitochondrial polarization following A23187 or thapsigargin exposure compared with control and the rapid loss of mitochondrial polarization following FCCP exposure in all groups.

We then explored the effects of two pharmacological agents known to alter ER  $\text{Ca}^{2+}$ . RPTC were treated with the SERCA inhibitor, thapsigargin, and the  $\text{Ca}^{2+}$  ionophore, A23187. Thapsigargin decreased ER  $\text{Ca}^{2+}$  over several min (Fig. 2A), which was more indicative of ER  $\text{Ca}^{2+}$  leakage than receptor-dependent ER  $\text{Ca}^{2+}$

release that is characterized by DAG, ryanodine, and  $\text{IP}_3$  receptor sensitive stores [1,21,22]. In contrast, A23187 caused a more rapid release of ER  $\text{Ca}^{2+}$ , an anticipated finding due to its ionophore activity.



Temporal and incremental changes in ER  $\text{Ca}^{2+}$  release and mitochondrial membrane potential were quantified (Fig. 3). A23187 decreased fluorescence by 50% and 70% at 5 and 15 min after treatment, respectively, and FCCP addition did not decrease fluorescence further. Thapsigargin decreased fluorescence by 25% and 55% at 5 and 15 min after treatment, respectively. In contrast to A23187, FCCP addition further decreased fluorescence after thapsigargin, providing evidence that thapsigargin did not completely release all ER  $\text{Ca}^{2+}$  stores under these conditions.

The advantage of this approach is that ER  $\text{Ca}^{2+}$  stores are measured directly, allowing observation of spatiotemporal differences between A23187- and thapsigargin-induced ER  $\text{Ca}^{2+}$  releases that would otherwise be masked by measuring cytosolic free  $\text{Ca}^{2+}$ . Interestingly, Jiang et al. reported a higher rate and magnitude of cytosolic  $\text{Ca}^{2+}$  increase in response to A23187 over thapsigargin using Fluo-3, but this approach required chelation of extracellular  $\text{Ca}^{2+}$  prior to treatment [12].

The effect of ER  $\text{Ca}^{2+}$  release on mitochondrial polarization was monitored using TMRM (Fig. 4). A23187- and thapsigargin-induced ER  $\text{Ca}^{2+}$  release decreased TMRM fluorescence by approximately 10%. In contrast, FCCP, the mitochondrial uncoupler added at the end of each experiment, decreased TMRM fluorescence by 60%, reflecting complete mitochondrial depolarization. Together, these data reveal that ER  $\text{Ca}^{2+}$  release produced by A23187 or thapsigargin partially decreases mitochondrial polarization and is consistent with previous reports [23].

In conclusion, confocal microscopy and Fluo5F are valuable tools for visualization of spatiotemporal dynamics of ER  $\text{Ca}^{2+}$  release in complex cellular systems in real time and will aid in our understanding of cellular dynamics and death.

## Disclaimer

The contents of this manuscript do not represent the views of the Department of Veteran Affairs or the United States Government.

## Acknowledgments

This work was supported by an NIH fellowship (F30 ES015964) to AE and NIH grant (DK-062028) to RGS, and by the Biomedical Laboratory Research and Development Program of the Department of Veterans Affairs. Animal facilities were funded by NIH grant C06 RR-015455 and Cell and Molecular Imaging (CMI) facility was supported, in part, by grant P30 CA138313 to the Hollings Cancer Center, Medical University of South Carolina. The authors also thank Dr. Venkat Ramshesh and Dr. John Lemasters (Medical University of South Carolina) for their advice and constructive discussions during the execution of these experiments.

## References

- [1] M.J. Berridge, Inositol trisphosphate and calcium signaling, *Ann. NY Acad. Sci.* 766 (1995) 31–43.

- [2] B.S. Cummings, G.R. Kinsey, L.J. Bolchoz, R.G. Schnellmann, Identification of caspase-independent apoptosis in epithelial and cancer cells, *J. Pharmacol. Exp. Ther.* 310 (2004) 126–134.
- [3] J.J. Lemasters, A.L. Nieminen, T. Qian, L.C. Trost, S.P. Elmore, Y. Nishimura, et al., The mitochondrial permeability transition in cell death: a common mechanism in necrosis, Apoptosis and autophagy, *Biochim. Biophys. Acta* 1366 (1998) 177–196.
- [4] J.J. Lemasters, T. Qian, S.P. Elmore, L.C. Trost, Y. Nishimura, B. Herman, et al., Confocal microscopy of the mitochondrial permeability transition in necrotic cell killing, apoptosis and autophagy, *BioFactors* 8 (1998) 283–285.
- [5] R.G. Schnellmann, Intracellular calcium chelators and oxidant-induced renal proximal tubule cell death, *J. Biochem. Toxicol.* 6 (1991) 299–303.
- [6] S.L. Waters, S.S. Sarang, K.K. Wang, R.G. Schnellmann, Calpains mediate calcium and chloride influx during the late phase of cell injury, *J. Pharmacol. Exp. Ther.* 283 (1997) 1177–1184.
- [7] S.L. Waters, J.K. Wong, R.G. Schnellmann, Depletion of endoplasmic reticulum calcium stores protects against hypoxia- and mitochondrial inhibitor-induced cellular injury and death, *Biochem. Biophys. Res. Commun.* 240 (1997) 57–60.
- [8] T. Pozzan, R. Rizzuto, P. Volpe, J. Meldolesi, Molecular and cellular physiology of intracellular calcium stores, *Physiol. Rev.* 74 (1994) 595–636.
- [9] X. Liu, T. Van Vleet, R.G. Schnellmann, The role of calpain in oncotic cell death, *Annu. Rev. Pharmacol. Toxicol.* 44 (2004) 349–370.
- [10] J.F. Harriman, X.L. Liu, M.D. Aleo, K. Machaca, R.G. Schnellmann, Endoplasmic reticulum  $\text{Ca}^{2+}$  signaling and calpains mediate renal cell death, *Cell Death Differ.* 9 (2002) 734–741.
- [11] S. Muruganandan, A.E. Cribb, Calpain-induced endoplasmic reticulum stress and cell death following cytotoxic damage to renal cells, *Toxicol. Sci.* 94 (2006) 118–128.
- [12] Y. Jiang, A.M. Cypess, E.D. Muse, C.R. Wu, C.G. Unson, R.B. Merrifield, et al., Glucagon receptor activates extracellular signal-regulated protein kinase 1/2 via cAMP-dependent protein kinase, *Proc. Natl. Acad. Sci. USA* 98 (2001) 10102–10107.
- [13] V. Robert, F. De Giorgi, M.L. Massimino, M. Cantini, T. Pozzan, Direct monitoring of the calcium concentration in the sarcoplasmic and endoplasmic reticulum of skeletal muscle myotubes, *J. Biol. Chem.* 273 (1998) 30372–30378.
- [14] M. Montero, M. Brini, R. Marsault, J. Alvarez, R. Sitia, T. Pozzan, et al., Monitoring dynamic changes in free  $\text{Ca}^{2+}$  concentration in the endoplasmic reticulum of intact cells, *EMBO J.* 14 (1995) 5467–5475.
- [15] G. Nowak, R.G. Schnellmann, Improved culture conditions stimulate gluconeogenesis in primary cultures of renal proximal tubule cells, *Am. J. Physiol.* 268 (1995) C1053–C1061.
- [16] G. Nowak, R.G. Schnellmann, L-ascorbic acid regulates growth and metabolism of renal cells: improvements in cell culture, *Am. J. Physiol.* 271 (1996) C2072–C2080.
- [17] B.S. Cummings, J. McHowat, R.G. Schnellmann, Role of an endoplasmic reticulum  $\text{Ca}^{2+}$ -independent phospholipase A2 in cisplatin-induced renal cell apoptosis, *J. Pharmacol. Exp. Ther.* 308 (2004) 921–928.
- [18] B.S. Cummings, J. McHowat, R.G. Schnellmann, Role of an endoplasmic reticulum  $\text{Ca}^{2+}$ -independent phospholipase A(2) in oxidant-induced renal cell death, *Am. J. Physiol. Renal Physiol.* 283 (2002) F492–F498.
- [19] A. Takahashi, P. Camacho, J.D. Lechleiter, B. Herman, Measurement of intracellular calcium, *Physiol. Rev.* 79 (1999) 1089–1125.
- [20] J.J. Lemasters, V.K. Ramshesh, Imaging of mitochondrial polarization and depolarization with cationic fluorophores, *Methods Cell Biol.* 80 (2007) 283–295.
- [21] J.H. Bian, T.K. Ghosh, J.C. Wang, D.L. Gill, Identification of intracellular calcium pools. Selective modification by thapsigargin, *J. Biol. Chem.* 266 (1991) 8801–8806.
- [22] K. Hirose, S. Kadowaki, M. Tanabe, H. Takeshima, M. Iino, Spatiotemporal dynamics of inositol 1,4,5-trisphosphate that underlies complex  $\text{Ca}^{2+}$  mobilization patterns, *Science* 284 (1999) 1527–1530.
- [23] J.J. Lemasters, T.P. Theruvath, Z. Zhong, A.L. Nieminen, Mitochondrial calcium and the permeability transition in cell death, *Biochim. Biophys. Acta* 1787 (2009) 1395–1401.