



Monoubiquitination of Tob/BTG family proteins competes with degradation-targeting polyubiquitination

Toru Suzuki^{a,*}, Minsoo Kim^b, Hiroko Kozuka-Hata^c, Masato Watanabe^d, Masaaki Oyama^c, Kouhei Tsumoto^{c,d}, Tadashi Yamamoto^{a,e,*}

^a Division of Oncology, Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan

^b Division of Bacterial Infection, Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan

^c Medical Proteomics Laboratory, Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan

^d Department of Medical Genome Science, School of Frontier Sciences, University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa 277-8562, Japan

^e Cell Signal Unit, Okinawa Institute of Science and Technology, 1919-1 Onna-son, Kunigami, Okinawa 904-0412, Japan

ARTICLE INFO

Article history:

Received 20 April 2011

Available online 28 April 2011

Keywords:

Tob
Monoubiquitination
Polyubiquitination
Degradation

ABSTRACT

Tob belongs to the anti-proliferative Tob/BTG protein family. The expression level of Tob family proteins is strictly regulated both transcriptionally and through post-translational modification. Ubiquitin (Ub)/proteasome-dependent degradation of Tob family proteins is critical in controlling cell cycle progression and DNA damage responses. Various Ub ligases (E3s) are responsible for degradation of Tob protein. Here, we show that Tob family proteins undergo monoubiquitination even in the absence of E3s *in vitro*. Determination of the ubiquitination site(s) in Tob by mass spectrometric analysis revealed that two lysine residues (Lys48 and Lys63) located in Tob/BTG homology domain are ubiquitinated. A mutant Tob, in which both Lys48 and Lys63 are substituted with alanine, is more strongly polyubiquitinated than wild-type Tob *in vivo*. These data suggest that monoubiquitination of Tob family proteins confers resistance against polyubiquitination, which targets proteins for degradation. The strategy for regulating the stability of Tob family proteins suggests a novel role for monoubiquitination.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Tob, a member of the anti-proliferative Tob/BTG family of proteins, is involved in the regulation of cell cycle progression at the G1 phase [1–3]. Tob's expression levels and phosphorylation status change during the cell cycle [2], and decreased expression and/or phosphorylation relative to normal tissues have been detected in several human cancers, including lung and thyroid cancer [3,4]. These observations suggest that the level of Tob expression and its phosphorylation status are strictly regulated to maintain normal cell growth. Whereas the expression of Tob family genes is regulated at the transcriptional level during cell differentiation or DNA damage response [1], it has been reported that post-translationally the Ub/proteasome pathway is also involved in controlling the

Abbreviations: Ub, ubiquitin; E1, ubiquitin-activating enzyme; E2, ubiquitin-conjugating enzyme; E3, ubiquitin ligase; UBD, ubiquitin-binding domain; PCR, polymerase chain reaction; GST, glutathione S-transferase; SDS, sodium dodecyl sulfate; PAGE, polyacrylamide gel electrophoresis.

* Corresponding authors at: Division of Oncology, Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. Fax: +81 3 5449 5413 (T. Yamamoto).

E-mail addresses: toru@ims.u-tokyo.ac.jp (T. Suzuki), tyamamot@ims.u-tokyo.ac.jp (T. Yamamoto).

amounts of these proteins [5]. Indeed, several E3 Ub ligases including SCF^{Skp2} regulate the level of Tob protein [6,7]. Furthermore, proteasome-mediated degradation of Tob is important for regulating cell cycle progression as well as DNA damage-induced apoptosis [6,8].

Proteins targeted for degradation by the Ub/proteasome system are marked by covalent linkage to Ub. Initially, Ub is activated by a Ub-activating enzyme (E1) in an ATP-dependent manner. The activated Ub is subsequently transferred to a Ub-conjugating enzyme (E2), and ultimately binds to a lysine residue on the substrate in a process mediated by an E3 ligase [9]. Different types of Ub modifications have distinct cellular functions [10–12]. A polyUb chain linked to Lys48 is associated with proteasome-dependent degradation, whereas a polyUb chain linked to Lys63 affects endocytosis or DNA repair processes. Moreover, covalent modification of proteins by a single Ub (monoubiquitination) plays important roles in regulation of signal transduction, endocytosis, and DNA repair processes [13,14]. Recent reports have shown a number of Ub-binding domain (UBD)-containing proteins to be monoubiquitinated *in vitro* independently of the activity of E3 ligases [15]. Monoubiquitination of ubiquitin-binding proteins inhibits their capacity to bind to and control the functions of ubiquitinated targets *in vivo* [16].

In this study, we found that Tob/BTG family proteins are mono-ubiquitinated in the absence of E3s *in vitro*. We further showed that the monoubiquitination event correlates with lower levels of poly-ubiquitination. Therefore, monoubiquitination of Tob/BTG family proteins appears to regulate their stability by affecting polyubiquitination and concomitant proteasome-dependent degradation.

2. Materials and methods

2.1. DNA constructs

The cDNAs encoding Tob family genes were inserted into pET-26b expression vector (Novagen) for protein expression in *Escherichia coli*. Since full-length Tob protein could not be prepared due to its instability and insolubility, Tob constructs lacking several amino acids in the carboxy-terminal region were prepared. The Tob deletion constructs (amino acids 1–332, 1–232 and 1–112) were constructed by polymerase chain reaction (PCR). The cDNAs of E2 proteins were cloned by PCR and inserted into pGEX6P-1 vector (GE Healthcare). To express Tob in mammalian cells, we used the SR α promoter-driven mammalian expression vector pME18S [2].

2.2. Preparation of recombinant proteins

6xHis-tagged proteins of the Tob/BTG family were expressed in *E. coli* BL21, and the cells were lysed by sonication in buffer A (50 mM Tris–HCl [pH 8.0], 200 mM NaCl, 1 mM PMSF, 5 mM imidazole). The proteins were purified using Ni–NTA Sepharose beads (Qiagen), and eluted from the beads with buffer A containing 125 mM imidazole. Glutathione S-transferase (GST)-fused E2 proteins expressed in BL21 *E. coli* were purified with glutathione-Sepharose beads (GE Healthcare). The GST portion of each protein was cleaved with precision protease (GE Healthcare). All purified proteins were dialyzed in buffer B (20 mM Tris–HCl [pH 7.5], 50 mM NaCl, 5% glycerol).

2.3. Antibodies

Anti-Tob monoclonal antibody (4B1) was obtained from IBL. Anti-ubiquitin (FK2) and anti-His (OGHis) antibodies were purchased from MBL.

2.4. Cell culture, transfection, and immunoblotting

Cos7 cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum in 5% CO₂ at 37 °C. The cells were transfected with the expression plasmids using FuGENE 6 transfection reagent (Roche). Immunoblot analysis was performed as described previously [2].

2.5. Detection of polyubiquitination *in vivo*

Cos7 cells were co-transfected with Tob expression vectors and 6xHis-Ub expression vector. The cells were treated the next day with 50 μ M MG132 (Peptide Institute) for 3 h. Cell lysates were prepared using denaturing buffer (6 M guanidine–HCl, 10 mM phosphate buffer [pH 8.0], 10 mM imidazole). His-Ub-conjugated proteins were purified with Ni–NTA agarose and analyzed by sodium dodecyl sulfate (SDS)–polyacrylamide gel electrophoresis (PAGE) followed by immunoblotting.

2.6. *In vitro* ubiquitination assay

Bovine ubiquitin (15 μ g, Sigma), recombinant E1 (200 ng, Boston Biochem), recombinant E2 protein (1 μ g), and recombinant

Tob family proteins (100 ng) were mixed in 30 μ l of buffer C (40 mM Tris–HCl [pH 7.4], 5 mM MgCl₂, 2 mM dithiothreitol, 2 μ M ATP). The mixtures were incubated for 1.5 h at 30 °C, and the reaction products were analyzed by SDS–PAGE followed by immunoblotting.

2.7. Mass spectrometric analysis

The *in vitro* ubiquitination of Tob was performed as described above. In total, 5 μ g of recombinant Tob was subjected to the reaction. After the reaction, Tob was purified with Ni–NTA Sepharose in buffer (20 mM Tris–HCl [pH 7.5], 500 mM NaCl, 1% Triton X-100, 5 mM 2-mercaptoethanol). The purified Tob was then separated by SDS–PAGE and visualized using Silver Staining Kit MS (Wako). For mass spectrometric analysis, bands corresponding to ubiquitinated Tob were excised from the gel and digested in the gel with trypsin (1 pmol/ μ l) for 12 h at 37 °C. The eluted peptides were loaded onto the automated nanoflow liquid chromatograph (Dina) and tandem mass spectrometer (QSTAR Elite). The peptide masses obtained by LC–MS/MS analysis were searched against the non-redundant protein sequence database of the National Center for Biotechnology Information using the Mascot search engine (Matrix Science, London, UK).

3. Results

3.1. *In vitro* monoubiquitination of Tob

Tob/BTG family proteins are polyubiquitinated and subsequently degraded by the 26S proteasome [5,8]. Several E3 Ub ligases are responsible for Tob ubiquitination [6,7]. To detect the activity of Tob ubiquitination in cytoplasmic extracts, we set up an *in vitro* ubiquitination assay using recombinant E1, E2 and Tob-His (amino acids 1–332) as well as Ub. Unexpectedly, a single slow migrating form of Tob was detected by immunoblotting even in reactions lacking cytoplasmic extracts (Fig. 1, left and middle panels). The slow migrating form was recognized by the anti-Ub antibody and was produced only when all the protein components were present (Fig. 1, right panel). We did not detect any obvious signals other than the slow migrating form. These data suggest that Tob is monoubiquitinated *in vitro* in a manner independent of E3s.

3.2. Tob/BTG family proteins are monoubiquitinated *in vitro*

We next carried out an *in vitro* ubiquitination assay using the two shorter forms of His-tagged Tob, Tob-His (1–232) and Tob-His (1–112), and found both forms were also monoubiquitinated (Fig. 2A). As the amino acid sequence (amino acids 1–112) is well conserved among the Tob/BTG family proteins, we tested whether the propensity to undergo monoubiquitination is conserved among protein family members. *In vitro* ubiquitination assay followed by immunoblot analysis revealed that both Tob2 and BTG2 also became monoubiquitinated (Fig. 2B). Interestingly, UBCH4 and UBCH5a but not UBCH3 were able to mediate the monoubiquitination of Tob/BTG family proteins.

3.3. Determination of monoubiquitination site(s)

The data shown in Fig. 2A indicate that monoubiquitination of Tob occurs within its N terminal half. To determine the monoubiquitination site(s), we performed mass spectrometry. We prepared large amounts of *in vitro* ubiquitinated Tob proteins and subjected them to SDS–PAGE (Fig. 3A). After silver staining of the gels, the corresponding band was cut out from the gel and analyzed by peptide mass fingerprinting, and we identified Lys48 and Lys63 of Tob




Fig. 1. E3-independent ubiquitination of Tob *in vitro*. *In vitro* ubiquitination assay of recombinant Tob was performed in the presence of the indicated proteins. Recombinant UbcH5b was used as an E2 protein. Immunoblot analysis of the reaction products using anti-Tob antibody (left and middle) and anti-ubiquitin antibody (right) are shown.




Fig. 2. E3-independent ubiquitination of Tob family proteins *in vitro*. *In vitro* ubiquitination assay of the indicated constructs of proteins belonging to the Tob/BTG family were performed. E2 proteins were used as indicated. (A) The reaction products of Tob-His (1–332) and Tob-His (1–230) are detected with anti-Tob antibody and those of Tob-His (1–112) are detected with anti-His antibody. (B) The reaction products of Tob2-His and Btg2-His are detected with anti-His antibody.

as *in vitro* monoubiquitinated sites. It is noteworthy that the amino acid sequence around Lys48 and Lys63 of Tob are located in the A domain, which is highly conserved among Tob/BTG family proteins [1] (Fig. 3B).

To confirm that Tob is ubiquitinated at Lys48 and Lys63 *in vitro*, we constructed Tob mutants in which Lys48, Lys63, or both Lys48 and Lys63 are substituted with alanine (TobK48A, TobK63A, or TobK48,63A, respectively). *In vitro* ubiquitination assay revealed that only a small amount of a slow migrating form was produced from the TobK48A mutant, whereas a slow migrating form was clearly detected in the reaction products of the TobK63A mutant (Fig. 3C), albeit slightly less than that seen in wild-type Tob. Furthermore, TobK48,63A yielded no slow migrating form. These data suggest that Lys48 is a major monoubiquitination site in Tob and that the contribution of Lys63 to monoubiquitination of Tob is small. Lys48 but not Lys63 is conserved among the corresponding amino acid sequences of all Tob/BTG family proteins.

3.4. Tob monoubiquitination affects Tob polyubiquitination

To examine the biological significance of Tob monoubiquitination *in vivo*, wild-type Tob or Tob mutant (TobK48A, TobK63A, or TobK48,63A) vectors were transfected into Cos7 cells together with 6xHis-tagged Ub expression vector. His-Ub-conjugated proteins were then purified with Ni-NTA agarose under denaturing

conditions. Immunoblot analysis of the purified fractions showed that polyubiquitination of TobK48A and TobK63A is enhanced relative to wild-type Tob (Fig. 4). These data suggest that monoubiquitination of Tob at either Lys48 or Lys63 suppresses polyubiquitination of Tob *in vivo*. It should be noted that the degree of polyubiquitination of TobK63A is greater than that of TobK48A. Although Lys48 is a major Tob ubiquitination site *in vitro* (Fig. 3C), monoubiquitination at Lys63 contributed more strongly than monoubiquitination at Lys48 toward suppression of Tob polyubiquitination *in vivo*. Nevertheless, since TobK48,63A is much more strongly polyubiquitinated than either TobK48A or TobK63A (Fig. 4), we concluded that tandem monoubiquitination at both Lys48 and Lys63 downregulated polyubiquitination of Tob.

4. Discussion

In this study, we found that Tob/BTG family proteins undergo monoubiquitination in the absence of E3s *in vitro*. The lysine residues universally conserved among members were identified as major monoubiquitination sites. Elevated polyubiquitination of TobK48,63A in comparison with wild-type Tob (Fig. 4) suggests that monoubiquitination somehow suppresses polyubiquitination. Just as monoubiquitination of UBD-containing proteins triggers intramolecular interactions with the UBD that prevent them from binding *in trans* to ubiquitinated targets [15], monoubiquitination




Fig. 3. Determination of Tob monoubiquitination sites. (A) Silver staining of *in vitro* ubiquitinated Tob. The position of monoubiquitinated Tob is indicated with a red square. The asterisk indicates a non-specific protein band derived from *E. coli*. (B) Alignment of Tob/BTG family protein amino acid sequence surrounding candidate ubiquitination sites. (C) *In vitro* ubiquitination assay of Tob mutants. Immunoblot analysis of the reaction products of the indicated Tob mutants using anti-Tob antibody are shown. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. Monoubiquitination of Tob affects its polyubiquitination *in vivo*. Wild-type Tob or Tob mutant (TobK48A, TobK63A, or TobK48,63A) vectors were transfected into Cos7 cells together with 6xHis-tagged Ub expression vector. The next day, cells were treated with MG132 (+) or DMSO (–) for 3 h. His-Ub-conjugated proteins were purified on Ni-NTA agarose under denaturing conditions. Immunoblot analysis of the purified fractions using anti-Tob antibody is shown.

of Tob may induce a conformational change that restricts E3s' access, resulting in suppression of polyubiquitination. Alternatively, UBD-containing proteins might be involved. Given that UBD-containing proteins recognize monoubiquitinated Tob, they may prevent E3s from polyubiquitinating the monoubiquitinated Tob protein. Structural analysis of monoubiquitinated Tob will help in understanding whether the inhibition of polyubiquitination by monoubiquitination is caused by intramolecular or intermolecular interactions. In the latter case, identification of molecules that specifically interact with monoubiquitinated Tob would be an important task.

While UBD-containing proteins monoubiquitinate themselves through the interaction between UBD and Ub-conjugated E2 enzymes [15], proteins in the Tob/BTG family lack any obvious UBD. Nevertheless, we cannot exclude the possibility that the unique sequences conserved among proteins of the Tob/BTG family in their amino-terminal regions might serve as a novel domain for self-monoubiquitination.

Tob is responsible for keeping cells in the quiescent state in the absence of mitogenic stimuli and also for blocking DNA damage-induced apoptosis [2,8]. Monoubiquitination of Tob may modulate Tob's ability to perform these activities. Generation of antibodies specific to monoubiquitinated Tob is needed to analyze its subcellular localization and protein interactions and ultimately to understand the biological significance of Tob monoubiquitination. In conclusion our current finding that the degree of polyubiquitination of Tob is controlled by its monoubiquitination indicates a novel strategy for protein stabilization.

Acknowledgments

We thank members of the Division of Oncology, IMSUT, for scientific comments and valuable discussions. We also thank Dr. Robert Whittier for comments on the manuscript. This work was supported by a grant for scientific research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] F. Tirone, The gene PC3/TIS21/BTG2, prototype member of the PC3/BTG/TOB family: regulator in control of cell growth, differentiation, and DNA repair? *J. Cell. Physiol.* 187 (2001) 155–165.
- [2] T. Suzuki, J. Tsuzuku, R. Ajima, T. Nakamura, Y. Yoshida, T. Yamamoto, Phosphorylation of three regulatory serines of Tob by Erk1 and Erk2 is required for Ras-mediated cell proliferation and transformation, *Genes Dev.* 16 (2002) 1356–1370.

- [3] Y. Yoshida, T. Nakamura, M. Komoda, H. Satoh, T. Suzuki, J. Tsuzuku, T. Miyasaka, E.H. Yoshida, H. Umemori, R.K. Kunisaki, K. Tani, S. Ishii, S. Mori, M. Suganuma, T. Noda, T. Yamamoto, Mice lacking a transcriptional corepressor Tob are predisposed to cancer, *Genes Dev.* 17 (2003) 1201–1206.
- [4] Y. Ito, T. Suzuki, H. Yoshida, C. Tomoda, T. Uruno, Y. Takamura, A. Miya, K. Kobayashi, F. Matsuzuka, K. Kuma, T. Yamamoto, A. Miyauchi, Phosphorylation and inactivation of Tob contributes to the progression of papillary carcinoma of the thyroid, *Cancer Lett.* 220 (2005) 237–242.
- [5] H. Sasajima, K. Nakagawa, H. Yokosawa, Antiproliferative proteins of the BTG/Tob family are degraded by the ubiquitin–proteasome system, *Eur. J. Biochem.* 269 (2002) 3596–3604.
- [6] Y. Hiramatsu, K. Kitagawa, T. Suzuki, C. Uchida, T. Hattori, H. Kikuchi, T. Oda, S. Hatakeyama, K.I. Nakayama, T. Yamamoto, H. Konno, M. Kitagawa, Degradation of Tob1 mediated by SCF^{Skp2}-dependent ubiquitination, *Cancer Res.* 66 (2006) 8477–8483.
- [7] M. Watanabe, T. Suzuki, M. Kim, Y. Abe, Y. Yoshida, S. Sugano, T. Yamamoto, Coronin7 forms a novel E3 ubiquitin ligase complex to promote the degradation of the anti-proliferative protein Tob, *FEBS Lett.* 407 (2011) 108–112.
- [8] T. Suzuki, J. Tsuzuku, K. Kawakami, T. Miyasaka, T. Yamamoto, Proteasome-mediated degradation of Tob is pivotal for triggering UV-induced apoptosis, *Oncogene* 28 (2009) 401–411.
- [9] A. Herskho, A. Ciechanover, The ubiquitin system, *Annu. Rev. Biochem.* 67 (1998) 425–479.
- [10] C.M. Pickart, M.J. Eddins, Ubiquitin: structures, functions, mechanisms, *Biochim. Biophys. Acta* 1695 (2004) 55–72.
- [11] K. Haglund, I. Dikic, Ubiquitylation and cell signaling, *EMBO J.* 24 (2005) 3353–3359.
- [12] W. Harper, B.A. Schulman, Structural complexity in ubiquitin recognition, *Cell* 124 (2006) 1133–1136.
- [13] L. Sun, Z.J. Chen, The novel functions of ubiquitination in signaling, *Curr. Opin. Cell Biol.* 16 (2004) 119–126.
- [14] S. Sigismund, S. Polo, P.P. Di-Fiore, Signaling through monoubiquitination, *Curr. Top. Microbiol. Immunol.* 286 (2004) 149–185.
- [15] D. Hoeller, C.M. Hecker, S. Wagner, V. Rogov, V. Dotsch, I. Dikic, E3-independent monoubiquitination of ubiquitin-binding proteins, *Mol. Cell* 26 (2007) 891–898.
- [16] D. Hoeller, N. Crosetto, B. Blagoev, C. Raiborg, R. Tikkanen, S. Wagner, K. Kowanetz, R. Breitling, M. Mann, H. Stenmark, I. Dikic, Regulation of ubiquitin-binding proteins by monoubiquitination, *Nat. Cell Biol.* 8 (2006) 163–169.