



Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: [www.elsevier.com/locate/ybbrc](http://www.elsevier.com/locate/ybbrc)



# An insight into the interaction mode between CheB and chemoreceptor from two crystal structures of CheB methyltransferase catalytic domain

Kwang-Hwi Cho<sup>a</sup>, Brian R. Crane<sup>b</sup>, SangYoun Park<sup>a,\*</sup>

<sup>a</sup> School of Systems Biomedical Science, Soongsil University, Seoul, Republic of Korea

<sup>b</sup> Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY, USA

## ARTICLE INFO

### Article history:

Received 1 June 2011

Available online 21 June 2011

### Keywords:

CheB  
Methyltransferase  
*Thermotoga maritima*  
Chemoreceptor  
Methyl-accepting chemotaxis proteins  
Protein–protein interaction

## ABSTRACT

We have determined 2.2 Å resolution crystal structure of *Thermotoga maritima* CheB methyltransferase domain to provide insight into the interaction mode between CheB and chemoreceptors. *T. maritima* CheB methyltransferase domain has identical topology of a modified doubly-wound  $\alpha/\beta$  fold that was observed from the previously reported *Salmonella typhimurium* counterpart, but the analysis of the electrostatic potential surface near the catalytic triad indicated considerable charge distribution difference. As the CheB demethylation consensus sites of the chemoreceptors, the CheB substrate, are not uniquely conserved between *T. maritima* and *S. typhimurium*, such surfaces with differing electrostatic properties may reflect CheB regions that mediate protein–protein interaction. Via the computational docking of the two *T. maritima* and *S. typhimurium* CheB structures to the respective *T. maritima* and *Escherichia coli* chemoreceptors, we propose a CheB:chemoreceptor interaction mode.

© 2011 Elsevier Inc. All rights reserved.

## 1. Introduction

Bacterial chemotaxis is a signal transduction cascade which allows bacteria to swim towards an attractant or away from a repellent chemical. Bacterial chemotaxis begins in response to the sensing of extracellular chemicals by the membrane-spanning chemoreceptors (also known as methyl-accepting chemotaxis proteins; MCPs), and bacteria use their flagella to either smooth-swim forward, or to tumble resulting in a randomly changing swimming direction. In bacteria such as *Escherichia coli*, the respective smooth-swimming and tumbling behavior is the result of alternation between a counter-clockwise (CCW) and a clockwise (CW) flagellar rotation [1–10]. In *E. coli*, a chemical ligand binding to the extracellular domain of the chemoreceptor is transmitted to the cytoplasmic CheA kinase, thus triggering kinase activity. CheA-bound ATP is auto-phosphorylated to a histidine of CheA, and is subsequently transferred to an aspartate of the two response-regulators, CheY and CheB. CheY and CheB bind competitively to CheA [11], and consequently determine two responses: “excitation” and “adaptation”. In the “excitation” process mediated by CheY, phospho-CheY detaches from CheA to interact with the switch protein FlhM at the flagellar motor complex, which alters the rotational direction of the flagella.

However, after the CheY-mediated response, bacteria return to the pre-stimulus state via a slower “adaptation” process mediated by CheB. Adaptation can be viewed as a negative feedback mechanism that allows the bacteria to detect temporally increasing gradients of chemicals, by preventing saturation. In general, two chemoreceptor modifying enzymes – CheR and CheB – each catalyze the methylation and demethylation of glutamates in the chemoreceptors, and the level of reversible methylation during the “adaptation” behavior enables the chemoreceptor’s attenuated sensitivity to the ligand. Methyltransferase CheR constantly methylates the chemoreceptors via S-adenosyl-L-methionine, and methyltransferase CheB functions in accordance with the activated CheA. CheB binds to CheA via an N-terminal CheY-like response-regulator domain which becomes phosphorylated by CheA. As is seen in CheY after CheA phosphorylation, phospho-CheB detaches from CheA, resulting in a concomitant opening of the C-terminal catalytic domain (CheB<sub>c</sub>) for catalytic methyltransferase activity [12–15]. The activated CheB<sub>c</sub> subsequently hydrolyzes the methyl glutamates of the chemoreceptor, and results in the deactivation of the chemoreceptor, thus setting a new threshold compared to the previous chemical concentration.

Glutamates that undergo CheR-mediated methylation and CheB-mediated demethylation are located at the cytoplasmic regions of the membrane-spanning chemoreceptors. The CheR-mediated methylation consensus sites have been defined for chemoreceptors of *Salmonella enterica* Tar [16] and *E. coli* Tsr [17], and the protein–protein interaction surface harboring the glutamate methylation sites has been predicted via docking studies

\* Corresponding author. Address: School of Systems Biomedical Science, College of Natural Science, Soongsil University, 511 Sangdo-Dong, Dongjak-Gu, Seoul 156-743, Republic of Korea. Fax: +82 2 824 4383.

E-mail address: [psy@ssu.ac.kr](mailto:psy@ssu.ac.kr) (S. Park).

[18] using the structures of the *E. coli* Tsr chemoreceptor (PDB code: 1QU7) [19] and CheR (PDB code: 1BC5) [20]. A recent study has also defined the CheR-mediated methylation consensus sites of *Thermotoga maritima* chemoreceptors, and the results demonstrate that the sites are not uniquely conserved between distant bacterial species of *S. enterica* or *E. coli* [22].

Despite the known crystal structures of full-length *Salmonella typhimurium* CheB (PDB code: 1A2O) [15] and CheB<sub>c</sub> (PDB code: 1CHD) [21], only limited attempts have been made to model the CheB<sub>c</sub> and the chemoreceptor interaction. Provided that the interaction modes are conserved between distant bacterial species, an improved model may be generated by comparing the docking results from more than two sets of interacting protein pairs. For this approach, we have determined the crystal structure of *T. maritima* CheB<sub>c</sub>, and by using this structure together with the structures of *S. typhimurium* CheB<sub>c</sub> [15,21], *T. maritima* chemoreceptor [23] and *E. coli* chemoreceptor [19], we propose a CheB<sub>c</sub> and chemoreceptor interaction model.

## 2. Materials and methods

### 2.1. Protein preparation

The genes encoding *T. maritima* methyltransferase domain of CheB (CheB<sub>c</sub>; residues 153–344) were PCR cloned into the vector pET28a (Novagen) using the *T. maritima* genomic DNA (ATCC). The protein was expressed with N-terminal His<sub>6</sub>-tag in *E. coli* strain BL21 (DE3) (Stratagene) using kanamycin selection (25 µg/mL). Using a conventional shaker, the transformed cells were grown at 37 °C in 2 L of Terrific Broth medium to OD<sub>600</sub> = ~0.6. The recombinant protein expression was induced with 0.5 mM isopropyl β-D-thiogalactopyranoside (IPTG), and the cells were further grown at 25 °C for 16 h. The cell pellets were harvested using centrifugation at 4500g for 10 min at 4 °C, and were re-suspended in ice-cold lysis buffer (20 mM TRIS pH 7.5, 500 mM sodium chloride and 5 mM imidazole) prior to homogenization by sonication.

The cell lysate was centrifuged at 70000g for 30 min at 4 °C. The supernatants were loaded onto Nickel-NTA columns, washed with wash buffer (20 mM TRIS pH 7.5, 500 mM sodium chloride and 20 mM imidazole) and the recombinant proteins were eluted with elution buffer (20 mM TRIS pH 7.5, 500 mM sodium chloride and 200 mM imidazole). The His<sub>6</sub>-tag was removed by treating the eluted protein with human thrombin (Roche) for 16 h at 4 °C. The protein was further purified using a Superdex 200 sizing column (GE Healthcare) equilibrated in gel-filtration buffer (50 mM TRIS pH 7.5 and 150 mM NaCl). The eluting fractions corresponding to CheB<sub>c</sub> were concentrated to ~50 mg/mL by centrifugation (Amicon Millipore). Protein purity determined on SDS-PAGE was >95% and the concentration was estimated by absorption at λ = 280 nm by employing the calculated molar extinction coefficient of 7680 M<sup>-1</sup> cm<sup>-1</sup> (SWISS-PROT; <http://www.expasy.ch/>).

### 2.2. Crystallization and data collection

Conditions for growing *T. maritima* CheB<sub>c</sub> crystals were found in commercial screening solutions (Hampton Research). Crystallization screenings were performed at 25 °C by using hanging drop vapor diffusion method where the protein and the well solutions were mixed in 1:1 ratio of total 2 µL volumes. Diffraction quality crystals grew after 2 months against a reservoir of 0.2 M ammonium sulfate, 0.1 M cacodylate pH 6.5 and 30% (w/v) PEG 8 K. Diffraction data were collected under a 100 K nitrogen stream at a NSLS beamline (X25) on a CCD detector (ADSC Quantum 315). The crystal belongs to the space group P2<sub>1</sub>2<sub>1</sub>2<sub>1</sub> and contains one molecule per asymmetric unit. Data were processed by DENZO and SCALEPACK [24].

### 2.3. Structure determination and refinement

The position of CheB<sub>c</sub> was determined by AMoRe molecular replacement [25] with free catalytic domain of *S. typhimurium* CheB as a search model (PDB code: 1CHD) and the diffraction data of 4.0–10.0 Å resolution. The entire model of CheB<sub>c</sub> was automatically built with ARP/wARP and SHELX [26] in CCP4 suite [27]. The final model (one CheB<sub>c</sub> molecule in asymmetric unit, residues 156–344) was further refined in CNS (*R*-factor = 0.204, *R*<sub>free</sub> = 0.223 where *R*<sub>free</sub> is the *R*-factor for 10% of randomly selected reflections excluded from the refinement) [28].

### 2.4. Computational modeling of CheB and chemoreceptor interaction

The structures of the *S. typhimurium* CheB methyltransferase domain (PDB code: 1CHD) and the *E. coli* chemoreceptor Tsr (PDB code: 1QU7; edited only to include α-helices containing the methylation Glu/Gln of residues 294–320 and 464–493 in each subunit), both lacking all heteroatoms, were docked using the ZDOCK program [29]. This program carries out rigid-body docking at all possible orientations between the two proteins, CheB<sub>c</sub> and Tsr. Among them, 1000 predicted complexes were selected based on the scoring function implemented in the program. The RDOCK program [30] was used for refinement and energy minimization of the complexes with a CHARMM force field [31]. The docked complexes were ranked based on the shape complementarities and electrostatic considerations implemented in the program, and only the top 40% were considered for the subsequent filtering processes. The top 40% complexes were filtered further on the basis of the constraints of the distance measured from three residues of the CheB catalytic triad (Ser164, His190 and Asp286) and each of the methylation Glu/Gln residues (Gln297, Glu304, Gln311). Cα-to-Cα distance constraints of at least one triad residue to Glu (or Gln) within 9 Å, and the sum of three triad residues to Glu (or Gln) within 32 Å were applied. This filtering step yielded two complexes for the methylation site Gln297, two complexes for site Glu304, and six complexes for site Gln311.

Similar steps were applied for the docking of *T. maritima* CheB<sub>c</sub> and the *T. maritima* chemoreceptor TM1143 (PDB code: 2CH7; edited to only to include α-helices containing the methylation Glu/Gln of residues 262–297 and 454–489 in each subunit). The two structures were docked using ZDOCK and minimized with RDOCK. The docked complexes were ranked and only the top 40% were considered for the subsequent filtering process by Cα-to-Cα distance. Distance constraints were based on the distance measured from three residues of the CheB catalytic triad (Ser166, His193 and Asp289) and each of the methylation Glu/Gln residues (Gln281 and Glu294). Cα-to-Cα distance constraints of at least one triad residue to Glu/Gln within 10 Å, and the sum of three triad residues to Glu/Gln within 30 Å were applied. This filtering step yielded only two complexes (No. 400 and No. 554) for the Gln281 methylation site. The 10 complexes resulting in the case of *E. coli* (or *S. typhimurium*) proteins and two complexes for the *T. maritima* proteins were assessed and compared using the PyMol molecular graphics display program [31]. All the structural figures were also rendered with PyMol [32].

## 3. Results and discussion

### 3.1. Overall structure of *T. maritima* CheB<sub>c</sub>

The crystal structure of *T. maritima* CheB methyltransferase domain was determined to a resolution of 2.2 Å using a *S. typhimurium* CheB structure (PDB code: 1CHD) as a molecular replacement model. The refined model harbors one α/β sandwich-fold CheB<sub>c</sub> molecule

within the crystal asymmetric unit, which is composed of nine  $\beta$ -strands, six  $\alpha$ -helices, and one  $3_{10}$ -helix (Fig. 1). Among the 192 residues designed in the protein construct, 189 residues were built in the model. The three N-terminal CheB residues (153–KPA–155) were not included in the model, as their electron densities were not clearly discernable. One hundred and six water molecules were also included. Data collection and refinement statistics are provided in Tables 1 and 2.

The fold of *T. maritima* CheB<sub>c</sub> is identical to that of *S. typhimurium* CheB<sub>c</sub>, which is a modified, doubly wound  $\alpha/\beta$  fold containing the central seven parallel  $\beta$ -strands flanked by six  $\alpha$ -helices and an anti-parallel  $\beta$ -hairpin (Fig. 2). In brief, the fold starts from  $\beta$ 1 and the subsequently alternating  $\alpha$ -helices and  $\beta$ -strands ( $\alpha$ 1- $\beta$ 2/ $\alpha$ 2- $\beta$ 3/ $\beta$ 4) form one core of the central parallel  $\beta$ -sheet ( $\beta$ 1/ $\beta$ 2/ $\beta$ 4/ $\beta$ 3). The following two  $\beta$ -strands ( $\beta$ 5- $\beta$ 6) form an anti-parallel  $\beta$ -hairpin distal to the central parallel  $\beta$ -sheet, which result in a modified  $\alpha/\beta$  fold. Finally, the three following sets of alternating  $\alpha$ -helix and the  $\beta$ -strand ( $\alpha$ 3- $\beta$ 7/ $\alpha$ 4- $\beta$ 8/ $\alpha$ 5- $\beta$ 9) complete the second half of the central seven-stranded  $\beta$ -sheet ( $\beta$ 9/ $\beta$ 8/ $\beta$ 7/ $\beta$ 1/ $\beta$ 2/ $\beta$ 4/ $\beta$ 3). The catalytic Ser-His-Asp triad which generates the methylesterase activity lie at the surface created mainly by loops that connect the  $\beta$ -strands and the  $\alpha$ -helices ( $\beta$ 1- $\alpha$ 1 loop,  $\beta$ 2- $\alpha$ 2 loop,  $\beta$ 6- $\alpha$ 3 loop,  $\beta$ 7- $\alpha$ 4 loop). Two other loops ( $\beta$ 3- $\beta$ 4 and  $3_{10}$  helix- $\alpha$ 5 loop) are also in the vicinity of the catalytic triad (Fig. 2).

### 3.2. Surface electrostatic potential difference between *T. maritima* and *S. typhimurium* CheB<sub>c</sub>

As *T. maritima* and *S. typhimurium* CheB<sub>c</sub>s are conserved with 36% identity (54% similarity) in their protein sequences (Fig. 1), the superposition of *T. maritima* and *S. typhimurium* CheB<sub>c</sub>s (PDB code: 1CHD) is clearly indicative of an overlap of secondary structural elements ( $C_{\alpha}$  RMSD = 1.032 Å aligning 182 residues). Partial sequence conservation also occurs near the catalytic surface of

**Table 1**

Data collection and phasing statistics.

Wavelength (Å)	1.100
Resolution (Å)	30–2.15
Highest shell	(2.23–2.15)
Completeness (%)	99.9 (99.9)
$R_{\text{merge}}^a$	0.106 (0.382)
$I/\sigma$ (I)	35 (8.4)

$$^a R_{\text{merge}} = \sum_j |I_j - \langle I \rangle| / \sum_j I_j$$

**Table 2**


Refinement statistics.

Space group	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub> ( $a = 52.47$ Å, $b = 65.08$ Å, $c = 78.35$ Å)
Resolution range (Å)	30–2.2
Unique reflections (test set)	13,555 (1382)
Wilson B ( $\text{\AA}^2$ )	28.51
R-factor <sup>a</sup> ( $R_{\text{free}}^b$ )	0.204 (0.223)
No. of scatters (No. of residues)	2928 (189)
No. of water molecules	106
RMSD bonds (Å)	0.00624
RMSD angles ( $^\circ$ )	1.405
Average B-factor (main chain) ( $\text{\AA}^2$ )	24.29
Average B-factor (side chain) ( $\text{\AA}^2$ )	26.40
Average B-factor (water) ( $\text{\AA}^2$ )	38.31

$$^a R\text{-factor} = \sum (|F_{\text{obs}}| - |F_{\text{calc}}|) / \sum |F_{\text{obs}}|$$


<sup>b</sup> R-factor for 10% of randomly selected reflections excluded from the refinement.

the protein, which is created by the contiguous loops that connect the  $\beta$ -strands and the  $\alpha$ -helices (Fig. 1,  $\beta$ 1- $\alpha$ 1 loop,  $\beta$ 2- $\alpha$ 2 loop,  $\beta$ 6- $\alpha$ 3 loop,  $\beta$ 7- $\alpha$ 4 loop). However, analysis of the electrostatic potential surface near the catalytic triad reflects considerable differences in the charge distribution between the two *T. maritima* and *S. typhimurium* CheB<sub>c</sub>s. The region of conspicuous differences is outlined along the arrows near the catalytic triad surface (Fig. 2C).



**Fig. 1.** Sequence alignment of CheB<sub>c</sub> from five organisms. CheB<sub>c</sub> sequences from *T. maritima*, *E. coli*, *S. typhimurium*, *Bacillus subtilis* and *Myxococcus xanthus* were aligned on the basis of the amino acid sequences. Secondary structural elements are indicated as block rectangles ( $\alpha$ -helix) and arrows ( $\beta$ -sheet) on top of the corresponding amino acid residues. Conserved residues are boxed around the amino acids. Catalytically critical residues (Ser-His-Asp triad) are indicated with an asterisk.






**Fig. 2.** Similarities in the overall fold do not account for the dissimilarities in the electrostatic potential surface of *T. maritima* and *S. typhimurium* CheB<sub>c</sub>. (A and B) Schematic ribbon diagrams of the superimposed structures of *T. maritima* and *S. typhimurium* CheB<sub>c</sub> are shown in two different orientations which are rotated 90° with respect to each other along the horizontal arrow. Three side chains of the catalytic triad are shown in sticks. (C) Differences in the electrostatic potential surface near the catalytic triad are highlighted using colors by charge. The surface was generated from the 90°-rotated orientation (B) which looks at the catalytic triad from the top. Dissimilar regions of charge distribution in *T. maritima* and *S. typhimurium* are mostly located around regions indicated with arrows. Since the consensus sequences adjacent to the methylation/demethylation glutamates of the chemoreceptors are not exactly conserved between the two species, such surface with electrostatic difference may be the sites of chemoreceptor interaction. (For interpretation of references to color in this figure legend, the reader is referred to the web version of this article.)

The CheR-mediated methylation consensus sites neighboring the modified glutamate have been defined for chemoreceptors of *S. enterica* Tar [17], *E. coli* Tsr [16], and several *T. maritima* chemoreceptors [22], and the results demonstrate that the sites are not uniquely conserved. As perfect complementarity of proteins, the fundamental driver of biological specificity, within biologically relevant complexes results during and requires co-evolution [33], such surfaces with differing electrostatic properties may reflect CheB<sub>c</sub> regions that mediate interactions with chemoreceptors.

Methylation-undergoing glutamates of chemoreceptors, which are also the substrates for methyltransferase CheB, are located in the cytoplasmic regions of the membrane-spanning chemoreceptor. The methylation (or demethylation) consensus sequences (Glx-**Glx**-X-X-Ala-Ser/Thr, methylation residue is in bold) of *S. enterica* (Tar) and *E. coli* (Tsr) chemoreceptors [16,17] have been shown to differ slightly from that of the *T. maritima* receptors

(Ala/Ser-sm-X-**Glx**-Glu-X-sm-Ala/Ser, methylation residues are in bold; sm indicates small residues which are either Ala, Gly, Ser or Thr) [22]. Two CheB<sub>c</sub>s recognizing the different consensus sequence of the chemoreceptors can be mapped to the different surface complementarities which have co-evolved between the chemoreceptor substrate and the CheB enzyme. Electrostatic potential surfaces near the *T. maritima* and *S. typhimurium* CheB<sub>c</sub> catalytic triad with substantial differences suggested that two CheB<sub>c</sub> structures in conjunction with the chemoreceptor's structural information may assist in generating a reliable CheB and chemoreceptor interaction model. A systematic docking study has been carried out using two sets of CheB<sub>c</sub> and chemoreceptor structures originating from *T. maritima* and *S. typhimurium*. Provided that the general interaction modes are conserved between the two distant bacterial species, an interaction model has been generated by comparing the overlapping docking results.




**Fig. 3.** Proposed chemoreceptor docking site for *T. maritima* and *S. typhimurium* CheB<sub>c</sub>. (A) Docking models were generated from the PDB coordinates of the chemoreceptors and the CheB<sub>c</sub>s from *T. maritima* and *S. typhimurium* using ZDOCK and RDOCK. Overall docking modes are shown with the chemoreceptor positions relative to the superimposed CheB<sub>c</sub> (only *T. maritima* CheB<sub>c</sub> is shown for clarity). (B) The methylation residue for the *T. maritima* chemoreceptor (Gln281) is shown with sticks in relation to the catalytic triad. Distances indicate the measured bonding distance between atoms in the model.

### 3.3. Prediction of chemoreceptor and CheB interaction mode

*E. coli* and *S. typhimurium* both belong to the Enterobacteriaceae family and are closely related in the bacterial phylogeny tree. As the CheBs from the two species also share 95% sequence identity, the structures of the cytoplasmic regions of the *E. coli* Tsr chemoreceptor (PDB code: 1QU7) and *S. typhimurium* CheB<sub>c</sub> (PDB code: 1CHD) were docked using ZDOCK and RDOCK. Docking was carried out solely using the truncated four-helix bundle of the chemoreceptor, which includes the methylation Glu/Gln (Gln297, Glu304, and Gln311). The docking results were sorted on the basis of the relevant complementarities and electrostatic considerations, and further filtered using distance constraints between the catalytic residues and methylation glutamates. The final result yielded 10 CheB<sub>c</sub> and chemoreceptor complexes in three Tsr methylation Glu/Gln residues (two complexes for site Gln297, two complexes for site Glu304, and six complexes for site Gln311). However, when the 10 complexes were superposed relative to CheB<sub>c</sub>, the positions of the chemoreceptor relative to CheB<sub>c</sub> – i.e. the directionality of the chemoreceptor four-helix bundle – in the 10 complexes were heterogeneous, which indicates that generating a single interaction mode is not possible (results not shown).

Similar docking steps were applied for the case of *T. maritima* CheB<sub>c</sub> and the TM1143 chemoreceptor (PDB code: 2CH7). A recently reported cytoplasmic structure of the *T. maritima* chemore-

ceptor TM0014 (PDB code: 3G67) [34] was not included in the docking set due to the absence of the methylation consensus sequences. Similar to the previous case, docking was carried out solely using the truncated four-helix bundle of the chemoreceptor which includes the methylation Glu/Gln (Gln294 and Gln281). After sorting and filtering processes using distance constraints, the final docking result with the *T. maritima* proteins yielded two CheB<sub>c</sub> and chemoreceptor complexes (No. 400 and No. 554) around only the Gln281 methylation residue. The positions of the chemoreceptor relative to CheB<sub>c</sub> in the two resultant complexes are practically identical, with only a slight  $\sim 1^\circ$  shift in the four-helix bundle around Gln281 (results not shown). When the two docking complexes from *T. maritima* proteins were overlaid with the 10 docking complexes from *E. coli* proteins, only one *E. coli* docking pair (No. 640 from site Gln311) matched exactly with an identical helical arrangement of the chemoreceptor substrate relative to CheB<sub>c</sub> (Fig. 3). Surprisingly, when the  $\alpha$ -helix of the chemoreceptor containing the demethylation site was observed outside the context of the four-helix bundle, it sits within a groove created at the adjunction of the four CheB<sub>c</sub>  $\alpha$ -helices ( $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ , and  $\alpha 4$ , Fig. 4). The interaction surface is created by the contacting loops that connect the  $\beta$ -strands to the  $\alpha$ -helices ( $\beta 1$ - $\alpha 1$  loop,  $\beta 2$ - $\alpha 2$  loop,  $\beta 6$ - $\alpha 3$  loop,  $\beta 7$ - $\alpha 4$  loop), and is the region where majority of the electrostatic potential surface differed between the two *T. maritima* and *S. typhimurium* CheB<sub>c</sub>s (Figs. 2C and 4). It is worth



**Fig. 4.** View of the proposed docking site with chemoreceptor  $\alpha$ -helix shown over the *T. maritima* CheB<sub>c</sub> electrostatic potential surface. View of the truncated  $\alpha$ -helical substrate of the *T. maritima* and *E. coli* chemoreceptor overlaid on top of the *T. maritima* CheB<sub>c</sub> electrostatic potential surface are shown in two different orientations, which are rotated 90° with respect to each other along the indicated axis (only *T. maritima* CheB<sub>c</sub> is shown for clarity). The  $\alpha$ -helical substrate of the chemoreceptor sits within a groove created at the active site.

noting that this CheB<sub>c</sub> groove has also been previously predicted to be the chemoreceptor interface, which was concluded from the manual inspection of a 10 Å – diameter channel that accommodates an  $\alpha$ -helix [21].

We have succeeded in generating a probable model between the two interacting proteins using the systematic docking approach that defines the helical orientation of the chemoreceptor relative to CheB<sub>c</sub>. However, since *T. maritima* CheB demethylates on at least two different methyl glutamates of the chemoreceptor, more than one interaction mode – an interface covering methylation residues Gln294 as well as Gln281 – should be predicted from the docking methods. As no docking results satisfy the constraints on Gln294, we are unable to suggest multiple modes of CheB<sub>c</sub> interaction between the different demethylation glutamates.

In summary, we have determined the crystal structure of *T. maritima* CheB<sub>c</sub> and used it to generate a chemoreceptor interaction model by carrying out docking studies. We have clearly demonstrated that the limited efficiencies of the docking processes can be enhanced in cases in which two or more structure sets of interacting proteins are known from different species. Although we had aimed to crystallize *T. maritima* CheB<sub>c</sub> in complex with the TM1143 chemoreceptor, to date this has not proven successful. A more complete picture of methyltransferase CheB<sub>c</sub> and the chemoreceptor substrate interaction thus awaits a successful determination of the complex structure.

#### 4. Accession code

Worldwide Protein Data Bank coordinate for *T. maritima* CheB<sub>c</sub> has been deposited with an accession code 3SFT.

#### Acknowledgments

This work was supported by the Human Resources Development of the Korea Institute of Energy Technology Evaluation and Planning (KETEP) grant funded by the Ministry of Knowledge Economy, Republic of Korea (No. 20104010100610) to K.W.C., and NIH

grant GM066775 to B.R.C. We thank J. Y. Park for graphical assistance during figure preparation. This work was supported by the Research carried out (in part) at the National Synchrotron Light Source, Brookhaven National Laboratory, which is supported by the US Department of Energy, Division of Materials Sciences and Division of Chemical Sciences, under Contract No. DE-AC02-98CH10886.

#### References

- [1] K.A. Borkovich, M.I. Simon, The dynamics of protein phosphorylation in bacterial chemotaxis, *Cell* 63 (1990) 1339–1348.
- [2] J.S. Parkinson, E.C. Kofoed, Communication modules in bacterial signaling proteins, *Annu. Rev. Genet.* 26 (1992) 71–112.
- [3] J.L. Appleby, J.S. Parkinson, R.B. Bourret, Signal transduction via the multi-step phosphorelay: not necessarily a road less travelled, *Cell* 86 (1996) 845–848.
- [4] P.N. Goudreau, A.M. Stock, Signal transduction in bacteria: molecular mechanisms of stimulus-response coupling, *Curr. Opin. Microbiol.* 1 (1998) 160–169.
- [5] H.C. Berg, The rotary motor of bacterial flagella, *Annu. Rev. Biochem.* 72 (2003) 19–54.
- [6] S. Kojima, D.F. Blair, The bacterial flagellar motor: structure and function of a complex molecular machine, *Int. Rev. Cytol.* 233 (2004) 93–134.
- [7] G.H. Wadhams, J.P. Armitage, Making sense of it all: bacterial chemotaxis, *Nat. Rev. Mol. Cell Biol.* 5 (2004) 1024–1037.
- [8] V. Sourjik, Receptor clustering and signal processing in *E. coli* chemotaxis, *Trends Microbiol.* 12 (2004) 569–576.
- [9] J.S. Parkinson, P. Ames, C.A. Studdert, Collaborative signaling by bacterial chemoreceptors, *Curr. Opin. Microbiol.* 8 (2005) 116–121.
- [10] M. Inouye, R. Dutta, Histidine Kinases in Signal Transduction, Academic Press, San Diego, 2002.
- [11] J. Li, R.V. Swanson, M.I. Simon, R.M. Weis, The response regulators CheB and CheY exhibit competitive bind to the kinase CheA, *Biochemistry* 34 (1995) 14626–14636.
- [12] A. Lupas, J. Stock, Phosphorylation of an N-terminal regulatory domain activates the CheB methyltransferase in bacterial chemotaxis, *J. Biol. Chem.* 264 (1989) 17337–17342.
- [13] R.C. Stewart, Activating and inhibitory mutations in the regulatory domain of CheB, the methyltransferase in bacterial chemotaxis, *J. Biol. Chem.* 268 (1993) 1921–1930.
- [14] G.S. Anand, P.N. Goudreau, A.M. Stock, Activation of methyltransferase CheB: evidence of a dual role for the regulatory domain, *Biochemistry* 37 (1998) 14038–14047.
- [15] S. Djordjevic, P.N. Goudreau, Q. Xu, A.M. Stock, A.H. West, Structural basis for methyltransferase CheB regulation by a phosphorylation-activated domain, *Proc. Natl. Acad. Sci. USA* 95 (1998) 1381–1386.



- [16] T.C. Terwilliger, D.E. Koshland Jr., Sites of methyl esterification and deamination on the aspartate receptor involved in chemotaxis, *J. Biol. Chem.* 259 (1984) 7719–7725.
- [17] M.R. Kehry, M.W. Bond, M.W. Hunkapiller, F.W. Dahlquist, Enzymatic deamidation of methyl-accepting chemotaxis proteins in *Escherichia coli* catalyzed by the cheB gene product, *Proc. Natl. Acad. Sci. USA* 80 (1983) 3599–3603.
- [18] E. Perez, A.H. West, A.M. Stock, S. Djordjevic, Discrimination between different methylation states of chemotaxis receptor Tar by receptor methyltransferase CheR, *Biochemistry* 43 (2004) 953–961.
- [19] K.K. Kim, H. Yokota, S.H. Kim, Four-helical-bundle structure of the cytoplasmic domain of a serine chemotaxis receptor, *Nature* 400 (1999) 787–792.
- [20] S. Djordjevic, A.M. Stock, Chemotaxis receptor recognition by protein methyltransferase CheR, *Nat. Struct. Biol.* 5 (1998) 446–450.
- [21] A.H. West, E. Martinez-Hackert, A.M. Stock, Crystal structure of the catalytic domain of the chemotaxis receptor methyltransferase, CheB, *J. Mol. Biol.* 250 (1995) 276–290.
- [22] E. Perez, H. Zheng, A.M. Stock, Identification of methylation sites in *Thermotoga maritima* chemotaxis receptors, *J. Bacteriol.* 188 (2006) 4093–4100.
- [23] S.-Y. Park, P.R. Borbat, G. Gonzalez-Bonet, J. Bhatnagar, A.M. Pollard, J.H. Freed, A.M. Bilwes, B.R. Crane, Reconstruction of the chemotaxis receptor-kinase assembly, *Nat. Struct. Mol. Biol.* 13 (2006) 400–407.
- [24] Z. Otwinowski, W. Minor, Processing of X-ray diffraction data in oscillation mode, *Methods Enzymol.* 276 (1997) 307–325.
- [25] J. Navaza, *AMoRe*: an automated package for molecular replacement, *Acta Cryst. A* 50 (1994) 157–163.
- [26] A. Perrakis, R.M. Morris, V.S. Lamzin, Automated protein model building combined with iterative structure refinement, *Nat. Struct. Biol.* 6 (1999) 458–463.
- [27] Collaborative Computational Project Number 4, The CCP4 suite: programs for protein crystallography, *Acta Cryst. D* 50 (1994) 760–763.
- [28] A.T. Brünger, P.D. Adams, G.M. Clore, W.L. DeLano, P. Gros, R.W. Grosse-Kunstleve, J.S. Jiang, J. Kuszewski, M. Nilges, N.S. Pannu, R.J. Read, L.M. Rice, T. Simonson, G.L. Warren, Crystallography and NMR system: a new software suite for macromolecular structure determination, *Acta Crystallogr. D Biol. Crystallogr.* 54 (1998) 905–921.
- [29] R. Chen, L. Li, Z. Weng, ZDOCK: an initial-stage protein-docking algorithm, *Proteins* 52 (2003) 80–87.
- [30] L. Li, R. Chen, Z. Weng, RDOCK: refinement of rigid-body protein docking predictions, *Proteins* 53 (2003) 693–707.
- [31] B.R. Brooks, R.E. Bruccoleri, B.D. Olafson, D.J. States, S. Swaminathan, M. Karplus, CHARMM: a program for macromolecular energy, minimization, and dynamics calculations, *J. Comput. Chem.* 4 (1983) 187–217.
- [32] W.L. DeLano, The PyMOL Molecular Graphics System, DeLano Scientific, San Carlos, CA, USA (2002) <<http://www.pymol.org>>.
- [33] S.Y. Park, W. Jin, J.R. Woo, S.E. Shoelson, Crystal structures of Human TBC1D1 and TBC1D4 (AS160) RabGAP domains reveal critical elements for GLUT4 translocation, *J. Biol. Chem.* 286 (2011) 18130–18138.
- [34] A.M. Pollard, A.M. Bilwes, B.R. Crane, The structure of a soluble chemoreceptor suggests a mechanism for propagating conformational signals, *Biochemistry* 48 (2009) 1936–1944.